

Widespread concerns still exist in relation to discrimination towards women and girls and FGM

It is heartening that the number of children presenting with female genital mutilation (FGM) in the UK is less than expected from the estimates.¹ This is against a decade of changes to legal frameworks, including introducing FGM Protection Orders (FGMPOs) to protect girls and women at risk, alongside educational efforts to increase public awareness by a range of stakeholders.

It is possible that FGM is abandoned after migration to the UK¹ and that anti-FGM measures need to be proportionate to the empirical evidence of the risk of FGM in the UK. However, there remain widespread concerns connected to discrimination towards women and girls of which FGM is one facet.


A criminal conviction for FGM in England in 2019 focused on witchcraft, rather than the injury and the harm that the girl sustained. In immigration cases, the home office appears to expect medical examinations to prove whether women and girls have had FGM or not, and in family law cases there are concerns that the approach to FGMPOs might have led to disengagement from some individuals, with concerns of a lack of state support to accompany legal measures. The most successful changes appear to be through education and public health prevention strategies. However, we note this with cautionary optimism due to a lack of empirical research, which is clearly imperative.

Using a disease progression model, we suggest that the study¹ is visualising the interconnection of legislation and health measures *with* education campaigns.² Interventions can include 'upstream' or 'downstream' approaches. Downstream approaches are individualised, catering to specific needs. Upstream approaches are system-wide, such as the introduction of FGMPOs. While often seen as a heavy-handed approach by policy-makers, the effectiveness of upstream interventions improves over time, as resistance to change fades and subsequent generations grow up with changes normalised.³

Upstream interventions tend to be cheaper with better health outcomes.⁴ This may offer reassurance to the longitudinal

effectiveness of anti-FGM measures. However, it would be unwise to suggest that effectiveness increases over time without additional interventions.

The use of legislation and government policies to change individuals' and communities' behaviours, attitudes and practices towards FGM is under-researched. The distance between on-the-ground impact through education and a legislative response can affect the success not just of eradication of FGM but also the wider necessary reduction in gender inequality. While the improvement of law was useful in 2012, the inflammatory and hyperbolic language centring on one conviction means the application of legal mechanisms risks becoming a barrier to reaching the United Nations 'Sustainable Development Goals to eradicate FGM'. Upstream interventions are considered more effective in terms of their longevity and equitability. Recognising that legislation alone is not sufficient to achieve the desired outcome but that health, education and law combined have the potential to end FGM in our generation is important.

Felicity Gerry,^{1,2} Charlotte Proudman,^{3,4}
Hoda Ali,⁵ Joseph Home,^{6,7}
Andrew Graeme Rowland ^{6,7}

¹Libertas Chambers, London, UK

²Deakin University, Burwood, Victoria, Australia

³Queens' College, Cambridge, UK

⁴Goldsmith Chambers, London, UK

⁵Perivale Primary School, Perivale, UK

⁶School of Health and Society, University of Salford, Salford, UK

⁷Emergency Department, Pennine Acute Hospitals NHS Trust, Manchester, UK

Correspondence to Professor Felicity Gerry, Libertas Chambers, London EC4M 7AN, UK; fgerryqc@libertaschambers.com

Twitter Andrew Graeme Rowland @DrAndrewRowland

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Provenance and peer review Not commissioned; internally peer reviewed.



OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Gerry F, Proudman C, Ali H, et al. *Arch Dis Child* 2021;**106**:929.

Accepted 10 January 2021

Published Online First 29 January 2021

Arch Dis Child 2021;**106**:929.

doi:10.1136/archdischild-2020-321187

ORCID iD

Andrew Graeme Rowland <http://orcid.org/0000-0001-9564-0032>

REFERENCES

- Hodes D, Ayadi O'Donnell N, Pall K, et al. Epidemiological surveillance study of female genital mutilation in the UK. *Arch Dis Child* 2021;106:372–6.
- Pandve HT. Quaternary prevention: need of the hour. *J Family Med Prim Care* 2014;3:309–10.
- Capewell S, Capewell A. An effectiveness hierarchy of preventive interventions: neglected paradigm or self-evident truth? *J Public Health* 2018;40:350–8.
- World Health Organization. EPHO5: disease prevention, including early detection of illness. Available: <http://www.euro.who.int/en/health-topics/Health-systems/public-health-services/policy/the-10-essential-public-health-operations/epho5-disease-prevention,-including-early-detection-of-illness2> [Accessed 11 Nov 2020].