

## Ameliorative Effect of *Ocimum Sanctum* on Meloxicam Induced Toxicity in Wistar Rats

R. Mahaprabhu, A. G. Bhandarkar, Babu Lal Jangir, S. P. Rahangadale, N. V. Kurkure

Department of Pathology, Nagpur Veterinary College, Maharashtra Animal and Fishery Sciences University, Nagpur- 440 006, India

### ABSTRACT

An ameliorating effect of *Ocimum sanctum* on the toxic effect of meloxicam, a new non-steroidal anti-inflammatory drug was studied by evaluating haemato-biochemical parameters, oxidative stress, gross and histopathological changes in various organs of Wistar rats. A total of thirty-six male rats were divided in six experimental groups each comprising of six rats and numbered from G<sub>1</sub> to G<sub>6</sub>. Meloxicam toxicity was induced by oral feeding of meloxicam at 1.2 mg/kg and 2.4 mg/kg body weight in G<sub>2</sub> and G<sub>3</sub> respectively for 28 days. Group G<sub>4</sub> and G<sub>5</sub> were fed with 1.2-mg/kg body weight and 2.4-mg/kg body weight of meloxicam along with 200 mg/kg body weight of aqueous extract of *Ocimum sanctum*. Group G<sub>1</sub> serve as control while group G<sub>6</sub> was kept as treatment control and fed only aqueous extract of *Ocimum sanctum* at 200 mg/kg body weight. Clinical finding showed mild diarrhea from 23<sup>rd</sup> day onwards in-group treated with 2.4-mg/kg body of meloxicam. Significant reduction of hemoglobin and packed cell volume (PCV) was observed in both the group treated with 1.2 mg/kg and 2.4-mg/kg body wt. of meloxicam. *Ocimum sanctum* could restore the hemoglobin and PCV value in-group treated with meloxicam at low dose level. Serum alkaline phosphatase, serum glutamic pyruvic transaminase, Serum glutamic oxaloacetic transaminase and total bilirubin were found elevated in meloxicam treated groups and indicated hepatotoxic activity of meloxicam. *Ocimum sanctum* could reduce hepatotoxic activity of meloxicam in group G<sub>4</sub> receiving meloxicam at lower dose rate along with *Ocimum sanctum* failed to regulate creatinine level in meloxicam treated groups. In meloxicam toxicity elevated Lipid peroxidation values was noticed in liver and kidneys, while superoxide dismutase and glutathione did not revealed any change. Stomach and intestine revealed hemorrhagic gastroenteritis and ulcers. Perivascular necrosis with infiltration with inflammatory cells was evident in liver. Interstitial nephritis, myocardial necrosis and spongiform encephalopathy were important lesions. The *Ocimum sanctum* could only counteract the toxic effect of meloxicam in liver and gastrointestinal tract.

**Key words:** Meloxicam toxicity, *Ocimum sanctum*, oxidative stress, Wistar rats

#### Access this article online

Quick Response Code:



Website:

[www.toxicologyinternational.com](http://www.toxicologyinternational.com)

DOI:

10.4103/0971-6580.84265

### INTRODUCTION

Meloxicam, a new non-steroidal anti-inflammatory drug (NSAID) is widely used in humans, cattle, buffalo, goats and dogs. Meloxicam is a COX-2 (cyclo-oxygenase) inhibitor at its lowest therapeutic dose and is an anti-inflammatory by inhibiting prostanoid synthesis in inflammatory cells<sup>[1]</sup>.

**Address for correspondence:** Dr. Nitin Kurkure, Department of Pathology, Nagpur Veterinary College, Maharashtra Animal and Fishery Sciences University, Nagpur- 440 006, India. E- mail: nitinkurkureatrediffmail.com

Meloxicam inhibits COX-2 about 12 times more selectively than COX-1.<sup>[2]</sup> Meloxicam is reported to be safer as it produce significantly lower incidence of gastrointestinal adverse effects as compared to diclofenac sodium.<sup>[3]</sup> Most NSAIDs are absorbed well from gastric and intestinal mucosa. They are bound to plasma albumin and distributed all over the body evenly. Most NSAIDs are metabolized in the liver by oxidation and converted to inactive metabolites, which are then typically excreted in the urine while some drugs are partially excreted in bile. Meloxicam, as NSAID, either alone or with antimicrobial drugs, is indicated for use in ruminants for the treatment of pneumonia, pleuritis, laminitis, myositis, sprain, mastitis, prolapse of uterus, premature labour etc. The principle side effect of meloxicam is gastrointestinal irritation (vomiting, diarrhea and ulceration). Rare but important side effects include liver and kidney toxicity<sup>[4]</sup>.

The plant *Ocimum sanctum* belongs to family "Labiatae" and is a well-known pot herb regarded as sacred by Hindus. *Ocimum sanctum* (Tulsi) is a medicinal plant commonly grown in India. Different parts of this plant have been reported to exhibit several medicinal properties.<sup>[5]</sup> *Ocimum sanctum* has been reported to possess antihepatotoxicity property,<sup>[6]</sup> and two triterpenes from the leaves have been shown to possess hepatoprotective effect against carbon tetrachloride induced damage in rats.<sup>[7]</sup> The present study is formulated to study sub acute toxicity of meloxicam in Wistar rats. An attempt is also made to evaluate protective property of *Ocimum sanctum* during meloxicam toxicity.

## MATERIALS AND METHODS

### Chemicals

Meloxicam pure base powder was obtained from M/s. Cadila Pharmaceutical Ltd., Dholka and Sodium Carboxy Methyl Cellulose pure base powder was obtained from M/s Fischer scientific, Mumbai. All the biochemical kits and chemicals used in the study were of analytical grade.

### Herbal medicine

Aqueous extract of dried leaves of *Ocimum sanctum* was prepared as follows and was used as herbal medicine. The leaves of *Ocimum sanctum* were collected from neighborhood and identified based on morphological characters. The leaves were dried in shadow at room temperature and powdered. The powder was stored in glass bottle in cool and dry place away from direct sunlight and used for preparation of aqueous extract. The aqueous extract was prepared by the method described earlier.<sup>[8]</sup> The extractability percentage was 6.25 and extract was dark brown in color. Extract was finally stored in desiccators in a cool and dry place.

### Experimental animals

Thirty-six male Wistar rats were divided into six groups of 6 animals each. All the animals were maintained in

polypropylene cages, (47 × 34 × 18 cm). The maximum number of animals in each cage was six. Dried rice husk was used as a bedding material. Bedding material was changed every alternate day. All the animals were kept under standard managemental conditions as per the norms of Committee for the purpose of control and supervision on experiments on animals (CPCSEA), and were maintained under a controlled environment with temperature at 23 ± 2°C, relative humidity at 55 ± 5%, and a 12-hours/12 hours light/dark cycle throughout the experiment. The animals were fed on a standard pellet diet. The pellet diet consisted of 22.02% crude protein, 4.25% crude oil, 3.02% crude fiber, 7.5% ash and 1.38% sand silica. They were given *ad-lib* feed and wholesome drinking water throughout the experiment. The necessary Institute Animal Ethical committee approval was obtained.

### Design of experiment

The rats were acclimatized for 15 days to the environment, before the start of the experiment. Each group of rat was given separate identification mark by using picric acid. A total of thirty-six male rats, were divided into six groups (G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub>, G<sub>5</sub> and G<sub>6</sub>) with six rats in each group. Group 'G<sub>1</sub>' served as control and was treated with vehicle, 0.5 per cent sodium carboxy methyl cellulose at 1 ml. Group 'G<sub>2</sub>' and 'G<sub>3</sub>', which received meloxicam at 1.2 mg/kg body weight, and 2.4 mg /kg body weight, respectively. Group 'G<sub>4</sub>' and G<sub>5</sub> received meloxicam at 1.2 mg/kg body weight and 2.4 mg /kg body weight, respectively along with 200 mg/kg of aqueous extract of *Ocimum sanctum* and Group G<sub>6</sub> was treated with 200 mg of aqueous extract of *Ocimum sanctum* per oral. Vehicle used for diluting meloxicam to obtain the desired concentration was 0.5 per cent Sodium Carboxy Methyl Cellulose in distilled water.

### Collection of material for hematological and biochemical parameters

The rats from all experimental groups were sacrificed on 28<sup>th</sup> day of experiment. Prior to sacrifice the rats were fasted for 12 hours. Blood was collected on the day of sacrifice (28<sup>th</sup> day) from retro orbital plexus with the help of capillary tube in two vials, one coated with Ethylenediaminetetraacetic acid (EDTA) as an anticoagulant and in second vial without anticoagulant, later used for serum separation. Haematology parameters like Hemoglobin (Hb), PCV, Total Erythrocyte Count (TEC), Total Leukocyte Count (TLC), Differential Leukocyte Count (DLC) and haematocrit values were studied by standard routine procedures.<sup>[9]</sup> Blood samples collected in vials without anticoagulant and were kept undisturbed. The serum was separated and stored at -20°C for subsequent analysis. SGOT, SGPT, ALP, Glucose, Total Protein, Albumin, Globulin, Creatinine, Cholesterol and Triglyceride estimations were carried out by analytical kits supplied by RFCL, Haridwar, Uttarakhand, India on clinical chemistry semi auto analyzer.

Animals were sacrificed by cervical dislocation at the end of experiment. A detailed post mortem examination was carried out. The representative tissues - liver, kidney, heart lung, spleen, brain and spinal cord were fixed in 10% buffered formalin immediately on removal. They were subjected for histopathological examination following standard procedure.<sup>[10]</sup> Tissues such as liver and kidney were collected in ice cold containers for studying oxidative stress and lipid peroxidation parameters viz. LPO,<sup>[11]</sup> Reduced glutathione (GSH)<sup>[12]</sup> and SOD.<sup>[13]</sup>

### Statistical analysis

The data generated was analyzed with one way analysis of variance (ANOVA).<sup>[14]</sup> The level of significance was set at  $P \leq 0.05$ .

## RESULTS AND DISCUSSION

### Clinical observations

The rats from group  $G_2$  receiving 1.2 mg/kg of meloxicam did not exhibit any clinical symptoms up to the end of experiment, while the rats receiving 2.4 mg/kg body weight of meloxicam showed mild diarrhea on 23<sup>rd</sup> day of experiment which continued till the end of experiment. The group  $G_4$  and  $G_5$  receiving 1.2 mg/kg and 2.4-mg/kg body weight of meloxicam respectively, along with aqueous extract of *Ocimum sanctum* did not show any clinical symptom. Group  $G_6$  also remained normal throughout the period of experiment. The gastrointestinal disorder due to subacute meloxicam toxicity has been reported earlier,<sup>[15]</sup> which could be the outcome of gastroenteritis as observed in present study.

### Hematological parameters

Hematological parameters like Hb concentration, TLC, PCV and TEC were estimated in control and treatment groups and are presented in Tables 1 and 2. The significant reduction in hemoglobin values of rats receiving of meloxicam indicated deleterious effect of meloxicam on blood cells. PCV values showed significant decrease in  $G_2$  and  $G_3$  group as compared to control. There was marked elevation in neutrophils count of rats from  $G_2$ ,  $G_3$  and  $G_5$  groups. Lymphocytes count showed relative decrease in group  $G_2$ ,  $G_3$  and  $G_5$  as a result of neutrophilia in these groups. The monocyte and eosinophils fluctuated within the normal range. Further, it was observed that the *Ocimum sanctum* could counteract untoward effect of meloxicam up to 1.2 mg/kg body weight. When meloxicam was fed orally at 2 mg/kg for 16 days, Alencar, et al.<sup>[16]</sup> reported anemia in meloxicam intoxicated dogs and substantiated the findings of present experiment. The increase in leucocyte count recorded in the present study could be attributed to ulcerogenic effect of meloxicam as observed in present study and as reported earlier.<sup>[16]</sup> *Ocimum sanctum* supplementation indicated protective role of *Ocimum sanctum* at low dose toxicity of meloxicam in the rats.

### Serum biochemical parameters

The serum biochemical parameter studied during the present experiment is presented in Table 3. The observations revealed that the rats treated with 1.2 mg ( $G_2$ ) and 2.4 mg ( $G_3$ ) of meloxicam resulted in two-fold elevation of alkaline phosphatase value, significantly increased serum bilirubin, serum creatinine when compared with control group. The group  $G_4$  and  $G_5$  receiving meloxicam and *Ocimum sanctum* showed relatively reduced values as compared to group

**Table 1: Mean hemoglobin, total erythrocyte count, packed cell volume and total leucocyte count of rats from various treatment groups**

Experimental groups	Hb (g/dl)	TEC ( $10^9$ /cumm)	PCV (%)	WBC ( $10^3$ /cumm)
$G_1$	13.16 <sup>c</sup> ± 0.35	8.84 ± 0.16	44.5 <sup>b</sup> 0.88	8.63 ± 0.69
$G_2$	9.80 <sup>a</sup> ± 0.57	7.68 0.71	36.5 <sup>a</sup> ± 0.71	10.33 ± 1.43
$G_3$	9.63 <sup>a</sup> ± 0.28	7.02 ± 0.72	35.83 <sup>a</sup> ± 1.27	12.08 ± 0.49
$G_4$	11.36 <sup>b</sup> ± 0.44	8.73 ± 0.25	42.33 <sup>b</sup> ± 0.84	8.72 ± 0.95
$G_5$	9.80 <sup>a</sup> ± 0.35	7.68 ± 0.71	37.65 <sup>a</sup> ± 0.65	10.05 ± 1.57
$G_6$	12.16 <sup>bc</sup> ± 0.57	8.14 ± 0.35	42.33 <sup>b</sup> ± 1.20	9.06 ± 0.81

Values are mean ± S.E. for 6 rats in each group. Values not sharing a common superscript in a column differ significantly ( $P < 0.05$ ).

**Table 2: Mean differential leucocyte count of rats from various treatment groups**

Experimental groups	Neutrophils	Lymphocytes	Monocytes	Eosinophils	Basophils
$G_1$	25.5 <sup>a</sup> ± 2.15	68.66 <sup>c</sup> ± 2.48	3.50 ± 0.61	2.33 ± 0.49	0.00 ± 0.00
$G_2$	55.5 <sup>c</sup> ± 2.43	38.00 <sup>a</sup> ± 3.09	3.66 ± 0.76	3.00 ± 1.54	0.00 ± 0.00
$G_3$	60.5 <sup>c</sup> ± 4.26	33.33 <sup>a</sup> ± 3.98	4.50 ± 1.11	1.66 ± 0.91	0.00 ± 0.00
$G_4$	37.66 <sup>b</sup> ± 4.16	51.66 <sup>b</sup> ± 6.84	6.50 ± 1.38	4.16 ± 3.02	0.00 ± 0.00
$G_5$	61.16 <sup>c</sup> ± 3.72	31.00 <sup>a</sup> ± 3.08	3.50 ± 0.88	4.33 ± 1.49	0.00 ± 0.00
$G_6$	33.5 <sup>ab</sup> ± 2.07	59.00 <sup>bc</sup> ± 2.9	4.66 ± 0.61	2.83 ± 1.07	0.00 ± 0.00

Values are mean ± S.E. for 6 rats in each group. Values not sharing a common superscript, in a column differ significantly ( $P < 0.05$ ).

**Table 3: Mean SGPT, SGOT, total bilirubin and alkaline phosphatase of rats from various treatment groups**

Experimental groups	SGPT (IU/L)	SGOT (IU/L)	Total bilirubin (IU/L)	Alkaline phosphatase (IU/L)	Creatinine (IU/L)
G <sub>1</sub>	81.53 ± 8.26	162.22 ± 15.17	0.32 <sup>a±0.03</sup>	62.31 <sup>a</sup> ± 3.52	0.37 <sup>a</sup> ± 0.02
G <sub>2</sub>	87.37 ± 15.19	245.90 ± 11.71	0.66 <sup>ab</sup> ± 0.06	114.82 <sup>b</sup> ± 28.74	0.44 <sup>a</sup> ± 0.08
G <sub>3</sub>	84.34 ± 16.33	231.09 ± 14.47	1.00 <sup>bc</sup> ± 0.29	142.55 <sup>c</sup> ± 26.99	0.86 <sup>b</sup> ± 0.13
G <sub>4</sub>	85.99 ± 8.10	238.06 ± 41.75	0.64 <sup>ab</sup> ± 0.02	65.87 <sup>a</sup> ± 14.91	0.30 <sup>a</sup> ± 0.05
G <sub>5</sub>	70.63 ± 3.2	210.16 ± 14.37	0.68 <sup>abc</sup> ± 0.05	67.26 <sup>a</sup> ± 9.37	0.41 <sup>a</sup> ± 0.09
G <sub>6</sub>	79.20 ± 3.98	183.83 ± 25.40	1.12 <sup>c</sup> ± 0.21	57.56 <sup>a</sup> ± 6.91	0.36 <sup>a</sup> ± 0.04

Values are mean ± S.E. for 6 rats in each group. Values not sharing a common superscript in a column differ significantly ( $P < 0.05$ ).

**Table 4: Mean tissue lipid peroxidase on liver and kidneys of rats from various treatment groups**

Experimental groups	Lipid peroxidase (Nmoles of MDA/g tissue)	
	Liver	Kidney
G <sub>1</sub>	38.20 <sup>a</sup> ± 3.06	39.61 <sup>ab</sup> ± 7.12
G <sub>2</sub>	113.14 <sup>c</sup> ± 27.88	72.82 <sup>b</sup> ± 11.00
G <sub>3</sub>	119.87 <sup>c</sup> ± 28.47	91.15 <sup>c</sup> ± 26.27
G <sub>4</sub>	46.53 <sup>a</sup> ± 8.4	52.69 <sup>ab</sup> ± 9.40
G <sub>5</sub>	63.84 <sup>ab</sup> ± 14.0	60.06 <sup>ab</sup> ± 5.34
G <sub>6</sub>	59.03 <sup>ab</sup> ± 10.35	34.41 <sup>a</sup> ± 1.78

Values are mean ± S.E. for 6 rats in each group. Values not sharing a common superscript in a column differ significantly ( $P < 0.05$ ).

treated with meloxicam alone. The elevated levels suggests hepatotoxic and nephrotoxic effect of meloxicam. Similar serum biochemical changes in case of a female patient with rheumatoid arthritis who developed acute cytolytic hepatitis due to meloxicam was reported earlier.<sup>[17]</sup> Dogs treated with meloxicam at 1 mg and 2-mg/kg body weight had elevation of serum creatinine<sup>[18]</sup> and is in agreement with the present findings. Serum biochemical parameters of *Ocimum sanctum* treated rats were not elevated, in turn indicating its protective action.

The values of parameters like LPO, GSH and antioxidant enzymes like SOD, activities of experimental rats were shown in Tables 4 and 5. The values of LPO in liver and kidney in G<sub>2</sub> and G<sub>3</sub> groups were found significantly elevated as compared with control group showed significant difference. Present observation is in accordance with Villages *et al.*,<sup>[19]</sup> who also observed elevation in oxidative stress parameters at higher dose of meloxicam (15 mg/kg) in rats. The other parameters studied did not showed any significant difference between the groups. However, there was an increase in the glutathione content and SOD values of meloxicam toxicated rats.

### Gross findings

Grossly, rats from group G<sub>2</sub> showed congestion of glandular stomach. The mucus membrane of intestine was thickened and become rough. Rats from group G<sub>3</sub> showed hemorrhages

in glandular stomach [Figure 1], congestion and subserosal hemorrhages in intestine [Figure 2]. Congestion and small necrotic area was seen on liver surface. Similar changes were also recorded in-group G<sub>5</sub>. Group G<sub>4</sub> did not show any detectable macroscopic lesion. Control groups G<sub>1</sub> and G<sub>6</sub> also did not show any gross pathological lesions.

### Microscopic findings


Liver from group G<sub>2</sub> revealed varied degree of changes from granular degenerative changes to necrosis of hepatocytes [Figure 3]. Infiltration of inflammatory cells and fibrous tissue in perivascular, perilobular and periportal areas of liver was seen [Figure 4]. Liver from group G<sub>3</sub> revealed granular degenerative changes and necrosis of hepatocytes from perivascular areas. Mild congestion in liver and proliferation of Kupffer cell was pronounced in the liver from group G<sub>4</sub> and G<sub>5</sub>.

Stomach from Group G<sub>2</sub> revealed intact mucosal epithelium with mild infiltration of round cells in between the glands. Stomach from group G<sub>3</sub> showed ulceration, hemorrhages and desquamation of mucosal epithelium at many places [Figure 5]. Stomach from group G<sub>4</sub> and G<sub>5</sub> revealed similar changes as that of group G<sub>2</sub>. Intact intestinal mucosa with mild submucosal infiltration was seen in rats from group G<sub>2</sub> [Figure 6], while intestine group G<sub>3</sub> showed hydropic degenerative changes in intestinal epithelium, hemorrhages and desquamation of the epithelial mucosa [Figure 7]. The group G<sub>4</sub> and G<sub>5</sub> did not show any detectable abnormality in intestine. Kidney from group G<sub>2</sub> showed granular degenerative changes in the tubular epithelium, while kidneys from group G<sub>3</sub> revealed extensive degenerative changes of tubular epithelium and interstitial nephritis [Figure 8], Hemorrhages were the prominent findings. The kidneys from G<sub>4</sub> rats did not show any changes other than mild degenerative changes of tubular epithelium. In-group G<sub>5</sub>, severe granular degenerative changes were observed in tubular epithelium leading occlusion of tubular lumen. Glomerular tuft of kidney from group G<sub>5</sub> showed hemorrhages. The section of brain from group G<sub>3</sub> showed perivascular edema, gliosis [Figure 9], and necrosis of neurons. Section of brain from other groups did not reveal any histological abnormalities. Focal myocardial necrosis and infiltration of inflammatory cells in between the cardiac muscles was evident in hear of

**Table 5: Mean reduced glutathione and super oxide dismutase on liver and kidneys of rats from various treatment groups**

Experimental groups	Reduced glutathione (Mmoles/g of wet tissue)		Super oxide dismutase (Units/mg protein)	
	Liver	Kidney	Liver	Kidney
G <sub>1</sub>	2.05 ± 1.39	1.46 ± 0.09	11.92 ± 0.58	8.58 ± 1.63
G <sub>2</sub>	3.26 ± 9.53	2.44 ± 0.33	153.27 ± 58.70	49.63 ± 6.49
G <sub>3</sub>	2.67 ± 7.8	2.72 ± 0.64	152.65 ± 62.54	118.24 ± 67.40
G <sub>4</sub>	3.71 ± 1.40	1.66 ± 0.19	171.63 ± 58.02	115.94 ± 68.21
G <sub>5</sub>	3.82 ± 7.73	1.94 ± 0.21	160.36 ± 59.80	279.65 ± 119.10
G <sub>6</sub>	2.13 ± 2.16	1.17 ± 0.14	80.49 ± 7.42	60.89 ± 14.08


Values are mean ± S.E. for 6 rats in each group. Values not sharing a common superscript in a column differ significantly ( $P < 0.05$ ).



**Figure 1:** Hemorrhagic spot in the glandular stomach of meloxicam treated group G<sub>3</sub>




**Figure 2:** Subserosal hemorrhages in duodenum of group



**Figure 3:** Section of liver (Group G<sub>2</sub>) showing vacuolar degenerative changes and necrosis of hepatocytes (H and E, × 200)


rats from group G<sub>3</sub> [Figure 10]. Section of heart from other groups did not reveal any histological abnormalities.

Histomorphological observations from group G<sub>4</sub> indicated hepatoprotective role of *Ocimum sanctum* against meloxicam toxicity at low dose level and confirms the findings of hepatoprotective action of *Ocimum sanctum* as reported




**Figure 4:** Section of liver (Group G<sub>2</sub>) showing infiltration of inflammatory cells and fibrous tissue in perivascular, perilobular and periportal areas of liver. (H and E, × 200)


earlier.<sup>[20]</sup> The findings of gastric hemorrhages and intestinal ulceration due to meloxicam toxicity is reported earlier<sup>[16]</sup> and confirmed the findings of present study. Castro, *et al.*<sup>[21]</sup> also reported mononuclear infiltration in stomach of meloxicam treated dogs. The stomach and intestinal sections from groups receiving meloxicam and extract of *Ocimum sanctum* did not show any microscopic lesions in the stomach




**Figure 5:** Section of glandular stomach (Group G<sub>3</sub>) showing ulcer (H and E, ×100)




**Figure 6:** Section of intestine (Group G<sub>2</sub>) showing infiltration of inflammatory cells and hydropic degeneration in mucosal epithelium (H and E, ×100)




**Figure 7:** Section of intestine (Group G<sub>2</sub>) showing hydropic degeneration in mucosal epithelium of intestine and sloughing (H and E, ×100)



**Figure 8:** Section of kidney (Group G<sub>3</sub>) showing interstitial nephritis and hemorrhages (H and E, ×100)



**Figure 9:** Section of brain (Group G<sub>3</sub>) showing gliosis and perivascular edema (H and E, ×100)



**Figure 10:** Section of heart (Group G<sub>3</sub>) showing myocardial infarct (H and E, ×100)

and intestine. This points out towards a protective action of *Ocimum sanctum*, which confirms the earlier findings of Mandal *et al.*<sup>[22]</sup> who reported antitumorogenic properties of *Ocimum sanctum* in rats induced by aspirin. Inhibition of gastric secretion and ulceration in albino rats by *Ocimum sanctum* was reported earlier,<sup>[23]</sup> and could be the possible mechanism of protection by *Ocimum sanctum*. Khan *et al.*<sup>[24]</sup> reported interstitial nephritis and nephritic syndrome as side effect of NSAIDs due to transient interference in electrolytic homeostasis. The *Ocimum sanctum* could not counteract nephrotoxic effect of meloxicam in present experiment. Sections of heart from group G<sub>3</sub> revealed focal myocardial necrosis and infiltration of inflammatory cells, which could be the outcome of sub acute meloxicam toxicity. Group G<sub>5</sub> revealed myocardial infarcts and focal necrotic area in the section of heart which might be due to cumulative toxic action of meloxicam on cardiac muscle.

From the present study we could conclude that Meloxicam at both dose levels causes anemia, hepatotoxicity, nephrotoxicity and produces lesion in GI tract. The lesions can also be noticed in the heart and brain. Supplementation with *Ocimum sanctum* could counteract the toxic effect only at lower dose level. Even though many previous studies observed the toxicity of meloxicam, the basic mechanism of the toxicity is not understood thoroughly. A further detailed study at molecular level is needed to know the exact mechanism of meloxicam toxicity, and the protective mechanism of *Ocimum sanctum* could be explored.

## ACKNOWLEDGMENT

Research presented in this manuscript was supported by grant from Department of Veterinary Pathology, Nagpur Veterinary College, Nagpur, India.

## REFERENCES

- Fleischmann R, Iqbal I, Slobodin G. Meloxicam. Expert Opin Pharmacother 2002;3:1501-12.
- Ogino K, Harada Y, Hatanaka K. Inhibitory effect of Meloxicam on COX-2. Jap Pharmacol Soc. The 69<sup>th</sup> Annu. Meet Nagasaki, Japan; 1996.
- Hawkey C, Kahan A, Steinbrück K, Alegre C, Baumelou E, Bégaud B, *et al.* Gastrointestinal tolerability of meloxicam compared to diclofenac in osteoarthritis patients. International MELISSA Study Group. Meloxicam Large-scale International Study Safety Assessment. Br J Rheumatol 1998;37:937-45.
- Vane JR, Botting RM. Mechanism of action of aspirin-like drugs. Semin Arthritis Rheum 1997;26 (6 Suppl 1):2-10.
- Kirtikar KR, Basu BD. In :*Ocimum sanctum* in Indian Medicinal Plants. Published by LB Basu, Allahabad.1965.
- Bhargava KP, Singh N. Anti-stress activity of *Ocimum sanctum* Linn. Ind J Med Res 1981;73:443-51.
- Rajasekaran M. Antifertility and some pharmacological studies on Triterpines. Ph.D. Thesis. India: University of Madras; 1988.
- Rosenthaler. The Chemical investigation of plants. 1<sup>st</sup>ed. London: Bell and sons;1930. p. 36.
- Benjamin MM. Outline of Veterinary Clinical Pathology. Kalyani Publishers New Delhi, 1985.
- Luna AG. Manual of histological staining methods of the Armed Force Institute of Pathology. 3<sup>rd</sup> ed. London: Mc Graw Hill Book Co; 1968. p. 24-5.
- Shafiq-Ur-Rehman. Lead induced regional lipid peroxidation in brain. Toxicol Lett 1984;21:333-7.
- Sedlak J, Lindsay RH. Estimation of total, protein bound and non-protein sulfhydryl groups in tissue with Ellmans reagent. Anal Biochem 1968;25:192-205.
- Marklund S, Marklund G. Involvement of the super oxide anion radical in the autooxidation of pyrogallol and a convenient assay for super oxide dismutase. Eur J Biochem 1974;47:469-74.
- Snedecor GW, WG Cochran. Statistical Method. 8<sup>th</sup> ed. New Delhi: Oxford and IBH Publishing Company; 1995.
- MacDonald TM, Morant SV, Goldstein JL, Burke TA, Pettitt D. Channeling bias and the incidence of gastrointestinal hemorrhage in users of meloxicam, coxibs, and older, non-specific non-steroidal anti-inflammatory drugs. Gut 2003;52:1265-70.
- Alencar MM, Pinto MT, Oliveira DM, Pessoa AW, De P, Candido IA, *et al.* Margin of safety of Meloxicam in dogs: deleterious effects on blood cells and gastrointestinal tract. Cienc Rural 2003;33: 525-32.
- Staerckel P, Horsmans Y. Meloxicam-induced liver toxicity. Acta Gastroenterol Belg 1999;62:255-6.
- Alencar MM, Pinto MT, Oliveira DM, Pessoa AW, De P, Candido IA, *et al.* Evaluation of the pharmacological activity of meloxicam on renal function in dogs. Cienc Animal 2002;12:25-33.
- Villegas I, Alarcón de la Lastra C, La Casa C, Motilva V, Martín MJ. Effect of food intake and oxidative stress on intestinal lesions caused by meloxicam and piroxicam in rats. Eur J Pharmacol 2001;414:79-86.
- Sankaran JR. An all India multicentric clinical survey on a herbal clue Tefroli for Hepatitis. J National Integ Med Assoc 1984;26:255-61.
- Castro MB, Develey FF, Azarias RE, Mourao GB, Marcos HR, Barbosa AM, *et al.* Clinical, gastroscopical and histopathological findings in dogs treated with meloxicam. Ars Veterinaria 1999;15:154-9.
- Mandal S, Das DN, De K, Ray K, Roy G, Chaudhuri SB, *et al.* *Ocimum sanctum* Linn-a study on gastric ulceration and gastric secretion in rats. Indian J Physiol Pharmacol 1993;37:91-2.
- Sen P. Therapeutic potentials of Tulsi: from experience to facts. Drugs News & Views 1993;1:15-21.
- Khan KN, Venturini CM, Bunch RT, Brassard JA, Koki AT, Morris DL, *et al.* Interspecies differences in renal localization of cyclooxygenase isoforms: Implications in nonsteroidal anti-inflammatory drug-related nephrotoxicity. Toxicol Pathol 1998;26:612-20.

**How to cite this article:** Mahaprabhu R, Bhandarkar AG, Jangir BL, Rahangadale SP, Kurkure NV. Ameliorative effect of *Ocimum Sanctum* on meloxicam induced toxicity in wistar rats. Toxicol Int 2011;18:130-6.

**Source of Support:** Grant from Department of Veterinary Pathology, Nagpur Veterinary College, Nagpur, India. **Conflict of Interest:** None declared.