

## Research Article

# Antitussive Activity of the Water-Extracted Carbohydrate Polymer from *Terminalia chebula* on Citric Acid-Induced Cough

Gabriela Nosalova,<sup>1</sup> Ludovit Jurecek,<sup>1</sup> Udipta Ranjan Chatterjee,<sup>2</sup>  
Sujoy Kumar Majee,<sup>2</sup> Slavomir Nosal,<sup>3</sup> and Bimalendu Ray<sup>2</sup>

<sup>1</sup> Department of Pharmacology, Jessenius Faculty of Medicine in Martin, Sklabinska 26, 036 01 Martin, Slovakia

<sup>2</sup> Natural Products Laboratory, Department of Chemistry, The University of Burdwan, Burdwan, West Bengal 713 104, India

<sup>3</sup> Clinic of Pediatric Anesthesiology and Intensive Medicine, Jessenius Faculty of Medicine and Martin University Hospital, Kollárova 2, 036 01 Martin, Slovakia

Correspondence should be addressed to Bimalendu Ray; [bimalendu.ray@gmail.com](mailto:bimalendu.ray@gmail.com)

Received 8 March 2013; Revised 16 May 2013; Accepted 18 May 2013

Academic Editor: Jairo Kenupp Bastos

Copyright © 2013 Gabriela Nosalova et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Terminalia chebula*, a medicinal plant, is widely used in the management of various diseases. As the water extract of its dried ripe fruit is a frequently used preparation, we decided to look for bioactive polysaccharide in this extract. We demonstrate that the obtained polysaccharide fraction, CP, contained a highly branched arabinogalactan protein having a (1 → 3)-, (1 → 6)- and (1 → 3, 6)-linked β-D-Galp together with (1 → 5)- and (1 → 3)-linked α-L-Araf and nonreducing end units of α-L-Araf. This polymer possesses strong antitussive property. Our results showed that the number of citric acid-induced cough efforts decreased significantly after the oral application of polysaccharide fraction in a dose of 50 mg kg<sup>-1</sup> body weight. Its antitussive efficacy was higher than cough suppressive effect of standard drug codeine. Therefore, traditional aqueous extraction method provides a major polysaccharide, which induces a pharmacological effect: this could represent an attractive approach in phytotherapeutic managements.

## 1. Introduction

The cough reflex represents the most important defensive reflex of the airways, which together with mucociliary transport system forms the main mechanism for the cleaning of respiratory tract [1]. Coughing protects the breathing passages from blocking, thereby preventing the infected mucus from falling into lungs and bronchial tubes, which could be very dangerous. The cough reflex belongs to the most frequent symptoms of respiratory system diseases and is the most common reason why sick patients visit physicians [2, 3]. Pathological cough has significant impact on patient's quality of life observed either in physical activity or psychosocial domain. The present therapies are often limited for the lack of effective medications. Moreover, most of the existed medicines could bring about inevitable side effects. Thus, there is

a need to develop new compounds with strong antitussive activity accompanied by favourable pharmacological and clinical properties. Natural products have been invaluable as biologically validated platforms for drug development [4–6]. Indeed, a series of review articles demonstrated the continuing and valuable contributions of nature as a source of lead compounds that have provided the basis and inspiration for the synthesis of new drugs [7, 8].

*Terminalia chebula*, a medicinal plant belonging to genus *Terminalia* (family, Combretaceae), is cultivated in Tibet, Taiwan, China, and India [9, 10]. The dried ripe fruit of *T. chebula*, locally known as haritaki, has been widely used for the treatment of fever, sore throat, cough, vomiting, hiccough, bleeding, piles, diarrhoea, gout, and heart and bladder diseases [11]. Haritaki fruits are reported to have free radical

scavenging activities [12]. It is active against different gram-positive and gram-negative bacteria [13]. Phytochemical analysis of *T. chebula* shows the presence of gallic acid, chebulagic acid, corilagin, mannitol, ascorbic acid (vitamin C), and other compounds [14, 15]. The extract of *T. chebula* contained 24–64% tannins, 4.4% gallic acid, 22.2% total carbohydrate, and 5.3% uronic acid [15, 16]. Despite the general interest in phytoconstituents, it remains ironic that research on the chemical and biological aspects of the high molecular weight bioactive compounds has been neglected, although the aqueous extract of this fruit, which no doubt contains high molecular weight compound, is used in traditional medicine. Notably, many polysaccharides from medicinal plants exhibit a large range of pharmacological effects [17–20]. Hence, the aim of the present study was, first of all, to characterize the polysaccharide from *T. chebula* fruits. In addition, the antitussive activity of the isolated polysaccharide in terms of the number of cough efforts and specific airway resistance *in vivo* conditions in awaken male TRIK strain guinea pigs was evaluated.

## 2. Materials and Methods

**2.1. Plant Materials.** Dry fruit powder (batch no. HTC008) of *Terminalia chebula* was purchased from Divya Pharmacy (Unit A-1, Haridwar, Uttarakhand, India).

**2.2. Chemicals.** The chemicals used were of analytical grade or the best available.

Codeine phosphate was purchased from Lachema (Brno, Czech Republic). Codeine phosphate, citric acid, and all samples were dissolved in water for injection.

**2.3. General Methods.** Ultraviolet-visible (UV-VIS) spectra were recorded on a UV-24500 spectrophotometer (Shimadzu, Japan). IR spectra were obtained on a FT IR spectrophotometer (PerkinElmer FT/IR Spectrum RX 1) using a KBr discs containing finely ground samples. The  $^1\text{H}$  NMR spectrum of the water-extracted carbohydrate polymer (CP) was recorded on a Bruker 500 spectrometer (Bruker Biospin AG, Fallanden, Switzerland) operating at 500 MHz. These macromolecules were deuterium-exchanged by lyophilization with  $\text{D}_2\text{O}$  ( $\times 3$ ) and then examined as 1% solutions in  $\text{D}_2\text{O}$  (99.96 atom % D). GC was carried out with a Shimadzu GC-17A chromatograph fitted with a flame ionization detector and a DB-225 column (30 m  $\times$  0.53 mm i.d.), using a program that maintained an isocratic temperature of 210°C for 18 min and helium as gas vector. GC-MS was performed with a Shimadzu QP 5050A GC-MS instrument at 70 eV. Conditions for GC-MS were as described previously [21]. Dialysis (molecular cutoff 12 kDa; Sigma-Aldrich) against distilled  $\text{H}_2\text{O}$  were performed with continuous stirring. Evaporations were carried out under reduced pressure at around 50°C (SB 1100 Rotary Evaporator; Eyela, Tokyo, Japan), and small volumes of sample solutions were freeze-dried (Cool Safe 55-F freeze drier; Scanvac, Lynge, Denmark). Moisture content was determined by drying the ground material in an oven at 110°C for 3 h. Total carbohydrate content was estimated by

the phenol-sulfuric acid assay [22] using glucose as standard. Total uronic acid was assayed as anhydrogalacturonic acid using *m*-hydroxydiphenyl color reagent [23]. Estimation of soluble protein content was performed by the Folin-Ciocalteu method. Amino acids were released by hydrolysis with 6 M HCl at 110°C for 22 h in a sealed tube and were analyzed as described previously [24]. Methylation was carried out by the method of Sims and Bacic [25]. In the methylation procedure, free hydroxyl groups in the carbohydrates were deprotonated and methylated, then the glycosidic linkages were hydrolyzed, and the partially methylated monosaccharides were reduced to alditols and acetylated. The partially methylated alditol acetates were analysed by GC and GC-MS using a DB-225 column (J&W Scientific) and using conditions as described previously [21]. The partially methylated alditol acetates were identified by the measurement of relative retention times, methoxyl substitution pattern as obtained from GC-MS, and carbohydrate composition of the nonmethylated polymers.

**2.4. Isolation of Polysaccharide.** Extraction of polysaccharide was conducted by stirring a suspension of *Terminalia chebula* fruit powder (20 g) in water (300 mL, pH 6.0) at 25–32°C for 12 h. Toluene was added to inhibit microbial growth. Separation of the residue from the extract was performed by filtration through a glass filter (G-2). The insoluble material was extracted twice more under similar condition at a solute to solvent ratio of 1:100 (w/v). The combined liquid extract was dialyzed extensively against water and lyophilized. The recovered material was dissolved in water, precipitated by the addition of ethanol (4 volumes), and then collected by centrifugation ( $\times 3$ ). The final pellet was dissolved in water and lyophilized to yield the water-extracted carbohydrate polymer, named CP (0.563 g).

The arabinogalactan protein (AGP) was isolated according to Schultz et al. [26]. Briefly, to a solution of CP in 1% NaCl (w/v) was added an equal volume of  $\beta$ -glucosyl Yariv reagent also in 1% NaCl. The mixture was kept at 4°C for 18 h and then centrifuged. The pellet was washed with 1% NaCl followed by pure methanol ( $\times 3$ ), dried, and treated with sodium metabisulfite (10%). The resulting solution was dialyzed and freeze-dried to yield AGP.

**2.5. Animals.** Adult healthy awaken male TRIK strain guinea pigs, weighing 200–350 g, supplied by the department of experimental pharmacology, Slovak Academy of Science, Dobra Voda, Slovakia, were kept in faculty animal house with food and water *ad libitum* and with a standard air conditioning system. The animals were kept one week in quarantine before starting the experiment.

The carbohydrate polymer (CP) from *T. chebula* fruit was tested on individual group of animals consisting of 20 guinea pigs, whereas the control agents (“positive” control codeine, “negative” control water for injection) were tested on 8 guinea pigs. All tested compounds were applied by peroral route of administration: carbohydrate polymer (CP) in the dose of 50 mg kg<sup>-1</sup>, codeine in the dose 10 mg kg<sup>-1</sup>, and water for injection in the dose 1 mL kg<sup>-1</sup> body weight.

The experimental protocols were approved by the Institutional Ethics Committee of the Jessenius Faculty of Medicine, Comenius University in Martin, Slovakia, registered in Institutional Review Board/Institutional Ethic Board Office (IRB 00005636), complied with Slovakian and European Community regulations for use of laboratory animals, and followed the criteria of experimental animal's welfare.

**2.5.1. Assessment of Chemically Induced Cough Reflex and Airways Defense Reflexes.** Antitussive activity and reactivity of the airway smooth muscle *in vivo* were determined as described [27, 28]. Awaken guinea pigs were individually placed in a body plethysmograph box (HSE type 855, Hugo Sachs Elektronik, Germany) and restricted so that the head protrudes into the nasal chamber and the neck were sealed with a soft diaphragm.

The cough reflex was induced by aerosol of citric acid in a concentration 0.3 M. The citric acid aerosol, generated by a jet nebulizer (PARI jet nebulizer, Paul Ritzau, Pari-Werk GmbH, Germany, output  $5 \text{ L s}^{-1}$ , particles mass median diameter  $1.2 \mu\text{m}$ ), was delivered to the head chamber of the plethysmograph for 3 min. The cough effort was defined as sudden PC-recorded enhancement of expiratory flow associated with typical cough motion and sound followed by two trained observers. Both influence on citric acid-induced cough and specific airway resistance were registered before any agent application (values labeled as N in graphs) and after that in 30, 60, 120, and 300 min time intervals. The minimal time interval between two measurements was 2 h for cough receptors adjustment as well as adaptation of laboratory animals on that kind of irritation [20, 29].

The reactivity of the airway smooth muscle *in vivo* conditions was expressed as values of specific airway resistance calculated according to Pennock et al. [30] by time difference between pressure changes in head and chest parts of body plethysmograph during normal breathing pattern. This noninvasive plethysmograph technique is commonly used for evaluation of bronchoactive substances effect [31]. The value of specific airway resistance is proportional to phase difference between nasal and thoracic respiratory airflow. The bigger the phase difference the higher is the value of specific airway resistance and hence, higher degree of bronchoconstriction. The values of sRaw were measured consecutively after the citric acid exposure and the cough response registration during 1 min interval.

**2.5.2. Statistics.** The changes of number of the citric acid-induced cough efforts and values of specific airways resistance in 30, 60, 120, and 300 min intervals were compared and statistically evaluated with initial data (N means number of cough efforts before application of agents). Student's *t*-test was used for the statistical analysis of the obtained results. Data are presented as mean  $\pm$  standard error of the mean (SEM).  $P < 0.05$  was considered statistically significant. Significance of  $P < 0.05$ ,  $P < 0.01$ , and  $P < 0.001$  is shown by one, two, or three asterisks, respectively.

### 3. Results and Discussion

In Indian ayurvedic system of medicine, a decoction of *Terminalia chebula* fruit powder in water is used as herbal remedy for chronic coughs and breathlessness [11]; therefore dry fruit powder of this medicinal plant was extracted with water. The yield of the water-extracted carbohydrate polymer (named as CP), after fractional precipitation with ethanol, was 29 mg per gram of dry fruit powder. Fraction CP contains 64% (w/w) neutral sugar along with 13% (w/w) protein. Aspartic acid (4 mol%), threonine (4 mol%), serine (5 mol%), glutamic acid (37 mol%), proline (9 mol%), glycine (10 mol%), alanine (6 mol%), cysteine (1 mol%), valine (3 mol%), isoleucine (2 mol%), leucine (5 mol%), tyrosine (1 mol%), phenylalanine (2 mol%), histidine (2 mol%), lysine (6 mol%), and arginine (3 mol%) were the amino acids associated with the protein present in fraction CP. The uronide content of this fraction was 2.3% (w/w). Sugar compositional analysis revealed that fraction CP consists mainly of arabinose, galactose, and glucose as the major neutral sugar in the molar ratio of 61 : 21 : 18. Therefore, this fraction may contain arabinogalactan. The high amount of glucose probably arises from starch, the presence of which has been confirmed by the blue coloration it developed with iodine reagent. Considering that the water extracted carbohydrate polymer (CP) contained galactosyl and arabinosyl residues as the major sugars and its protein content is 13%, we have tested its reactivity with  $\beta$ -glucosyl Yariv reagent that specifically precipitates arabinogalactan proteins (AGP). We found that the major part of CP was Yariv soluble. Sugar compositional analysis of this precipitate shows that it consisted mainly of galactose (77 mol%) residues and, to a lesser extent, arabinose (23 mol%) residues, confirming the presence of AGP.

We used FT-IR spectroscopy to assess the arabinogalactan protein structure. FT-IR spectroscopy is a vibrational spectroscopic method that can quantitatively detect, without derivatization, the relative amounts of functional groups, including  $-\text{OH}$ ,  $\text{NH}_2$ , amides, methyl, methylene, carbohydrate fingerprint region, with absorbance at certain wavenumber. FT-IR spectrum of the CP, as shown in Figure 1, contained bands characteristics of polysaccharide and protein. The stretching vibrations of  $-\text{OH}$  and  $-\text{NH}_2$  groups in sugar and protein moieties were indicated by a broad band around  $3400 \text{ cm}^{-1}$ . Low-intensity bands around  $2926 \text{ cm}^{-1}$  derived from stretching resonances of  $\text{CH}$ ,  $\text{CH}_2$ , or  $\text{CH}_3$  groups in both monosaccharide and amino acid residues were also observed. It also contained bands at 1098 and  $1066 \text{ cm}^{-1}$  characteristic of carbohydrate. The presence of protein component was demonstrated by the occurrence of two bands of decreasing intensity: amide I towards  $1643 \text{ cm}^{-1}$  and amide II towards  $1542 \text{ cm}^{-1}$  with  $\text{N-H}$  and  $\text{C-N}$  links as reported previously for other proteins [32]. The chemical composition of CP and its IR spectrum suggests the presence of arabinogalactan protein.

Methylation analysis showed that CP is a highly branched polysaccharide, containing nonreducing end units of Ara<sub>f</sub> (2,3,5-Me<sub>3</sub>-Ara) (16%), and Gal<sub>p</sub> (2,3,4,6-Me<sub>4</sub>-Gal) (4%).




FIGURE 1: FT-IR spectrum of the carbohydrate polymer (CP) from *T. chebula*. The spectrum shows amide-I and amide-II bands at 1643 and 1542  $\text{cm}^{-1}$  and carboxylate ion band at 1375  $\text{cm}^{-1}$  together with bands at 1066 and 1098  $\text{cm}^{-1}$  characteristics of carbohydrate.

The arabinosyl units were substituted at O-5 (Araf) and/or O-4 (Arap) and O-3, in accord with 2,3-Me<sub>2</sub>-Ara (6%) and 2,5-Me<sub>2</sub>-Ara (4%) derivatives, respectively. The glucosyl units were substituted at O-4 and O-4,6, as demonstrated by the presence of 2,3,6-Me<sub>3</sub>-Glc (7%) and 2,3-Me<sub>2</sub>-Glc (1%) derivatives, respectively. The galactopyranosyl units are, mainly, 3-O- and 3,6-di-O-substituted, in accord with 2,4,6-Me<sub>3</sub>-Gal (19%) and 2,4-Me<sub>2</sub>-Gal (36%) methylated derivatives, respectively. Also present were 6-O-substituted galactopyranosyl units, as shown by the 2,3,4-Me<sub>3</sub>-Gal (7%) derivatives. Taken together, the glycosidic linkage composition of CP suggests the presence of arabinogalactan.

The anomeric signals in the <sup>1</sup>H NMR spectrum (Figure 2) of CP were assigned according to sugar composition, glycosidic linkage makeup, and data in the literature [33, 34]. The signal at  $\delta$  5.05 was assigned to anomeric protons (H1) of the nonreducing terminal  $\alpha$ -Araf and the signals in between  $\delta$  5.23 and  $\delta$  5.32 originated from the anomeric protons (H1) of 1,5- and 1,3-linked Araf, respectively. This arabinogalactan also showed one group of H1 signals from  $\delta$  4.44 to 4.48 and a second group of H1 signals from  $\delta$  4.5 to 4.52 originating from resonances of anomeric protons (H1) of different  $\beta$ -Galp residues of the  $\beta$ -1,3-linked galactan backbone. The signal at  $\delta$  4.3 was assigned to H4 protons of  $\beta$ -Galp residues. This spectrum also contained resonances characteristic of ring protons (H2–H5) between 3.5 and 4.2 ppm. The signal that appeared around 5.35 ppm can be attributed to anomeric proton of the Glcp moieties in CP. Signals for protein and deuterated water were designated as P and HOD, respectively. On the basis of the data obtained, it can be concluded that the carbohydrate polymer of *Terminalia chebula* fruit powder is an arabinogalactan protein.

The effect of polysaccharide component (CP) on the citric acid-induced cough reflex and reactivity of the airways smooth muscle using double chamber body plethysmograph




FIGURE 2: <sup>1</sup>H NMR spectrum at 500 MHz of the carbohydrate polymer (CP) from *T. chebula*. The spectrum of the macromolecule was recorded at 70°C in D<sub>2</sub>O solution. Insets: expansion of  $\delta_H$  6 to 7 and 2 to 3 regions. Numerical values are in  $\delta$  (ppm). The signals for protein and deuterated water are designated as P and HOD, respectively.




FIGURE 3: The influence of the carbohydrate polymer (CP) from *T. chebula*, codeine (positive control) and water for injection (negative control), on the citric acid-induced cough efforts in guinea pigs recorded at 30, 60, 120, and 300 min time intervals. The number of cough efforts before the application of the polysaccharides and codeine is given by N.  $P < 0.05$  was considered statistically significant, and the significance of  $P < 0.05$ ;  $P < 0.01$ ; and  $P < 0.001$  is shown by \*, \*\*, or \*\*\*, respectively. All used substances were applied by peroral route of administration: plant polysaccharides in the dose of 50  $\text{mg kg}^{-1}$ , codeine in the dose 10  $\text{mg kg}^{-1}$  and water in the dose 1  $\text{mL kg}^{-1}$ , body weight.

were tested *in vivo* conditions. The cough reflex was chemically induced by exposing the guinea pigs to an aerosol of 0.3 M citric acid for 3 minutes. Cough efforts and specific airways resistances were registered before (N) and subsequently 30, 60, 120, and 300 minutes after oral administrations of CP as well as controls. The polysaccharide component CP was administered orally to 20 healthy male awoken guinea pigs in the dose of 50  $\text{mg kg}^{-1}$  body weight. Efficiency of tested fraction was compared with antitussive activity of codeine (positive control) and water for injection (negative control). As shown in Figure 3, peroral administration of polysaccharide from *T. chebula* brought a highly significant ( $P < 0.001$ ) decrease in number of cough efforts in the dose




FIGURE 4: Total antitussive activity on guinea pigs of the carbohydrate polymer (CP) from *T. chebula* along with standard antitussive compound codeine (administered in the dose of  $50 \text{ mg kg}^{-1}$  and  $10 \text{ mg kg}^{-1}$  body weight, resp.).

of  $50 \text{ mg kg}^{-1}$  body weight. This significant reduction of the cough efforts observed at 30 minutes after administration indicates prompt onset of action of the polysaccharide component. Furthermore, the positive effect on this parameter was observed during all the study time intervals. These results supported our previous finding that several naturally occurring polysaccharides possess antitussive activity [35–38]. Importantly, the total antitussive activity of CP was quantitatively higher than that of codeine, the strongest antitussive agent used in clinical practice (Figure 4). Narcotics such as codeine are not suitable for long-term treatment due to their habit-forming properties and unfavorable side effects including depressive activities on respiratory center and expectoration of mucus from the airways. These results showed not only the validity of the use of this plant in ayurvedic medicine but also the possibility to substitute codeine with more safer and effective antitussive agent from natural source.

Airway resistance is a concept used in respiratory physiology to describe mechanical factors, which limit the access of inspired air to the pulmonary alveoli and thus determine airflow. It is dictated by, *inter alia*, the diameter of the airways. At present, the relationship between cough and bronchoconstriction is not known with certainty, although it is generally accepted that bronchodilating substances can cause cough suppression. Therefore, we have evaluated the changes of specific airway resistance as indicator of this activity. Our result suggests that the application of arabinogalactan from *T. chebula* in the dose, which provoked cough suppressive activities, did not significantly change the values of specific airway resistance (Figure 5). Therefore, this parameter do not modulates the antitussive activity of polysaccharide.

The mechanism(s) responsible for the antitussive activity of the polysaccharide from *T. chebula* cannot be defined from the present study. It may be due to spasmolytic activity, protective effects on mucous, and other mechanisms. Further studies in this area would be needed.




FIGURE 5: The influence of the carbohydrate polymer (CP) from the medicinal plant *T. chebula* and control agents (water for injection and codeine) on citric acid-induced changes of specific airway resistance (sRaw) *in vivo* conditions recorded at 0 (values labeled as *N* in graph), 30, 60, 120, and 300 min time intervals.

Arabinogalactan structures have been associated with various biological activities [18, 39–46]. Investigations of biological activities of arabinogalactan isolated from various medicinal plants emphasized the importance of highly branched side chains for the expression of the observed activities [18, 39–41, 45, 46]. Samuelsson and coworker [45] employing multivariate statistical analysis suggested that the magnitude of the biological activity of arabinogalactan is influenced by the content of certain side chains in the polymer. It has been shown that high activity correlates to large neutral side chains with high amounts of (1 → 6)- and (1 → 3,6)-linked Gal and low amounts of (1 → 4)-linked GalA. The arabinogalactan of the present study contains side chains possessing these distinctive features, and hence, these portions might be considered to be important as functional sites.

#### 4. Conclusions

The findings of this study highlight several novel and important aspects of *T. chebula*-derived polysaccharide with regard to its structure and antitussive properties. A highly branched arabinogalactan protein could be isolated from the fruit powder of this medicinal plant using traditional water extraction protocol. This water-soluble macromolecule demonstrated strong antitussive activity. Moreover, administration of this compound did not change specific airways resistance significantly. The highly branched side chains of the arabinogalactan might be the functional sites. Finally, the biological activity observed in *T. chebula* provides a scientific basis for the use of the plant in traditional medicines.

#### Conflict of Interests

The authors declare that they have no conflict of interests.

## Acknowledgments

This work was supported by the Science and Technology Assistance Agency (APVV no. 0030-07, LPP-0317-09), Center of Experimental and Clinical Respiratory cofinanced from EC sources, and Department of Science & Technology (project number SR/S1/OC-38/2012).

## References

- [1] A. B. Goldsobel and B. E. Chipps, "Cough in the pediatric population," *Journal of Pediatrics*, vol. 156, no. 3, pp. 352–358, 2010.
- [2] S. M. Schappert and C. W. Burt, "Ambulatory care visits to physician offices, hospital outpatient departments, and emergency departments: United States, 2001-02," *Vital and Health Statistics*, vol. 13, no. 159, pp. 1–66, 2006.
- [3] K. F. Chung and J. G. Widdicombe, "Pharmacology and therapeutics of cough. Preface," *Handbook of Experimental Pharmacology*, no. 187, pp. 23–49, 2009.
- [4] I. A. Ramallo, M. O. Salazar, L. Mendez, and R. L. E. Furlan, "Chemically engineered extracts: source of bioactive compounds," *Accounts of Chemical Research*, vol. 44, no. 4, pp. 241–250, 2011.
- [5] J. Clardy and C. Walsh, "Lessons from natural molecules," *Nature*, vol. 432, no. 7019, pp. 829–837, 2004.
- [6] D. J. Newman and G. M. Cragg, "Natural products as sources of new drugs over the last 25 years," *Journal of Natural Products*, vol. 70, no. 3, pp. 461–477, 2007.
- [7] A. L. Harvey, "Natural products in drug discovery," *Drug Discovery Today*, vol. 13, no. 19-20, pp. 894–901, 2008.
- [8] G. M. Cragg, P. G. Grothaus, and D. J. Newman, "Impact of natural products on developing new anti-cancer agents," *Chemical Reviews*, vol. 109, no. 7, pp. 3012–3043, 2009.
- [9] C. L. Chang and C. S. Lin, "Phytochemical composition, antioxidant activity, and neuroprotective effect of *Terminalia chebula* Retzius extracts," *Evidence-Based Complementary and Alternative Medicine*, vol. 2012, Article ID 125247, 7 pages, 2012.
- [10] P. Kannan, S. R. Ramadevi, and H. Waheeta, "Antibacterial activity of *Terminalia chebula* fruit extract," *African Journal of Microbiology Research*, vol. 3, no. 4, pp. 180–184, 2009.
- [11] N. N. Barthakur and N. P. Arnold, "Nutritive value of the chebulic myrobalan (*Terminalia chebula* retz.) and its potential as a food source," *Food Chemistry*, vol. 40, no. 2, pp. 213–219, 1991.
- [12] P. Khazaeli, R. Goldoozian, and F. Shariffar, "An evaluation of extracts of five traditional medicinal plants from Iran on the inhibition of mushroom tyrosinase activity and scavenging of free radicals," *International Journal of Cosmetic Science*, vol. 31, no. 5, pp. 375–381, 2009.
- [13] A. Agrawal, A. Gupta, N. K. Choudhary et al., "Antibacterial activity of hydroalcoholic extract of *Terminalia chebula* Retz. on different gram-positive and gram-negative bacteria," *International Journal of Pharmaceuticals & Biological Archives*, vol. 1, no. 4, pp. 485–488, 2010.
- [14] I. S. Grover and S. Bala, "Antimutagenic activity of *Terminalia chebula* (myroblan) in *Salmonella typhimurium*," *Indian Journal of Experimental Biology*, vol. 30, no. 4, pp. 339–341, 1992.
- [15] R. R. Chattopadhyay and S. K. Bhattacharyya, "Plant review *Terminalia chebula*: an update," *Pharmacognosy Reviews*, vol. 1, no. 1, pp. 151–156, 2007.
- [16] W. Panunto, K. Jaijoy, N. Lerdvuthisophon et al., "Acute and chronic toxicity studies of the water extract from dried fruits of *Terminalia chebula* Retz. in rats," *International Journal of Applied Research in Natural Products*, vol. 3, no. 4, pp. 36–43, 2010.
- [17] T. Ghosh, K. Chattopadhyay, M. Marschall, P. Karmakar, P. Mandal, and B. Ray, "Focus on antivirally active sulfated polysaccharides: from structure-activity analysis to clinical evaluation," *Glycobiology*, vol. 19, no. 1, pp. 2–15, 2009.
- [18] K. T. Inngjerdingen, A. Coulibaly, D. Diallo, T. E. Michaelsen, and B. S. Paulsen, "A complement fixing polysaccharide from *Biophytum petersianum* klotzsch, a medicinal plant from Mali, West Africa," *Biomacromolecules*, vol. 7, no. 1, pp. 48–53, 2006.
- [19] M. S. Mantovani, M. F. Bellini, J. P. F. Angeli, R. J. Oliveira, A. F. Silva, and L. R. Ribeiro, "β-Glucans in promoting health: prevention against mutation and cancer," *Mutation Research*, vol. 658, no. 3, pp. 154–161, 2008.
- [20] M. Šutovská, S. Fraňová, L. Prišežňaková et al., "Antitussive activity of polysaccharides isolated from the Malian medicinal plants," *International Journal of Biological Macromolecules*, vol. 44, no. 3, pp. 236–239, 2009.
- [21] S. Mazumder, C. Morvan, S. Thakur, and B. Ray, "Cell wall polysaccharides from chalkumra (*Benincasa hispida*) fruit. Part I. Isolation and characterization of pectins," *Journal of Agricultural and Food Chemistry*, vol. 52, no. 11, pp. 3556–3562, 2004.
- [22] M. Dubois, K. A. Gilles, J. K. Hamilton, P. A. Rebers, and F. Smith, "Colorimetric method for determination of sugars and related substances," *Analytical Chemistry*, vol. 28, no. 3, pp. 350–356, 1956.
- [23] A. Ahmed and J. M. Labavitch, "A simplified method for accurate determination of cell wall uronide content," *Journal of Food Biochemistry*, vol. 1, no. 4, pp. 361–365, 1977.
- [24] B. Ray, "Polysaccharides from *Enteromorpha compressa* isolation, purification and structural features," *Carbohydrate Polymers*, vol. 66, no. 3, pp. 408–416, 2006.
- [25] I. M. Sims and A. Bacic, "Extracellular polysaccharides from suspension cultures of *Nicotiana plumbaginifolia*," *Phytochemistry*, vol. 38, no. 6, pp. 1397–1405, 1995.
- [26] C. J. Schultz, K. L. Johnson, G. Currie, and A. Bacic, "The classical arabinogalactan protein gene family of Arabidopsis," *Plant Cell*, vol. 12, no. 9, pp. 1751–1767, 2000.
- [27] B. J. Canning, S. B. Mazzone, S. N. Meeker, N. Mori, S. M. Reynolds, and B. J. Udem, "Identification of the tracheal and laryngeal afferent neurons mediating cough in anaesthetized guinea-pigs," *Journal of Physiology*, vol. 557, no. 2, pp. 543–558, 2004.
- [28] B. J. Canning and Y. L. Chou, "Coughsensors. I. Physiological and pharmacological properties of the afferent nerves regulating cough," in *Pharmacology and Therapeutics of Cough*, F. Chung and J. G. Widdicombe, Eds., vol. 187, pp. 23–49, Springer, Berlin, Germany, 2009.
- [29] G. Nosalova, S. Franova, J. Mokry, and M. Sutovska, "Pharmacotherapy of cough," in *Cough from Lab to Clinic*, J. Korpas, A. S. Paintal, and A. Anand, Eds., pp. 253–316, ANE Books, New Delhi, India, 2007.
- [30] B. E. Pennock, C. P. Cox, and R. M. Rogers, "A noninvasive technique for measurement of changes in specific airway resistance," *Journal of Applied Physiology Respiratory Environmental and Exercise Physiology*, vol. 46, no. 2, pp. 399–406, 1979.
- [31] Y. Tohda, M. Muraki, T. Iwanaga, R. Haraguchi, M. Fukuoka, and S. Nakajima, "Dual-phase response model for bronchial asthma," *Allergy and Asthma Proceedings*, vol. 21, no. 2, pp. 79–84, 2000.

- [32] D. Bertrand and E. Dufour, "La spectroscopie infra-rouge et ses applications analytiques," *Technique & Documentation*, Lavoisier, Paris, France, 2000.
- [33] T. R. Cipriani, C. G. Mellinger, L. M. De Souza et al., "A polysaccharide from a tea (infusion) of *Maytenus ilicifolia* leaves with anti-ulcer protective effects," *Journal of Natural Products*, vol. 69, no. 7, pp. 1018–1021, 2006.
- [34] S. Ozaki, N. Oki, S. Suzuki, and S. Kitamura, "Structural characterization and hypoglycemic effects of arabinogalactan-protein from the tuberous cortex of the white-skinned sweet potato (*Ipomoea batatas* L.)," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 22, pp. 11593–11599, 2010.
- [35] N. Chattopadhyay, G. Nosál'ová, S. Saha, S. S. Bandyopadhyay, D. Flešková, and B. Ray, "Structural features and antitussive activity of water extracted polysaccharide from *Adhatoda vasica*," *Carbohydrate Polymers*, vol. 83, no. 4, pp. 1970–1974, 2011.
- [36] S. Saha, G. Nosál'ová, D. Ghosh, D. Flešková, P. Capek, and B. Ray, "Structural features and in vivo antitussive activity of the water extracted polymer from *Glycyrrhiza glabra*," *International Journal of Biological Macromolecules*, vol. 48, no. 4, pp. 634–638, 2011.
- [37] S. Sinha, G. Nosál'ová, S. S. Bandyopadhyay, D. Flešková, and B. Ray, "In vivo anti-tussive activity and structural features of a polysaccharide fraction from water extracted *Withania somnifera*," *Journal of Ethnopharmacology*, vol. 134, no. 2, pp. 510–513, 2011.
- [38] G. Nosál'ová, L. Prisenžňáková, E. Paulovičová et al., "Antitussive and immunomodulating activities of instant coffee arabinogalactan-protein," *International Journal of Biological Macromolecules*, vol. 49, no. 4, pp. 493–497, 2011.
- [39] S. Baek, J. G. Lee, S. Y. Park, O. N. Bae, D. Kim, and J. H. Park, "Pectic polysaccharides from *panax ginseng* as the antirotavirus principals in ginseng," *Biomacromolecules*, vol. 11, no. 8, pp. 2044–2052, 2010.
- [40] C. S. Nergard, H. Kiyohara, J. C. Reynolds et al., "Structure and Structure-activity relationships of three mitogenic and complement fixing pectic arabinogalactans from the malian antiulcer plants *Cochlospermum tinctorium* A. Rich and *Vernonia kotschyana* sch. bip. ex walp," *Biomacromolecules*, vol. 7, no. 1, pp. 71–79, 2006.
- [41] J. Lee, J. S. Shim, J. S. Lee, M. Kim, M. Chung, and K. H. Kim, "Pectin-like acidic polysaccharide from *Panax ginseng* with selective antiadhesive activity against pathogenic bacteria," *Carbohydrate Research*, vol. 341, no. 9, pp. 1154–1163, 2006.
- [42] Y. Xu, Q. Dong, H. Qiu, R. Cong, and K. Ding, "Structural characterization of an arabinogalactan from *Platycodon grandiflorum* roots and antiangiogenic activity of its sulfated derivative," *Biomacromolecules*, vol. 11, no. 10, pp. 2558–2566, 2010.
- [43] D. Ghosh, S. Ray, K. Ghosh et al., "Antioxidative carbohydrate polymer from *Enhydra fluctuans* and its interaction with bovine serum albumin," *Biomacromolecules*, vol. 14, no. 6, pp. 1761–1768, 2013.
- [44] B. S. Paulsen, "Biologically active polysaccharides as possible lead compounds," *Phytochemistry Reviews*, vol. 1, no. 3, pp. 379–387, 2002.
- [45] A. B. Samuelsen, B. Westereng, O. Yousif, A. K. Holtekjølen, T. E. Michaelsen, and S. H. Knutsen, "Structural features and complement-fixing activity of pectin from three Brassica oleracea varieties: white cabbage, kale, and red kale," *Biomacromolecules*, vol. 8, no. 2, pp. 644–649, 2007.
- [46] G. Nosalova, D. Fleskova, L. Jurecek, V. Sadlonova, and B. Ray, "Herbal polysaccharides and cough reflex," *Respiratory Physiology & Neurobiology*, vol. 187, no. 1, pp. 47–51, 2013.