



HHS Public Access

Author manuscript

Obesity (Silver Spring). Author manuscript; available in PMC 2016 May 18.

Published in final edited form as:

Obesity (Silver Spring). 2015 July ; 23(7): 1422–1428. doi:10.1002/oby.21119.

Resting State Brain Connectivity After Surgical and Behavioral Weight Loss

Rebecca J. Lepping, PhD¹, Amanda S. Bruce, PhD², Alex Francisco, BS³, Hung-Wen Yeh, PhD⁴, Laura E. Martin, PhD^{1,5}, Joshua N. Powell, AB⁶, Laura Hancock, PhD⁷, Trisha M. Patrician, PhD⁸, Florence J. Breslin, MS⁶, Niazy Selim, MD⁹, Joseph E. Donnelly, EdD^{10,11}, William M. Brooks, PhD^{1,12}, Cary R. Savage, PhD^{6,13}, W. Kyle Simmons, PhD^{14,15}, and Jared M. Bruce, PhD³

¹Hoglund Brain Imaging Center, University of Kansas Medical Center

²Department of Pediatrics, University of Kansas Medical Center and Children's Mercy Hospital

³Department of Psychology, University of Missouri-Kansas City

⁴Department of Biostatistics, University of Kansas Medical Center

⁵Department of Preventive Medicine, University of Kansas Medical Center

⁶Center for Health Behavior Neuroscience, University of Kansas Medical Center

⁷Department of Psychiatry and Human Behavior, Alpert Medical School, Brown University

⁸Minneapolis VA Health Care Center

⁹Department of Surgery-General, University of Kansas Medical Center

¹⁰Center for Physical Activity and Weight Management, University of Kansas Medical Center

¹¹Department of Medicine-Cardiology, University of Kansas Medical Center

¹²Department of Neurology, University of Kansas Medical Center

¹³Department of Psychiatry, University of Kansas Medical Center

¹⁴Laureate Institute for Brain Research

¹⁵Faculty of Community Medicine, The University of Tulsa

Abstract

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence to: Jared M. Bruce, Ph.D., Department of Psychology, The University of Missouri-Kansas City, 5030 Cherry Street, Kansas City, Missouri 64110, Phone: (816)235-5429, FAX: (816)235-1062, ; Email: brucejm@umkc.edu

Work performed in Kansas City, Kansas, USA.

CONFLICT OF INTEREST

The other authors declare no conflict of interest.

ETHICAL APPROVAL AND INFORMED CONSENT

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study.

OBJECTIVE—We previously reported changes in food-cue neural reactivity associated with behavioral and surgical weight loss interventions. Resting functional connectivity represents tonic neural activity that may contribute to weight loss success. Here we explore whether intervention type is associated with differences in functional connectivity after weight loss.

METHODS—Fifteen obese participants were recruited prior to adjustable gastric banding surgery. Thirteen demographically matched obese participants were selected from a separate behavioral diet intervention. Resting state fMRI was collected three months after surgery/behavioral intervention. ANOVA was used to examine post-weight loss differences between the two groups in connectivity to seed regions previously identified as showing differential cue-reactivity after weight loss.

RESULTS—Following weight loss, behavioral dieters exhibited increased connectivity between left precuneus/superior parietal lobule (SPL) and bilateral insula pre- to post-meal and bariatric patients exhibited decreased connectivity between these regions pre- to post-meal ($p_{\text{corrected}} < .05$).

CONCLUSIONS—Behavioral dieters showed increased connectivity pre- to post-meal between a region associated with processing of self-referent information (precuneus/SPL) and a region associated with interoception (insula) whereas bariatric patients showed decreased connectivity between these regions. This may reflect increased attention to hunger signals following surgical procedures, and increased attention to satiety signals following behavioral diet interventions.

Keywords

resting state functional MRI; bariatric surgery; neural mechanisms; weight loss

Functional neuroimaging has improved our understanding of the hedonic brain systems associated with food motivation and obesity. While reports of resting state functional connectivity differences based on obesity status are emerging [1–6], differential connectivity associated with weight loss method has not been examined. Intrinsic resting brain connectivity may elucidate tonic neural activity, which may be critical in understanding the underlying neural mechanisms that lead to successful weight loss.

In task-based fMRI studies, bariatric surgery has been associated with decreased activation to food cues in both cognitive control and reward regions [7, 8]. Reduction in activation to food cues after surgery is also associated with decreased desire for calorically dense foods [9]. We recently showed [10] that behavioral weight loss is associated with increased activation to food images in self-referential processing, valuation (*e.g.*, medial prefrontal cortex), and salience (*e.g.*, precuneus/superior parietal) regions, whereas surgical weight loss is associated with increased cue-reactivity in sensory processing regions (*e.g.*, middle, inferior temporal cortex). This suggests that different methods of weight loss could affect brain responses to food-based stimuli.

The purpose of the present study was to explore whether those brain regions that show differential changes in food-cue reactivity after behavioral and bariatric weight loss interventions [10] also exhibit different resting state functional connectivity after weight loss. Based on our task results [10], we hypothesized that behavioral dieters would show greater connectivity with valuation and salience regions (*i.e.*, MPFC and precuneus/SPL),

and that bariatric patients would show greater connectivity with sensory processing regions (*i.e.*, middle and inferior temporal cortex). We also hypothesized that functional connectivity would be sensitive to hunger state, and would differ between intervention groups following weight loss.

PARTICIPANTS AND METHODS

Participants/Recruitment

Participants were not randomized to treatment condition. They were selected from independent sources of candidates who had decided to undergo bariatric weight-loss surgery or enroll in a behavioral weight-loss intervention research study.

Surgical participants—Obese (body mass index (BMI) 30–45kg/m²) participants ($n=15$; 3 males; Age= 41.40 \pm 9.80; Education=13.85 \pm 1.95 years) planning to undergo adjustable gastric banding weight loss surgery (LapBand®) were recruited from two surgical sites.

Diet participants—13 obese diet participants (4 males; Age=40.23 \pm 8.01; Education=15.31 \pm 1.93 years [some college]) were selected to match demographically with the bariatric group, blind to imaging data, from a larger behavioral weight-loss clinical trial (N=120) [NIH DK080090; NCT02031848] recruited via advertisements, and a university-based weight management center. Participants underwent a three-month weight loss intervention of behavioral strategies, moderate calorie restriction with provided pre-packaged meals, and physical activity.

Special diets (*e.g.*, vegetarian and Atkins), appetite or metabolic medications (*e.g.*, thyroid, beta blockers, and Meridia), smoking and diabetes were exclusions for the diet participants. Because most patient presenting for bariatric surgery have comorbid health conditions, patients who had well-controlled diabetes (most recent hemoglobin A1c $<$ 7) and were not taking insulin or other injectable medications (*i.e.* GLP-1 agonists) were included in the bariatric group. Additional exclusion criteria for both groups included current eating disorder, current major depression, history of neurological disease, pregnancy within the past 6 months, cancer, heart disease, and contraindications for MRI (*e.g.*, metal implants). Participants taking selective serotonin reuptake inhibitors (SSRIs) were included in the bariatric group.

One bariatric participant was unable complete the resting-state scan and another participant's data were unusable due to excess movement (*i.e.*, greater than 50% censored) during the scan [11]. Diet participants with unusable resting-state data were not selected for this study. Therefore, 13 Bariatric participants and 13 Diet participants were included in the final analyses. No significant differences between the final bariatric and diet groups were observed for age [$t(24)=.49$; $p=.63$], education [$\chi^2(3)=3.93$; $p=.27$], sex [$\chi^2(1)=0.87$; $p=.35$], pre-intervention BMI [$t(24)=1.68$; $p=.11$], or percent weight lost [$t(24)=.99$; $p=.33$]. Additional demographic and anthropometric data are included in Table 1.

Procedures and Methods

Resting state data were collected three-months post intervention. To investigate changes that accompany typical mealtime eating behavior, participants were scanned while hungry (Premeal; following at least a 4-hour fast) and after eating a small, standardized (500 kcal) lunch (Postmeal). As dietary restrictions on certain foods (*i.e.* bread products) are in place following bariatric surgery, the format of the meal was slightly different between the two groups. Specifically, the bariatric participants' meal included a lean meat (turkey or ham) wrap, while diet participants' meal included an equivalent sandwich. Both groups reported similar levels of satiation post-meal (0–100 visual analog scale, "How full do you feel right now?": Bariatric=67.92<pm>32.29; Diet=74.92<pm>22.63; $t(24)=-0.64$, $p=.53$). Order of scans (Premeal, Postmeal) was counterbalanced across participants (Figure 1). The resting-state scan (6 minutes 36 seconds) followed a structural scan and two functional scans while passively viewing food and animal pictures [7, 10]. Participants were instructed to close their eyes during the resting state scan. The entire scanning session lasted approximately 45 minutes.

fMRI Data Acquisition

Data were acquired with a 3-Tesla Siemens Allegra, head-only magnetic resonance imaging scanner. To minimize susceptibility artifact in ventromedial prefrontal regions, participants were positioned so that the angle of the AC-PC plane was between 17<deg> and 22<deg> in scanner coordinate space [12]. T1-weighted anatomic images were acquired with a 3D MPRAGE sequence (TR/TE = 23/4ms, flip angle=8<deg>, FOV=192mm, matrix=192×192mm, 208 slices, slice thickness=1mm). Task-based and resting-state gradient echo blood-oxygenation level dependent (BOLD) scans were acquired in 43 contiguous axial slices at an angle of 40<deg> to the AC-PC line (TR/TE 3000/30ms, slice thickness=3mm (0.5mm skip), in-plane resolution=3×3mm, 130 volumes).

Data Preprocessing

Data preprocessing and statistical analysis were conducted using Analysis of Functional NeuroImages (AFNI) [13], and a modified version of the ANATICOR method, developed by Jo, and colleagues, implemented in `afni_restproc.py` [14]. The first four volumes of the functional scans were removed and a de-spiking interpolation algorithm (*i.e.*, 3dDespike) was used to remove any transient signal spikes from the data. The volumes were slice time corrected and co-registered to the first volume (which was registered to the anatomical scan). Several nuisance variables were measured [*i.e.*, six motion parameters (3 translations, 3 rotations), average ventricle signal, and average local white matter signal (15mm spherical neighborhood, 3dLocalstat)]. These nuisance variables' predicted timecourse was constructed and then subtracted from each resting-state voxel timecourse using multiple regression, yielding a residual timecourse for each voxel. The residual images were smoothed with a 6mm FWHM Gaussian kernel, resampled to a 2×2×2mm grid, and spatially transformed to stereotaxic space conforming to the Talairach and Tournoux Atlas [15].

Further motion correction procedures (*i.e.*, scrubbing) were utilized to reduce false group differences due to uncontrolled subject motion [16, 17]. The six motion parameters from the image registration process were used to construct a time series reflecting the Euclidean

normalized derivative of the motion. This timeseries was thresholded so that any time point where the derivative was greater than 0.3 (roughly 0.3mm motion) was censored. We also censored any time point where more than 5% of brain voxels were considered outliers (3dToutcount). Time points censored by the union of both methods were removed in the subsequent regression analysis. The percentage of data removed in this manner did not differ between the two groups [$t(24)=1.21$; $p=.24$; Bariatric=11.9<pm>9.2%; Diet=8.0<pm>7.4%], nor did it differ between imaging sessions [Wilks' $\lambda(3,24)=1.12$; $p=.30$; $\eta^2=.05$].

Seed Region Selection and Identification—Seed regions were defined as 5mm radii spheres in regions identified in our previous food cue reactivity study: medial prefrontal cortex (Talairach (TAL)X,Y,Z=6,50,19), precuneus/SPL (-30,-67,40), right middle (48,-55,7) and left inferior temporal gyrus (-42,-64,-2) [10]. At the subject-level, the four seed timeseries (MPFC, precuneus/SPL, right middle and left inferior temporal gyrus) were constructed by calculating the average time series over the voxels within each of the seed regions. Using multiple regression, we produced for each seed a map of the correlations (r -values) between the seed timeseries and each voxel in the brain. These r -values were transformed to z -scores. While the motion scrubbing procedure described above removes time points most affected by motion artifact, datasets with more motion may still contain residual effects. These effects may induce spurious correlations in an individual data set and increase noise in the overall sample, reducing power. As an additional conservative approach to minimize effects of motion, z -score maps for each dataset were weighted by multiplying the participant's percentage of TRs remaining after censoring by the z -score at each voxel (e.g., 100% TRs included, weighting factor=1; 65% included, weighting factor=.65). This additional step ensures that participants who moved least contribute more to the final group analyses, while those who moved more contribute less [18].

We implemented two-way mixed effects ANOVA comparing weighted z -scores for fixed-effects of Group (Diet, Bariatric) \times Satiety (Premeal, Postmeal) with Participants as random effects. To elucidate differences in functional connectivity to those regions that were previously identified as showing cue-reactivity changes after weight loss, these analyses focus on data collected *after* the 3-month weight loss intervention (Post-intervention) that each group underwent (either a diet/behavioral intervention, or bariatric surgery), as well as the effect of satiety (Premeal vs. Postmeal). These F -statistic maps were corrected for multiple comparisons at $p < .05$ using a voxel-wise threshold of $p < .005$, combined with Monte Carlo simulations of minimum cluster size (616mm³) determined for the whole brain [19].

RESULTS

For the ANOVA seeded in the left precuneus/SPL (Table 2), there was a significant main effect of intervention type (Bariatric vs. Diet) in the middle temporal gyrus ($p_{\text{corrected}} < .01$). Collapsed across intervention groups, there was no main effect of satiety (Premeal vs. Postmeal) in resting state correlations with the left precuneus/SPL. However, there were significant interactions in functional connectivity between satiety (Pre to Postmeal) and intervention type (Diet vs. Bariatric) between the left precuneus/SPL and the following

regions: right precentral gyrus, spreading into insula ($p_{\text{corrected}} < .01$), left middle occipital gyrus ($p_{\text{corrected}} < .01$), right superior temporal gyrus ($p_{\text{corrected}} < .01$), and left insula ($p_{\text{corrected}} < .02$). Figure 2 illustrates the directionality of these interactions. Post intervention, bariatric participants showed greater functional connectivity between left precuneus/SPL and right precentral gyrus and insula, left middle occipital gyrus, right superior temporal gyrus and left insula prior to eating compared to the diet participants. Additionally, resting state functional connectivity changed differentially between the groups after going from a fasted state to a fed state, such that correlations between left precuneus/SPL and each of these regions *increased* from premeal to postmeal for those in the diet intervention, and *decreased* for those in the bariatric intervention.

No significant main effects or interactions were found for the ANOVA seeded from the medial PFC; however, some trends were observed (Table 3). There was a subthreshold main effect of satiety in the posterior cingulate ($p_{\text{corrected}} < .09$). Subthreshold interactions between satiety and intervention type were observed in the left dorsolateral prefrontal cortex ($p_{\text{corrected}} < .08$), and the left superior frontal cortex ($p_{\text{corrected}} < .08$; Figure 3). No significant effects were found for the ANOVAs seeded from either temporal cortex region (all $p_{\text{corrected}} > .10$).

DISCUSSION AND CONCLUSION

The goal of the current study was to determine whether regions of the brain that showed differential changes in food cue reactivity after weight loss dependent on weight loss method (*i.e.* surgical or behavioral) also exhibited group differences after weight loss in functional connectivity with the rest of the brain. Consistent with our hypotheses, we found that the precuneus/SPL and bilateral insula connectivity changed differentially pre- to post-meal depending on whether the participants had completed a behavioral or surgical weight loss intervention. We used the dorsal and lateral part of the precuneus/SPL as a seed for these connectivity analyses, a subregion previously shown to be differentially associated with reactivity to food cues in surgical versus behavioral weight loss [10]. This region is also involved in mental imagery involving motor planning and self-awareness [20–23]. Differential connectivity was found between the precuneus/SPL and insular regions associated with interoception [24, 25]. The groups not only show differential connectivity between these two regions of the brain when fasted, but also demonstrate oppositional change in connectivity strength after a meal.

This pattern of connectivity could reflect differences in interoceptive signaling and awareness that may have led to weight loss. Hunger and satiety are components of interoceptive signaling that lead to initiation and cessation of food intake. Though only speculative, greater functional connectivity between the precuneus/SPL and insula could indicate greater interoceptive self-awareness. If true, those individuals who have lost weight through surgery may be more aware of internal bodily signals of *hunger*, while signals of satiety may be more automatic due to the physical restrictions placed on the stomach through surgery. Alternatively, individuals who have successfully lost weight through dieting may have greater awareness of bodily signals monitoring feelings of *fullness* to know when to stop eating. These differences in connectivity are not due to differences in weight loss

success between groups, as there were no differences in the amount of weight lost between the two groups.

Contrary to expectations, this study found no significant differences in connectivity to either medial prefrontal or temporal cortex. A marginal effect of satiety was found between MPFC and posterior cingulate, and a marginal interaction of satiety and intervention type was found between MPFC and middle and superior frontal cortex. The lack of significance may be due to our modest sample size. Accordingly, our subthreshold results should be interpreted cautiously. One of the strengths of this study is that the analysis was focused on functional connectivity with a priori regions that showed differential responsiveness to food cues *in these same subjects* [10]. Due to limited sample size, we have avoided further exploratory analyses. Nevertheless, it is possible that intervention type may differentially impact connectivity patterns with the default mode or salience networks, as has been shown in other recent work in obesity [1-6].

This study is limited by the fact that the groups were not selected via random assignment. Preexisting factors could lead individuals to consider surgical versus behavioral weight loss. Also, since the bariatric surgery group all underwent laparoscopic gastric banding, results cannot be extended directly to gastric sleeve or gastric bypass patients. Additionally, the groups were not matched for comorbid conditions, such as diabetes, however, in the bariatric surgery group, only those participants with well-controlled diabetes were included. That said, we carefully matched the groups on demographics, pre-intervention BMI, as well as the total weight lost during the intervention. Therefore, results are not simply due to differential treatment effectiveness. Other factors that were not measured in this study, such as menstrual cycle phase for female participants, could also lead to differences in brain connectivity patterns [26-28]. In this study, the intervention groups were matched for gender and age, and both men and women were included in the analysis. Therefore, it is unlikely that the results were systematically biased by menstrual cycle phase. It is also unknown if possible physical activity differences between the groups could influence functional connectivity [4]. All surgical participants were under the care of a physician, and physical activity recommendations were monitored on an individual level. Those in the behavioral diet intervention were under standardized physical activity guidelines according to the intervention. As the two groups received different instructions for physical activity, we cannot confidently conclude that physical activity levels were the same for both groups. Care should be taken in future studies to control for these additional variables.

In conclusion, method of weight loss seems to be related to differential connectivity between regions of the brain involved in self-imagery and interoception, as well as differences in whether that connectivity emerges during states of hunger or satiety. Surgery may lead individuals to increase attention to bodily signals of hunger, whereas successful dieting may require more attention to signals of fullness. While both methods are effective for initial weight loss, patterns of functional connectivity in the brain suggest differences in the underlying mechanisms associated with weight loss approaches. Future research should examine whether these neurofunctional differences are maintained over time in extended longitudinal studies, and how they may be related to successful weight loss maintenance.

ACKNOWLEDGMENTS

This project was supported in part by a grant from the University of Missouri Research Board to Dr. J. Bruce, pilot funding to Dr. A. Bruce from a CTSA grant from NCATS awarded to the University of Kansas Medical Center for Frontiers: The Heartland Institute for Clinical and Translational Research (UL1 TR000001), and by R01DK080090 to Dr. Savage. The Hoglund Brain Imaging Center is supported by a generous gift from Forrest and Sally Hoglund and funding from the National Institutes of Health (UL1 TR000001). The contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH or any individual institute. All authors were involved in writing the paper and had final approval of the submitted and published versions.

Dr. Bruce is a consultant to the National Hockey League and Princeton University, a member of the Novartis Unbranded Speaker's Bureau, and a member of the Novartis Cognition in Multiple Sclerosis Medical Advisory Board. He has received funding unrelated to the current work, from Cephalon and the National Multiple Sclerosis Society.

Special characters

pm	plus minus \pm
gx	Greek Chi, uppercase
deg	degrees
gl	Greek lambda, lowercase
ge	Greek eta, lowercase
ga	Greek alpha, lowercase

Abbreviations

rs-fMRI	resting state functional magnetic resonance imaging
BMI	body mass index
PFC	prefrontal cortex
SPL	superior parietal lobule
MPFC	medial prefrontal cortex
ROI	region of interest
BOLD	blood-oxygenation level dependent

References

1. Kullmann S, et al. The obese brain: association of body mass index and insulin sensitivity with resting state network functional connectivity. *Hum Brain Mapp.* 2012; 33(5):1052–1061. [PubMed: 21520345]
2. Garcia-Garcia I, et al. Alterations of the salience network in obesity: a resting-state fMRI study. *Hum Brain Mapp.* 2013; 34(11):2786–2797. [PubMed: 22522963]
3. Lips MA, et al. Resting-state functional connectivity of brain regions involved in cognitive control, motivation, and reward is enhanced in obese females. *Am J Clin Nutr.* 2014; 100(2):524–531. [PubMed: 24965310]
4. McFadden KL, et al. Effects of exercise on resting-state default mode and salience network activity in overweight/obese adults. *Neuroreport.* 2013; 24(15):866–871. [PubMed: 24022176]

5. Frank S, et al. Altered brain activity in severely obese women may recover after Roux-en-Y gastric bypass surgery. *Int J Obes (Lond)*. 2014; 38(3):341–348. [PubMed: 23711773]
6. Kullmann S, et al. Functional network connectivity underlying food processing: disturbed salience and visual processing in overweight and obese adults. *Cereb Cortex*. 2013; 23(5):1247–1256. [PubMed: 22586138]
7. Bruce JM, et al. Changes in brain activation to food pictures after adjustable gastric banding. *Surg Obes Relat Dis*. 2012; 8(5):602–608. [PubMed: 21996599]
8. Ochner CN, et al. Neural responsivity to food cues in fasted and fed states pre and post gastric bypass surgery. *Neurosci Res*. 2012; 74(2):138–143. [PubMed: 22921709]
9. Ochner CN, et al. Relation between changes in neural responsivity and reductions in desire to eat high-calorie foods following gastric bypass surgery. *Neuroscience*. 2012; 209:128–135. [PubMed: 22406414]
10. Bruce AS, et al. A comparison of functional brain changes associated with surgical versus behavioral weight loss. *Obesity (Silver Spring)*. 2014; 22(2):337–343. [PubMed: 24115765]
11. Siegel JS, et al. Statistical improvements in functional magnetic resonance imaging analyses produced by censoring high-motion data points. *Hum Brain Mapp*. 2014; 35(5):1981–1996. [PubMed: 23861343]
12. Bruce AS, et al. Branding and a child's brain: an fMRI study of neural responses to logos. *Soc Cogn Affect Neurosci*. 2014; 9(1):118–122. [PubMed: 22997054]
13. Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res*. 1996; 29(3):162–173. [PubMed: 8812068]
14. Jo HJ, et al. Effective Preprocessing Procedures Virtually Eliminate Distance-Dependent Motion Artifacts in Resting State FMRI. *J Appl Math*. 2013; 2013
15. Talairach, J.; Tournoux, P. *Co-planar Stereotaxic Atlas of the Human Brain*. New York: Thieme Medical Publishers Inc.; 1988.
16. Power JD, et al. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage*. 2012; 59(3):2142–2154. [PubMed: 22019881]
17. Power JD, et al. Steps toward optimizing motion artifact removal in functional connectivity MRI; a reply to Carp. *Neuroimage*. 2013; 76:439–441. [PubMed: 22440651]
18. Harris JL, et al. High-field proton magnetic resonance spectroscopy reveals metabolic effects of normal brain aging. *Neurobiol Aging*. 2014; 35(7):1686–1694. [PubMed: 24559659]
19. Forman SD, et al. Improved assessment of significant activation in functional magnetic resonance imaging (fMRI): use of a cluster-size threshold. *Magn Reson Med*. 1995; 33(5):636–647. [PubMed: 7596267]
20. Lamm C, Decety J, Singer T. Meta-analytic evidence for common and distinct neural networks associated with directly experienced pain and empathy for pain. *Neuroimage*. 2011; 54(3):2492–2502. [PubMed: 20946964]
21. Legrand D, Ruby P. What is self-specific? Theoretical investigation and critical review of neuroimaging results. *Psychol Rev*. 2009; 116(1):252–282. [PubMed: 19159156]
22. Gruber O, et al. Brain mechanisms associated with background monitoring of the environment for potentially significant sensory events. *Brain Cogn*. 2009; 69(3):559–564. [PubMed: 19135767]
23. Kircher TT, et al. The neural correlates of intentional and incidental self processing. *Neuropsychologia*. 2002; 40(6):683–692. [PubMed: 11792407]
24. Simmons WK, et al. Keeping the body in mind: insula functional organization and functional connectivity integrate interoceptive, exteroceptive, and emotional awareness. *Hum Brain Mapp*. 2013; 34(11):2944–2958. [PubMed: 22696421]
25. Simmons WK, et al. Category-specific integration of homeostatic signals in caudal but not rostral human insula. *Nat Neurosci*. 2013; 16(11):1551–1552. [PubMed: 24077565]
26. Arelin K, et al. Progesterone mediates brain functional connectivity changes during the menstrual cycle—a pilot resting state MRI study. *Front Neurosci*. 2015; 9:44. [PubMed: 25755630]
27. Petersen N, et al. Oral contraceptive pill use and menstrual cycle phase are associated with altered resting state functional connectivity. *Neuroimage*. 2014; 90:24–32. [PubMed: 24365676]

28. Weis S, et al. Dynamic changes in functional cerebral connectivity of spatial cognition during the menstrual cycle. *Hum Brain Mapp.* 2011; 32(10):1544–1556. [PubMed: 20814961]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

WHAT IS ALREADY KNOWN ABOUT THIS SUBJECT

- Resting state functional connectivity differs based on obesity status
- Bariatric surgery is associated with decreased brain activation to food cues in cognitive control and reward regions
- Changes in brain activation to food cues after weight loss in self-referential processing, valuation, and salience regions are different if weight loss is achieved through behavioral or surgical means

WHAT DOES THIS STUDY ADD

- Evidence for different resting state functional connectivity after behavioral and bariatric weight loss interventions in regions that show differential changes in food-cue reactivity between groups
- Evidence that connectivity between regions involved in self-awareness and interoceptive signaling is different between the two groups after weight loss
- Evidence that functional connectivity between these two regions differs by group dependent on hunger state and could reflect differences in interoceptive signaling and awareness that may have led to weight loss




Figure 1.

Participants were randomized to receive one of two counterbalanced scanning orders. Half of the participants were scanned pre-meal, then at a 500 kcal meal, and immediately were scanned post-meal. The other half ate the 500 kcal meal upon arrival for their appointment and were immediately scanned post-meal, then waited for four hours and were scanned pre-meal.




Figure 2. Maps show the interaction between intervention type and satiety on voxel-wise correlations with the precuneus/SPL seed. At three months post-intervention, correlations between left precuneus/SPL and right precentral gyrus, right insula, and left insula increased from premeal to postmeal for those in the diet intervention, and decreased for those in the bariatric intervention ($p_{\text{corrected}} < .05$). Error bars denote standard error.




Figure 3.

Maps show the interaction between intervention type and satiety on voxel-wise correlations with the MPFC seed. Although subthreshold, at three months post-intervention, correlations between right MPFC and left middle and superior prefrontal cortex increased from premeal to postmeal for those in the diet intervention, and decreased for those in the bariatric intervention ($p_{\text{corrected}} < .10$). Error bars denote standard error.

Demographic and anthropometric characteristics for participants included in the analyses. Body Mass Index (BMI) is calculated as body weight (in kg) divided by height (in meters) squared.

Table 1

Group	Male/Female	Age [years] (Mean <pm>SD)	BMI-baseline [kg/m ²] (Mean <pm>SD)	BMI-post-intervention (Mean <pm>SD)	BMI-percent change (Mean <pm>SD)
Bariatric Surgery Intervention	2/11	42.00 <pm>10.35	41.35 <pm>1.97	37.43 <pm>2.73	-9.50% <pm>4.32%
Behavioral Diet Intervention	4/9	40.23 <pm>8.01	40.10 <pm>1.80	35.62 <pm>2.22	-11.16% <pm>4.19%

Resting State Functional Connectivity with Left Precuneus/SPL 3 Months Post-intervention (Behavioral or Bariatric Surgery)

Table 2

Region	L/R	BA	X	Y	Z	<i>P</i>_{corrected}	mm³
<i>Intervention Main Effect</i>							
Middle temporal gyrus	L	39	-39	-69	18	< .01	944
<i>Satiety Main Effect</i>							
None							
<i>Intervention × Satiety Interaction</i>							
Precentral gyrus/ Insula	R	4	53	-9	24	< .01	1624
Middle occipital gyrus	L	19	-45	-77	8	< .01	1384
Superior temporal gyrus	R	22	55	-37	20	< .01	1128
Insula	L	13	-41	1	12	< .02	784

Table 3 Resting State Functional Connectivity with Right Medial Prefrontal Cortex 3 Months Post-intervention (Behavioral or Bariatric Surgery) – Subthreshold Effects

Region	L/R	BA	X	Y	Z	<i>P</i> _{corrected}	mm ³
<i>Intervention Main Effect</i>							
None							
<i>Satiety Main Effect</i>							
Posterior cingulate	R	7	11	-49	32	<.09	568
<i>Intervention × Satiety Interaction</i>							
Middle frontal	L	6	-39	1	50	<.08	576
Superior frontal	L	6	-13	15	58	<.08	584