

# Genome-wide identification and expression analysis of the *14-3-3* gene family in soybean (*Glycine max*)

Yongbin Wang<sup>1,2</sup>, Lei Ling<sup>3</sup>, Zhenfeng Jiang<sup>2</sup>, Weiwei Tan<sup>1</sup>, Zhaojun Liu<sup>1</sup>, Licheng Wu<sup>1</sup>, Yuanling Zhao<sup>4</sup>, Shanyong Xia<sup>4</sup>, Jun Ma<sup>4</sup>, Guangjin Wang<sup>5</sup> and Wenbin Li<sup>2</sup>

<sup>1</sup> Biotechnology Research Institute, Heilongjiang Academy of Agricultural Sciences, Harbin, Heilongjiang, China

<sup>2</sup> Key Laboratory of Soybean Biology in Chinese Ministry of Education, Key Laboratory of Soybean Biology and Breeding/Genetics of Chinese Agriculture Ministry, Northeast Agricultural University, Harbin, Heilongjiang, China

<sup>3</sup> Harbin Normal University, Harbin, Heilongjiang, China

<sup>4</sup> Heilongjiang Academy of Agricultural Sciences, Harbin, Heilongjiang, China

<sup>5</sup> Soybean Research Institute, Heilongjiang Academy of Agricultural Sciences, Harbin, Heilongjiang, China

## ABSTRACT

In eukaryotes, proteins encoded by the *14-3-3* genes are ubiquitously involved in the plant growth and development. The *14-3-3* gene family has been identified in several plants. In the present study, we identified 22 *GmGF14* genes in the soybean genomic data. On the basis of the evolutionary analysis, they were clustered into  $\varepsilon$  and non- $\varepsilon$  groups. The *GmGF14s* of two groups were highly conserved in motifs and gene structures. RNA-seq analysis suggested that *GmGF14* genes were the major regulator of soybean morphogenesis. Moreover, the expression level of most *GmGF14s* changed obviously in multiple stress responses (drought, salt and cold), suggesting that they have the abilities of responding to multiple stresses. Taken together, this study shows that soybean *14-3-3s* participate in plant growth and can respond to various environmental stresses. These results provide important information for further understanding of the functions of *14-3-3* genes in soybean.

**Subjects** Agricultural Science, Bioinformatics, Plant Science

**Keywords** Soybean, *14-3-3* genes, Expression, Abiotic stress

## INTRODUCTION

The *14-3-3* genes are first isolated from brain tissue, and they are ubiquitously found in eukaryotes (Li *et al.*, 2015; Yang *et al.*, 2017; Kumar *et al.*, 2015; Takahashi, 2006). *14-3-3* proteins are highly conserved small acidic proteins in different organisms, encoded by a large gene family (27–32 kDa) (Ferl, Lu & Bowen, 1994; Cao & Tan, 2018). These proteins can form dimers (homo- or hetero- dimers) and have approximately nine antiparallel  $\alpha$ -helices (Ferl, Manak & Reyes, 2002; Rodriguez & Guan, 2010). These structures work as binding sites and interact with *14-3-3* proteins and their target, and they also bring two proteins together as a protein complex based on their dimeric properties (Sijbesma *et al.*, 2017; Valente *et al.*, 2012; Li & Dhaubhadel, 2012). *14-3-3s* are involved in several protein-protein interactions, such as responding to biotic/abiotic stress, participating in

Submitted 18 March 2019

Accepted 25 September 2019

Published 6 December 2019

Corresponding authors

Guangjin Wang, gjw1962@yeah.net,

wenbinli@yahoo.com

Wenbin Li, wenbinli@neau.edu.cn

Academic editor

Shalu Jhanwar

Additional Information and  
Declarations can be found on  
page 17

DOI 10.7717/peerj.7950

© Copyright  
2019 Wang et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

plant hormone signaling and regulating tissue development in various plants (Roberts, Salinas & Collinge, 2002; Camoni et al., 2018; Zhang et al., 2010).

To date, more 14-3-3s have been reported in several plants, such as Arabidopsis, rice, tobacco, populus and *Medicago truncatula* (Chen et al., 2006; Rosenquist et al., 2001; Xu & Shi, 2006; Tian et al., 2015; Cheng et al., 2016). In plants, the 14-3-3 proteins were named as GF14 or GRF due to they are a part of protein/G-box complex (De Vetten & Ferl, 1994; Rosenquist et al., 2001). 14-3-3s are distributed in different organelles, such as cytoplasm, cell membrane, nucleus, chloroplast and mitochondria (Bihn et al., 2010; Sehnke et al., 2000; Ferl, Manak & Reyes, 2002). In plants, they were regulated by several biological processes (Cheng et al., 2016; Tian et al., 2015); for example, multiple mutant analysis suggested that Arabidopsis 14-3-3 genes regulate root growth, chloroplast division, photosynthesis and leaf longevity (Liesbeth et al., 2015). GhGRFs were found involving in plant development and signaling transduction in cotton fiber (Zhang et al., 2010). In addition, an increasing number of works were carried out to investigate the roles of 14-3-3s in plants under multiple stresses (Roberts, Salinas & Collinge, 2002). Most of the OsGRF genes' expression changed under heat, cold and salt stresses (Yashvardhini et al., 2017). The results of overexpression of AtGRF6 in transgenic cotton showed a stay-green phenotype, indicating that they can improve plant tolerance to drought stress (Juqiang et al., 2004).

Soybean is an important cash crop in the world, while its production is often influenced by various environmental stresses (Masuda & Goldsmith, 2009). However, not enough attention is focused on soybean GmGF14s. Previous studies identified 18 GmGF14 genes in soybean, which was different from our study (Li & Dhaubhadel, 2011). In this study, we identified a total of 22 GmGF14 genes in soybean genome. Phylogenetic relationship, gene structures, protein motifs and expression patterns of all the GmGF14 genes were analyzed, together with their responses to various stresses in soybean. These results will provide important information for further study to understand the regulating mechanism of GmGF14 genes during their growth and their responding abilities to various environmental stresses.

## MATERIALS & METHODS

### Identification of 14-3-3 genes

The Hidden Markov Model (HMM) profiles of the 14-3-3 motif (PF00244) were downloaded from the Pfam database (Punta et al., 2004). HMM searched 14-3-3 motif (PF00244) from the *Glycine max* protein database with values (*e*-value) cut-off at 0.1 (Punta et al., 2004). The integrity of the 14-3-3 motif was determined using the online program SMART (<http://smart.embl-heidelberg.de/>) with an *e*-value < 0.1 (Ivica, Tobias & Peer, 2012). In addition, the three fields (length, molecular weight, and isoelectric point) of each 14-3-3 protein was predicted by the online ExPASy program (<http://www.expasy.org/tools/>) (Johana et al., 2015).

### Phylogenetic analysis

To investigate the phylogenetic relationship of the 14-3-3 gene families in *Arabidopsis thaliana*, *Oryza sativa*, *Medicago truncatula* and *Glycine max*, 14-3-3 protein sequences were

downloaded from Phytozome v12.1 (<http://www.phytozome.org>) (Goodstein *et al.*, 2012). 14-3-3s were aligned using the BioEdit program. A neighbor-joining (NJ) phylogenetic tree was constructed using the MEGA 5.0 program (Tamura *et al.*, 2011). Bootstrapping was performed with 1000 replications. Genes were classified according to the distance homology with *Arabidopsis thaliana* genes (Feri, Lu & Bowen, 1994).

### Sequence alignment, motif prediction and gene structure of 14-3-3 genes

The 3D structure of 14-3-3 proteins were predicted by using Phyre<sup>2</sup> and ESPript 3.0 software (Gouet, Robert & Courcelle, 2003; Kelley *et al.*, 2015). Multiple alignments of proteins were conducted using Jalview software with ClustalW method (Clamp *et al.*, 2004). The online MEME analysis is used to identify the unknown conserved motifs (<http://meme.ebi.edu.au/>) by using the following parameters: site distribution: zero or one occurrence (of a contributing motif site) per sequence, maximum number of motifs: 20, and optimum motif width  $\geq 6$  and  $\leq 200$  (Bailey *et al.*, 2015). A gene structure display server program (<http://gsds.cbi.pku.edu.cn/>) was used to display the *G. max* 14-3-3 gene structures.

### Gene duplication and collinearity analysis

The physical locations of the *GmGF14* genes on the soybean chromosomes were mapped by using MG2C website ([http://mg2c.iask.in/mg2c\\_v2.0/](http://mg2c.iask.in/mg2c_v2.0/)). The analysis of synteny among the soybean genomes was conducted locally using a method similar to that developed for the PGDD (<http://chibba.agtec.uga.edu/duplication/>) (Krzywinski & Schein, 2009). First, BLASTP, OrthoMCL software (<http://orthomcl.org/orthomcl/about.do#release>) and MCScanX software (Wang *et al.*, 2012) were used to search for potential homologous gene pairs ( $E < 1 \times 10^{-5}$ , top 5 matches) across multiple genomes. Then, these homologous pairs were used as the input for the PGDD database (<http://chibba.agtec.uga.edu/duplication/>). Ideograms were created by using Circos (Krzywinski & Schein, 2009).

### Calculating $K_a$ and $K_s$

The  $K_a$  and  $K_s$  were used to assess selection history and divergence time (Li, Gojobori & Nei, 1981). The number of synonymous ( $K_s$ ) and nonsynonymous ( $K_a$ ) substitutions of duplicated 14-3-3 genes were computed by using the KaKs\_Calculator 2.0 with the NG method (Wang *et al.*, 2010). The divergence time ( $T$ ) was calculated using the formula  $T = Ks / (2 \times 6.1 \times 10^{-9}) \times 10^{-6}$  million years ago (MYA) (Kim *et al.*, 2013).

### 14-3-3 genes expression analysis of soybean

The expression data of 14-3-3 genes in different tissues, including root, root hair, flower, nodule, pod, stem, leaf, SAM and seed, were available in Phytozome v12.1 database (<https://phytozome.jgi.doe.gov/pz/portal.html>). We retrieved the fragments per kilobase per million reads (FPKM) value representing the expression level of *GmGF14* genes, then generated the heatmap and k-means clustering by using R 3.2.2 software.

## Plant material and treatments

*Glycine max* (Williams 82) was used in this study. Seeds were planted in a 3:1 (w/w) mixture of soil and sand, germinated, and irrigated with half-strength Hoagland solution once every 2 days. The seedlings were grown in a night temperature of 20 °C and day temperature of 22 °C, relative humidity of 60%, and a 16/8 h photoperiod (daytime: 05:00–21:00). After 4 weeks, the germinated seedlings were treated with 20% PEG6000 (drought), 250 mM NaCl solution (salt), and 4 °C (cold). Control and treated seedlings were harvested 1 h, 6 h, and 12 h after treatment. All samples were frozen in liquid nitrogen and stored at –80 °C until use.

## RNA extraction and Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted from the root of soybean using RNAiso Plus (TaKaRa, Toyoto, Japan) according to manufacturer's instructions. 2 µg RNA was extracted using PrimeScript RT reagent Kit with gDNA Eraser (TaKaRa, Toyoto, Japan). The cDNA samples were diluted to 2.5 ng/L. Quantitative Real-time PCR (qRT-PCR) was performed using SYBR *Premix Ex Taq* II (TaKaRa, Toyoto, Japan) on an ABI Prism 7000 sequence detection system (Applied Biosystems, USA) with the primers listed in Table S1. PCR amplification was performed in accordance with SYBR *Premix Ex Taq* (TaKaRa, Toyoto, Japan) response system. For each sample, three biological replicates were conducted. Relative expression was calculated by the  $2^{-\Delta\Delta C_t}$  method (Livak & Schmittgen, 2001). The *actin* and *GAPDH* genes were used as internal control.

## Gene ontology enrichment

Once the sequences were obtained ran a BLASTX search against the Uniref100 database at a 1e–30 significance level. The matches were extracted and compared to the GO annotation generated against Uniref100 hits located at EBI. The GO annotation of the *GmGF14* genes by using WEGO 2.0 website (<http://wego.genomics.org.cn/>).

## RESULTS

### Identification and multiple sequences alignment of *GmGF14* genes

We identified 22 *GmGF14* genes, which were named from *GmGF14a* to *GmGF14v* based on their physical locations on chromosomes. ExpASy predicted that 22 *GmGF14* proteins have different physical and chemical properties as their amino acid lengths ranged from 71 aa (*GmGF14v*) to 754 aa (*GmGF14j*), with an average of 295 aa, and their molecular weights ranged from 7.92 kDa (*GmGF14v*) to 81.75 kDa (*GmGF14j*), the isoelectric points ranged from 4.67 (*GmGF14c/e*) to 5.7 (*GmGF14v*). Detail information of *GmGF14* proteins is provided in Table 1. Besides, we found that most of the *GmGF14s* contain highly conserved domains, and ten  $\alpha$ -helices were identified in their secondary structures (Fig. 1). In addition, the C-terminal end of 14-3-3 proteins are quite unique in sequence and length.

### Phylogenetic analysis of the *GmGF14s*

We constructed a phylogenetic tree to show the phylogenetic and evolutionary relationships of *GmGF14* genes among *A. thaliana*, *O. sativa*, *M. truncatula* and *G. max* (Fig. 2). The


**Table 1** List of all *GmGF14* genes identified in the *Glycine max* genome.

Gene name	Gene locus	Chromosome location	Length (aa)	pI	Molecular weight (Da)	Group
GmGF14a	Glyma.01G058000	Chr01:7642485-7646277	260	4.83	29,487.17	$\epsilon$
GmGF14b	Glyma.02G115900	Chr02:11280858-11285984	260	4.83	29,498.19	$\epsilon$
GmGF14c	Glyma.02G208700	Chr02:39388574-39391014	263	4.67	29,353.86	non- $\epsilon$
GmGF14d	Glyma.04G092600	Chr04:8158031-8160711	251	4.81	28,208.84	non- $\epsilon$
GmGF14e	Glyma.04G099900	Chr04:9132954-9135203	289	4.67	32,432.54	non- $\epsilon$
GmGF14f	Glyma.04G183400	Chr04:45129363-45133820	727	4.94	79,220.24	$\epsilon$
GmGF14g	Glyma.05G158100	Chr05:35025422-35029392	260	4.8	29,249.8	$\epsilon$
GmGF14h	Glyma.06G094400	Chr06:7432085-7434388	251	4.81	28,365.07	non- $\epsilon$
GmGF14i	Glyma.06G101500	Chr06:8052625-8054939	280	5.46	31,708.21	non- $\epsilon$
GmGF14j	Glyma.06G182800	Chr06:15705290-15709591	754	5.06	81,749.77	$\epsilon$
GmGF14k	Glyma.07G226000	Chr07:40298318-40302692	260	4.79	29,579.23	$\epsilon$
GmGF14l	Glyma.08G115800	Chr08:8877809-8881104	260	4.9	29,247.74	$\epsilon$
GmGF14m	Glyma.08G363800	Chr08:47528826-47532060	261	4.81	29,384.49	non- $\epsilon$
GmGF14n	Glyma.12G210400	Chr12:36943077-36946491	262	4.73	29,461.02	$\epsilon$
GmGF14o	Glyma.12G229200	Chr12:38919217-38923409	266	4.85	30,493.26	$\epsilon$
GmGF14p	Glyma.13G270600	Chr13:37265741-37269626	264	4.84	30,207.93	$\epsilon$
GmGF14q	Glyma.13G290900	Chr13:39120795-39124124	262	4.77	29,518.07	$\epsilon$
GmGF14r	Glyma.14G176900	Chr14:43637893-43642553	315	4.71	35,233.85	non- $\epsilon$
GmGF14s	Glyma.17G208100	Chr17:34108328-34108849	160	5.61	18,687.61	non- $\epsilon$
GmGF14t	Glyma.18G298300	Chr18:57587135-57590454	258	4.7	29,063.69	non- $\epsilon$
GmGF14u	Glyma.20G025900	Chr20:2845106-2852380	261	4.79	29,640.22	$\epsilon$
GmGF14v	Glyma.20G043700	Chr20:7939112-7939943	71	5.7	7920.14	$\epsilon$

22 GmGF14 proteins were composed of  $\epsilon$  group or non- $\epsilon$  group, 13 GmGF14 proteins (GmGF14a/b/f/g/j/k/l/n/o/p/q/u/v) belonged to the former group, while the other 9 GmGF14 proteins (GmGF14c/d/e/h/i/m/r/s/t) belonged to the non- $\epsilon$  group.

### Gene structure and motif analysis

Exon/intron pattern divergence plays a crucial role during evolution. We analyzed the exon/intron pattern of *GmGF14s* and found that genes of soybean contained 1–6 introns. Among them, non- $\epsilon$  group *GmGF14* genes contained 1–4 introns, whereas  $\epsilon$  group genes had 1–6 introns. The exon/intron pattern were obviously different in the two groups of *GmGF14* genes, suggesting the diversity of *GmGF14* genes during the evolution (Fig. 3A, Tables S2 and S3). A total of 15 conserved motifs in *GmGF14* genes were identified by MEME software. As shown in Fig. 3B, 5 motifs (motifs 1–5) were annotated as 14-3-3 domains, and most of GmGF14 proteins contained these motifs. All non- $\epsilon$  group GmGF14 proteins shared motifs 3, 4 and 15, whereas most  $\epsilon$  group soybean 14-3-3 proteins contained the motifs 1–7 and motif 15. In addition, the GmGF14f/j in  $\epsilon$  group contained motifs 8-14, and GmGF14v only had motif 6.



**Figure 1** Multiple sequence alignment of all 14-3-3 proteins from soybean.


Full-size DOI: 10.7717/peerj.7950/fig-1

## Chromosomal location and duplication analysis

A chromosomal location map of *GmGF14* genes on each chromosome was drawn. As shown in Fig. 4, 22 *GmGF14* genes were mapped to thirteen of twenty chromosomes unevenly, and they were densely distributed on chromosome 4 and chromosome 6, containing 3 members, respectively (Fig. 4). Most of them were distributed on the two ends of the chromosomes. To better understand the evolution of soybean 14-3-3 genes, we checked genome duplication events in this gene family. The *GmGF14* gene pairs had 19 segmental duplication events without tandem duplication (Table S4). Among the duplication events, genes on chromosome 8 had the largest number of that (Fig. 4).

## Evolution and divergence of the 14-3-3 gene family

We found 19 pairs of paralogous in soybean, 27 orthologous pairs between soybean and Arabidopsis, 15 orthologous pairs in soybean and *M. truncatula* (Table 2). Additionally, we found that two 14-3-3 genes (*GmGF14i* and *GmGF14s*) did not have any homology genes. Two or more *GmGF14* genes matched to one *AtGRF* gene or *Mt14-3-3* gene, implying that these genes might play key roles in the *GmGF14* genes' expansion during evolution. In addition, to examine the evolutionary selection process, we calculated *Ka/Ks* ratios of 19 *GmGF14* paralogous pairs (Table 3). All the *Ka/Ks* value were under 0.3, indicating that they had evolved mainly in strong purifying selection. The gene differentiation of the 19 gene pairs were approximately occurred in the 5–20 MYA.



**Figure 2** Phylogenetic tree analysis of the 14-3-3 genes in *Glycine max*, *Arabidopsis thaliana*, *Medicago truncatula* and *Oryza sativa*. The phylogenetic tree was constructed using MEGA 5.0 by the neighbor-joining method. The Bootstrap value was 1,000 replicates. The two clusters were designated as non- $\epsilon$  group and  $\epsilon$  group, and indicated them in a specific color.


Full-size DOI: 10.7717/peerj.7950/fig-2

### **Cis-elements in *GmGF14s* promoters**

*Cis*-elements involved in transcriptional regulation and can response to variety stresses. We isolated the sequence which is 1.5 kb upstream of the *GmGF14* genes to explore their potential function (Table 4). We found nine potential elements, such as ABRE, AuxRR-core, GARE-motif, CGTCA/TGACG-motif, P-box, TATC-box, TCA-element and TGA-element, were involved in ABA (abscisic acid), IAA (auxin), GA (gibberellin), MeJA (methyl jasmonate) and SA (salicylic acid) regulating mechanism. Additionally, there were four elements (TC-rich repeats, ARE, MBS and LTR) involved in defense/stress, anaerobic induction, drought and low-temperature responses, respectively. In the *GmGF14* promoters, we found different types and numbers of *cis*-elements, indicating that they participated in different regulatory mechanisms during plant growth and development.

### **Expression analysis of *GmGF14* genes in different tissues**

We analyzed the expression level of *GmGF14* genes in different soybean tissues and organs (e.g., root, root hair, flower, nodule, pod, stem, leaf, SAM and seed) based on RNA-seq



**Figure 3** Conserved motifs and gene structure in *GmGF14s*. (A) Exon/intron structures of *GmGF14* genes. (B) Conserved motifs of the *GmGF14s*. Each motif is represented by a number in colored box.


Full-size [DOI: 10.7717/peerj.7950/fig-3](https://doi.org/10.7717/peerj.7950/fig-3)

data (Fig. 5). Results showed that the expression level of most *GmGF14* genes varies in different tissues, suggesting the diversity of their roles. Significantly, most *GmGF14s*' expression level in vegetative organs (e.g., root, root hair, stem, leaf, and SAM) were higher than that of reproductive organs (e.g., flower, pod and seed). Ten *GmGF14* genes (*GmGF14e/i/h/c/r/m/n/q/g/t*) were highly expressed in all tested tissues, suggesting that they regulated the growth and development of soybean. *GmGF14k* was specifically expressed in root, *GmGF14p* was highly expressed in pod and stem, *GmGF14o* was highly expressed in pod. In addition, *GmGF14s* and *GmGF14v* can't be detected in these tissues.

### Expression patterns of *GmGF14s* under abiotic stress

Drought, salinity and cold are major factors affecting the production of soybean under natural conditions. We selected 19 genes from *GmGF14* which were differentially expressed (except *GmGF14s*, *GmGF14u* and *GmGF14v*) to further explore their expression pattern by qRT-PCR under abiotic stresses (Figs. 6–8, Tables S5–S8). The expression levels of them were changed over time during the stresses, showing that there were dynamic processes in the *GmGF14s* when they responding to stresses. In addition, we found that different gene duplication pairs have different expression patterns under the abiotic stresses (Table S5). During drought treatment, the expression pattern of three genes (*GmGF14a/b/g*) were all up-regulated in all the time point (Fig. 6, Table S6). Five genes (*GmGF14a/b/e/g/t*) were highly induced at 1 h, while the expression levels of *GmGF14c/d/q/r* did not significantly changed at all the time after drought treatment. Conversely, under drought treatment, four *GmGF14* genes (*GmGF14c/d/q/r*) were obviously down-regulated. Under salt stress, the expression levels of *GmGF14a/b/c/e/g/i/q/r/t* at 1 h time point up-regulated considerably, then the expression levels decreased (Fig. 7, Table S7). The expression levels of three *GmGF14* genes (*GmGF14f/h/p*) were generally down-regulated at one time point. 5 of 19 *GmGF14* genes (*GmGF14g/h/l/m/t*) were expressed essentially identical, with expression peaking at the first time point (1h) under cold stress (Fig. 8, Table S8). Eight genes





**Figure 4** Chromosome location and duplication events analysis in *Glycine max*.

Full-size  DOI: [10.7717/peerj.7950/fig-4](https://doi.org/10.7717/peerj.7950/fig-4)

(*GmGF14a/b/c/d/e/n/q/r*) were not significantly changed at first time points (1h), followed by a strongly decrease under cold stress.

### Gene ontology enrichment

To further understand the functions of the GmGF14s, we performed GO annotation and GO enrichment analyses (Fig. S2 and Table S9). Usually, the GO terms included biological process, molecular function and cellular component. Within the biological process, most genes were assigned to the cellular process (16/22), single-organism process (16/22), biological regulation (14/22), regulation of biological process (14/22) and response to stimulus (14/22). In terms of cellular process, most GmGF14 proteins were predicted to be involved in cell (20/22), cell part (20/22), membrane (16/22), organelle (16/22) and organelle part (14/22). Few genes had predicted distributions in cell junction (2) and symplast (2). Within the molecular function category, only 20 GmGF14 proteins were

**Table 2** List of paralogous and orthologous pairs between soybean and *Arabidopsis thaliana* and *Medicago truncatula*. Paralogous (Gm-Gm) and orthologous (Gm-Mt and Gm-At) gene pairs.

Gm-Gm	Gm-Mt	Gm-At
GmGF14a/GmGF14b	GmGF14c/Mt14-3-3i	GmGF14f/AtGRF16
GmGF14a/GmGF14k	GmGF14r/Mt14-3-3i	GmGF14j/AtGRF16
GmGF14a/GmGF14u	GmGF14a/Mt14-3-3h	GmGF14d/AtGRF8
GmGF14b/GmGF14k	GmGF14b/Mt14-3-3h	GmGF14d/AtGRF6
GmGF14b/GmGF14u	GmGF14d/Mt14-3-3f	GmGF14h/AtGRF8
GmGF14c/GmGF14e	GmGF14h/Mt14-3-3f	GmGF14h/AtGRF6
GmGF14c/GmGF14r	GmGF14l/Mt14-3-3j	GmGF14c/AtGRF1
GmGF14d/GmGF14 h	GmGF14g/Mt14-3-3j	GmGF14c/AtGRF4
GmGF14f/GmGF14j	GmGF14k/Mt14-3-3g	GmGF14c/AtGRF2
GmGF14g/GmGF14l	GmGF14u/Mt14-3-3g	GmGF14e/AtGRF1
GmGF14g/GmGF14n	GmGF14n/Mt14-3-3c	GmGF14e/AtGRF4
GmGF14g/GmGF14q	GmGF14q/Mt14-3-3c	GmGF14e/AtGRF2
GmGF14k/GmGF14u	GmGF14e/Mt14-3-3e	GmGF14r/AtGRF1
GmGF14l/GmGF14n	GmGF14o/Mt14-3-3b	GmGF14r/AtGRF4
GmGF14l/GmGF14q	GmGF14p/Mt14-3-3b	GmGF14r/AtGRF2
GmGF14m/GmGF14t		GmGF14g/AtGRF9
GmGF14n/GmGF14q		GmGF14l/AtGRF9
GmGF14o/GmGF14p		GmGF14n/AtGRF9
GmGF14r/GmGF14e		GmGF14q/AtGRF9
		GmGF14m/AtGRF3
		GmGF14m/AtGRF7
		GmGF14m/AtGRF5
		GmGF14m/AtGRF5
		GmGF14t/AtGRF3
		GmGF14t/AtGRF7
		GmGF14t/AtGRF5
		GmGF14a/AtGRF12
		GmGF14b/AtGRF12
		GmGF14k/AtGRF12
		GmGF14u/AtGRF12


predicted to be involved in binding. The number of GmGF14 proteins predicted to cell and binding were very high, suggesting that the *GmGF14* gene family may play a crucial role in protein binding and cell development.

## DISCUSSION

In eukaryotes, the *14-3-3s* were highly conserved and could form homo- or hetero- dimers, which then produced different proteins in a protein complex (*Takahashi et al., 2003; Ferl, Manak & Reyes, 2002*). They played important roles in various biological progresses and signal transduction process (*Yoon et al., 2012; Wilson, Swatek & Thelen, 2016*). Hence, we carried out genome-wide analysis of *GmGF14* genes by bioinformatics analysis and qRT-PCR to investigate their regulation during development processes and/or stress

**Table 3** List of *Ka*, *Ks* and *Ka/Ks* values calculated for paralogous *GmGF14* gene pairs.

Gene 1	Gene 2	<i>Ka</i>	<i>Ks</i>	<i>Ka/Ks</i> ratio
GmGF14a	GmGF14b	0.006657464	0.098400891	0.067656541
GmGF14a	GmGF14k	0.050954843	0.601979458	0.084645484
GmGF14a	GmGF14u	0.052668354	0.646064072	0.081521874
GmGF14b	GmGF14k	0.053326335	0.624443785	0.085398135
GmGF14b	GmGF14u	0.05474254	0.686620201	0.07972754
GmGF14c	GmGF14e	0.041987099	0.46411507	0.090467003
GmGF14c	GmGF14r	0.007777127	0.133330521	0.058329686
GmGF14d	GmGF14 h	0.014039603	0.130901237	0.1072534
GmGF14f	GmGF14j	0.038562958	0.163525641	0.235822089
GmGF14g	GmGF14l	0.016897364	0.089125506	0.189590661
GmGF14g	GmGF14n	0.095210598	1.445569001	0.065863752
GmGF14g	GmGF14q	0.096117116	1.432594886	0.067093019
GmGF14k	GmGF14u	0.006698342	0.14254911	0.046989715
GmGF14l	GmGF14n	0.08659209	1.528599239	0.056648
GmGF14l	GmGF14q	0.087490577	1.514174801	0.057781028
GmGF14m	GmGF14t	0.052328317	0.201934457	0.259135156
GmGF14n	GmGF14q	0.008325325	0.127514764	0.065289105
GmGF14o	GmGF14p	0.015822932	0.069501128	0.227664399
GmGF14r	GmGF14e	0.037835791	0.428494041	0.088299457


**Figure 5** Expression analysis of *GmGF14* genes in different tissues. The gene expression values are square-root transformed fragments per kilo-bases per million mapped reads (FPKM). Different colors in map represent FPKM values as shown in bar at top of figure.Full-size DOI: [10.7717/peerj.7950/fig-5](https://doi.org/10.7717/peerj.7950/fig-5)

responses. Li identified 18 *GmGF14* genes in soybean (Li & Dhaubhadel, 2011), however, we found 22 *GmGF14* genes. This may due to the fact that we used a newer version database compared with previous study. Recently, the *14-3-3s* has been reported in several plants, such as Arabidopsis (13), tobacco (17), rice (18), Populus (12), cotton (6), banana (25) and

**Table 4** The number and composition of *cis*-acting regulatory elements of each *GmGF14* gene.

Gene	ABRE (ABA)	AuxRR-core (IAA)	TGA-element (IAA)	CGTCA-motif (MeJA)	TGACG-motif (MeJA)	GARE-motif (GA)	P-box (GA)	TATC-box (GA)	TCA-element (SA)	TC-rich repeats (Defense/stress)	LTR (cold)	ARE (anaerobic)	MBS (drought)
GmGF14a								1			1	1	
GmGF14b								1			1		
GmGF14c	1			2	2								
GmGF14d	3		1	2	2					1			
GmGF14e	4			1	1						1	1	
GmGF14f	2	1		1	1	2			1			1	1
GmGF14g	3			1	2		1						
GmGF14 h	7		1							1			
GmGF14i	3			1	1		1				1		
GmGF14j		1	1			1						2	1
GmGF14k				1	1		1	1					
GmGF14l	2			1	1		1						
GmGF14m	3			1	1								
GmGF14n						1	2						
GmGF14o						1	1						
GmGF14p						1							
GmGF14q				1	1	1	1						
GmGF14r	3			2	2								1
GmGF14s				1	1								
GmGF14t	1			2	2								
GmGF14u				2	2		2	1					
GmGF14v													

## drought




**Figure 6** qRT-PCR analysis reveals *GmGF14* genes under PEG (drought) treatment compared to the controls. Stress treatments and time course are described in 'Materials and Methods'. (A–S) represent different genes which were used in qRT-PCR analysis. Asterisks on top of the bars indicating statistically significant differences between the stress and counterpart controls (\* $p < 0.05$ , \*\* $p < 0.01$ ). Error bars represent SD of biologic replicates.

Full-size DOI: [10.7717/peerj.7950/fig-6](https://doi.org/10.7717/peerj.7950/fig-6)

grape (11) (Saalbach et al., 1997; Ferl, Lu & Bowen, 1994; Yashvardhini et al., 2017; Tian et al., 2015; Zhang et al., 2010; Li et al., 2012; Cheng et al., 2018).

In soybean, 14-3-3s were divided into two groups,  $\epsilon$  group (13 members) and non- $\epsilon$  group (9 members) based on their phylogenetic analysis. Besides, there is a very close relationship between soybean and *M. truncatula*, suggesting that the 14-3-3 family members in legumes are relatively conserved. In addition,  $\epsilon$  group *GmGF14* genes have much more exons/intron than non- $\epsilon$  group genes, while the first intron of non- $\epsilon$  group was longer than that of  $\epsilon$  group. Besides, the members in  $\epsilon$  group contained eight motifs, while non- $\epsilon$  group members had less, usually 3-4 motifs. Furthermore, protein structure analysis show that compared with other species such as banana, grape and rice, the members of 14-3-3s had ten typical antiparallel  $\alpha$ -helices (Yashvardhini et al., 2017; Cheng et al., 2018; Li et al., 2012). The result of 14-3-3 proteins was different from that in other species, that might

## salt




**Figure 7** qRT-PCR analysis reveals. Stress treatments and time course are described in ‘Materials and Methods’. (A–S) represent different genes which were used in qRT-PCR analysis. Asterisks on top of the bars indicating statistically significant differences between the stress and counterpart controls (\* $p < 0.05$ , \*\* $p < 0.01$ ). Error bars represent SD of biologic replicates.

Full-size DOI: [10.7717/peerj.7950/fig-7](https://doi.org/10.7717/peerj.7950/fig-7)

due to the soybean genome has undergone two gene duplication events and has more gene diversity in the process of evolution (Wang *et al.*, 2017).

Gene duplication events is important in gene family expansion and could gain functional diversity during evolution, including tandem, transposition and segment duplication events (Kaessmann, 2010). There were 19 gene pairs involved in segment duplication, while no tandem duplication event occurred in *GmGF14s*, indicating that the segment duplication maybe the major gene duplication for this gene family’s expansion (Cheng *et al.*, 2018). Among them,  $\epsilon$  group had more gene duplication events (14/19; 73.68%) than non- $\epsilon$  group (5/19; 26.32%) in soybean. In addition, we calculated the  $K_s$  value of each paralogous pairs, and found the most recent duplication event in soybean appeared between 5 to 20 MYA, which is consisting with the recent whole genome duplication (WGD) event in soybean (Wang *et al.*, 2017). The  $K_a/K_s$  of all the *GmGF14* gene pairs were less than 0.3, suggesting that they were evolved in mainly under strong purifying selection. This result was similar to other plants, meaning that 14-3-3 genes evolved more slowly at the protein level in plants, and they have a conserved evolutionary pattern in *GmGF14* genes.

## cold



**Figure 8** qRT-PCR analysis reveals. *GmGF14* genes under cold treatment compared to the controls. Stress treatments and time course are described in 'Materials and Methods'. (A–S) represent different genes which were used in qRT-PCR analysis. Asterisks on top of the bars indicating statistically significant differences between the stress and counterpart controls (\* $p < 0.05$ , \*\* $p < 0.01$ ). Error bars represent SD of biologic replicates.

Full-size DOI: 10.7717/peerj.7950/fig-8

It has been reported that the *14-3-3* genes were expressed in different tissues in many plants. *PvGRF1r* might involve in flower development based on the expression patterns in switchgrass (Wu *et al.*, 2016). In banana, the expression quantity of most *MaGRFs* were accumulated during fruit ripening obviously (Li *et al.*, 2012). The expression levels of most *GmGF14* genes in vegetative organs were higher than that of reproductive organs in plants, suggesting that *14-3-3* genes may participate in morphogenesis directly or indirectly. In soybean, *14-3-3* genes involved in nodule mature, they can affect the formation of the early nodule development when the expression levels of *SGF14c* and *SGF14l* reduced (Radwan *et al.*, 2012). In *M. truncatula*, *Mt14-3-3* genes were involved in rhizobium infections, *Mt14-3-3c* was involved in the early stage of nodule formation (Chen *et al.*, 2006). These results suggesting that the *14-3-3* gene family has various functions, they were similar to the results of the GO enrichment. For the GO enrichment, *GmGF14* genes were mainly concentrated in cell development and protein binding. Except this, different *GmGF14* genes had similar expression patterns in different tissues. For example, paralogous pairs

*GmGF14a/b*, *GmGF14k/u*, and *GmGF14o/p* had similar expression patterns in most tested tissues, meanwhile, they also had gene duplication relationship, indicating that they might have similar functions in different tissues, in accordance with the results of GO enrichment.

More and more evidences had suggested that 14-3-3 genes response to environmental stimuli in many plants (Xu & Shi, 2006; Chen et al., 2006; Li et al., 2015). Plant 14-3-3 genes are signal moderators, playing an important role in response to abiotic stress (Li et al., 2015). Overexpression of *AtGRF9* resulted in more carbon distribution from the shoot to the root, and enhanced the drought tolerance of plant by increasing proton secretion in the root growing zone (He et al., 2015). As homologous gene of *AtGRF9*, *GmGF14g* was up regulated during the treatment, and the 1 h drought stress treatment caused a threefold increase of its expression level (its expression increased 3-fold with 1 h drought stress treatment). In tomato, transcription level of four 14-3-3 genes were significantly up-regulated under salt stress (Xu & Shi, 2006). In this study, nine genes (*GmGF14a/b/c/e/g/i/q/r/t*) were first up-regulated and then decreased after salt treatment, indicating that soybean 14-3-3 genes have different regulatory mechanisms under stress. In addition, many 14-3-3 genes (e.g., *GmGF14b/c/g/j*) of soybean changed distinctly under cold treatment, most of the gene expression levels decreased at 6 h and 12 h treatment time points, suggested that they might play a potential role in responding to cold stress. The results of GO enrichment suggested that *GmGF14* genes can respond to stimuli.

The ABA signaling pathway is a major pathway in response to the drought, salt and cold stresses (Zhang et al., 2006; Yu & Qi, 2017). 14-3-3s promoter region contain ABRE promoters, and they could response to stresses directly or indirectly by involving in the ABA signal pathway. In addition, four *cis*-elements of *GmGF14* genes (TC-rich repeats, ARE, MBS and LTR) involved in responding to different abiotic stresses, while other eight *cis*-elements were involved in multiple plant hormone stress responses. Taken together, these results reported that *GmGF14s* may have various functions, including regulate plant growth and response to abiotic stresses.

## CONCLUSIONS

All 22 *GmGF14s* were classified into the  $\epsilon$  group and non- $\epsilon$  group based on their phylogenetic relationship among *A. thaliana*, *O. sativa* and *M. truncatula*. Gene structure and duplication event showed that 14-3-3 gene family was relatively conserved. RNA-seq and qRT-PCR were used to explore the function of *GmGF14s*. The expression levels of most *GmGF14s* showed that they could response to multiple stresses. In summary, these results suggest the potential roles of *GmGF14* genes in plant development and multiple stress responses, therefore provide scientific references for the further study of *GmGF14* genes' function.

## ACKNOWLEDGEMENTS

The authors would like to thank the Key Laboratory of Crop and Livestock Molecular Breeding of Heilongjiang Province for providing plenty of helpful manpower and material support.



## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the National Key R&D Program for Crop Breeding (2016YFD0102105), the Science and Technology Program for Innovation Talents of Harbin (2014RFQYJ016, 2014RFXJY011) and the Project of the Heilongjiang Academy of Agricultural Sciences (HAAS) (2017BZ12). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

National Key R&D Program for Crop Breeding: 2016YFD0102105.

Science and Technology Program for Innovation Talents of Harbin: 2014RFQYJ016, 2014RFXJY011.

Project of the Heilongjiang Academy of Agricultural Sciences (HAAS): 2017BZ12.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Yongbin Wang performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Lei Ling analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Zhenfeng Jiang performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Weiwei Tan Zhaojun Liu performed the experiments, prepared figures and/or tables, approved the final draft.
- Licheng Wu, Shanyong Xia and Jun Ma performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Yuanling Zhao performed the experiments, prepared figures and/or tables, approved the final draft.
- Guangjin Wang conceived and designed the experiments, prepared figures and/or tables, approved the final draft.
- Wenbin Li conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

The raw measurements are available in the [Supplementary Files](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.7950#supplemental-information>.

## REFERENCES

- Bailey TL, Johnson J, Grant CE, Noble WS. 2015. The MEME suite. *Nucleic Acids Research* 43:W39–W49 DOI 10.1093/nar/gkv416.
- Bihn EA, Paul AL, Wang SW, Erdos GW, Ferl RJ. 2010. Localization of 14-3-3 proteins in the nuclei of arabidopsis and maize. *Plant Journal for Cell & Molecular Biology* 12(6):1439–1445 DOI 10.1046/j.1365-313x.1997.12061439.x.
- Camoni L, Visconti S, Aducci P, Marra M. 2018. 14-3-3 proteins in plant hormone signaling: doing several things at once. *Frontiers in Plant Science* 9:Article 297 DOI 10.3389/fpls.2018.00297.
- Cao J, Tan X. 2018. Comparative and evolutionary analysis of the 14-3-3 family genes in eleven fishes. *Gene* 662:76–82 DOI 10.1016/j.gene.2018.04.016.
- Chen F, Li Q, Sun L, He Z. 2006. The rice 14-3-3 gene family and its involvement in responses to biotic and abiotic stress. *DNA Research* 13(2):53–63 DOI 10.1093/dnares/dsl001.
- Cheng C, Yi W, Chai F, Li S, Xin H, Liang Z. 2018. Genome-wide identification and characterization of the 14-3-3 family in *Vitis vinifera* L. during berry development and cold- and heat-stress response. *BMC Genomics* 19(1):579 DOI 10.1186/s12864-018-4955-8.
- Cheng Q, Cheng L, Shen J, Zhang Y, Cao H, Dan L, Shen C. 2016. Genome-wide identification and expression analysis of the 14-3-3 family genes in *Medicago truncatula*. *Frontiers in Plant Science* 7:Article 320 DOI 10.3389/fpls.2016.00320.
- Clamp M, Cuff J, Searle SM, Barton GJ. 2004. The jalview java alignment editor. *Bioinformatics* 20(3):426–427 DOI 10.1093/bioinformatics/btg430.
- De Vetten NC, Ferl RJ. 1994. Two genes encoding GF14 (14-3-3) proteins in *Zea mays*. Structure, expression, and potential regulation by the G-box binding complex. *Plant Physiology* 106(4):1593–1604 DOI 10.1104/pp.106.4.1593.
- Ferl RJ, Lu G, Bowen BW. 1994. Evolutionary implications of the family of 14-3-3 brain protein homologs in *Arabidopsis thaliana*. *Genetica* 92(2):129–138 DOI 10.1007/bf00163762.
- Ferl RJ, Manak MS, Reyes MF. 2002. The 14-3-3s. *Genome Biology* 3(7):1–7 DOI 10.1186/gb-2002-3-7-reviews3010.
- Goodstein DM, Shengqiang S, Russell H, Rochak N, Hayes RD, Joni F, Therese M, William D, Uffe H, Nicholas P. 2012. Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Research* 40(D1):D1178–D1186 DOI 10.1093/nar/gkr944.
- Gouet P, Robert X, Courcelle E. 2003. ESPript/ENDscript: extracting and rendering sequence and 3D information from atomic structures of proteins. *Nucleic Acids Research* 31(13):3320–3323 DOI 10.1007/s10404-008-0309-1.

- He Y, Wu J, Lv B, Li J, Gao Z, Xu W, Baluška F, Shi W, Pang CS, Zhang J. 2015. Involvement of 14-3-3 protein GRF9 in root growth and response under polyethylene glycol-induced water stress. *Journal of Experimental Botany* **66**(8):2271–2281 DOI 10.1093/jxb/erv149.
- Ivica L, Tobias D, Peer B. 2012. SMART 7: recent updates to the protein domain annotation resource. *Nucleic Acids Research* **40**(D1):302–305 DOI 10.1093/nar/gkr931.
- Johana R, Teresa R, Gloria Isabel M, Elsa Z, Juan Carlos R, Beatriz Eugenia F, Jaime R. 2015. Genotypic analysis of genes associated with independent resistance and cross-resistance to isoniazid and ethionamide in mycobacterium tuberculosis clinical isolates. *Antimicrobial Agents & Chemotherapy* **59**(12):7805–7810 DOI 10.1128/AAC.01028-15.
- Juqiang Y, Cixin H, Jing W, Zhehui M, Holaday SA, Allen RD, Hong Z. 2004. Overexpression of the Arabidopsis 14-3-3 protein GF14 lambda in cotton leads to a stay-green phenotype and improves stress tolerance under moderate drought conditions. *Plant & Cell Physiology* **45**(8):1007–1014 DOI 10.1093/pcp/pch115.
- Kaessmann H. 2010. Origins, evolution, and phenotypic impact of new genes. *Genome Research* **20**(10):1313–1326 DOI 10.1101/gr.101386.109.
- Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre<sup>2</sup> web portal for protein modeling, prediction and analysis. *Nature Protocols* **10**(6):845–858 DOI 10.1038/nprot.2015.053.
- Kim MY, Yang JK, Lee T, Lee SH. 2013. Divergence of flowering-related genes in three legume species. *Plant Genome* **6**(3):841–856 DOI 10.3835/plantgenome2013.03.0008.
- Krzywinski M, Schein JI. 2009. Circos: an information aesthetic for comparative genomics. *Genome Research* **19**(9):1639–1645 DOI 10.1101/gr.092759.109.
- Kumar K, Muthamilarasan M, Bonthala VS, Roy R, Prasad M. 2015. Unraveling 14-3-3 proteins in C4 panicoids with emphasis on model plant *Setaria italica* reveals phosphorylation-dependent subcellular localization of RS splicing factor. *PLOS ONE* **10**:e0123236 DOI 10.1371/journal.pone.0123236.
- Li MY, Xu BY, Liu JH, Yang XL, Zhang JB, Jia CH, Ren LC, Jin ZQ. 2012. Identification and expression analysis of four 14-3-3 genes during fruit ripening in banana (*Musa acuminata* L. AAA group, cv. Brazilian). *Plant Cell Reports* **31**(2):369–378 DOI 10.1007/s00299-011-1172-1.
- Li R, Jiang X, Jin D, Dhaubhadel S, Bian S, Li X. 2015. Identification of 14-3-3 family in common bean and their response to abiotic stress. *PLOS ONE* **10**(11):e143280 DOI 10.1371/journal.pone.0143280.
- Li WH, Gojobori T, Nei M. 1981. Pseudogenes as a paradigm of neutral evolution. *Nature* **292**(5820):237–239 DOI 10.1038/292237a0.
- Li X, Dhaubhadel S. 2011. Soybean 14-3-3 gene family: identification and molecular characterization. *Planta* **233**(3):569–582 DOI 10.1007/s00425-010-1315-6.
- Li X, Dhaubhadel S. 2012. 14-3-3 proteins act as scaffolds for GmMYB62 and GmMYB176 and regulate their intracellular localization in soybean. *Plant Signal Behav* **7**(8):965–968 DOI 10.4161/psb.20940.

- Liesbeth V, Tognetti VB, Nathalie G, Judith VD, Liesbeth DM, Agnieszka B, Riet DR, Frank VB, Dirk I. 2015. Growth regulating factor 5 stimulates Arabidopsis chloroplast division, photosynthesis, and leaf longevity. *Plant Physiology* **167**(3):817–832 DOI [10.1104/pp.114.256180](https://doi.org/10.1104/pp.114.256180).
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* **25**:402–408.
- Masuda T, Goldsmith PD. 2009. World soybean production: area harvested, yield, and long-term projections. *International Food & Agribusiness Management Review* **12**:233–236.
- Punta M, Coggill PC, Eberhardt RY, Mistry J, Tate J, Bournsnel C, Pang N, Forslund K, Ceric G, Clements J. 2004. The Pfam protein families database. *Nucleic Acids Research* **28**(1):263–266 DOI [10.1093/nar/gkh121](https://doi.org/10.1093/nar/gkh121).
- Radwan O, Wu X, Govindarajulu M, Libault M, Neece DJ, Oh MH, Berg RH, Stacey G, Taylor CG, Huber SC. 2012. 14-3-3 proteins SGF14c and SGF14l play critical roles during soybean nodulation. *Plant Physiology* **160**(4):2125–2136 DOI [10.1104/pp.112.207027](https://doi.org/10.1104/pp.112.207027).
- Roberts MR, Salinas J, Collinge DB. 2002. 14-3-3 proteins and the response to abiotic and biotic stress. *Plant Molecular Biology* **50**(6):1031–1039 DOI [10.1023/a:1021261614491](https://doi.org/10.1023/a:1021261614491).
- Rodriguez LG, Guan JL. 2010. 14-3-3 regulation of cell spreading and migration requires a functional amphipathic groove. *Journal of Cellular Physiology* **202**(1):285–294 DOI [10.1002/jcp.20122](https://doi.org/10.1002/jcp.20122).
- Rosenquist M, Alsterfjord M, Larsson C, Sommarin M. 2001. Data mining the arabidopsis genome reveals fifteen 14-3-3 genes. Expression is demonstrated for two out of five novel genes. *Plant Physiology* **127**(1):142–149 DOI [10.1104/pp.127.1.142](https://doi.org/10.1104/pp.127.1.142).
- Saalbach G, Schwerdel M, Natura G, Buschmann P, Christov V, Dahse I. 1997. Over-expression of plant 14-3-3 proteins in tobacco: enhancement of the plasma membrane K<sup>+</sup> conductance of mesophyll cells. *Febs Letters* **413**(2):294–298 DOI [10.1016/S0014-5793\(97\)00865-X](https://doi.org/10.1016/S0014-5793(97)00865-X).
- Sehnke PC, Henry R, Cline K, Ferl RJ. 2000. Interaction of a plant 14-3-3 protein with the signal peptide of a thylakoid-targeted chloroplast precursor protein and the presence of 14-3-3 isoforms in the chloroplast stroma. *Plant Physiology* **122**(1):235–241 DOI [10.2307/4279094](https://doi.org/10.2307/4279094).
- Sijbesma E, Skora L, Leysen S, Brunsveld L, Koch U, Nussbaumer P, Jahnke W, Ottmann C. 2017. Identification of two secondary ligand binding sites in 14-3-3 proteins using fragment screening. *Biochemistry* **56**(30):3972–3982 DOI [10.1021/acs.biochem.7b00153](https://doi.org/10.1021/acs.biochem.7b00153).
- Takahashi Y. 2006. Handbook of Neurochemistry and Molecular Neurobiology. 249–270 DOI [10.1007/978-0-387-30381-9\\_12](https://doi.org/10.1007/978-0-387-30381-9_12).
- Takahashi Y, Fukazawa J, Matushita A, Ishida S. 2003. Involvement of RSG and 14-3-3 proteins in the transcriptional regulation of a GA biosynthetic gene. *Journal of Plant Growth Regulation* **22**(2):195–204 DOI [10.1007/s00344-003-0035-6](https://doi.org/10.1007/s00344-003-0035-6).

- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011.** MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology & Evolution* **28(10)**:2731–2739.
- Tian F, Wang T, Xie Y, Zhang J, Hu J. 2015.** Genome-wide identification, classification, and expression analysis of 14-3-3 gene family in *Populus*. *PLOS ONE* **10(4)**:e123225 DOI [10.1371/journal.pone.0123225](https://doi.org/10.1371/journal.pone.0123225).
- Valente C, Turacchio G, Mariggìo S, Pagliuso A, Gaibisso R, Tullio GD, Santoro M, Formiggini F, Spanò S, Piccini D. 2012.** A 14-3-3[ $\gamma$ ] dimer-based scaffold bridges CtBP1-S/BARS to PI(4)KIII[ $\beta$ ] to regulate post-Golgi carrier formation. *Nature Cell Biology* **14(4)**:343–354 DOI [10.1038/ncb2445](https://doi.org/10.1038/ncb2445).
- Wang D, Zhang Y, Zhang Z, Zhu J, Yu J. 2010.** KaKs\_Calculator 2.0: a toolkit incorporating gamma-series methods and sliding window strategies. *Genomics Proteomics & Bioinformatics* **8(1)**:77–80 DOI [10.1016/S1672-0229\(10\)60008-3](https://doi.org/10.1016/S1672-0229(10)60008-3).
- Wang J, Sun P, Li Y, Liu Y, Yu J, Ma X, Sun S, Yang N, Xia R, Lei T. 2017.** Hierarchically aligning 10 legume genomes establishes a family-level genomics platform. *Plant Physiology* **174(1)**:284–300 DOI [10.1104/pp.16.01981](https://doi.org/10.1104/pp.16.01981).
- Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Lee T, Jin H, Marler B, Guo H, Kissinger JC. 2012.** MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Research* **40(7)**:e49 DOI [10.1093/nar/gkr1293](https://doi.org/10.1093/nar/gkr1293).
- Wilson RS, Swatek KN, Thelen JJ. 2016.** Regulation of the regulators: post-translational modifications, subcellular, and spatiotemporal distribution of plant 14-3-3 proteins. *Frontiers in Plant Science* **7**:Article 611 DOI [10.3389/fpls.2016.00611](https://doi.org/10.3389/fpls.2016.00611).
- Wu S, Yan HD, Zhang AL, Huang LK, Yin GH, Lee S. 2016.** Identification and characterization of the 14-3-3 gene family in switchgrass. *Genetics & Molecular Research* **15(4)**:Article gmr15048688 DOI [10.4238/gmr15048688](https://doi.org/10.4238/gmr15048688).
- Xu W, Shi W. 2006.** Expression profiling of the 14-3-3 gene family in response to salt stress and potassium and iron deficiencies in young tomato (*Solanum lycopersicum*) roots: analysis by real-time RT-PCR. *Annals Of Botany* **98(5)**:965–974 DOI [10.1093/aob/mcl189](https://doi.org/10.1093/aob/mcl189).
- Yang L, You J, Wang Y, Li J, Quan W, Yin M, Wang Q, Chan Z. 2017.** Systematic analysis of the G-box Factor 14-3-3 gene family and functional characterization of GF14a in *Brachypodium distachyon*. *Plant Physiology & Biochemistry* **117**:1–11 DOI [10.1016/j.plaphy.2017.05.013](https://doi.org/10.1016/j.plaphy.2017.05.013).
- Yashvardhini N, Bhattacharya S, Chaudhuri S, Sengupta DN. 2017.** Molecular characterization of the 14-3-3 gene family in rice and its expression studies under abiotic stress. *Planta* **247(1)**:1–25 DOI [10.1007/s00425-017-2779-4](https://doi.org/10.1007/s00425-017-2779-4).
- Yoon BC, Zivraj KH, Strohlic L, Holt CE. 2012.** 14-3-3 proteins regulate retinal axon growth by modulating ADF/cofilin activity. *Developmental Neurobiology* **72(4)**:600–614 DOI [10.1002/dneu.20955](https://doi.org/10.1002/dneu.20955).
- Yu F, Qi X. 2017.** Ubiquitination modification precisely modulates the ABA signaling pathway in plants. *Hereditas* **39(8)**:692–706 DOI [10.16288/j.ycz.17-043](https://doi.org/10.16288/j.ycz.17-043).

**Zhang J, Jia W, Yang J, Ismail AM. 2006.** Role of ABA in integrating plant responses to drought and salt stresses. *Field Crops Research* **97**(1):111–119  
[DOI 10.1016/j.fcr.2005.08.018](https://doi.org/10.1016/j.fcr.2005.08.018).

**Zhang ZT, Ying Z, Yang L, Shao SQ, Li BY, Shi HY, Li XB. 2010.** Interactome analysis of the six cotton 14-3-3s that are preferentially expressed in fibres and involved in cell elongation. *Journal of Experimental Botany* **61**(12):3331–3344  
[DOI 10.1093/jxb/erq155](https://doi.org/10.1093/jxb/erq155).