









RESEARCH ARTICLE

Modifications to student quarantine policies in K–12 schools implementing multiple COVID-19 prevention strategies restores in-person education without increasing SARS-CoV-2 transmission risk, January–March 2021

Patrick Dawson^{1,2}^{*}, Mary Claire Worrell¹^{*}, Sara Malone³, Stephanie A. Fritz³^{*}, Heather P. McLaughlin¹^{*}, Brock K. Montgomery³, Mary Boyle³, Ashley Gomel⁴, Samantha Hayes⁴, Brett Maricque³, Albert M. Lai³^{*}, Julie A. Neidich³^{*}, Sarah C. Tinker¹, Justin S. Lee¹, Suxiang Tong¹, COVID-19 Response Fieldwork Laboratory Teams[¶], Rachel C. Orscheln³, Rachel Charney⁴, Terri Rebmann⁴^{*}, Missouri School District Data Coordination Group[¶], Jon Mooney⁵, Catherine Rains⁵, Nancy Yoon⁵, Mabelle Petit⁵, Katie Towns⁵, Clay Goddard⁵, Spring Schmidt⁶, Lisa C. Barrios¹, John C. Neatherlin¹, Johanna S. Salzer¹, Jason G. Newland³

1 COVID-19 Response Team, Centers for Disease Control and Prevention, Atlanta, GA, United States of America, **2** Epidemic Intelligence Service, Centers for Disease Control and Prevention, Atlanta, GA, United States of America, **3** Washington University in St. Louis, St. Louis, MO, United States of America, **4** Saint Louis University, St. Louis, MO, United States of America, **5** Springfield-Greene County Health Department, Springfield, MO, United States of America, **6** Saint Louis County Department of Public Health, Berkeley, MO, United States of America

 These authors contributed equally to this work.

[¶] Membership of the COVID-19 Response Fieldwork and Laboratory Teams and Missouri School District Data and Coordination Group is provided in the Acknowledgments.

* yds5@cdc.gov



OPEN ACCESS

Citation: Dawson P, Worrell MC, Malone S, Fritz SA, McLaughlin HP, Montgomery BK, et al. (2022) Modifications to student quarantine policies in K–12 schools implementing multiple COVID-19 prevention strategies restores in-person education without increasing SARS-CoV-2 transmission risk, January–March 2021. PLoS ONE 17(10): e0266292. <https://doi.org/10.1371/journal.pone.0266292>

Editor: Leonid Chindelevitch, Imperial College London, UNITED KINGDOM

Received: March 16, 2022

Accepted: September 15, 2022

Published: October 20, 2022

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: A limited dataset has been added as [Supporting Information](#) file 1. School data are provided in the supplemental file. Individual-level data cannot be shared publicly because of participation of students under the age of 18. Data are available from the Washington University in St. Louis Study PI (contact co-author Jason Newland [jnewland@wustl.edu] or Institutional PoC Cindy Terrill [terrill@wustl.edu])

Abstract

Objective

To determine whether modified K–12 student quarantine policies that allow some students to continue in-person education during their quarantine period increase schoolwide SARS-CoV-2 transmission risk following the increase in cases in winter 2020–2021.

Methods

We conducted a prospective cohort study of COVID-19 cases and close contacts among students and staff (n = 65,621) in 103 Missouri public schools. Participants were offered free, saliva-based RT-PCR testing. The projected number of school-based transmission events among untested close contacts was extrapolated from the percentage of events detected among tested asymptomatic close contacts and summed with the number of detected events for a projected total. An adjusted Cox regression model compared hazard rates of school-based SARS-CoV-2 infections between schools with a modified versus standard quarantine policy.

for researchers who meet the criteria for access to confidential data.

Funding: The Missouri Department of Health and Senior Services (<https://health.mo.gov/>) provided funding to Washington University in St. Louis to support this study (no grant #, authors: SM, SAF, BKM, MB, BM, AML, LAN, RCO, JGN). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Results

From January–March 2021, a projected 23 (1%) school-based transmission events occurred among 1,636 school close contacts. There was no difference in the adjusted hazard rates of school-based SARS-CoV-2 infections between schools with a modified versus standard quarantine policy (hazard ratio = 1.00; 95% confidence interval: 0.97–1.03).

Discussion

School-based SARS-CoV-2 transmission was rare in 103 K–12 schools implementing multiple COVID-19 prevention strategies. Modified student quarantine policies were not associated with increased school incidence of COVID-19. Modifications to student quarantine policies may be a useful strategy for K–12 schools to safely reduce disruptions to in-person education during times of increased COVID-19 community incidence.

Introduction

The coronavirus disease 2019 (COVID-19) pandemic substantially impacted the operation of kindergarten through grade 12 (K–12) schools in the United States, with many schools switching from in-person instruction to part- or full-time virtual learning during the 2020–2021 school year [1]. Virtual instruction can be an effective learning model for some students, but unequal access to computers, internet services, and childcare services may put some children at an educational disadvantage [2]. Disruptions to in-person instruction may also affect students beyond educational attainment, as K–12 schools often provide critical services (e.g., nutritional, physical, and mental health support) to students and their families [3]. Therefore, schools nationwide have sought strategies to safely operate and maintain in-person learning during the COVID-19 pandemic.

The U.S. Food and Drug Administration (FDA) has authorized the Comirnaty/ Pfizer-BioNTech COVID-19 Vaccine for use in children ages 5 years and older through an Emergency Use authorization (ages 5–15 years) and a full authorization for ages 16 and above [4]. As of March 14, 2022, 34% of the population 5–11 years and 70% of the population 12–17 years had received at least one dose of a COVID-19 vaccine [5]. Vaccine hesitancy and inequitable access to vaccine mean not all communities have high coverage of COVID-19 vaccination [6, 7]. Thus, layered prevention strategies continue to be essential to prevent school-based SARS-CoV-2 transmission for K–12 school students when there are increased COVID-19 Community Levels [8]: implementing a universal masking policy, ensuring physical distancing in classrooms, increasing classroom ventilation with outdoor air, screening testing, having a robust case identification and contact tracing system, and following local isolation and quarantine guidance [9]. Under a standard quarantine policy, students typically must forfeit all in-person activities including in-person instruction for 7–14 days after their last exposure [10]. In Greene County, a 10-day standard quarantine was implemented during the 2020–2021 school year with the option to complete quarantine on day 7 with a negative PCR test collected between 5–7 days post-exposure or a negative antigen test on day 7 post-exposure. St. Louis County implemented a 14-day standard quarantine without the option for completing quarantine early during the same school year.

However, some schools have implemented modifications to standard quarantine guidance to minimize disruptions to in-person learning for students if the exposure is deemed low risk.

In Greene County, Missouri, K–12 schools implemented a modified quarantine policy permitting student close contacts of a person having COVID-19 to attend school in person during their quarantine period if: they were aged ≤ 18 years, their only exposure was in a classroom, they did not have direct physical contact for ≥ 15 minutes in one 24-hour period with the person having COVID-19, and both the student close contact and the person having COVID-19 had worn masks appropriately during the exposure event [11]. In a two-week investigation in Greene County in December 2020, an estimated 240 days of in-person instruction were saved for 30 students participating in the modified quarantine [12]. Additionally, none of the students participating in the modified quarantine were identified as having a SARS-CoV-2 infection during the 14 days following exposure through testing or symptom monitoring [12].

Layered COVID-19 prevention strategies in K–12 schools can limit school-based SARS-CoV-2 transmission despite concurrent high community incidence [9, 12–15], unless there are lapses in implementation or adherence [16, 17]. However, the effect of individual prevention strategies on school-based transmission has not been thoroughly assessed in previous reports. Furthermore, less is known about the impact that modifications to quarantine policies have on school-based transmission risk and schoolwide COVID-19 incidence.

To better understand the effects of COVID-19 prevention strategies and modifications to quarantine policies on school-based SARS-CoV-2 transmission, we conducted a two-month investigation in Missouri public schools. The objectives were to 1) measure the frequency of school-based SARS-CoV-2 transmission; 2) quantify the relative risks of school-based transmission among schoolwide COVID-19 mitigation policies (e.g., quarantine, masking, ventilation, classroom distancing); and 3) compare the schoolwide incidence between schools implementing a modified quarantine and schools following standard quarantine.

Methods

During January 25–March 21, 2021, we worked with school officials in 103 schools across six Missouri public school districts in Greene County (57 schools; districts A–C) and St. Louis County (46 schools; districts D–F), with an estimated 65,621 students and staff (S1 Table). During the investigation, districts A–C schools had implemented a modified quarantine policy while districts D–F schools followed standard quarantine [11, 18, 19]. The project was reviewed and approved by the Washington University in St. Louis Institutional Review Board and conducted consistent with applicable federal law and U.S. Centers for Disease Control and Prevention (CDC) policy (45 C.F.R. part 46, 21 C.F.R. part 56; 42 U.S.C. Sect. 241(d); 5 U.S.C. Sect. 552a; 44 U.S.C. Sect. 3501 et seq.). The project was deemed by the IRB to be a non-research public health surveillance activity [45 CFR 46.102(l)(2)]. Therefore, the need for informed consent was waived. However, participants provided oral agreement to participate, and parents/guardians provided oral agreement for their children aged < 18 years.

Participants

School officials were notified of students and staff members who received a positive SARS-CoV-2 nucleic acid amplification test (NAAT) or antigen test by parents/guardians, local health departments, or school-based testing, typically ≤ 2 days after the laboratory result. For persons having COVID-19 who had been physically present in school, at a school-associated event (e.g., school athletics, extracurricular activities), or on a school bus while potentially infectious (starting two days before symptom onset or collection of their first positive test specimen), school officials conducted contact tracing to identify their school-based close contacts. A close contact was defined as someone who was ≤ 6 feet away from a person with COVID-19 for ≥ 15 minutes in one 24-hour period. In districts A–C, school officials determined if student

close contacts (hereafter, contacts) met criteria for a modified quarantine. Student close contacts of a person having COVID-19 were eligible for modified quarantine if: they were aged ≤ 18 years, their only exposure was in a classroom, they did not have direct physical contact for ≥ 15 minutes in one 24-hour period with the person having COVID-19, and both the student close contact and the person having COVID-19 had worn masks appropriately during the exposure event. School officials followed everyone through completion of their isolation or quarantine period, including whether contacts received a positive NAAT or antigen test.

Case notification

After the school case notification and contact tracing is completed, we asked eligible individuals to participate in an enhanced investigation consisting of a telephone interview and free saliva-based testing. Persons with COVID-19 were eligible for the enhanced investigation if they had been physically present in school, at a school-associated event, or on a school bus while potentially infectious within 14 days of recruitment by the investigation team; contacts were eligible if their most recent school-based exposure was within 14 days of recruitment. Contacts were ineligible if they lived with the person with COVID-19 from the school-based exposure. Participants provided oral agreement to participate, and parents/guardians provided oral agreement for children aged < 18 years.

Interview

For the enhanced investigation, a trained interviewer conducted a standardized telephone interview addressing clinical symptoms; school, community, and household exposures; and demographics. For children aged 12–17 years, a parent/guardian and/or the child was interviewed; for children aged < 12 years, only a parent/guardian was interviewed. Upon completion of their quarantine period, participating contacts were reinterviewed about symptoms, additional exposures and activities, and SARS-CoV-2 testing.

Laboratory testing

Saliva specimens were collected from persons with COVID-19 soon after recruitment and from contacts 5–14 days after their last school-based exposure. Contacts were offered testing regardless of presence of symptoms. Specimens were tested for SARS-CoV-2 by real-time reverse transcription polymerase chain reaction (RT-PCR) as previously described [12]. Full genome sequences were generated from RT-PCR-positive saliva specimens at CDC [20].

Case determination

For each school-based contact who received a positive test result from the enhanced investigation or elsewhere reported to school officials, we conducted a case determination process to assess the likelihood of the infection being from school-based transmission. Infections were classified as probable, possible, or unlikely school-based transmission using epidemiologic and sequencing data from case-contact pairs. The following criteria were used to make the classifications for school-based transmission:

1. Unlikely: the close contact lived in the same household as another person with COVID-19 ≤ 14 days before symptom onset or date of collection of their first positive specimen; their exposure, symptom, or testing timeline was not consistent with the known epidemiology of COVID-19; or the sequence generated from their specimen had > 5 single nucleotide polymorphisms (SNPs) compared to the sequence generated from their school-based index case's specimen.

2. Possible: the close contact had non-household community exposure to a person with COVID-19 ≤ 14 days before symptom onset or date of collection of their first positive specimen and was
3. Probable: the close contact had non-household community exposure to a person with COVID-19 ≤ 14 days before symptom onset or date of collection of their first positive specimen and their only identified close contact was with the school-based person with COVID-19. OR the sequence generated from the close contact's specimen had ≤ 5 SNPs compared to the sequence generated from their school-based index case's specimen.

Classifications were made by at least two members of the investigation team. Discordant classifications were resolved by group discussion with at least one additional team member. In the absence of these data, classification defaulted to probable as this was the most conservative approach in evaluating the effectiveness of COVID-19 mitigation strategies.

Analysis

Data on school- and district-level student enrollment, staffing, demographics, and COVID-19 prevention measures were collected from school officials using a standardized survey. All contact tracing data from school officials were collected in Microsoft Excel and entered into a REDCap database (hosted by Washington University in St. Louis; version 9.5.5) along with data from the enhanced investigation. Data were cleaned and analyzed using R (R Core Team; version 3.6.1) and SAS (SAS Institute; version 9.4).

Descriptive analyses were conducted and two-sided tests for statistical significance were measured using Fisher's exact test where appropriate (including comparing the frequency of transmission events among students in modified quarantine versus standard quarantine). Relative risks of school-based SARS-CoV-2 infections by schoolwide COVID-19 policies were computed using log-binomial regression and accounted for school-level cluster-correlated observations using generalized estimating equations with an independent correlation matrix. Adjusted models included a variable indicating whether the contact was a student or a staff member.

Since not all close contacts received SARS-CoV-2 testing, the percentage of asymptomatic contacts we tested who received a positive test result was extrapolated to contacts who were never tested (contacts who did not receive testing were presumed to be asymptomatic) to create a projected total number of contacts who were infected with SARS-CoV-2: the detected number of positive test results among those who received testing + the extrapolated number of positive test results among those who did not receive testing. Schoolwide COVID-19 crude incidence rates and Cox proportional hazard rates (using observed and projected total case numbers) were compared between schools with a modified versus standard quarantine policy. The approximate number of students and staff at each school whose attendance was 100% virtual or had COVID-19 ≤ 90 days before the start of the study were subtracted from the denominator. Those presumed to have been infected outside of school were censored at the day of their positive test specimen collection. Hazard ratios were adjusted for potential school-level confounding factors: quartiles of the percentage of students eligible to receive free or reduced-price lunch (as a proxy for school resources) and the school's total number of persons with COVID-19 that attended school or a school-related event during the study period.

Results

School characteristics

Of 103 participating schools, COVID-19 prevention strategy survey data were available for 100 (97%). [S1A–S1D Fig](#) describes the mitigation strategies being utilized in the schools. Virtual

instruction was offered by 92% of schools, universal masking policies (face masks required for all students, teachers, staff, and visitors on school grounds) were implemented in 97% of schools, 94% of schools reported efforts to increase ventilation in classrooms, and desks were spaced at least three feet apart in all classrooms in 84% of schools.

School-based secondary transmission

From January 25 to March 21, 2021, a total of 1,864 students, teachers, and staff were identified by school officials through case identification and contact tracing, including 228 eligible persons with COVID-19 and 1,636 contacts (Table 1). Among contacts, 16 (1%) had two school-based exposures within the same 14-day window and 12 (1%) had a second school-based exposure after completion of their first quarantine period. Eligible cases and contacts were identified at 68 (66%) schools. The median number of contacts identified per school index person with COVID-19 was 6 (interquartile range, 2–11).

Of 1,636 contacts, 496 (30%) were tested for SARS-CoV-2, and 14 (3% of tested; 1% overall) received a positive test result. Among tested contacts, we tested 372 (75%) and 124 (25%) were tested elsewhere. Case determinations for the 14 contacts with a positive test result classified 11 as resulting from probable school-based transmission, one as possible, and two as unlikely. For subsequent analyses, probable and possible were combined for a total of 12 school-based transmission events. Six contacts who received a positive test result had paired whole genome sequencing data with their school-based index case, which confirmed epidemiologic linkages for four (67%; all identical or nearly identical sequences) and ruled out epidemiologic linkages for two (33%; classified as unlikely). One pair's sequences, both assigned to the B.1.2 lineage, differed by 15 single nucleotide polymorphisms (SNPs). The other pair's sequences, assigned to lineages B.1.2 and B.1.1.416, differed by 35 SNPs. The 12 school-based transmission events were generated by 11 school index cases (5%) in 10 different schools; a cluster in one school involved a transmission chain with one primary, one secondary, and two tertiary cases (Table 2).

Of 307 contacts we tested who reported no symptoms during the 14 days after their school-based exposure, 3 (1%) were positive. Therefore, among 1,140 contacts with no reported test results (and presumed to be asymptomatic), a projected 11 additional asymptomatic infections may have occurred for a projected total of 23 (1%) school-based transmission events.

Relative risk of school-based SARS-COV-2 infection by schoolwide COVID-19 policy

Unadjusted and adjusted relative risks of school-based SARS-CoV-2 infection among school contacts by schoolwide COVID-19 policy were statistically underpowered, primarily due to the low number of observed school-based infections and the homogeneity of prevention strategies implemented by schools (S2 Table).

Modified versus standard quarantine policies

In schools with a modified quarantine policy, 336 (49%) of 681 contacts were eligible for modified quarantine. Among 345 contacts who school officials deemed ineligible for modified quarantine, 29 were not eligible because they were not students. The primary reasons cited for ineligibility among 316 students include exposure during lunch ($n = 143$; 45%), unmasked exposure ($n = 103$; 33%), exposure during athletic activities ($n = 64$; 20%), and prolonged direct contact ($n = 34$; 11%); multiple reasons were cited for some students (Fig 1).

A total of 66 (20%) students in a modified quarantine had available test results; three students with positive tests (5% of tested; 1% of all students in modified quarantine) were

Table 1. Characteristics of school-based persons with index cases of COVID-19 and close contacts from 103 K–12 schools, Greene and St. Louis Counties, Missouri, January 25–March 21, 2021.

	Persons with index case of COVID-19 (N = 228)	Close contacts (N = 1,636)	Total (N = 1,864)
	Median (Range)		
Age of students, years*	13 (5–18)	13 (4–18)	13 (4–18)
Age of teachers/staff, years†	38 (22–58)	42 (21–64)	40 (21–64)
	N (%)		
School location			
Greene County	90 (39)	671 (41)	761 (41)
District A	60 (26)	485 (30)	545 (29)
District B	14 (6)	92 (6)	106 (6)
District C	16 (7)	94 (6)	110 (6)
St. Louis County	138 (61)	965 (59)	1,103 (59)
District D	100 (44)	719 (44)	819 (44)
District E	31 (14)	183 (11)	214 (11)
District F	7 (3)	63 (4)	70 (4)
School role‡			
Student	163 (71)	1,553 (95)	1,716 (92)
Elementary grade (grades K–5)	49 (21)	492 (30)	541 (29)
Middle school grade (grades 6–8)	41 (18)	387 (24)	428 (23)
High school grade (grades 9–12)	73 (32)	632 (39)	705 (38)
Staff	47 (21)	83 (5)	130 (7)
Teacher	20 (9)	40 (2)	60 (3)
Non-teaching staff	27 (12)	43 (3)	70 (4)
Gender			
Female	80 (35)	360 (22)	440 (24)
Male	48 (21)	345 (21)	393 (21)
Transgender, non-binary, or other gender	0 (–)	5 (<1)	5 (<1)
Unknown	100 (44)	926 (57)	1,026 (55)
Race			
American Indian or Alaska Native	0 (–)	1 (<1)	1 (<1)
Asian	0 (–)	32 (2)	32 (2)
Black or African American	3 (1)	97 (6)	100 (5)
Native Hawaiian or Other Pacific Islander	0 (–)	4 (<1)	4 (<1)
White	117 (51)	515 (31)	632 (34)
Multiracial	8 (4)	35 (2)	43 (2)
Prefer not to say or unknown	100 (44)	952 (58)	1,052 (56)
Ethnicity			
Hispanic/Latino	5 (2)	54 (3)	59 (3)
Non-Hispanic/Latino	123 (54)	648 (40)	771 (41)
Prefer not to say or unknown	100 (44)	934 (57)	1,034 (55)
Preexisting medical conditions§			
Yes	34 (15)	145 (9)	179 (10)
No	87 (38)	528 (32)	615 (33)

(Continued)

Table 1. (Continued)

	Persons with index case of COVID-19	Close contacts	Total
	(N = 228)	(N = 1,636)	(N = 1,864)
Unknown	107 (47)	963 (59)	1,070 (57)

Abbreviations: COVID-19 = coronavirus disease 2019; K–12 = kindergarten through grade 12.

* Among students, age was unknown for 64 (39%) cases and 897 (58%) contacts.

† Among teachers and staff, age was unknown for 18 (38%) cases and 36 (43%) contacts.

‡ For 18 cases, school role is unknown. For 42 student contacts enrolled in K–8 intermediate schools, grade is unknown.

§ Preexisting medical conditions included, but were not limited to, type 1 or type 2 diabetes, hypertension, cardiovascular disease, chronic renal disease, chronic lung disease, immunosuppressive conditions, autoimmune conditions, and premature birth.

<https://doi.org/10.1371/journal.pone.0266292.t001>

determined to have been infected through school-based transmission. All three students discontinued modified quarantine upon receiving a positive test result to begin isolation. None infected another person in the school environment; therefore, there was no observed onward transmission from students in modified quarantine. Applying the same extrapolation approach previously described, a projected additional three cases (1% of 270) would be expected among the 270 student close contacts in modified quarantine without test results for a total of six (2%) transmission events among students in modified quarantine. A projected additional eight cases (1% of 835) would be expected among the 835 student close contacts in standard quarantine without test results for a total of 14 (1%) transmission events among students in standard quarantine.

The difference in frequency of transmission events between student contacts in modified versus standard quarantine was not different when comparing observed cases ($P = 0.41$) or projected cases ($P = 0.50$); thus, students selected for modified quarantine were no more likely to test positive or develop disease and pose a risk for onward school-based transmission than students in standard quarantine.

Using observed cases, the average crude incidence rate of school-based SARS-CoV-2 infections was 1.94 per 100,000 per week in schools that implemented a modified quarantine policy and 4.00 per 100,000 per week in schools following standard quarantine ($P = 0.24$). The adjusted hazard rates of school-based SARS-CoV-2 infections were not different between schools that implemented a modified quarantine policy and schools that did not when using observed cases or total projected cases (for both, hazard ratio, HR = 1.00; 95% confidence interval, CI: 0.97–1.03). The adjusted probability of school-based SARS-CoV-2 infections based on total projected cases reached a maximum of 0.83% (95% CI: 0.75–0.91%) by the end of the study (Fig 2).

Discussion

From January 25 to March 21, 2021, school-based SARS-CoV-2 transmission was rare across 103 Missouri public schools, with a projected total of 23 transmission events. All schools implemented multiple COVID-19 prevention strategies, such as universal masking policies, spacing desks three feet apart, increasing ventilation in classrooms, and contact tracing with implementation of quarantine and isolation policies. These measures have been shown to limit school-based SARS-CoV-2 transmission in some specific settings [9, 12–15], but this report demonstrates this in urban, suburban, and rural public school districts; in elementary, middle, and high school grades; in schools that have implemented a modified quarantine and those that have not; during periods of moderate- to high- COVID-19 Community levels [8, 21, 22]; and for a combined estimated school population of >65,000 students, teachers, and staff. Since

Table 2. Characteristics of school-based persons with index cases of COVID-19 and close contacts who received a positive SARS-CoV-2 test result from K–12 schools, Greene and St. Louis Counties, Missouri, January 25–March 21, 2021.

Transmission event	School district	School type*	Person with index case of COVID-19	Close contact with positive SARS-CoV-2 test
N/A†	A	Elementary school	• Staff	• Student
			• Symptomatic 4d before exposure event‡	• Standard quarantine • Classroom exposure (testing session)
			• 34 contacts	• Unlikely school-based transmission + sequencing§
1	A	Middle school	• Student	• Student
			• Symptomatic 24h after exposure event	• Modified quarantine • Classroom exposure
			• 20 contacts	• Probable school-based transmission
2	A	Intermediate school	• Student	• Student
			• Symptomatic 7d before exposure event	• Modified quarantine • Classroom exposure
			• 20 contacts	• Probable school-based transmission
3	B	Middle school	• Student	• Student
			• Symptomatic 6d before exposure event	• Modified quarantine • Classroom exposure
			• 6 contacts	• Probable school-based transmission
N/A	C	Elementary school	• Student	• Student
			• Symptomatic day of exposure event	• Modified quarantine • Classroom exposure
			• 15 contacts	• Unlikely school-based transmission + sequencing
4	C	High school	• Staff	• Staff
			• Symptomatic 24h after exposure event	• Standard quarantine School kitchen exposure
			• 2 contacts	• Probable school-based transmission + sequencing
5	D	Elementary school	• Staff	• Staff
			• Unknown symptom status	• Standard quarantine • Exposure circumstances unknown
			• 2 contacts	• Probable school-based transmission
6	D	Middle school	• Student	• Student
			• Unknown symptom status	• Standard quarantine • Exposure circumstances unknown
			• 6 contacts	• Probable school-based transmission
7	D	High school	• Student	• Student
			• Symptomatic 48h after exposure event	• Standard quarantine • Classroom exposure + community exposure
			• 4 contacts	• Possible school-based transmission
8	D	High school	• Unknown role	• Student
			• Unknown symptom status	• Standard quarantine • Exposure circumstances unknown
			• Unknown # contacts	• Probable school-based transmission
9	D	High school	• Student	• Student
			• Symptomatic day of exposure event	• Standard quarantine • Exposure circumstances unknown
			• 11 contacts	• Probable school-based transmission

(Continued)

Table 2. (Continued)

Transmission event	School district	School type*	Person with index case of COVID-19	Close contact with positive SARS-CoV-2 test
10 [‡]	F	Elementary school	• Staff	• Teacher
			• Symptomatic day of exposure event	• Standard quarantine
			• 2 contacts	• Unclear exposure location (the staff member’s office is located next to the teacher’s classroom)
11–12	F	Elementary school	• Teacher	• Probable school-based transmission + sequencing
			• Student	• Student
			• Standard quarantine	• Standard quarantine
			• Classroom exposure	• Classroom exposure
			• Symptomatic day of exposure event [¶]	• Probable school-based transmission + sequencing
			• 28 contacts	• Student
• Standard quarantine	• Standard quarantine			
• Classroom exposure	• Classroom exposure			
			• Probable school-based transmission + sequencing	

Abbreviations: COVID-19 = coronavirus disease 2019; d = days; h = hours; K–12 = kindergarten through grade 12.

* School types are elementary school (grades K–5), middle school (grades 6–8), intermediate school (grades K–8), and high school (grades 9–12).

† The close contact received a positive SARS-CoV-2 test result, but school-based transmission was determined to be unlikely.

‡ Symptom onset dates may reflect a person’s retrospective recognition of earlier minor symptoms prior to development of more noticeable symptoms that prompted testing and/or their diagnosis of COVID-19. Therefore, some people may have continued to attend school without considering themselves as having symptoms of COVID-19 until after their diagnosis.

§ Epidemiologic data supported by sequencing results (identical or nearly identical sequences with ≤5 single nucleotide polymorphisms (SNPs) between the specimens from the close contact and school-based person with index case of COVID-19 supported probable school-based transmission; sequences with >5 SNPs supported unlikely school-based transmission).

|| Transmission events 10, 11, and 12 are hypothesized to be part of a single transmission chain at the same school, with a primary case generating a secondary case (event 10), and the secondary case generating two tertiary cases (events 11 and 12).

¶ The teacher became symptomatic before learning of their exposure to the staff member having COVID-19 in transmission event 10.

<https://doi.org/10.1371/journal.pone.0266292.t002>

this investigation, there have been two waves of increased incidence of COVID-19 in the U.S. driven by the Delta and Omicron variants [23, 24]. During future times of increased COVID-19 Community Levels [8], implementation of layered COVID-19 prevention strategies in schools will continue to play a pivotal role in preventing school-based SARS-CoV-2 transmission.

Modifications to student quarantine policies to allow students to continue in-person learning following a low-risk exposure to a person with COVID-19 at school did not result in onward SARS-CoV-2 transmission by generation of tertiary cases. These policies also did not result in increased schoolwide incidence relative to schools that did not implement a modified quarantine policy, even after adjusting for potential confounding factors. Student close contacts deemed eligible for modified quarantine were no more likely to receive a positive test result than their peers in standard quarantine and regained a combined estimated 2,664 days of in-person schooling (assuming each of the 333 students in modified quarantine who did not test positive would have missed eight school days if they had not been eligible for modified quarantine), enough school days for one student to attend kindergarten through their high school graduation. Other school districts with multiple prevention strategies in place have eliminated quarantine when masks are reliably used among persons with COVID-19 and their contacts without observing untoward effects [13]. Beyond the educational benefits of restoring




Fig 1. Primary reasons close contacts of persons with COVID-19 (n = 345) were deemed ineligible for modified quarantine by K-12 school officials, Greene County, Missouri, January 25–March 21, 2021. Abbreviations: COVID-19 = coronavirus disease 2019; K-12 = kindergarten through grade 12. The number inside a rectangle corresponds to the number of close contacts who were deemed ineligible for a modified quarantine based on the category below the rectangle. Rectangles that overlap two reasons indicate both reasons were cited in the decision to deem the individual ineligible for a modified quarantine. *Under Greene County's modified quarantine policy, only students aged ≤ 18 years were eligible for a modified quarantine. † Prolonged direct contact was defined as direct physical contact with the person having COVID-19 for ≥ 15 minutes. ‡ Includes 7 contacts who were also ineligible due to extracurricular activities and 2 contacts who were also ineligible due to contact outside of school.

<https://doi.org/10.1371/journal.pone.0266292.g001>

in-person learning, consideration of modified quarantine policies should factor in the psychosocial impacts on students and families.

There are several limitations to this report. First, school contact tracing may have not identified all persons exposed to someone with COVID-19 in a school setting. Second, we did not test all identified contacts for SARS-CoV-2 following exposure, and therefore the observed number and incidence of school-based SARS-CoV-2 infections is possibly an underestimation. Not all participants or parents/guardians of participants agreed to testing and some participants chose to be tested using other resources within the community. However, positive test results outside of the enhanced investigation were reported to school officials and we projected the number of asymptomatic infections among those without test results to minimize underestimation. Third, due to low variability in school-level prevention strategies and the low number of identified school-based transmission events, analyses of the effect of specific prevention strategies on SARS-CoV-2 transmission were underpowered. Fourth, sequencing data and interview data were not available for all identified school index case-positive contact pairs, and in these instances, all persons who received a positive test result during the 14-day window were presumed to have been infected in school. For the six contacts with a positive test result who had paired sequencing data, it was determined that two (33%) were not infected from their school-based index case; it is possible that if sequencing data were available, school-based transmission may have been ruled out for some of the other contacts with a positive test result.

Public health implications

In this two-month investigation of SARS-CoV-2 transmission in 103 schools implementing layered COVID-19 prevention strategies and modifications to quarantine policies, school-based SARS-CoV-2 transmission was rare and schools that implemented a modified




Fig 2. Adjusted probability of school-based SARS-CoV-2 infections based on total projected cases ($n = 23$) in schools with and without a modified quarantine policy, Greene and St. Louis Counties, Missouri, January 25–March 21, 2021. Top-right inset shows zoomed-in view of adjusted probability curves over the study period. The adjusted* probability curve of school-based SARS-CoV-2 infection in schools with a standard quarantine policy is shown by the teal line; in schools with a modified quarantine policy, it is shown by the dotted black line. The adjusted probability curves and 95% confidence intervals were the same for schools with and without a modified quarantine policy. * The Cox regression model was adjusted for potential school-level confounding factors: quartiles of the percentage of students eligible to receive free or reduced-price lunch (as a proxy for school resources) and the school's total number of cases that attended school or a school-related event during the study period.

<https://doi.org/10.1371/journal.pone.0266292.g002>

quarantine policy did not have greater incidence of school-based SARS-CoV-2 infections. Given these findings and the benefits of restoring in-person learning for students, schools implementing multiple COVID-19 prevention strategies might consider adopting a similar policy during times of increased community incidence. This should be done in conjunction with reinforcement of public health messaging to promote quarantine adherence outside of school. K–12 schools implementing universal face mask policies, promoting physical distancing, and increasing ventilation in classrooms continue to experience low rates of school-based SARS-CoV-2 transmission. As vaccination become available for the majority of the school-aged population since the completion of this investigation, additional consideration and adaptation to modified quarantine policies could be considered. Modifications to student quarantine policies may reduce disruptions to in-person education while maintaining a safe environment for the students and staff.

Supporting information

S1 Table. Characteristics of 103 public K–12 schools participating in SARS-CoV-2 transmission investigation, Greene and St. Louis Counties, Missouri, January 25–March 21, 2021. Abbreviations: K–12 = kindergarten through grade 12; NR = not reported. Note: Data are from surveys completed by school and district officials unless otherwise noted. * Data

from: <https://dese.mo.gov/school-data>. † Includes participating schools only. ‡ Race and ethnicity categories at the school level differed from those collected from individuals as part of the investigation.

(DOCX)

S2 Table. Unadjusted and adjusted relative risks of school-based SARS-CoV-2 infection among school close contacts by schoolwide COVID-19 policy in K–12 schools, Greene and St. Louis Counties, Missouri, January 25–March 21, 2021. Abbreviations: COVID-19 = coronavirus disease 2019; K–12 = kindergarten through grade 12; RR = relative risk; CI = confidence interval. Note: All regression models accounted for cluster-correlated observations at the school level using generalized estimating equations (GEE) with an independent correlation matrix. * Regression models were adjusted for school role (student versus staff). † Surveyed ventilation strategies included opening doors when possible, opening windows when possible, using fans to circulate air, and updating heating, ventilation, and air conditioning (HVAC) systems specifically to prevent COVID-19.

(DOCX)

S1 Fig. (A–D) Schoolwide COVID-19 policies reported in 100 K–12 schools, Greene and St. Louis Counties, Missouri, January 25–March 21, 2021. Abbreviations: COVID-19 = coronavirus disease 2019; HVAC = heating, ventilation, and air conditioning; K–12 = kindergarten through grade 12. * School buildings also include the areas listed in the subsequent two categories: hallways, stairways, gymnasiums, cafeterias, and other special use rooms.

(TIF)

S1 File. Limited dataset.

(XLSX)

Acknowledgments

All the students, families, educators, nurses, administrators, and staff members from participating schools and school districts in Greene and St. Louis Counties, Missouri; Dr. James Blaine; Pam A. Miller; Dr. Matt Pearce; Dr. Shawn Randles; CDC COVID-19 Response Team (Jennifer Frazier, Elizabeth Haller, Laura Hughes-Baker, Catherine N. Rasberry, Hailey Reid); Jordan Valley Community Health Center (Shelley Hall, Shannel Johnson, Mandy Jones, Jennifer Logan, Dr. Matthew Stinson, Amber Tenorio); Mercy Hospital Springfield and St. Louis; Ozarks Technical Community College (Katelyn Antrim, Dr. Daniela Brink, Alane Cordray, Katherine Ebersole, Chanté Whittle); Saint Louis University (Mary Beal, Adrienne Beckett-Ansa, Janaki Bhave, Allie Bodin, Kajal Dholakia, Andrea Hoppert, Rachel Leimkuehler, Elena Dalleo Locascio, Ruband Mahmood, Rachel L. Mazzara, Margaret O'Brien, Komal Patel, Lydia Thomas, Audrey Yao); Springfield-Greene County Health Department (Sean Barnhill, Jordan Coiner, Brad Stulce, Kathryn Wall); University of Missouri School of Medicine (Hosea Covington, Evan Garrad, Dr. David Haustein, Tricia Haynes, Spencer Blake Price, Wyatt Whitman); 4ES Corporation (Dr. Bre Peeler); and Washington University in St. Louis (Lori Barganier, Caleb Gentry, Ian T. Lackey, Savannah Low, Dr. Suong T. Nguyen, Jenna Rideout, Tejas Sekhar, Cindy Terrill, Cole Tipton).

COVID-19 response fieldwork and laboratory teams

Thu-ha Dinh, M.D., Catherine V. Donovan, Ph.D., Victoria Foltz, B.S., Jessica L. Halpin, M.S., Sooji Lee, M.S.P.H., Michelle O'Hegarty, Ph.D., Jonathan Steinberg, M.P.H., Elaine Stevens-Emilien, M.S., Shaniece C. Theodore, Ph.D., Caitlin M. Worrell, M.P.H. (COVID-19 School

Fieldwork Team, CDC); Yan Li, M.Sc., Ying Tao, Ph.D., Anna Uehara, Ph.D., Jing Zhang, Ph.D. (COVID-19 Surge Laboratory Group, CDC); Sarah Greene, M.D., Ph.D., Jaimee Hall, D.O., Alex S. Plattner, M.D. (COVID-19 School Fieldwork Team, Washington University in St. Louis).

Group lead author: Catherine V. Donovan, Ph.D. (phz2@cdc.gov).

Missouri school district data and coordination group

Tammy Fitzpatrick, B.S.N., R.N., Mandy Williams, R.N., Logan-Rogersville School District, Rogersville, MO; Amanda Fields, B.S.N., R.N., Tim Pecoraro, Ed.D., Pattonville School District, St. Ann, MO; Natalie Botkin, B.S.N., R.N., Republic School District, Republic, MO; Glenn Hancock, Ed.S., Amy Wehr, B.S.N., R.N., Rockwood School District, Eureka, MO; Jean Grabeel, M.Ed., R.N., Lee Ann Neill, M.S.N., R.N., Springfield Public Schools, Springfield, MO; Kashina Bell, Ed.D., Sharonica Hardin-Bartley, Ph.D., The School District of University City, University City, MO.

Group lead author: Jean Grabeel, M.Ed., R.N. (jgrabeel@spsmail.org)

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Author Contributions

Conceptualization: Patrick Dawson, Mary Claire Worrell, Sara Malone, Stephanie A. Fritz, Sarah C. Tinker, Rachel C. Orscheln, Jon Mooney, Katie Towns, Clay Goddard, Spring Schmidt, Lisa C. Barrios, John C. Neatherlin, Johanna S. Salzer, Jason G. Newland.

Data curation: Patrick Dawson, Mary Claire Worrell, Sara Malone, Johanna S. Salzer.

Formal analysis: Patrick Dawson, Mary Claire Worrell, Justin S. Lee, Johanna S. Salzer.

Funding acquisition: Jason G. Newland.

Investigation: Patrick Dawson, Mary Claire Worrell, Sara Malone, Stephanie A. Fritz, Heather P. McLaughlin, Brock K. Montgomery, Mary Boyle, Ashley Gomel, Samantha Hayes, Brett Maricque, Albert M. Lai, Julie A. Neidich, Sarah C. Tinker, Justin S. Lee, Suxiang Tong, Rachel C. Orscheln, Rachel Charney, Terri Rebmann, Jon Mooney, Catherine Rains, Nancy Yoon, Mabelle Petit, Katie Towns, Clay Goddard, Spring Schmidt, John C. Neatherlin, Johanna S. Salzer, Jason G. Newland.

Methodology: Patrick Dawson, Mary Claire Worrell, Sara Malone, Stephanie A. Fritz, Heather P. McLaughlin, Brock K. Montgomery, Sarah C. Tinker, Justin S. Lee, Rachel C. Orscheln, Lisa C. Barrios, Johanna S. Salzer, Jason G. Newland.

Project administration: Patrick Dawson, Mary Claire Worrell, Sara Malone, Stephanie A. Fritz, Heather P. McLaughlin, Brock K. Montgomery, Mary Boyle, Samantha Hayes, Brett Maricque, Sarah C. Tinker, Katie Towns, Johanna S. Salzer, Jason G. Newland.

Resources: Patrick Dawson, Mary Claire Worrell, Brett Maricque, Justin S. Lee, Suxiang Tong, Jon Mooney, Catherine Rains, Mabelle Petit, Katie Towns, Johanna S. Salzer, Jason G. Newland.

Software: Sara Malone.

Supervision: Patrick Dawson, Mary Claire Worrell, Sara Malone, Stephanie A. Fritz, Heather P. McLaughlin, Brock K. Montgomery, Ashley Gomel, Samantha Hayes, Brett Maricque,

Sarah C. Tinker, Justin S. Lee, Rachel Charney, Terri Rebmann, Jon Mooney, Lisa C. Barrios, Johanna S. Salzer, Jason G. Newland.

Validation: Patrick Dawson, Mary Claire Worrell, Sara Malone, Heather P. McLaughlin, Sarah C. Tinker, Catherine Rains, Johanna S. Salzer, Jason G. Newland.

Visualization: Patrick Dawson, Mary Claire Worrell, Sara Malone.

Writing – original draft: Patrick Dawson, Mary Claire Worrell, Johanna S. Salzer.

Writing – review & editing: Patrick Dawson, Mary Claire Worrell, Sara Malone, Stephanie A. Fritz, Heather P. McLaughlin, Brock K. Montgomery, Mary Boyle, Ashley Gomel, Samantha Hayes, Brett Maricque, Albert M. Lai, Julie A. Neidich, Sarah C. Tinker, Justin S. Lee, Suxiang Tong, Rachel C. Orscheln, Rachel Charney, Terri Rebmann, Jon Mooney, Catherine Rains, Nancy Yoon, Mabelle Petit, Katie Towns, Clay Goddard, Spring Schmidt, Lisa C. Barrios, John C. Neatherlin, Johanna S. Salzer, Jason G. Newland.

References

1. Honein MA, Barrios LC, Brooks JT. Data and policy to guide opening schools safely to limit the spread of SARS-CoV-2 infection. *JAMA*. 2021; 325(9):823–4. <https://doi.org/10.1001/jama.2021.0374> PMID: 33497433
2. Black E, Ferdig R, Thompson LA. K-12 Virtual Schooling, COVID-19, and Student Success. *JAMA Pediatrics*. 2020; 175(2):119–20.
3. Masonbrink AR, Hurley E. Advocating for Children During the COVID-19 School Closures. *Pediatrics*. 2020; 146(3):e20201440.
4. FDA. COVID-19 Vaccines 2022 [cited 2022 3/5/2022]. Available from: <https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/covid-19-vaccines>.
5. Centers for Disease Control and Prevention. Demographic Trends of People Receiving COVID-19 Vaccinations in the United States 2022 [updated 3/5/2022; cited 2022 3/15/2022]. Available from: <https://covid.cdc.gov/covid-data-tracker/#vaccination-demographics-trends>.
6. Centers for Disease Control and Prevention. Vaccine Hesitancy for COVID-19: County and local estimates 2021 [3 August 2021]. Available from: <https://data.cdc.gov/Vaccinations/Vaccine-Hesitancy-for-COVID-19-County-and-local-es/q9mh-h2tw>.
7. Centers for Disease Control and Prevention. Increasing COVID-19 Vaccine Uptake among Members of Racial and Ethnic Minority Communities: A Guide for Developing, Implementing, and Monitoring Community-Driven Strategies 2021 [3 August 2021]. Available from: <https://www.cdc.gov/vaccines/covid-19/downloads/guide-awardees-community-driven-strategies.pdf>.
8. Centers for Disease Control and Prevention. COVID-19 Community Levels 2022 [cited 2022 3/15/2022]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/science/community-levels.html>.
9. Centers for Disease Control and Prevention. Operational Strategy for K-12 Schools through Phased Prevention: US Department of Health and Human Services, CDC; 2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/community/schools-childcare/operation-strategy.html>.
10. Centers for Disease Control and Prevention. Science Brief: Options to Reduce Quarantine for Contacts of Persons with SARS-CoV-2 Infection Using Symptom Monitoring and Diagnostic Testing 2020 [3 August 2021]. Available from: cdc.gov/coronavirus/2019-ncov/science/science-briefs/scientific-brief-options-to-reduce-quarantine.html.
11. Springfield-Greene County Health Department. Modified Quarantine 2021. Available from: <https://www.springfieldmo.gov/5369/Modified-Quarantine>.
12. Dawson P, Worrell MC, Malone S, Tinker SC, Fritz S, Maricque B, et al. Pilot Investigation of SARS-CoV-2 Secondary Transmission in Kindergarten Through Grade 12 Schools Implementing Mitigation Strategies—St. Louis County and City of Springfield, Missouri, December 2020. *Morbidity and Mortality Weekly Report*. 2021; 70(12):449–55. <https://doi.org/10.15585/mmwr.mm7012e4> PMID: 33764961
13. Hershov RB, Wu K, Lewis NM, Milne AT, Currie D, Smith AR, et al. Low SARS-CoV-2 Transmission in Elementary Schools—Salt Lake County, Utah, December 3, 2020–January 31, 2021. *Morbidity and Mortality Weekly Report*. 2021; 70(12):442–8. <https://doi.org/10.15585/mmwr.mm7012e3> PMID: 33764967

14. Falk A, Benda A, Falk P, Steffen S, Wallace Z, Høeg TB. COVID-19 Cases and Transmission in 17 K–12 Schools—Wood County, Wisconsin, August 31–November 29, 2020. *Morbidity and Mortality Weekly Report*. 2021; 70(4):136–40. <https://doi.org/10.15585/mmwr.mm7004e3> PMID: 33507890
15. Macartney K, Quinn HE, Pillsbury AJ, Koirala A, Deng L, Winkler N, et al. Transmission of SARS-CoV-2 in Australian educational settings: a prospective cohort study. *The Lancet Child & Adolescent Health*. 2020; 4(11):807–16. [https://doi.org/10.1016/S2352-4642\(20\)30251-0](https://doi.org/10.1016/S2352-4642(20)30251-0) PMID: 32758454
16. Stein-Zamir C, Abramson N, Shoob H, Libal E, Bitan M, Cardash T, et al. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020. *Euro Surveillance*. 2020; 25(29):pii = 2001352.
17. Gold JAW, Gettings JR, Kimball A, Franklin R, Rivera G, Morris E, et al. Clusters of SARS-CoV-2 Infection Among Elementary School Educators and Students in One School District—Georgia, December 2020–January 2021. *Morbidity and Mortality Weekly Report*. 2021; 70(8):289–92. <https://doi.org/10.15585/mmwr.mm7008e4> PMID: 33630823
18. Springfield-Greene County Health Department. Quarantine and Isolation 2021. Available from: <https://www.springfieldmo.gov/5251/Quarantine-and-Isolation>.
19. St. Louis County Department of Public Health. Home Quarantine Instructions for Close Contacts to People With COVID-19 2021. Available from: <https://stlouiscountymo.gov/st-louis-county-departments/public-health/covid-19/advisories-safety-recommendations/quarantine/>.
20. Paden CR, Tao Y, Queen K, Zhang J, Li Y, Uehara A, et al. Rapid, sensitive, full-genome sequencing of severe acute respiratory syndrome coronavirus 2. *Emerging Infectious Diseases*. 2020; 26(10):2401–5. <https://doi.org/10.3201/eid2610.201800> PMID: 32610037
21. Springfield-Greene County Health Department. COVID-19 Dashboard: Springfield-Greene County 2021. Available from: <https://www.springfieldmo.gov/5147/Recovery-Dashboard>.
22. St. Louis County Department of Public Health. COVID-19 Statistics 2021. Available from: <https://stlcorona.com/resources/covid-19-statistics/>.
23. Centers for Disease Control and Prevention. Trends in Number of COVID-19 Cases and Deaths in the US Reported to CDC, by State/Territory 2022. Available from: https://covid.cdc.gov/covid-data-tracker/#trends_dailycases.
24. Centers for Disease Control and Prevention. Variant Proportions 2022 [cited 2022 3/5/2022]. Available from: <https://covid.cdc.gov/covid-data-tracker/#variant-proportions>.