



RESPONSE TO COMMENT ON CEFALU ET AL.

Update and Next Steps for Real-World Translation of Interventions for Type 2 Diabetes Prevention: Reflections From a *Diabetes Care* Editors' Expert Forum.

Diabetes Care 2016;39:1186–1201

Diabetes Care 2017;40:e23–e24 | DOI: 10.2337/dci16-0036

William T. Cefalu,¹ John B. Buse,²
Jaakko Tuomilehto,^{3,4,5,6}
G. Alexander Fleming,⁷
Ele Ferrannini,⁸ Hertz C. Gerstein,⁹
Peter H. Bennett,¹⁰
Ambady Ramachandran,¹¹
Itamar Raz,¹² Julio Rosenstock,¹³ and
Steven E. Kahn¹⁴

We thank Dr. Garvey for his letter (1) in response to our 2015 *Diabetes Care* Editors' Expert Forum on the prevention of type 2 diabetes (2). We appreciate his comments regarding the importance of the topic and the need for effective preventive strategies. Dr. Garvey also pointed out two shortcomings in our paper that we would like to address.

Dr. Garvey first stated that “there was a relative lack of emphasis on medication-assisted weight loss as a primary treatment approach, whereas conventional diabetes medications were afforded greater discussion” (1). He further stated, “All five medications approved for chronic management of obesity are highly effective for treating type 2 diabetes, and published clinical trials for three of these medications also demonstrate high efficacy for type 2 diabetes prevention, namely, orlistat, liraglutide 3 mg, and phentermine/topiramate extended release (ER) [3], which was not referenced in the forum” (1). We fully agree that weight loss in the range of 10% does appear

to be maximally effective and can indeed reduce progression to type 2 diabetes. However, our goal in this Expert Forum was to specifically target studies whose primary aim was primary prevention of type 2 diabetes. For instance, the study evaluating phentermine/topiramate ER trials (4) was not primarily a type 2 diabetes prevention trial but rather reported type 2 diabetes incidence as a “subanalysis of a phase 3, randomized, placebo-controlled, double-blind study of overweight/obese subjects (BMI ≥ 27 to ≤ 45 kg/m²) with two or more comorbidities.” Thus, although we appreciate Dr. Garvey's point and do not dispute the evidence he presented for weight loss per se driving diabetes remission, our intent was to feature studies with a main goal of primary prevention of type 2 diabetes. In addition, although it is clear that weight reduction in obese subjects with prediabetes and metabolic syndrome reduces the risk of type 2 diabetes, this is not true in Asian populations with normal BMI and impaired glucose tolerance (IGT). Both Indian and Chinese prevention

trials have shown that we can achieve significant reduction in the incidence of type 2 diabetes without weight loss by lifestyle management.

The second shortcoming Dr. Garvey suggested was our “focus only on prediabetes as a condition conferring high diabetes risk and to ignore metabolic syndrome.” Again, for this Expert Forum, we tried to identify those cohorts within the broader category of “prediabetes” for which there were specific reported interventions. Clearly, the most consistently studied group of individuals with prediabetes has been those with IGT. We recognize that the “prediabetic” state encompasses impaired fasting glucose (IFG), IGT, and combined IFG/IGT. These three subcategories may represent somewhat different pathophysiologies, and individuals in all three—with or without metabolic syndrome—have been shown to have increased diabetes risk (5). Furthermore, it is understood that individuals at the lower end of the glycemic range in all three groups are at lower risk for progressing to diabetes than those at

¹Pennington Biomedical Research Center, Louisiana State University, Baton Rouge, LA

²University of North Carolina School of Medicine, Chapel Hill, NC

³Chronic Disease Prevention Unit, National Institute for Health and Welfare, Helsinki, Finland

⁴Dasman Diabetes Institute, Dasman, Kuwait

⁵Saudi Diabetes Research Group, King Abdulaziz University, Jeddah, Saudi Arabia

⁶Center for Vascular Prevention, Danube University Krems, Krems, Austria

⁷Kinexum, Harper's Ferry, WV

⁸CNR Institute of Clinical Physiology, Pisa, Italy

⁹McMaster University and Hamilton Health Sciences, Hamilton, Ontario, Canada

¹⁰National Institutes of Health, Phoenix, AZ

¹¹India Diabetes Research Foundation and Dr. A. Ramachandran's Diabetes Hospitals, Chennai, India

¹²Diabetes Unit, Department of Internal Medicine, Hadassah Hebrew University Hospital, Jerusalem, Israel

¹³Dallas Diabetes Research Center at Medical City and The University of Texas Southwestern Medical Center, Dallas, TX

¹⁴VA Puget Sound Health Care System and University of Washington, Seattle, WA

Corresponding author: William T. Cefalu, william.cefalu@pbrc.edu.

© 2017 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

the higher end. Although we are not ignoring the fact that other components of metabolic syndrome are also associated with an increased risk of type 2 diabetes, there are many other factors that increase risk in people with prediabetes. In addition, very few studies have evaluated the primary prevention potential specifically in people with metabolic syndrome.

We do not disagree with Dr. Garvey's statements; rather, we want to clarify that our approach was primarily to focus on major primary prevention programs. We agree that heightened awareness and further evaluation are required for individuals in all three subgroups of prediabetes (IFG, IGT, and combined IFG/IGT) and especially for those in each subgroup who are at the higher end of the glycemic range and thus at higher risk. We also recognize Dr. Garvey's work on weight loss and diabetes prevention and agree that weight loss in obese people may be the most important factor in type 2 diabetes prevention.

Acknowledgments. W.T.C. is supported in part by National Institutes of Health (NIH) grant 1U54-GM-104940, which funds the Louisiana Clinical and Translational Science Center, and NIH grant P50-AT-002776.

Duality of Interest. W.T.C. has served as principal investigator on clinical research grants received by his institutions from AstraZeneca, Janssen, MannKind Corp., and Sanofi and has served as a consultant for Intarcia Therapeutics and Sanofi. J.B.B. has been an investigator and/or consultant under contracts between his employer and the following companies: Adocia, Andromeda, Astellas, AstraZeneca, Boehringer Ingelheim, Bristol-Myers Squibb, Dance Biopharm, Elcelyx, Eli Lilly and Co., F. Hoffman-La Roche, Gl Dynamics, GlaxoSmithKline, Halozyme, Intarcia Therapeutics, Johnson & Johnson, Lexicon, MacroGenics, Medtronic Minimed, Merck, Metavention, Novo Nordisk, Orexigen, Osiris, Quest, Sanofi, Scion NeuroStim, Takeda, Tolerelex, and vTv Therapeutics. He is a

consultant to and has stock options from PhaseBio. J.T. has received grant support from AstraZeneca, Boehringer Ingelheim, Merck Sharp & Dohme, Merck Serono, Regeneron, and Sanofi. He has received honoraria for speaking engagements from AstraZeneca, Eli Lilly and Co., Novo Nordisk, and Sanofi and consulting fees from AstraZeneca, Bayer Pharma, and Novo Nordisk. He is a stock shareholder of Orion Pharma. G.A.F. is a panel member/consultant for Acosti, Adocia, Arisaph, Artemis, Bio-Cancer Treatment International Ltd., Becton Dickinson, BioCon, Diasome, Dompé, Emperra, Exsulin, Thermo Fisher Scientific, Gilead, Hyperion, IMTherapeutics, Innoneo, Intarcia Therapeutics, Islet Sciences, Lexicon, Locemia, Johnson & Johnson, MannKind Corp., Mars, MediWound, Melior, N-Gene, NuSirt, Royalty Pharma, Pfizer, Rhythm, Sanofi, SkyePharma, Strongbridge Biopharma, SynAgile, Takeda, Teva, Thermalin, Thetis, ThromboGenics, Tolerion, VeroScience, and Versartis. He is a stock/shareholder of Ammonett Pharma, Exsulin Corporation, Locemia, Innoneo, Synagile, and Thetis. E.F. is an advisory board member for Boehringer Ingelheim/Eli Lilly and Company and Merck Sharp & Dohme. He is a consultant for AstraZeneca, GlaxoSmithKline, and Janssen and a speaker for AstraZeneca, Boehringer Ingelheim, Eli Lilly and Co., Merck Sharp & Dohme, Mitsubishi, Novo Nordisk, Sanofi, and Takeda. He receives research grant support from Eli Lilly and Co. H.C.G. has received grant support from AstraZeneca, Eli Lilly and Co., Merck, and Sanofi; honoraria for speaking engagements from AstraZeneca, Berlin Chemie, Boehringer Ingelheim, Novo Nordisk, and Sanofi; and consulting fees from Abbott Pharmaceuticals, Amgen, AstraZeneca, Boehringer Ingelheim, Eli Lilly and Co., Kaneq Bioscience, Merck, Novo Nordisk, and Sanofi. A.R. is an advisory board member for AstraZeneca and Merck Sharp & Dohme. He has received honoraria for speaking engagements from Bayer, Eli Lilly and Co., Merck Sharp & Dohme, Novartis, Novo Nordisk, and Sanofi. The India Diabetes Research Foundation, of which he is president, has received research funding support from AstraZeneca and Novartis. I.R. has served on advisory boards for AstraZeneca/Bristol-Myers Squibb, Boehringer Ingelheim, Eli Lilly and Co., LabStyle Innovations, Merck Sharp & Dohme, Novo Nordisk, Orgenesis, Sanofi, and SmartZyme Innovation; has been a consultant for AstraZeneca/Bristol-Myers Squibb, FuturRx, Gili

Medical, Insuline Medical, Kamada, and NephroGenex; has been a speaker for AstraZeneca/Bristol-Myers Squibb, Boehringer Ingelheim, Eli Lilly and Co., Johnson & Johnson, Merck Sharp & Dohme, Novartis Pharma AG, Novo Nordisk, Sanofi, and Teva; and is a stock/shareholder in Glucome, Insuline Medical, LabStyle Innovations, Orgenesis, and SmartZyme Innovation. J.R. has served on advisory boards and received honoraria or consulting fees from AstraZeneca, Boehringer Ingelheim, Bristol-Myers Squibb, Daiichi Sankyo, Eli Lilly and Co., Intarcia Therapeutics, Janssen, Merck, Novo Nordisk, and Sanofi. He has received grants/research support from Asahi, AstraZeneca, Boehringer Ingelheim, Daiichi Sankyo, Eli Lilly and Co., GlaxoSmithKline, Hanmi, Intarcia Therapeutics, Janssen, Lexicon, MannKind Corporation, Merck, Novo Nordisk, Pfizer, Sanofi, and Takeda. S.E.K. is a consultant/advisory board member for Boehringer Ingelheim, Elcelyx, Eli Lilly and Co., GlaxoSmithKline, Intarcia Therapeutics, Janssen, Merck, Novo Nordisk, and Receptos and has received grant support from Eli Lilly and Co. No other potential conflicts of interest relevant to this article were reported.

References

1. Garvey WT. Comment on Cefalu et al. Update and next steps for real-world translation of interventions for type 2 diabetes prevention: reflections from a *Diabetes Care* Editors' Expert Forum. *Diabetes Care* 2016;39:1186–1201 (Letter). *Diabetes Care* 2017;40:e21–e22. DOI: 10.2337/dci16-0022
2. Cefalu WT, Buse JB, Tuomilehto J, et al. Update and next steps for real-world translation of interventions for type 2 diabetes prevention: reflections from a *Diabetes Care* Editors' Expert Forum. *Diabetes Care* 2016;39:1186–1201
3. Garvey WT, Mechanick JL, Brett EM, et al.; Reviewers of the AACE/ACE Obesity Clinical Practice Guidelines. American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract* 2016;22:842–884
4. Garvey WT, Ryan DH, Henry R, et al. Prevention of type 2 diabetes in subjects with prediabetes and metabolic syndrome treated with phentermine and topiramate extended release. *Diabetes Care* 2014;37:912–921
5. Cefalu WT. "Prediabetes": are there problems with this label? No, we need heightened awareness of this condition! *Diabetes Care* 2016;39:1472–1477