

Clinical guidelines in Hematology

Wanderley Marques Bernardo

Medicine School, Universidade de
Sao Paulo – USP, Sao Paulo, SP
Centro Universitário Lusfada – UNILUS,
Santos, SP, Brazil

Clinical guidelines are now a powerful tool in decision making in the complex process of healthcare. There is no absolute definition of its impact on the clinical outcomes and in different patient populations. Nevertheless, its role is unquestionable in the regulation and organization of the healthcare system as a whole.⁽¹⁻⁸⁾

Evidence-based clinical guidelines balance the diverse interests involved during the process to ensure that patients receive an adequate standard of healthcare. Through clinical guidelines we can compare our experience with recommendations, which not only teaches us and brings our knowledge up-to-date, but also allows us to reflect on the main issue: what level of uncertainty am I accepting with my current conduct of this patient?⁽⁹⁻¹⁵⁾

The central principle of the *Associação Médica Brasileira (AMB)/Conselho Federal de Medicina (CFM)* Guidelines Program is to prepare the physician to answer four basic questions: a) what do I do in my clinical practice; b) for whom do I do it; c) how do I do it and d) why do I do it?⁽¹⁶⁾

The development of recommendations can be interpreted as a way to limit medical autonomy, but in fact, it is to make our actions in healthcare in Brazil transparent, clearly stating the strength of scientific evidence that supports each of these conducts by estimating the level of uncertainty involved in decision making.⁽¹⁷⁻²²⁾

We established standards, providing conduct options focused on the patient in relation to what we do: recommendations for diagnosis, prevention, treatment and prognosis; for whom we do it: patients with indications to meet their expectations and individuality, and never forgetting the minorities; how we do it: defining the method by which to develop our detailed and explicit conduct; and why we do it: to support our decisions on the benefits, risks and harm to patients.⁽²³⁻²⁶⁾

The AMB-CFM Guidelines Program together with the societies of medical specialties, members of the AMB, has already prepared 500 clinical guidelines and today has about 120 in development. In addition, continuing medical education and participation in international networks that develop evidence-based guidelines is included in the Guidelines Programs.

In 2011, Brazilian hematology through its society (the *Associação Brasileira de Hematologia e Hemoterapia – ABHH*) started an unprecedented process of developing evidence-based protocols within the AMB-CFM Guidelines Program. The association initially chose six major hematological diseases: Sickle cell anemia, chronic myeloid leukemia, acute promyelocytic leukemia, non-promyelocytic leukemia, idiopathic thrombocytopenic purpura and multiple myeloma.

Each theme (Guideline) is composed of important clinical questions (on average 15) prepared by experts. These questions are structured using the acronym, PICO (P: patient, I: Intervention C: Comparison, O: Outcome) as a guide to search available evidence by an extensive systematic review of the literature to find evidence to support the recommendations for each clinical question. The recommendations are based on the strongest scientific evidence and aim to help hematologists make their decisions on each individual patient.^(3,16,19,21)

In mid-2012, the first six issues will be completed initiating a series of feasible guidelines developed using a rigorous methodology written in a clear and objective language. Without doubt, participants at all levels of the healthcare system will benefit, but mostly these benefits will be reflected in the care provided to hematology patients in Brazil.

We recognize the difficulties of obtaining and critically analyzing the evidence, the pressure of interest that are not always directed to the care of patients and the difficulties of the National Health System in relation to its structure, diversity and inequality. But we also know the effort and determination of many, who, through the guidelines, will establish a discerning, flexible, ethical, and reflective language, based on evidence that meets the basic needs and expectations of patients.^(2,21,22)

Conflict-of-interest disclosure:
The author declares no competing
financial interest

Submitted: 10/27/2011
Accepted: 10/28/2011

Corresponding author:

Wanderley Marques Bernardo
Rua São Carlos do Pinhal, 324
01333-903 – São Paulo, SP, Brazil
wmbernardo@usp.br

www.rbhh.org or www.scielo.br/rbhh

DOI:

References

1. Andrews EJ, Redmond HP. A review of clinical guidelines. *Br J Surg*. 2004;9(8):1:956-64.
2. Norheim OF. Healthcare rationing - are additional criteria needed for assessing evidence based clinical practice guidelines? *BMJ*. 1999; 319(7222):1426-9.
3. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ*. 1996;312(7023):71-2. Comment in: *J Psychiatr Ment Health Nurs*. 2005;12(6):739-44. *Aust Health Rev*. 2008;32(2):204-7. *BMJ*. 1996;313(7050):169-70; author reply 170-1.
4. Hopkins A. Some reservations about clinical guidelines. *Arch Dis Child*. 1995;72(1):70-5.
5. Woolf SH. Practice guidelines: a new reality in medicine. I. Recent developments. *Arch Intern Med*. 1990;150(9):1811-8.
6. Petrie JC, Grimshaw JM, Bryson A. The Scottish Intercollegiate Guidelines Network Initiative: getting validated guidelines into local practice. *Health Bull (Edinb)*. 1995;53(6):345-8.
7. Horton R. NICE: a step forward in the quality of NHS care. National Institute for Clinical Excellence. National Health Service. *Lancet*. 1999;353(9158):1028-9. Comment on: *Lancet*. 1999;353(9158): 1079-82.
8. Atkins D, Fink K, Slutsky J; Agency for Healthcare Research and Quality; North American Evidence-based Practice Centers. Better information for better health care: the Evidence-based Practice Center program and the Agency for Healthcare Research and Quality. *Ann Intern Med*. 2005;142(12 Pt 2):1035-41.
9. Bernardo WM, Jatene FB, Nobre MRC. Apesar das evidências, por que persiste a variação nos cuidados ao paciente cirúrgico? *Rev Assoc Med Bras*. 2005;51(4):183-4.
10. Lassen K, Hannemann P, Ljungqvist O, Fearon K, Dejong CH, von Meyenfeldt MF, Hausel J, Nygren J, Andersen J, Revhaug A; Enhanced Recovery After Surgery Group. Patterns in current perioperative practice: survey of colorectal surgeons in five northern European countries. *BMJ*. 2005;330(7505):1420-1. Comment in: *BMJ*. 2005;330(7505):1401-2.
11. Garber AM. Evidence-based guidelines as a foundation for performance incentives. *Health Aff (Millwood)*. 2005;24(1):174-9. Comment in: *Health Aff (Millwood)*. 2005;24(3):886.
12. Boyd CM, Darer J, Boulton C, Fried LP, Boulton L, Wu AW. Clinical practice guidelines and quality of care for older patients with multiple comorbid diseases: implications for pay for performance. *JAMA*. 2005;294(6):716-24. Comment in: *JAMA*. 2005;294(6): 741-3. *JAMA*. 2006;295(1):33; author reply 34-5.
13. Berger JT, Rosner F. The ethics of practice guidelines. *Arch Intern Med*. 1996;156(18):2051-6. Comment in: *Arch Intern Med*. 1996; 156(18):2038-40.
14. Choudhry NK, Stelfox HT, Detsky AS. Relationships between authors of clinical practice guidelines and the pharmaceutical industry. *JAMA*. 2002;287(5):612-7
15. Waclawski ER, Madan I. In the current era of evidence-based guidelines, do consensus-based guidelines still have a place? *Occup Med (Lond)*. 2005; 55(5):343-4. Comment on: *Occup Med (Lond)*. 2005;55(5):345-8. *Occup Med (Lond)*. 2005;55(5):369-70.
16. Projeto Diretrizes. Associação Médica Brasileira. Available at: <http://www.projetodiretrizes.amb.org.br>.
17. EPC Evidence Reports. Agency for Healthcare Research and Quality. Available at: <http://www.ahrq.gov/clinic/epcindex.htm>.
18. NICE guidance. National Health Institute. Available at: <http://www.nice.org.uk/>
19. Levels of Evidence and Grades of Recommendation. Centre for Evidence-Based Medicine. Oxford. Available at: <http://www.cebm.net/>.
20. Cabana MD, Rand CS, Powe NR, Wu AW, Wilson MH, Abboud PA, et al. Why don't physicians follow clinical practice guidelines? A framework for improvement. *JAMA*. 1999;282(15):1458-65. Comment in: *JAMA*. 200;283(13):1685; author reply 1686.
21. Grimshaw JM, Thomas RE, MacLennan G, Fraser C, Ramsay CR, Vale L, et al. Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health Technol Assess*. 2004;8 (6):iii-iv,1-72.
22. Woolf SH, Grol R, Hutchinson A, Eccles M, Grimshaw J. Clinical guidelines: potential benefits, limitations, and harms of clinical guidelines. *BMJ*. 1999; 318(7182):527-30. Comment in: *BMJ*. 2001;322(7297):1258-9.
23. The AGREE Collaboration. Appraisal of Guidelines Research and Evaluation. Agree Instrument. Available at: <http://www.agreecollaboration.org>.
24. Guidelines International Network (GIN).[Internet]. Scotland; c2002-2011. Available at: <http://www.g-i-n.net>
25. Hurwitz B. Legal and political considerations of clinical practice guidelines. *BMJ*.1999;318(7184):661-4.
26. Haycox A, Bagust A, Walley T. Clinical guidelines-the hidden costs. *BMJ* 1999;318(7180):391-3.