

Review Article

Asymptotic Identity in Min-Plus Algebra: A Report on CPNS

Ming Li¹ and Wei Zhao²

¹ School of Information Science & Technology, East China Normal University, No. 500 Dong-Chuan Road, Shanghai 200241, China

² Department of Computer and Information Science, University of Macau, Avenue Padre Tomas Pereira, Taipa, Macau

Correspondence should be addressed to Ming Li, ming_lihk@yahoo.com

Received 23 March 2011; Accepted 20 April 2011

Academic Editor: Carlo Cattani

Copyright © 2012 M. Li and W. Zhao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Network calculus is a theory initiated primarily in computer communication networks, especially in the aspect of real-time communications, where min-plus algebra plays a role. Cyber-physical networking systems (CPNSs) are recently developing fast and models in data flows as well as systems in CPNS are, accordingly, greatly desired. Though min-plus algebra may be a promising tool to linearize any node in CPNS as can be seen from its applications to the Internet computing, there are tough problems remaining unsolved in this regard. The identity in min-plus algebra is one problem we shall address. We shall point out the confusions about the conventional identity in the min-plus algebra and present an analytical expression of the asymptotic identity that may not cause confusions.

1. Introduction

We use the term cyber-physical networking systems (CPNS) instead of cyber-physical systems (CPS) as that in Song et al. [1] for the meaning of Internet of Things (IoT) that was stated by Commission of the European Communities [2] or Networks of Things (NoT) as discussed by Ferscha et al. [3], intending to emphasize the point that we are interested in the networking theory in CPS. Communication networks in CPNS include, but are never limited to, the Internet. Physical systems considered in CPNS are heterogeneous, ranging from telemedicine systems to geophysical ones, see, for example, Clifton et al. [4], Traynor [5], Chang [6]. Obviously, data in various physical systems are heterogeneous, see, for example, Chang [6], Goodchild [7], Lai and Xing [8], Mandelbrot [9–11], Hainaut and Devolder [12], Cattani [13–17], Chen et al. [18–22], Mikhael and Yang [23], Bakhoum and Toma [24–26], Li [27–32], Li et al. [33–39], Messina et al. [40], Humi [41], Dong [42], Liu [43], Toma [44], Abuzeid et al. [45], [46–49], Werner [50], and West [51], just naming a few.

There are two challenge issues in CPNS. On the one hand, data models that are irrelevant of statistics of a random function $x(t)$ are greatly desired. On the other hand, theory that may be used to linearize nonlinear data transmission systems but irrelevant of their nonlinearity is particularly

expected, because communication systems, including the Internet, are, in nature, nonlinear due to queuing, see, for example, Akimaru and Kawashima [52], Yue et al. [53], Gibson [54], Cooper [55], Pitts and Schormans [56], McDysan [57], and Stalling [58]. In short, we are interested in *data models that are irrelevant of their statistics and system theory that is irrelevant of the nonlinearity of systems*.

The early work regarding the above in italic may refer to Cruz [59–61], Zhao and Ramamritham [62], Raha et al. [63], Chang [64, 65], Boudec [66], Boudec and Patrick [67], Firoiu et al. [68], and Agrawal et al. [69]. Following Cruz [59, 60], the theory for the above in italic is called network calculus, see, for example, [66, 67], Jiang and Liu [70]. Chang [71] uses the term (σ, ρ) calculus, which is taken as the synonym of network calculus of Cruz in this paper.

The main application area of network calculus is conventionally to computer science, the Internet in particular, see, for example, Wang et al. [72, 73], Li and Zhao [74, 75], Fidler [76], Jiang [77], Jiang et al. [78], Liu et al. [79], Li et al. [80], Li and Kinghtly [81], Burchard et al. [82], Ng et al. [83], Raha et al. [84, 85], Starobinski and Sidi [86], Fukás et al. [87], Jia et al. [88], Golestani [89], and Lenzini et al. [90]. However, we have to emphasize the point that its applications are never limited to computer science. Rather, it is a theory to model data irrelevant of their statistics and to deal with data transmission without the necessity in principle to consider

the nonlinearity of transmission systems, as we shall explain in the next section. Therefore, it may be a promising tool to deal with data and systems in CPNS.

Basically, the fundamental theory of network calculus consists of three parts as described below.

- (i) (σ, ρ) model of arrival data $x(t)$,
- (ii) relationship between $x(t)$, single system (or node or server) $S(t)$ that is usually called service curve, and departure data $y(t)$,
- (iii) departure data $y(t)$ of a series of systems (nodes or servers) $S_n(t)$ ($n = 1, 2, \dots$), driven by arrival data $x(t)$,

where min-plus algebra plays a role, see, for example, [66, 67, 70, 71, 76].

The contributions of this paper are in the following three aspects:

- (i) the problem statement,
- (ii) the proof of the existence of the identity in the min-plus algebra in the domain of generalized functions,
- (iii) the asymptotic expression of the identity.

The rest of paper is organized as follows. Research background is discussed in Section 2. In Section 3, we will brief the min-plus algebra and state the problem regarding the identity in this algebra system. In Section 4, we shall address the existence of the identity in the min-plus algebra. The asymptotic expression of the identity is presented in Section 5. Discussions are given in Section 6, which is followed by conclusions.

2. Research Background

Data in CPNS are heterogeneous. They may be from sensors like radio-frequency identification (RFID), see, for example, [91], Ilie-Zudor et al. [92], Ahuja and Potti [93], data traffic in the Internet [38], transportation traffic (see [94–98]), ocean waves (see [31]), sea level (see [36, 99]), medical signals (see [14]), hydrological data (see [100]), financial data (see [101]), and so on. They may be Gaussian (see [29, 31]) or non-Gaussian (see [12, 102]). They may be in fractional order or integer order. In the case of fractional order, they may be unifractal or multifractal. The sample size of data of interest may be long enough for statistical analysis or very short, for example, a short conversation in mobile phone networks. On the other side, systems are also heterogeneous. Therefore, CPNS challenges us two tough issues. One is in data modeling and the other system modeling. We shall exhibit that the min-plus algebra in network calculus may yet serve as a tool in this regard.

2.1. Network Model. We first explain a single node in CPNS. Then, a model of tandem network is mentioned.

2.1.1. Nonlinearity of Node in CPNS. Denote by N a node in CPNS, see Figure 1. Suppose there are m clients arriving at




FIGURE 1: Single node in CPNS.

the input of N at time t , see, for example, Starobinski et al. [103].

Without confusions, we use N to represent the operator of node N such that

$$y_i(t) = Nx_i(t), \quad 1 \leq i \leq m. \quad (1)$$

Recall that queuing is a phenomenon often occurring in CPNS. For instance, cars in highways are often queued. Clients in a library for borrowing or returning books need queuing. Suppose client $x_i(t)$ suffers from delay $d_i(t)$. Then,

$$y_i(t) = x_i(t + d_i(t)), \quad 1 \leq i \leq m. \quad (2)$$

Note that $d_i(t)$ is a random variable in two senses. One is

$$d_i(t) \neq d_j(t), \quad 1 \leq i \leq m, 1 \leq j \leq m, i \neq j. \quad (3)$$

The other is

$$d_i(t_1) \neq d_i(t_2), \quad 1 \leq i \leq m, t_1 \neq t_2. \quad (4)$$

Therefore, we have the following remark.

Remark 1 (nonlinearity). A node N in CPNS is usually nonlinear. That is,

$$\sum y_i(t) \neq \sum Nx_i(t), \quad 1 \leq i \leq m. \quad (5)$$

2.1.2. Number of Arrivals is Random. The number of arrivals, denoted by m in Figure 1, is random.

Note 1. We need theory to deal with a nonlinear node N with m arrival clients, where m is a random variable.

2.1.3. Tandem Network Model. A single node previously described is not enough in CPNS since a client may be served by a series of n nodes, which we call tandem network, see Figure 2.

According to Remark 1, each node in Figure 2 is nonlinear. In addition, considering Note 1, we see that the number of arrival clients at the input of each node is random. Some clients may go through from $N1$ to Nn while others may not. For instance, client $x_{1i1}(t)$ leaves the tandem network when it passes through $N1$. Further more, some clients, for example, $x_{21}(t)$, arrive at this tandem network




FIGURE 2: Tandem network.

at the input of $N2$. In general, how many clients leave the tandem network at the output of a specific node and how many clients arrive at the input of another specific node are uncertain.

Note 2. We need theory to handle a nonlinear system that is a tandem network as that in Figure 2 to assure the quality of service (QoS) of a specific client or of a specific class of clients within a given period of time.

The above Note 1 and Note 2 propose two challenge tasks in system theory. We shall explain how min-plus algebra is capable of dealing with those tasks late.

2.2. Data Modeling. We consider two classes of data flow. One is arrival data in the aggregated case, or aggregated clients, and the other arrival data of a specific client. In terms of network communications, the former is usually called aggregated arrival traffic while later arrival traffic at connection level. Without confusions, we use the term traffic rather than client.

One of radical properties of arrival traffic (traffic for short) is remarked below.

Remark 2 (positive). Traffic $x_i(t)$ is positive. That is,

$$x_i(t) \geq 0, \quad t \in \mathbf{R}, \quad (6)$$

where \mathbf{R} is the set of real numbers.

Another radical property of traffic is that the maximum of $x_i(t)$ is finite. More precisely, the value of $x_i(t)$ may never be infinite. Thus, we have the following remark.

Remark 3 (finite range). The maximum of $x_i(t)$ is finite. That is,

$$0 \leq x_i(t) \leq x_{i,\max}. \quad (7)$$

Remark 4 (randomness). The function $x_i(t)$ is usually random. This implies that

$$x_i(t_1) \neq x_i(t_2) \quad \text{for } t_1 \neq t_2. \quad (8)$$

2.2.1. Traffic at Connection Level. At connection level, for instance, for the i th connection, traffic is $x_i(t)$. One particularity of $x_i(t)$ is that t for $x_i(t)$ usually lasts within a finite time interval, say, $[0, T]$. The width of the interval may be

short, such as a short conversation like a word “hello” or long, such as a long speech over a network. In any case, it is finite. Modeling $x_i(t)$ with short interval is particularly desired and challenging.

Note 3. In the discrete case, the length of $x_i(t)$ may be too short to the proper statistical analysis of $x_i(t)$ in practice.

Note 4. Without confusions, we use $[0, T]$ to represent the interval in both the continuous case and the discrete one. In the continuous case, $[0, T] \in \mathbf{R}$. In the discrete case, $[0, T] \in \mathbf{Z}$, where \mathbf{Z} is the set of integer numbers, implying $t = 0, 1, \dots, T$. We use $[t_1, t_2]$ to represent an interval the starting point of which is nonzero.

2.2.2. Aggregated Traffic. We adopt Figure 1 to discuss aggregated traffic. At time t , aggregated traffic denoted by $x(t)$ at a node is expressed by

$$x(t) = \sum x_i(t), \quad i = 1, \dots, m. \quad (9)$$

In contrary to $x_i(t)$, the particularity of $x(t)$ is that t for $x(t)$ usually lasts within an interval longer than that of $x_i(t)$. As a matter of fact, if $x_i(t)$ passes through a node, another arrival flow $x_j(t)$ ($j = 1, \dots, m$) may arrive at the node. Consequently, in general, we should consider $t \in (0, \infty)$ for $x(t)$.

2.3. Accumulated Traffic. Traffic, either $x_i(t)$ or $x(t)$, discussed previously is instantaneous one. Data modeling of instantaneous traffic is essential, as we need understanding what its behaviors are at instantaneous time t at the input of a node. However, from the point of view of the service of a node, we also need data modeling of accumulated traffic within a time interval, say, $[0, T]$, without loss of generality, because it is desired for us to understand what the service performance of the node is for the purpose of proper design of a buffer size as well as scheduling policy of the node.

2.3.1. Accumulated Traffic at Connection Level. In the continuous case, the accumulated traffic of $x_i(t)$ within the interval $[0, T]$ is denoted by $X_i(T)$. It is given by

$$X_i(T) = \int_0^T x_i(t) dt, \quad t \in \mathbf{R}. \quad (10)$$

In the discrete case,

$$X_i(T) = \sum_{t=0}^{T-1} x_i(t), \quad t \in \mathbf{Z}. \quad (11)$$

2.3.2. Accumulated Traffic in the Aggregated Case. Denote by $X(T)$ the accumulated traffic in the aggregated case within the interval $[0, T]$. Then, in the continuous case, we have

$$X(T) = \int_0^T x(t)dt, \quad t \in \mathbf{R}. \quad (12)$$

In the discrete case,

$$X(T) = \sum_{t=0}^{T-1} x(t), \quad t \in \mathbf{Z}. \quad (13)$$

The mathematical expressions of $X(T)$ and $X_i(T)$ appear similar except the subscript i . However, $X(T)$ differs from $X_i(T)$ substantially in analysis in methodology. On the one hand, T for $X_i(T)$ should be assumed to be short such that conventional methods in statistics fail to its statistical analysis. On the other hand, T for $X(T)$ may be large enough such that it may be sectioned for the statistical analysis, see, for example, Li et al. [104].

2.3.3. A Basic Property of Accumulated Traffic. One property of accumulated traffic, either $X(T)$ or $X_i(T)$, is the wide sense increasing. By wide sense increasing, we mean that

$$X_i(T_1) \leq X_i(T_2) \quad \text{for } T_1 \leq T_2, \quad (14)$$

or

$$X(T_1) \leq X(T_2) \quad \text{for } T_1 \leq T_2. \quad (15)$$

Therefore, the data functions or series we face with are increasing ones in the wide sense.

2.3.4. (σ, ρ) Model of Data. For $\sigma \geq 0$ and $\rho \geq 0$, the following is called the (σ, ρ) model of data $x_i(t)$,

$$X_i(T) = \int_0^T x_i(u)du \leq \sigma_i + \rho_i T. \quad (16)$$

Note 5. The model expressed by (16) is irrelevant of any information of statistics of $x_i(t)$. The advantage of this model is at the cost of using inequality instead of equality.

Note 6. The model of (16) is simple in computation. Thus, it may be effective in practice, particularly in environments of CPNS, where simple computations are always expected.

For accumulated traffic $X(T)$, we have

$$X(T) = \int_0^T x(u)du \leq \sigma + \rho T. \quad (17)$$

Due to sufficiently large T , we may set the starting time by T_0 . In this case, we have

$$\int_{T_0}^T x(u)du \leq \sigma(T_0) + \rho(T - T_0). \quad (18)$$

Moreover, we are allowed to section the above integral such that

$$\int_{nT}^{(n+1)T} x(u)du \leq \sigma(nT) + \rho(T), \quad n = 0, 1, \dots \quad (19)$$

Without loss of generality, we use (17) to explain σ and ρ .

Remark 5. The parameter σ represents the bound of the burstness or local irregularity of $x(t)$, because

$$0 \leq \lim_{T \rightarrow 0} \int_0^T x(u)du \leq \sigma. \quad (20)$$

Note that the above integral does not make sense if $\lim_{T \rightarrow 0} \int_0^T x(t)dt \neq 0$ for the continuous $x(t)$ even in the field of the Lebesgue's integrals, see Dudley [105], Bartle and Sherbert [106], and Trench [107] for the contents of the Lebesgue's integrals. However, it makes sense when it is considered in the domain of generalized functions, which we shall brief in the following section. A simple way to explain (20) is

$$\lim_{T \rightarrow 0} \int_0^T x(t)dt = \int_0^T \sigma_1 \delta(t)dt, \quad (21)$$

where $\sigma_1 \leq \sigma$ and $\delta(t)$ is the Dirac- δ function.

Remark 6. The parameter ρ represents the bound of the average rate of $X(T)$, because

$$0 \leq \lim_{T \rightarrow \infty} \frac{\int_0^T x(t)dt}{T} \leq \rho = \text{constant}. \quad (22)$$

Remark 7. The parameter σ measures the local property of $x(t)$ while ρ is a measure of global property of $x(t)$.

3. Min-Plus Algebra and Problem Statement

Min-plus convolution is essential in the min-plus algebra. In this section, we first briefly review the conventional convolution in linear systems. Then, we shall visit min-plus convolution. Finally, we shall state the problem in the aspect of identity in the min-plus algebra.

3.1. Conventional Convolution. Denote by p a real number that satisfies $1 \leq p < \infty$. If a function $f(t)$ defined on $[a, b]$, where a is allowed to be $-\infty$ and b is allowed to be ∞ , is measurable and

$$\int_a^b |f(u)|^p du < \infty, \quad (23)$$

we say that $f(t) \in L^p(a, b)$.

Suppose that two functions $f_1(t), f_2(t) \in L^1(-\infty, \infty)$. Then, one says that $f_1(t)$ convolutes $f_2(t)$ if

$$f_1(t) * f_2(t) = \int_{-\infty}^{\infty} f_1(u)f_2(t-u)du, \quad (24)$$

where $*$ is the symbol implying the operation of convolution. We call it conventional convolution so as to distinguish

it from the min-plus convolution we are discussing in this paper.

The conventional convolution is crucial for linear systems, see, for example, Gibson [54], Box et al. [108], Mitra and Kaiser [109], Papoulis [110], Harris [111], Mikusinski [112], Fuller [113], and Bendat and Piersol [114], just naming a few. It has the properties described by the following lemmas.

Lemma 1. *In the algebra system $(L^1; *)$, the conventional convolution is commutative.*

Lemma 2 (closure of $*$). *If $f_1(t), f_2(t) \in L^1$, then $f_1(t) * f_2(t) \in L^1$.*

Lemma 3. *In the algebra system $(L^1; +, *)$, where $+$ implies the ordinary addition, $*$ with respect to $+$ is distributive.*

Lemma 4. *For $a \in \mathbf{R}$, $[af_1(t)] * f_2(t) = f_1(t) * [af_2(t)] = a[f_1(t) * f_2(t)]$.*

Lemma 5. *The identity in $(L^1; *)$ is the Dirac- δ function $\delta(t)$ that is defined by*

$$f(t) = \int_{-\infty}^{\infty} f(u)\delta(t-u)du, \quad (25)$$

where $f(t) \in L^1(-\infty, \infty)$ is continuous at t .

In fact, in the domain of generalized functions, we have

$$\int_{-\infty}^{\infty} \delta(u)du < \infty. \quad (26)$$

Thus, $\delta(t) \in L^1(-\infty, \infty)$ in the sense of generalize functions. Consequently, $\delta(t)$ is taken as the asymptotic identity in $(L^1; *)$ in the domain of generalized functions. Accordingly, the inverse of the conventional convolution discussed by, for instance, Mikusinski [112], Bracewell [115], Huang and Qiu [116], Abutaleb et al. [117], Rhoads and Ekstrom [118], Todoschuck and Jensen [119], and Moreau et al. [120], exists because the necessary and sufficient condition that the inverse of an operation exists is that there exists the identity in that system, see, for example, Korn and Korn [121], Zhang [122], Riley et al. [123], Bronshtein et al. [124], and Stillwell [125], but it should be in the sense of generalized functions. As a matter of fact, the conventional convolution itself is in that sense, see, for example, Smith [126].

Theorem 1. *The algebra system $(L^1; *)$ is a group.*

Proof. First, the operation $*$ is closed in L^1 . Second, $*$ is commutative because, for any $f_1(t), f_2(t), f_3(t) \in L^1(-\infty, \infty)$,

$$f_1(t) * [f_2(t) * f_3(t)] = [f_1(t) * f_2(t)] * f_3(t). \quad (27)$$

Finally, there exists the left identity denoted by $\delta(t)$ and the right one again denoted by $\delta(t)$ in $(L^1; *)$ such that

$$f(t) * \delta(t) = \delta(t) * f(t) \quad \text{for any } f(t) \in L^1(-\infty, \infty). \quad (28)$$

Thus, $(L^1; *)$ is a group. \square

3.2. Min-Plus Convolution. Considering the property of wide sense increasing of accumulated traffic mentioned in Section 2.3, we denote by \mathbb{S} the set that contains all functions that are greater than or equal to zero and that are wide sense increasing.

Definition 1. Let $X_1(t), X_2(t) \in \mathbb{S}$. Then, the following operation is called min-plus convolution:

$$X_1(t) \otimes X_2(t) = \inf_{0 \leq u \leq t} \{X_1(u) + X_2(t-u)\}, \quad (29)$$

where \otimes represents the operation of the min-plus convolution.

Example 1. Let $X(t) = t^2$ for $t > 0$ and 0 elsewhere. Then, $X(t) \otimes X(t) = t^2/2$.

Lemma 6 (closure of \otimes). *Let $X_1(t), X_2(t) \in \mathbb{S}$. Then, $X_1(t) \otimes X_2(t) \in \mathbb{S}$.*

Lemma 7. *The operation \otimes is commutative. That is,*

$$X_1(t) \otimes X_2(t) = X_2(t) \otimes X_1(t) \quad \text{for } X_1(t), X_2(t) \in \mathbb{S}. \quad (30)$$

Define another operation that is denoted by \wedge such that

$$X_1(t) \wedge X_2(t) = \inf[X_1(t), X_2(t)] \quad \text{for } X_1(t), X_2(t) \in \mathbb{S}. \quad (31)$$

Then, we have an algebra system denoted by $(\mathbb{S}, \wedge, \otimes)$ that follows the distributive law.

Lemma 8. *The operation \otimes with respect to \wedge is distributive. That is, for $X_1(t), X_2(t), X_3(t) \in \mathbb{S}$, one has*

$$[X_1(t) \wedge X_2(t)] \otimes X_3(t) = [X_1(t) \otimes X_3(t)] \wedge [X_2(t) \otimes X_3(t)]. \quad (32)$$

The following rule useful in this research is stated as follows.

Lemma 9. *Suppose $K \in \mathbf{R}$. Then, for $X_1(t), X_2(t) \in \mathbb{S}$, one has*

$$[X_1(t) + K] \otimes X_2(t) = X_1(t) \otimes X_2(t) + K, \quad (33)$$

where $+$ is the ordinary addition.

Denote by $I_1(t)$ the conventional identity in the min-plus algebra, which is defined by

$$I_1(t) = \begin{cases} \infty, & t > 0, \\ 0, & t < 0, \end{cases} \quad (34)$$

see [66–70].

It seems quite obvious when one takes $I_1(t)$ as the identity in the min-plus algebra since

$$X(t) \otimes I_1(t) = I_1(t) \otimes X(t) = X(t). \quad (35)$$

However, we shall soon point the contradictions of $I_1(t)$ below.

3.3. *Problem Statement.* Denote by $u(t)$ the Heavyside unit step function. That is,

$$u(t) = \begin{cases} 1, & t > 0, \\ 0, & t < 0. \end{cases} \quad (36)$$

Then, for $K \in \mathbf{R}$, we have

$$Ku(t) = \begin{cases} K, & t > 0, \\ 0, & t < 0. \end{cases} \quad (37)$$

Using (34), we have

$$\begin{aligned} I_1(t) + Ku(t) &= \begin{cases} \infty + K, & t > 0 \\ 0, & t < 0 \end{cases} \\ &= \begin{cases} \infty, & t > 0 \\ 0, & t < 0 \end{cases} \quad (\text{Contradiction 1}) \\ &= I_1(t). \end{aligned} \quad (38)$$

The above is an obvious contradiction regarding the conventional identity defined by (34).

In addition to the above contradiction, we now state another problem regarding (34). As a matter of fact, if we let $X_1(t) = I_1(t)$ and $Ku(t)$ in Lemma 9, then, on the left side of (33) in Lemma 9, we have

$$[I_1(t) + Ku(t)] \otimes X_2(t) = I_1(t) \otimes X_2(t) = X_2(t). \quad (39)$$

On the other side, on the right side of (33) in Lemma 9, we have

$$\begin{aligned} [I_1(t) + Ku(t)] \otimes X_2(t) &= I_1(t) \otimes X_2(t) + Ku(t) \\ &= X_2(t) + Ku(t). \end{aligned} \quad (40)$$

Comparing the right sides of (39) with that of (40) yields another contradiction expressed by

$$X_2(t) = X_2(t) + Ku(t), \quad (\text{Contradiction 2}) \quad (41)$$

The above discussions imply that the definition of the identity of (34) in the min-plus algebra, which is commonly used in literature, see, for example, [66–70], may not be rigorous at least. Therefore, the conventional representation of the identity, that is, (34), may be inappropriate since it may mislead computation results like those in (39) and (40). Consequently, rigorous definition of the identity needs studying.

4. Existence of Identity in Min-Plus Algebra

The problems regarding the definition of the conventional identity, which we stated in Section 3.3, give rise to a question whether or not the identity in the min-plus algebra exists. The answer to this question is rarely seen, to the best of our knowledge. Another question resulted from Section 3.3 is what the rigorous representation of the identity is. We shall provide the answer to the first question in this section. The answer to the second will be explained in the next section.

4.1. *Preliminaries.* We brief some results in generalized functions [127–129] for the purpose of discussing the existence of identity.

Definition 2. Let $\text{supp}(f)$ be the support of a function $f : \mathbf{R} \rightarrow \mathbf{C}$. It implies $\{t : f(t) \neq 0\}$. The function is said to have a bounded support if there exist $a, b \in \mathbf{R}$ such that $\text{supp}(f) \subset [a, b]$.

Definition 3. A function $f : \mathbf{R} \rightarrow \mathbf{C}$ is said to have n time continuous derivatives if its first n derivatives exist and are continuous. If its derivatives of all orders exist and are continuous, f is said to be infinitely differentiable. In this case, f is said to be smooth.

Definition 4. A test function is a smooth $\mathbf{R} \rightarrow \mathbf{C}$ with $\text{supp}(f) \subset [a, b]$. The set of all test functions is denoted by \mathbf{D} .

Definition 5. A linear functional f on \mathbf{D} is a map $f : \mathbf{D} \rightarrow \mathbf{C}$ such that, for $a, b \in \mathbf{C}$ and $\phi, \psi \in \mathbf{D}$, $f(a\phi + b\psi) = af(\phi) + bf(\psi)$.

Definition 6. Denote by (ϕ_n) a sequence of test functions and Φ another test function. We say that $\phi_n \rightarrow \Phi$ if the following holds:

- (1) there is an interval $[a, b]$ that contains $\text{supp}(\Phi)$ and $\text{supp}(\phi_n)$ for all n ,
- (2) $\lim_{n \rightarrow \infty} \phi_n^{(k)}(t) \rightarrow \Phi^{(k)}(t)$ uniformly for $t \in [a, b]$.

Definition 7. A functional f on \mathbf{D} is continuous if it maps every convergent sequence in \mathbf{D} into a convergent sequence in \mathbf{C} . A continuous linear functional f on \mathbf{D} is termed a generalized function. It is often called a distribution in the sense of Schwartz.

Definition 8. A function $f : \mathbf{R} \rightarrow \mathbf{C}$ is locally integrable if $\int_a^b f(t)dt < \infty$ for all a, b .

Lemma 10. Any continuous, including piecewise continuous, function is locally integrable.

Lemma 11 (regular). Any locally integrable function f is a generalized function defined by

$$\langle f, \phi \rangle = \int_{-\infty}^{\infty} f(t)\phi(t)dt < \infty. \quad (42)$$

In this case, f is called regular.

Lemma 12. Any generalized function has derivatives of all orders.

Lemma 13. There exists the Fourier transform of any generalized function.

Definition 9 (rapid function). A function of rapid decay is a smooth function $\phi : \mathbf{R} \rightarrow \mathbf{C}$ such that $t^n \phi^{(r)}(t) \rightarrow 0$ as $t \rightarrow \pm \infty$ for all $n, r \geq 0$, where \mathbf{C} is the space of complex numbers. The set of all functions of rapid decay is denoted by \mathbf{S} .

Lemma 14. *Every function belonging to \mathbb{S} is absolutely integrable.*

4.2. *Proof of Existence.* Define the norm and inner product of $X \in \mathbb{S}$ by

$$\|X\|^2 = \langle X, X \rangle = \int_0^\infty X^2(u)w(u)du, \quad (43)$$

where $w \in \mathbb{S}$. Combining any $X \in \mathbb{S}$ with its limit yields a Hilbert space that we denote again by \mathbb{S} without confusions.

Let $g \in \mathbb{S}$ be a system function such that it transforms its input $X \in \mathbb{S}$ to the output by

$$y = (X \otimes g) \in \mathbb{S}. \quad (44)$$

Denote the system by the operator L . Then, we purposely force the functionality of L such that it maps an element $X \in \mathbb{S}$ to another element $(X \otimes g) \in \mathbb{S}$. Note that L is a linear operator. In fact, according to Lemma 8, we have

$$L(X \wedge g) = L(X) \wedge L(g). \quad (45)$$

In addition, from Lemma 9, we have

$$L(X + K) = L(X) + K. \quad (46)$$

Therefore, L is a linear mapping from \mathbb{S} to \mathbb{S} .

Denote by \mathbf{L} the space consisting of all such operators by

$$\mathbf{L}(\mathbb{S}, \mathbb{S}) = \mathbf{L}(\mathbb{S}). \quad (47)$$

Then, from Lemmas 8 and 9, one can easily see that $\mathbf{L}(\mathbb{S})$ is a linear space.

Lemma 15 (archimedes criterion). *For any positive real numbers $a > 0$ and $b > 0$, there exists positive integer $n \in \mathbf{Z}$ such that $na > b$ (see [130]).*

Lemma 16 (archimedes). *If $b \in \mathbf{R}$, there exists $n \in \mathbf{Z}$ such that $b < n$ (see [106]).*

Lemma 17. *An operator $T : X \mapsto Y$ is invertible if and only if there exists constant $m > 0$ such that for all $x \in X$, $\|Tx\| \geq m\|x\|$, where X and Y are linear normed spaces (see [131]).*

From the above discussions, we obtain the following theorem.

Theorem 2 (existence). *For $X, g \in \mathbb{S}$ and $X(0) \neq 0$ and $g(0) \neq 0$, if $L(X) = X \otimes g$ or $L_1(g) = g \otimes X$, then both L and L_1 are invertible. Consequently, the identity in the min-plus algebra exists.*

Proof. Consider

$$\begin{aligned} \|LX\| &= \sqrt{\|X \otimes g\|} \\ &= \sqrt{\int_0^\infty \left[\inf_{0 \leq u \leq t} \{X(u) + g(t-u)\} \right]^2 w(u)du}. \end{aligned} \quad (48)$$

Since

$$\inf_{0 \leq u \leq t} \{X(u) + g(t-u)\} \geq \inf \{X(u)\} = X(0) \quad (49)$$

and $X(u) \in \mathbb{S}$, we have

$$0 < X(0) \leq X(u). \quad (50)$$

According to Lemmas 15 and 16, there exists $m > 0$ such that

$$X(0) \geq m^2 X(u). \quad (51)$$

Therefore,

$$\begin{aligned} \|LX\| &\geq \sqrt{\int_0^\infty [\inf \{X(u)\}]^2 w(u)du} \\ &= \sqrt{\int_0^\infty [X(0)]^2 w(u)du} \\ &\geq m \sqrt{\int_0^\infty X(u)^2 w(u)du} = m\|X\|. \end{aligned} \quad (52)$$

Similarly, if $L_1 \in \mathbf{L}(\mathbb{S})$ is such that $L_1(g) = g \otimes X$, we have $\|L_1 g\| \geq m_1 \|g\|$ since $g(0) \neq 0$, where $m_1 > 0$ is a constant. Thus, according to Lemma 17, Theorem 2 holds. \square

Note 7. In Theorem 2, we need the conditions of $X(0) \neq 0$ and $g(0) \neq 0$. Since $X(t)$ and $g(t)$ are wide sense increasing, we need in fact $X(0) > 0$ and $g(0) > 0$.

5. Representation of Identity in Min-Plus Algebra

Express the Dirac- δ function by

$$\delta(t) = \frac{1}{2\pi} + \frac{1}{\pi} \sum_{k=-\infty}^{\infty} \cos(kt). \quad (53)$$

For the purpose of distinguishing the identity we present from the conventional one, we denote $I(t)$ as the identity in what follows instead of $I_1(t)$ as used in Section 3.

Theorem 3 (representation). *The identity in the min-plus algebra is expressed by*

$$I(t) = \lim_{T \rightarrow 0} \left[\frac{2}{T} + \frac{4}{T} \sum_{n=1}^{\infty} \cos\left(\frac{2n\pi t}{T}\right) \right]. \quad (54)$$

Proof. Take the following into account

$$\sum_{n=0}^{\infty} \delta(t - nT) \quad (T > 0). \quad (55)$$

Then, the identity in the discrete case is given by

$$I(k) = \sum_{n=0}^{\infty} \delta(k - nT). \quad (56)$$




FIGURE 3: Single node with arrival and departure traffic.

FIGURE 4: N tandem nodes with arrival and departure traffic.

The identity in the continuous case is taken as the limit expressed by

$$I(t) = \lim_{T \rightarrow 0} \sum_{n=0}^{\infty} \delta(t - nT). \quad (57)$$

Considering the Poisson's summation formula, we have

$$I(k) = \frac{2}{T} + \frac{4}{T} \sum_{n=1}^{\infty} \cos\left(\frac{2n\pi k}{T}\right). \quad (58)$$

In the limit case,

$$I(t) = \lim_{T \rightarrow 0} \left[\frac{2}{T} + \frac{4}{T} \sum_{n=1}^{\infty} \cos\left(\frac{2n\pi t}{T}\right) \right]. \quad (59)$$

This completes the proof. \square

Remark 8. If one uses the representation in Theorem 3, the contradictions given in (38) and (41) vanish.

Note 8. The identity expressed by (59) is an asymptotic one.

6. Discussions

We mention an application of min-plus algebra to CPNS. Denote by $Y_i(t)$ the accumulated function characterizing the output of the i th node (Figure 3). Then, the min-plus convolution can be used to establish the relationship between $X_i(t)$, $S_i(t)$, and $Y_i(t)$ by

$$Y_i(t) \geq X_i(t) \otimes S_i(t) = \inf_{0 \leq u \leq t} \{S_i(u) + X_i(t - u)\}. \quad (60)$$

Suppose a traffic function passes through N tandem nodes from the first node with the service curve $S_1(t)$ to the N th node with the service curve $S_N(t)$ to reach the destination as indicated in Figure 4. Denote the departure traffic of the N th node by $Y_N(t)$. Then,

$$Y_N(t) \geq X_1(t) \otimes S_N^1(t) = \inf_{0 \leq u \leq t} \{S_N^1(u) + X_1(t - u)\}, \quad (61)$$

where (see [132])

$$S_N^1(t) = S_1(t) \otimes S_2(t) \otimes \dots \otimes S_i(t) \dots \otimes S_N(t). \quad (62)$$

Note 9. Min-plus algebra can be used to linearize a nonlinear system as can be seen from (62). Thus, it may yet be used as a theory in the aspect of data transmission systems in CPNS.

7. Conclusions

We have proposed the problem regarding the conventional identity in the min-plus algebra. In addition, we have presented the proof that the identity in the min-plus algebra exists in the domain of generalized function. Moreover, we have given the asymptotic expression of the identity in the system of min-plus algebra.

Acknowledgment

This work was partly supported by the National Natural Science Foundation of China (NSFC) under the project Grant nos. 60873264, 61070214, and the 973 plan under the Project no. 2011CB302801/2011CB302802.

References

- [1] Z. Song, Y. Q. Chen, C. R. Sastry, and N. C. Tas, *Optimal Observation for Cyber-Physical Systems*, Springer, 2009.
- [2] Commission of the European Communities, "Internet of things—an action plan for Europe," *Journal of International Wildlife Law and Policy*, vol. 12, no. 1-2, pp. 108–120, 2009.
- [3] A. Ferscha, M. Hechinger, A. Riener et al., "Peer-it: stick-on solutions for networks of things," *Pervasive and Mobile Computing*, vol. 43, no. 3, pp. 448–479, 2008.
- [4] G. D. Clifton, H. Byer, K. Heaton, D. J. Haberman, and H. Gill, "Provision of pharmacy services to underserved populations via remote dispensing and two-way videoconferencing," *American Journal of Health-System Pharmacy*, vol. 60, no. 24, pp. 2577–2282, 2003.
- [5] K. Traynor, "Navy takes telepharmacy worldwide," *American Journal of Health-System Pharmacy*, vol. 67, no. 14, pp. 1134–1136, 2010.
- [6] K. T. Chang, *Introduction to Geographical Information Systems*, McGraw-Hill, New York, NY, USA, 2008.
- [7] M. F. Goodchild, "Twenty years of progress: GIScience in 2010," *Journal of Spatial Information Science*, vol. 2010, no. 1, pp. 3–20, 2010.
- [8] T. L. Lai and H. Xing, *Statistical Models and Methods for Financial Markets*, Springer, 2008.
- [9] B. B. Mandelbrot, *Gaussian Self-Affinity and Fractals*, Springer, 2001.
- [10] B. B. Mandelbrot, *Multifractals and 1/f Noise*, Springer, 1998.
- [11] B. B. Mandelbrot, *The Fractal Geometry of Nature*, W. H. Freeman, New York, NY, USA, 1982.
- [12] D. Hainaut and P. Devolder, "Mortality modelling with Lévy processes," *Insurance*, vol. 42, no. 1, pp. 409–418, 2008.
- [13] C. Cattani, "Harmonic wavelet approximation of random, fractal and high frequency signals," *Telecommunication Systems*, vol. 43, no. 3-4, pp. 207–217, 2010.
- [14] C. Cattani, "Fractals and hidden symmetries in DNA," *Mathematical Problems in Engineering*, vol. 2010, Article ID 507056, 31 pages, 2010.
- [15] G. Mattioli, M. Scalia, and C. Cattani, "Analysis of large-amplitude pulses in short time intervals: application to neuron interactions," *Mathematical Problems in Engineering*, vol. 2010, Article ID 895785, 15 pages, 2010.

- [16] C. Cattani, "Shannon wavelets for the solution of integrodifferential equations," *Mathematical Problems in Engineering*, vol. 2010, 22 pages, 2010.
- [17] C. Cattani, "Shannon wavelets theory," *Mathematical Problems in Engineering*, vol. 2008, Article ID 164808, 24 pages, 2008.
- [18] S. Y. Chen, Y. F. Li, and J. Zhang, "Vision processing for realtime 3-D data acquisition based on coded structured light," *IEEE Transactions on Image Processing*, vol. 17, no. 2, pp. 167–176, 2008.
- [19] S. Y. Chen, H. Tong, Z. Wang, S. Liu, M. Li, and B. Zhang, "Improved generalized belief propagation for vision processing," *Mathematical Problems in Engineering*, vol. 2011, Article ID 416963, 12 pages, 2011.
- [20] S. Y. Chen and Y. F. Li, "Determination of stripe edge blurring for depth sensing," *IEEE Sensors Journal*, vol. 11, no. 2, pp. 389–390, 2011.
- [21] S. Y. Chen and Q. Guan, "Parametric shape representation by a deformable NURBS model for cardiac functional measurements," *IEEE Transactions on Biomedical Engineering*, vol. 58, no. 3, pp. 480–487, 2011.
- [22] S. Chen, J. Zhang, H. Zhang et al., "Myocardial motion analysis for determination of tei-index of human heart," *Sensors*, vol. 10, no. 12, pp. 11428–11439, 2010.
- [23] W. Mikhael and T. Yang, "A gradient-based optimum block adaptation ICA technique for interference suppression in highly dynamic communication channels," *EURASIP Journal on Advances in Signal Processing*, vol. 2006, Article ID 84057, 10 pages, 2006.
- [24] E. G. Bakhoun and C. Toma, "Dynamical aspects of macroscopic and quantum transitions due to coherence function and time series events," *Mathematical Problems in Engineering*, vol. 2010, Article ID 428903, 13 pages, 2010.
- [25] E. G. Bakhoun and C. Toma, "Mathematical transform of traveling-wave equations and phase aspects of quantum interaction," *Mathematical Problems in Engineering*, vol. 2010, Article ID 695208, 15 pages, 2010.
- [26] E. G. Bakhoun and C. Toma, "Relativistic short range phenomena and space-time aspects of pulse measurements," *Mathematical Problems in Engineering*, vol. 2008, Article ID 410156, 20 pages, 2008.
- [27] M. Li, "Change trend of averaged Hurst parameter of traffic under DDOS flood attacks," *Computers and Security*, vol. 25, no. 3, pp. 213–220, 2006.
- [28] M. Li, "A class of negatively fractal dimensional gaussian random functions," *Mathematical Problems in Engineering*, vol. 2011, Article ID 291028, 18 pages, 2011.
- [29] M. Li, "Generation of teletraffic of generalized cauchy type," *Physica Scripta*, vol. 81, no. 2, Article ID 025007, 2010.
- [30] M. Li, "Fractal time series-a tutorial review," *Mathematical Problems in Engineering*, vol. 2010, Article ID 157264, 26 pages, 2010.
- [31] M. Li, "A method for requiring block size for spectrum measurement of ocean surface waves," *IEEE Transactions on Instrumentation and Measurement*, vol. 55, no. 6, pp. 2207–2215, 2006.
- [32] M. Li, "Modeling autocorrelation functions of long-range dependent teletraffic series based on optimal approximation in hilbert space-a further study," *Applied Mathematical Modelling*, vol. 31, no. 3, pp. 625–631, 2007.
- [33] M. Li, W. Zhao, and S. Chen, "MBm-based scalings of traffic propagated in internet," *Mathematical Problems in Engineering*, vol. 2011, Article ID 389803, 21 pages, 2011.
- [34] M. Li and W. Zhao, "Variance bound of ACF estimation of one block of fGn with LRD," *Mathematical Problems in Engineering*, vol. 2010, Article ID 560429, 14 pages, 2010.
- [35] M. Li and W. Zhao, "Detection of variations of local irregularity of traffic under DDOS flood attack," *Mathematical Problems in Engineering*, vol. 2008, Article ID 475878, 11 pages, 2008.
- [36] M. Li, C. Cattani, and S.-Y. Chen, "Viewing sea level by a one-dimensional random function with long memory," *Mathematical Problems in Engineering*, vol. 2011, Article ID 654284, 13 pages, 2011.
- [37] M. Li and J. Y. Li, "On the predictability of long-range dependent series," *Mathematical Problems in Engineering*, vol. 2010, Article ID 397454, 9 pages, 2010.
- [38] M. Li and S. C. Lim, "Modeling network traffic using generalized cauchy process," *Physica A*, vol. 387, no. 11, pp. 2584–2594, 2008.
- [39] M. Li, S. C. Lim, and Sy. Chen, "Exact solution of impulse response to a class of fractional oscillators and its stability," *Mathematical Problems in Engineering*, vol. 2011, Article ID 657839, 9 pages, 2011.
- [40] A. R. Messina, P. Esquivel, and F. Lezama, "Time-dependent statistical analysis of wide-area time-synchronized data," *Mathematical Problems in Engineering*, vol. 2010, Article ID 751659, 17 pages, 2010.
- [41] M. Humi, "Assessing local turbulence strength from a time series," *Mathematical Problems in Engineering*, vol. 2010, Article ID 316841, 13 pages, 2010.
- [42] M. Dong, "A tutorial on nonlinear time-series data mining in engineering asset health and reliability prediction: concepts, models, and algorithms," *Mathematical Problems in Engineering*, vol. 2010, Article ID 175936, 22 pages, 2010.
- [43] Z. Liu, "Chaotic time series analysis," *Mathematical Problems in Engineering*, vol. 2010, Article ID 720190, 31 pages, 2010.
- [44] G. Toma, "Specific differential equations for generating pulse sequences," *Mathematical Problems in Engineering*, vol. 2010, Article ID 324818, 11 pages, 2010.
- [45] O. M. Abuzeid, A. N. Al-Rabadi, and H. S. Alkhalidi, "Fractal geometry-based hypergeometric time series solution to the hereditary thermal creep model for the contact of rough surfaces using the Kelvin-Voigt medium," *Mathematical Problems in Engineering*, vol. 2010, Article ID 652306, 22 pages, 2010.
- [46] M. Li and W. Zhao, "Visiting power laws in cyber-physical networking systems," *Mathematical Problems in Engineering*, vol. 2012, Article ID 302786, p. 13, 2012.
- [47] M. Li and S. C. Lim, "Power spectrum of generalized cauchy process," *Telecommunication Systems*, vol. 43, no. 3-4, pp. 219–222, 2010.
- [48] J. He, H. Qian, Y. Zhou, and Z. Li, "Cryptanalysis and improvement of a block cipher based on multiple chaotic systems," *Mathematical Problems in Engineering*, vol. 2010, Article ID 590590, 14 pages, 2010.
- [49] Z. Liao, S. Hu, and W. Chen, "Determining neighborhoods of image pixels automatically for adaptive image denoising using nonlinear time series analysis," *Mathematical Problems in Engineering*, vol. 2010, 2010.
- [50] G. Werner, "Fractals in the nervous system: conceptual implications for theoretical neuroscience," *Frontiers in Fractal Physiology*, vol. 1, p. 28, 2010.
- [51] B. J. West, "Fractal physiology and the fractional calculus: a perspective," *Frontiers in Fractal Physiology*, vol. 1, 2010.
- [52] H. Akimaru and K. Kawashima, *Teletraffic: Theory and Applications*, Springer, 1993.

- [53] W. Yue, H. Takagi, and Y. Takahashi, *Advances in Queueing Theory and Network Applications*, Springer, 2009.
- [54] J. D. Gibson, *The Communications Handbook*, IEEE Press, 1997.
- [55] R. B. Cooper, *Introduction to Queueing Theory*, Elsevier, 2nd edition, 1981.
- [56] J. M. Pitts and J. A. Schormans, *Introduction to ATM Design and Performance: With Applications Analysis Software*, John Wiley & Sons, New York, NY, USA, 2nd edition, 2000.
- [57] D. McDysan, *QoS & Traffic Management in IP & ATM Networks*, McGraw-Hill, New York, NY, USA, 2000.
- [58] W. Stalling, *High-Speed Networks: TCP/IP and ATM Design Principles*, Prentice Hall, 2nd edition, 2002.
- [59] R. L. Cruz, "A calculus for network delay—part 1: network elements in isolation part 2," *IEEE Transactions on Information Theory*, vol. 37, no. 1, pp. 114–141, 1991.
- [60] R. L. Cruz, "A calculus for network delay—part 2: network analysis," *IEEE Transactions on Information Theory*, vol. 37, no. 1, pp. 132–141, 1991.
- [61] R. L. Cruz, "Quality of service guarantees in virtual circuit switched networks," *IEEE Journal on Selected Areas in Communications*, vol. 13, no. 6, pp. 1048–1056, 1995.
- [62] W. Zhao and K. Ramamritham, "Virtual time csma protocols for hard real-time communication," *IEEE Transactions on Software Engineering*, vol. 13, no. 8, pp. 938–952, 1987.
- [63] A. Raha, S. Kamat, and W. Zhao, "Guaranteeing end-to-end deadlines in ATM networks," in *Proceedings of the 15th International Conference on Distributed Computing Systems*, pp. 60–68, June 1995.
- [64] C. S. Chang, "On deterministic traffic regulation and service guarantees: a systematic approach by filtering," *IEEE Transactions on Information Theory*, vol. 44, no. 3, pp. 1097–1110, 1998.
- [65] C. S. Chang, "Stability, queue length, and delay of deterministic and stochastic queueing networks," *IEEE Transactions on Automatic Control*, vol. 39, no. 5, pp. 913–931, 1994.
- [66] J. Y. Le Boudec, "Application of network calculus to guaranteed service networks," *IEEE Transactions on Information Theory*, vol. 44, no. 3, pp. 1087–1096, 1998.
- [67] J. Y. Le Boudec and T. Patrick, *Network Calculus, A Theory of Deterministic Queueing Systems for the Internet*, Springer, 2001.
- [68] V. Firoiu, J. Y. Le Boudec, D. Towsley, and Z. L. I. Zhang, "Theories and models for internet quality of service," *Proceedings of IEEE*, vol. 90, no. 9, pp. 1565–1591, 2002.
- [69] R. Agrawal, F. Baccelli, and R. Rajan, *An Algebra for Queueing Networks with Time Varying Service and Its Application to the Analysis of Integrated Service Networks*, INRIA, 1998, RR-3435.
- [70] Y. M. Jiang and Y. Liu, *Stochastic Network Calculus*, Springer, 2008.
- [71] C. S. Chang, *Performance Guarantees in Communication Networks*, Springer, 2000.
- [72] H. Wang, J. B. Schmitt, and I. Martinovic, "Dynamic demultiplexing in network calculus—theory and application," *Performance Evaluation*, vol. 68, no. 2, pp. 201–219, 2011.
- [73] S. Q. Wang, D. Xuan, R. Bettati, and W. Zhao, "Toward statistical QoS guarantees in a differentiated services network," *Telecommunication Systems*, vol. 43, no. 3-4, pp. 253–263, 2010.
- [74] C. Z. Li and W. Zhao, "Stochastic performance analysis of non-feedforward networks," *Telecommunication Systems*, vol. 43, no. 3-4, pp. 237–252, 2010.
- [75] M. Li and W. Zhao, "Representation of a stochastic traffic bound," *IEEE Transactions on Parallel and Distributed Systems*, vol. 21, no. 9, pp. 1368–1372, 2010.
- [76] M. Fidler, "Survey of deterministic and stochastic service curve models in the network calculus," *IEEE Communications Surveys and Tutorials*, vol. 12, no. 1, pp. 59–86, 2010.
- [77] Y. M. Jiang, "Per-domain packet scale rate guarantee for expedited forwarding," *IEEE/ACM Transactions on Networking*, vol. 14, no. 3, pp. 630–643, 2006.
- [78] Y. M. Jiang, Q. Yin, Y. Liu, and S. Jiang, "Fundamental calculus on generalized stochastically bounded bursty traffic for communication networks," *Computer Networks*, vol. 53, no. 12, pp. 2011–2019, 2009.
- [79] Y. Liu, C. K. Tham, and Y. M. Jiang, "A calculus for stochastic QoS analysis," *Performance Evaluation*, vol. 64, no. 6, pp. 547–572, 2007.
- [80] C. Li, A. Burchard, and J. Liebeherr, "A network calculus with effective bandwidth," *IEEE/ACM Transactions on Networking*, vol. 15, no. 6, pp. 1442–1453, 2007.
- [81] C. Li and E. Knightly, "Schedulability criterion and performance analysis of coordinated schedulers," *IEEE/ACM Transactions on Networking*, vol. 13, no. 2, pp. 276–287, 2005.
- [82] A. Burchard, J. Liebeherr, and S. D. Patek, "A min-plus calculus for end-to-end statistical service guarantees," *IEEE Transactions on Information Theory*, vol. 52, no. 9, pp. 4105–4114, 2006.
- [83] J. K. Y. Ng, S. Song, and W. Zhao, "Integrated end-to-end delay analysis for regulated ATM networks," *Real-Time Systems*, vol. 25, no. 1, pp. 93–124, 2003.
- [84] A. Raha, S. Kamat, X. Jia, and W. Zhao, "Using traffic regulation to meet end-to-end deadlines in ATM networks," *IEEE Transactions on Computers*, vol. 48, no. 9, pp. 917–935, 1999.
- [85] A. Raha, W. Zhao, S. Kamat, and W. Jia, "Admission control for hard real-time connections in ATM LANs," *IEEE Proceedings*, vol. 148, no. 4, pp. 1–12, 2001.
- [86] D. Starobinski and M. Sidi, "Stochastically bounded burstiness for communication networks," *IEEE Transactions on Information Theory*, vol. 46, no. 1, pp. 206–212, 2000.
- [87] H. Fukś, A. T. Lawnczak, and S. Volkov, "Packet delay in models of data networks," *ACM Transactions on Modeling and Computer Simulation*, vol. 11, no. 3, pp. 233–250, 2001.
- [88] X. Jia, W. Zhao, and J. Li, "An integrated routing and admission control mechanism for real-time multicast connections in ATM networks," *IEEE Transactions on Communications*, vol. 49, no. 9, pp. 1515–1519, 2001.
- [89] S. J. Golestani, "Network delay analysis of a class of fair queueing algorithms," *IEEE Journal on Selected Areas in Communications*, vol. 13, no. 6, pp. 1057–1070, 1995.
- [90] L. Lenzi, E. Mingozzi, and G. Stea, "A methodology for computing end-to-end delay bounds in FIFO-multiplexing tandems," *Performance Evaluation*, vol. 65, no. 11-12, pp. 922–943, 2008.
- [91] INFOS D.4 Networked Enterprise & RFID INFOS G.2 Micro & Nanosystems, in co-operation with the Working Group RFID of the ETP EPOSS, Internet of Things in 2020, Roadmap for the Future[R]. Version 1.1, May 2008.
- [92] E. Ilie-Zudor, Z. Kemény, F. van Blommestein, L. Monostori, and A. van Der Meulen, "A survey of applications and requirements of unique identification systems and RFID techniques," *Computers in Industry*, vol. 62, no. 3, pp. 227–252, 2011.
- [93] S. Ahuja and P. Potti, "An introduction to RFID technology," *Communications and Network*, vol. 2, no. 3, pp. 183–186, 2010.

- [94] R. Wang, L. Zhang, R. Sun, J. Gong, and L. Cui, "A pervasive traffic information acquisition system based on wireless sensor networks," *IEEE Transactions on Intelligent Transportation Systems*, vol. 12, no. 2, pp. 615–621, 2011.
- [95] W. H. K. Lam, S. C. Wong, and H. K. Lo, "Emerging theories in traffic and transportation; and emerging methods for transportation planning and operations," *Transportation Research Part C*, vol. 19, no. 2, pp. 169–171, 2011.
- [96] L. N. Emmanuel, R. T. Antonio, and L. M. Ernesto, "A modeling framework for urban traffic systems microscopic simulation," *Simulation Modelling Practice and Theory*, vol. 18, no. 8, pp. 1145–1161, 2010.
- [97] J. A. Laval, "Hysteresis in traffic flow revisited: an improved measurement method," *Transportation Research Part B*, vol. 45, no. 2, pp. 385–391, 2011.
- [98] B. G. Heydecker and J. D. Addison, "Analysis and modelling of traffic flow under variable speed limits," *Transportation Research Part C*, vol. 19, no. 2, pp. 206–217, 2011.
- [99] P. Santoro, M. Fernández, M. Fossati, G. Cazes, R. Terra, and I. Piedra-Cueva, "Pre-operational forecasting of sea level height for the Río de la Plata," *Applied Mathematical Modelling*, vol. 35, no. 5, pp. 2264–2278, 2011.
- [100] D. She and X. Yang, "A new adaptive local linear prediction method and its application in hydrological time series," *Mathematical Problems in Engineering*, vol. 2010, Article ID 205438, 15 pages, 2010.
- [101] S. V. Muniandy, S. C. Lim, and R. Murugan, "Inhomogeneous scaling behaviors in Malaysian foreign currency exchange rates," *Physica A*, vol. 301, no. 1–4, pp. 407–428, 2001.
- [102] G. Samorodnitsky and M. S. Taqqu, *Stable Non-Gaussian Random Processes, Stochastic Models with Infinite Variance*, Chapman and Hall, New York, NY, USA, 1994.
- [103] D. Starobinski, M. Karpovsky, and L. A. Zakrevski, "Application of network calculus to general topologies using turn-prohibition," *IEEE/ACM Transactions on Networking*, vol. 11, no. 3, pp. 411–421, 2003.
- [104] M. Li, S. Wang, and W. Zhao, "A real-time and reliable approach to detecting traffic variations at abnormally high and low rates," *Springer Lecture Notes in Computer Science*, vol. 4158, pp. 541–550, 2006.
- [105] R. M. Dudley, *Real Analysis and Probability*, Cambridge University Press, 2002.
- [106] R. G. Bartle and D. R. Sherbert, *Introduction to Real Analysis*, John Wiley & Sons, 3rd edition, 2000.
- [107] W. F. Trench, *Introduction to Real Analysis*, Pearson Education, 2003.
- [108] G. E. P. Box, G. M. Jenkins, and G. C. Reinsel, *Time Series Analysis: Forecasting and Control*, Prentice Hall, 1994.
- [109] S. K. Mitra and J. F. Kaiser, *Handbook for Digital Signal Processing*, John Wiley & Sons, 1993.
- [110] A. Papoulis, *The Fourier Integral and Its Applications*, McGraw-Hill, 1962.
- [111] C. M. Harris, *Shock and Vibration Handbook*, McGraw-Hill, 4th edition, 1995.
- [112] J. Mikusinski, *Operational Calculus*, Pergamon Press, 1959.
- [113] W. A. Fuller, *Introduction to Statistical Time Series*, Wiley, 2nd edition, 1996.
- [114] J. S. Bendat and A. G. Piersol, *Random Data: Analysis and Measurement Procedure*, John Wiley & Sons, 3rd edition, 2000.
- [115] R. N. Bracewell, *The Fourier Transform and Its Applications*, McGraw-Hill, New York, NY, USA, 2nd edition, 1978.
- [116] X. Huang and P. H. Qiu, "Blind deconvolution for jump-preserving curve estimation," *Mathematical Problems in Engineering*, vol. 2010, Article ID 350849, 9 pages, 2010.
- [117] A. S. Abutaleb, N. M. Elhamy, and M. E. S. Waheed, "Blind deconvolution of the aortic pressure waveform using the malliavin calculus," *Mathematical Problems in Engineering*, vol. 2010, Article ID 102581, 27 pages, 2010.
- [118] R. L. Rhoads and M. P. Ekstrom, "Removal of interfering system distortion by deconvolution," *IEEE Transactions, Instrumentation and Measurement*, vol. 17, no. 4, pp. 333–337, 1968.
- [119] J. P. Todoeschuck and O. G. Jensen, "Scaling geology and seismic deconvolution," *Pure and Applied Geophysics*, vol. 131, no. 1-2, pp. 273–287, 1989.
- [120] S. Moreau, G. Plantier, J. C. Valière, H. Bailliet, and L. Simon, "Estimation of power spectral density from laser Doppler data via linear interpolation and deconvolution," *Experiments in Fluids*, vol. 50, no. 1, pp. 179–188, 2011.
- [121] G. A. Korn and T. M. Korn, *Mathematical Handbook for Scientists and Engineers*, McGraw-Hill, 1961.
- [122] H. R. Zhang, *Elementary of Modern Algebra*, People's Education Press, China, 1978.
- [123] K. F. Riley, M. P. Hobson, and S. J. Bence, *Mathematical Methods for Physics and Engineering*, Cambridge Press, 2006.
- [124] I. N. Bronshtein, K. A. Semendyayev, G. Musiol, and H. Muehlig, *Handbook of Mathematics*, Springer, 2007.
- [125] J. Stillwell, *Mathematics and Its History*, Springer, 3rd edition, 2010.
- [126] D. C. Smith, "An introduction to distribution theory for signals analysis—part 2. The convolution," *Digital Signal Processing*, vol. 16, no. 4, pp. 419–444, 2006.
- [127] J. Lighthill, *An Introduction to Fourier Analysis and Generalised Functions*, Cambridge University Press, 1958.
- [128] R. P. Kanwal, *Generalized Functions: Theory and Applications*, Birkhauser, 3rd edition, 2004.
- [129] I. M. Gelfand and K. Vilenkin, *Generalized Functions*, vol. 1, Academic Press, New York, NY, USA, 1964.
- [130] A. D. Aleksandro, *Mathematics, Its Essence, Methods and Role*, vol. 3, USSR Academy of Sciences, 1952.
- [131] V. I. Istratescu, *Introduction to Linear Operator Theory*, Marcel Dekker, New York, NY, USA, 1981.
- [132] M. Li and W. Zhao, "Sufficient condition for min-plus deconvolution to be closed in the service-curve set in computer networks," *International Journal of Computers*, vol. 1, no. 3, pp. 163–166, 2007.