



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

***In vivo* study of interferon-alpha-secreting cells in pig foetal lymphohaematopoietic organs following *in utero* TGEV coronavirus injection**

I. Šplíchal ⁽¹⁾ (*), Z. Řeháková ⁽¹⁾, M. Šinkora ⁽¹⁾, J. Šinkora ⁽¹⁾, I. Trebichavský ⁽¹⁾,
H. Laude ⁽²⁾ and B. Charley ⁽²⁾

⁽¹⁾ *Division of Immunology and Gnotobiology, Institute of Microbiology of the Academy
of Sciences of the Czech Republic, 54922 Nový Hrádek (Czech Republic), and*

⁽²⁾ *Laboratoire de Virologie et Immunologie moléculaires, INRA, 78350 Jouy-en-Josas (France)*

SUMMARY

Non-infectious UV-inactivated transmissible gastroenteritis virus (TGEV) was previously shown to induce interferon alpha (IFN α) secretion following *in vitro* incubation with blood mononuclear cells. In this study, pig foetuses at different stages of gestation were injected *in utero* with (a) partially UV-inactivated wild TGEV or (b) fully UV-inactivated wild or dm49-4 mutant TGEV coronavirus. Nucleated cells from foetal liver, bone marrow, spleen and blood were isolated 10 or 20 h after injection and assayed *ex vivo* for IFN α secretion by ELISPOT and ELISA techniques. The administration of TGEV induced IFN α -secreting cells in foetal lymphohaematopoietic organs at mid-gestation. In contrast, IFN α was not detected in control sham-operated foetuses. A specific point mutation in the amino acid sequence of the viral membrane glycoprotein M of TGEV mutant dm49-4 was associated with lower or absent IFN α *in utero* inducibility by mutant virus as compared with wild virus. Flow cytometry analysis did not show differences in leukocyte surface marker expression between control and TGEV- or between dm49-4 and wild virus-treated foetus cells, with the exception of a reduction in percentages of polymorphonuclear cells in TGEV-treated lymphohaematopoietic tissues, which is probably due to IFN α secretion. The present data provided *in vivo* evidence of IFN α secretion at the cell level in foetal lymphohaematopoietic organs. Such IFN α -secreting cells in lymphohaematopoietic tissues may be the source of IFN α detected during foetal infections.

Key-words: Coronavirus, Transmissible gastroenteritis virus, IFN α ; ELISA, ELISPOT, Foetus, Pig.

INTRODUCTION

Interferon- α (IFN α) is a critical component of early, host non-specific immune defence against viral infections. It acts as both an antiviral agent

and an immunomodulator as well as a cell growth inhibitor. Several leukocyte populations are able to secrete IFN α , in response to virus stimuli, depending on the virus used and on whether viral infection of leukocytes is necessary or not.

Submitted January 9, 1997, accepted April 24, 1997.

(*) Corresponding author, present address : National Institute of Animal Health, Kannondai, Tsukuba, Ibaraki 305, Japan.

Monocytes are most often associated with production of IFN α in response to infectious viruses (Roberts *et al.*, 1979; Saksela *et al.*, 1984), whereas a distinct population of leukocytes among peripheral blood mononuclear cells (PBMC), referred to as "natural interferon-producing cells" (NIPCs), is able to secrete IFN α following exposure to non-infectious viral structures (Lebon *et al.*, 1982a; Capobianchi *et al.*, 1985; Charley and Laude, 1988). Human and porcine NIPCs have been characterized as highly infrequent, non-phagocytic, non-adherent, non-B, non-T, MHC II⁺ and CD4⁺ cells (Cederblad and Alm, 1990; Charley and Lavenant, 1990; Sandberg *et al.*, 1990; Fitzgerald-Bocarsly, 1993; Nowacki and Charley, 1993).

Viral glycoproteins were suggested as being responsible for triggering synthesis of IFN α in NIPC. Monoclonal antibodies (mAbs) to viral glycoproteins of herpes simplex virus type 1, transmissible gastroenteritis virus (TGEV) and Aujeszky's disease virus were shown to block virus-induced IFN α secretion (Lebon, 1985; Charley and Laude, 1988; Artursson, 1993). However, the precise nature of NIPCs as well as of the interactions between NIPCs and virus leading to IFN α production remains to be elucidated.

TGEV is an enteric coronavirus which causes acute and fatal diarrhoea, as well as intense and early IFN α production in newborn piglets (La Bonnardière and Laude, 1981). *In vitro* studies on IFN α induction by TGEV have shown that IFN α -secreting cells (IFN α -SCs) were detected among non-adherent porcine PBMCs after exposure to UV-inactivated TGEV or glutaraldehyde-fixed TGEV-infected cells (Charley and Lavenant, 1990; Nowacki and Charley, 1993). A specific point mutation in the amino acid sequence of the viral membrane glycoprotein M

in dm49-4 mutant TGEV was shown to be associated with a defect in *in vitro* induction of IFN α (Laude *et al.*, 1992).

The epitheliochorial nature of placentation in pig prevents transfer of immunoglobulins or antigens from mother to foetus, which precludes any immune activation of foetuses (Šterzl *et al.*, 1966). This type of placentation together with multiparity, relatively long term of gestation and size of foetuses makes this species suitable for studies on the development of the immune system. In an earlier study on the prenatal ontogeny of porcine IFN α -SCs, we detected *in vitro* inducible IFN α -SCs in pig foetal liver and other lymphohaematopoietic organs, at very early stages of gestation (Šplíchal *et al.*, 1994). The present study was undertaken to evaluate *in utero* viral induction of IFN α -SC in pig foetuses by TGEV, either partially or fully UV-inactivated, at different stages of gestation. We found that 10 or 20 h after *in utero* injection of TGEV in the umbilical cord of pig foetuses, IFN α -SCs were detected *ex vivo* in lymphohaematopoietic tissues by the ELISPOT assay.

MATERIALS AND METHODS

Animals

Healthy pregnant gilts of miniature pig bred in the Laboratory of gnotobiology in Nový Hrádek were used in our experiments. They had free access to water but were starved 12 hours before first surgery. The gilts were subcutaneously (s.c.) premedicated with 1 mg of atropin sulphate (Hoechst-Biotika, Slovakia) per 25 kg of body weight, and they were anaesthetized with 1.5-2.5% of halothane (Léčiva, Czech Republic) mixed with O₂ and N₂O. HCG (1,500 U) (Léčiva, Czech Republic) and acetate

FCM	=	flow cytometry (analysis).
FCS	=	foetal calf serum.
FSC	=	forward scatter.
HCG	=	human chorionic gonadotropin.
IFN	=	interferon.
IFN α -SC	=	IFN α -secreting cell.
mAb	=	monoclonal antibody.
MOC	=	mononuclear cell.
NIPC	=	natural interferon-producing cell.

PBL	=	peripheral blood leukocyte.
PBMC	=	peripheral blood mononuclear cell.
PBS	=	phosphate-buffered saline.
PFU	=	plaque-forming unit.
PMN	=	polymorphonuclear (cell).
SSC	=	side scatter.
SWC3	=	an antigen common for myelomonocytic lineages.
TGEV	=	transmissible gastroenteritis virus.

medroxyprogesterone (50 mg per 25 kg of body weight) (Upjohn, Netherlands) were intramuscularly (i.m.) injected. In the first series of experiments, fourteen foetuses of 52, 82 and 101 days of gestation were injected with partially UV-inactivated TGEV (500 PFU/ml in 50, 300 and 500 μ l of saline, respectively) via the umbilical vein when the umbilical cord was exteriorized after laparotomy and uterotomy of the gilts. Control foetuses were subjected to the same surgery but treated with equivalent volumes of the saline only. The uterine and abdominal walls were sutured, and gilts were placed in a postsurgical care unit. They had free access to water but a limited amount of food. Animals were treated with 1,500,000 U penicillin G (Spofa, Czech Republic) s.c. and 0.5 g streptomycin (Medexport, Russia) i.m. per 25 kg of body weight.

In the second series of experiments, twenty-one foetuses at 75, 77, 91 and 105 days of gestation were injected via the umbilical vein (300-1,500 μ l) with a fully UV-inactivated wild or dm49-4 mutant TGEV (initial titre of 6×10^7 PFU per ml before inactivation). The injected volume was proportionally adjusted to expected body weight. The treatment of animals was the same as described above.

Cell suspensions

The second uterotomy was performed 20 or 10 hours later, in the first or second series of experiments, respectively. The sows were anaesthetized, foetuses were bled via umbilical cord arteries, and blood samples containing 20 U of heparin/ml of blood (Léčiva, Czech Republic) were collected. Cell suspensions from liver, spleen, both femurs and sternum were prepared by cutting these organs with scissors in cold PBS. Supernatants containing cells were collected after 10 min sedimentation at 1 g to remove debris and clumps. PBMCs were prepared by centrifugation of diluted blood on "Ficoll" density gradient (Pharmacia, Sweden), and red cells were depleted by hypotonic lysis with deionized water as described (Šplíchal *et al.*, 1994). In the case of a non-adherent cell fraction isolation, cells were incubated for 90 min in RPMI-1640 with 20% foetal calf serum (FCS) (Seromed, Germany), L-glutamine (2 mM), penicillin (100 U/ml) and streptomycin (100 μ g/ml) (Gibco, UK) in 5% CO₂ atmosphere at 37°C to remove plastic-adherent cells. The number of nucleated cells and cell viability were calculated before assays.

IFN α assays

Cell suspensions (200 μ l) in RPMI with 10% FCS and antibiotics were log₃ diluted in 3-6 wells of 96-well cell culture microplate (Costar, UK). ELISPOT was performed as described (Nowacki and Charley, 1993). Cells in ELISPOT plates or in tissue

culture microplates were incubated in 5% CO₂ atmosphere for 16 and 18 h, respectively. IFN α -SC frequency was estimated by an ELISPOT assay using peroxidase-labelled anti-pig IFN α mAb F17. Spots were counted under binocular microscopy, and the frequency was calculated from the total number of living nucleated cells and number of spots. IFN α titres in supernatants from tissue culture microplates incubated for 18 h, and in plasma, were estimated by pig IFN α -specific ELISA using the same antibodies as in the case of ELISPOT (De Arce *et al.*, 1992; Nowacki and Charley, 1993). Results are expressed as IFN α unit/ml. Production of IFN α /IFN α -SC (yield) was estimated from the IFN α titre and the number of IFN α -specific spots.

Flow cytometry analysis

The following mouse mAbs directed against porcine leukocyte surface markers were used: K252.1E4 (anti-CD45), 74-22-15 (anti-SWC3), an antigen common for myelomonocytic lineages), 10-2H2 (anti-CD4) and MSA3 (anti-SLA-DR). Erythrocytes were removed from cell suspensions by hypotonic lysis of the cell pellet with water. Leukocytes were washed and stained as described recently (Cukrowska *et al.*, 1996): briefly, cells were treated with a primary mouse mAb and then with fluorescein-conjugated swine anti-mouse Ig polyclonal antibodies (Sevac, Czech Republic); for double staining, biotinylated mAbs were revealed by streptavidin-phycoerythrin conjugate (Immunotech, France). Cells were divided into two major populations on the basis of their size and internal complexity: polymorphonuclear cells (PMN) with higher SSC parameter gated separately from mononuclear leukocytes (MOC) with lower internal complexity. Flow cytometry data were obtained using a "FACSort" flow cytometer (Becton-Dickinson, CA). Propidium iodide was added to cells just before cytometry to prevent counting of dead and damaged cells, and at least 10,000 events were collected. Data were analysed using PC-LYSYS 1.0 software (Becton-Dickinson, CA).

Virus

High-passage Purdue 115 "wild strain" of TGEV and dm49-4 mutant TGEV (Laude *et al.*, 1992) were used as virus sources. Procedures for virus preparation have been described previously (Charley and Laude, 1988). Viruses were inactivated by UV irradiation. In the first series of experiments, TGEV was partially UV-irradiated to obtain a residual infectivity titre of 5×10^4 PFU/ml. In the second series of experiments, both wild and dm 49-4 mutant virus, at initial titres of 6×10^7 PFU/ml, were fully inactivated before injection.

RESULTS

Induction of IFN α -SC following *in utero* injection of partially inactivated wild TGEV

The optimal amount of partially inactivated wild TGEV allowing foetus survival after intravenous injection at 54 days of gestation was determined in preliminary experiments. Foetuses were thereafter infected with this amount of partially inactivated TGEV, and other foetuses were sham-operated as controls. At 52 days of gestation only three foetuses (two infected and one control) survived till the next day; dead foetuses infected by virus did not exhibit any macroscopic pathological lesions when compared with the dead control foetus. Only cell suspensions and plasma of surviving foetuses were used for IFN α determina-

tions. IFN α -SCs or IFN α were detected in liver and bone marrow cells or plasma in only one of two TGEV-treated foetuses already at mid-gestation (day 52) (table I). In the spleen they were found at later stages of gestation. The highest IFN α -SC frequency was observed in foetal liver. A slightly higher IFN α -SC frequency was detected in non-adherent cells compared with that in total cells (table I, 101 days). The absence of IFN α -SCs in spleen cells at 52 days may be due to the limited number of cells. The highest IFN α -SC frequency was observed by 101 days of gestation, whilst IFN α yield (production of IFN α per cell) was roughly constant (table II). No IFN α secretion in bone marrow cell culture supernatants could be detected at 52 days of gestation (table II), although low numbers of IFN α -specific spots were detected in one stimulated foetus (table I).

Table I. IFN α -secreting cells in pig foetal lymphohaematopoietic organs, and plasma IFN α levels 20 h after experimental *in utero* injection of partially UV-inactivated wild TGEV coronavirus.

Day of gestation	Liver	IFN α -SC frequency (spots per 10 ⁵ cells)		Bone marrow	IFN α level (units per ml) Plasma
		Spleen			
52nd (n = 2)	26 ; 0	0 ^(b)		2.1 ; 0	4 800 ; 0
82nd (n = 4)	84.5 \pm 57.7	2.2 \pm 1.1		14.8 \pm 13.5	9 200 \pm 2 870
101st (n = 3)	47.3 \pm 23.4	6.3 \pm 0.2		7.7 \pm 4.7	10 190 \pm 2 920
101st ^(a) (n = 3)	66.3 \pm 23.7	ND		ND	

Results are expressed as individual data or as means \pm standard deviation. Day of gestation=day of stimulation by TGEV; ^(a) non-adherent fraction; ^(b) low number of cells. ND=non-detected.

Numbers of sham-operated controls on 52nd, 82nd and 101st day of gestation were 1, 2 and 2, respectively. No IFN α -SC or IFN α titres were detected in lymphohaematopoietic organs or plasma of controls.

Table II. IFN α yield per cell in pig foetal lymphohaematopoietic cell cultures 20 h after experimental *in utero* injection of partially UV-inactivated wild TGEV coronavirus.

Day of gestation	Liver	IFN α secretion (units per IFN α -SC)	
		Spleen	Bone marrow
52 nd (n = 2)	0.7 ; 0	0 ^(b)	0
82nd (n = 4)	1.9 \pm 0.8	0.3 \pm 0.1	0.9 \pm 0.8
101st (n = 3)	0.3 \pm 0.2	0.2 \pm 0.1	0.2 \pm 0.1
101st ^(a) (n = 3)	0.2 \pm 0.1	ND	ND

^(a) Non-adherent fraction; ^(b) low number of cells. ND=non-detected.

No IFN α titres were detected in cell culture supernatants of lymphohaematopoietic organs of controls.

As controls, IFN α titres were determined in the plasma of TGEV-injected foetuses. High IFN α titres were found in foetal plasmas at 82 and 101 days of gestation (table I). Only one TGEV-injected foetus had plasma IFN α at 52 days of gestation.

No IFN α -specific spots or IFN α secretion was found in control sham-operated foetuses, which were subjected to *in utero* injection of saline only (data not shown).

Flow cytometry (FCM) analysis was performed in 52- and 101-day-old foetuses. SSC/FSC dot plot analysis showed a lower percentage of PMN cells in virus-injected foetal pig organs (fig. 1). FCM of leukocyte cell markers did not show significant differences between control and TGEV-injected foetuses (data not shown).

After evaluation of the results from this first series of experiments, we reduced the length of *in utero* stimulation from 20 to 10 hours in order to analyse IFN α secretion at its expected time of maximal production.

Induction of IFN α -SC following *in utero* injection of fully inactivated wild and dm49-4 mutant TGEV

In a second series of experiments using fully inactivated virus, differences in IFN α induction in pig foetuses stimulated by wild or dm49-4 mutant TGEV were analysed. In order to exclude any possible IFN α secretion by cells of monocytic/macrophage lineage, only non-adherent cells were used (as performed with liver cells at 101 days: table I). The main differences observed between the two series of experiments were a much lower plasma IFN α level, a reduced IFN α -SC frequency in liver and spleen and the absence of IFN α -SCs in foetal bone marrow (table III compared with table I). IFN α yields per cell were similar (tables II and IV). Neither plasma IFN α nor IFN α -SCs in organs were found in dm49-4-injected foetuses, with the exception of very low IFN α -SC numbers in liver and spleen of 75-day-old foetuses (tables III and IV).

Flow cytometry showed lower percentages of PMN cells in liver and bone marrow of wild TGEV-infected foetuses than in dm49-4-treated foetuses, with one exception – the liver of 105-day-old foetuses (fig. 2). No significant changes in leukocyte marker expression were observed (data not shown).

DISCUSSION

Intraamniotic infections provoke abortion, infertility, foetal death and abnormal foetal development and are associated with increased levels of amniotic inflammatory cytokines (Gravett *et al.*, 1994; Romero *et al.*, 1994; Dudley *et al.*, 1996). IFN was found not only in human foetuses during intraamniotic rubella infection (Lebon *et al.*, 1985) but also in the amniotic fluid of pregnant women without clinical signs of congenital virus infection (Lebon *et al.*, 1982b). Following our previous demonstration that *in vitro* inducible IFN α -SCs were present at early stages of gestation in porcine foetal lymphohaematopoietic organs (Šplíchal *et al.*, 1994), the aim of the present study was therefore to evaluate, at the cell level, the secretion of IFN α following *in utero* intravenous injection of TGEV in pig foetuses at different stages of gestation. The study was divided into two parts: (a) induction of IFN α secretion by partially inactivated wild TGEV, for which IFN α secretion was analysed in total cell populations (first series of experiments) and (b) induction of IFN α secretion in the non-adherent IFN α -SC fraction (presumably NIPC-like cells) after exposure to non-infectious TGEV. Wild and dm49-4 mutant TGEVs were compared in order to evaluate the influence of an amino acid point mutation in protein M on the IFN α -inducing properties *in vivo* (second series of experiments).

The experimental approach using “open” surgery is an efficient procedure for *in vivo* infection of foetuses via the umbilical cord vein. It has, however, several disadvantages such as loss of amniotic fluid, damage of foetal membranes and possible damage of the umbilical cord (Kovářů *et al.*, 1971). In the present study we observed high mortality of pig foetuses at 52 days of gestation




Fig. 1. FCM of umbilical cord blood, bone marrow and liver cells of 52- and 101-day-old foetuses after *in utero* injection of control medium or TGEV.

Dot plot analysis of foetal bone marrow cells (101 days of gestation) from control (a) or TGEV-injected (b) foetuses. Percentages of PMN cells in the different cell suspensions at 52 and 101 days of gestation (c). Number of foetuses as indicated in table I.

even in the case of one control foetus treated by saline only, which could be due to easier damage of the umbilical cord at this period of gestation than with older foetuses.

The main finding of the present study was that IFN α -SCs were found as early as 52 days of gestation in foetal liver and bone marrow following *in utero* injection of partially UV-inactivated

Table III. IFN α -secreting cells in pig foetal lymphohaematopoietic organs and plasma IFN α levels 10 h after experimental *in utero* injection of UV-inactivated wild and dm49-4 mutant TGEV coronavirus.

Day of gestation (wild/mutant)	IFN α -SC frequency (spots per 10 ⁵ cells)				IFN α levels (units per ml)	
	Liver		Spleen		Plasma	
	Wild virus	Mutant virus	Wild virus	Mutant virus	Wild virus	Mutant virus
75th (n = 3/3)	3.4 \pm 2.9	0.1 \pm 0.1	1.9 \pm 1.4	0.1 \pm 0.1	990 \pm 120	0
77th (n = 2/1)	0.7 ; 1.7	0	0 ; 2.4	0	ND	0
91st (n = 3/3)	3.7 \pm 1.4	0	28.8 \pm 11.5	0	1480 \pm 1180	0
105th (n = 3/3)	0.1 \pm 0.1	0	18.1 \pm 1.7	0	1400 \pm 780	0

IFN α ELISPOT assay was performed with non-adherent cells. ND = non-detected.

Table IV. IFN α yield per cell in pig foetal lymphohaematopoietic organs 10 h after experimental *in utero* injection of UV-inactivated wild and dm49-4 mutant TGEV coronavirus.

Day of gestation (wild/mutant)	IFN α secretion (units per IFN α -SC)			
	Liver		Spleen	
	Wild virus	Mutant virus	Wild virus	Mutant virus
75th (n = 3/3)	0.8 \pm 0.4	0.1 \pm 0.1	0.1 \pm 0.1	0
77th (n = 2/1)	0.2 ; 1.2	0	0 ; 0.3	0
91rd (n = 3/3)	0.8 \pm 0.4	0	0.4 \pm 0.1	0
105th (n = 3/3)	0.1 \pm 0.1	0	0.3 \pm 0.1	0

IFN α ELISPOT assay was performed with non-adherent cells.

virus. This is, to our knowledge, the first demonstration of the presence of IFN α -SCs in foetal lymphohaematopoietic tissues after *in vivo* viral induction. The fact that IFN α -SCs were found in only one virus-injected foetus may be due to an unsuccessful virus inoculation. At 52 days of gestation, a low frequency of spots but no IFN α in supernatants were observed in bone marrow cells of one foetus. This discrepancy may be due to a lower sensitivity of ELISA in comparison with that of ELISPOT.

Haematopoiesis in the bone marrow of pig foetuses starts at around 50 days of gestation (Šterzl and Kovářů, 1977), and the finding of IFN α -SCs at that time indicates the early appearance of these cells in lymphohaematopoietic organs. This finding is in accordance with our previous data on *in vitro* inducible foetal IFN α -

SCs (Šplíchal *et al.*, 1994) and supports the likely haematopoietic origin of IFN α -SCs (Charley *et al.*, 1995).

At later stages of gestation, *in utero* TGEV injection induced IFN α -SCs in foetal liver, spleen and bone marrow. When fully UV-inactivated TGEV was used, IFN α -SCs were not found in bone marrow, which may reflect the inability of non-replicating virus to reach that organ. A much lower IFN α production was observed after injection of fully inactivated virus compared with that after injection of infectious TGEV, as evidenced by both lower plasma IFN α titres and IFN α -SC frequency. This lower IFN α induction very probably reflects a lower amount of virus available when non-infectious TGEV is injected.

IFN α -SCs or IFN α was never found in blood cell suspensions (negative data not shown), but




Fig. 2. Percentages of PMN cells in bone marrow and liver cell suspensions of wild or dm49-4 mutant TGEV-injected foetuses of different ages. Numbers of foetuses as indicated in table III.

due to the low number of PBMC obtained, we cannot conclude whether these negative results could only be due to an insufficient number of cells tested or to a real absence of IFN α -SCs.

The lower IFN α inducibility by the UV-inactivated dm49-4 TGEV mutant characterized by a mutation in glycoprotein M, in comparison with inducibility by wild virus, has already been shown in *in vitro* experiments (Laude *et al.*, 1992). In the present work, IFN α was induced by the dm49-4 mutant to very low levels at only one period of gestation (75 days), while IFN α was observed in liver and spleen of foetuses at all ages tested, after induction by wild virus. This result implies that *in utero* IFN α induction by inactivated TGEV largely depends upon the presence of a native envelope protein M, as expected from *in vitro* studies (Laude *et al.*, 1992).

The presence of IFN α -SCs in pig lymphohaematopoietic tissues, together with plasma IFN α ,

confirms recent data obtained in mice within a broader framework: briefly, early and transient serum IFN α production was detectable as well as the presence of IFN α -producing cells localized in specific areas of spleen and lymph nodes, but not in bone marrow, after *in vivo* inactivated virus injection (Eloranta *et al.*, 1996; Riffault *et al.*, 1996).

The aim of FCM was to study changes in cell populations during induction of IFN α production. NIPC size and granularity as well as surface markers have been described (Sandberg *et al.*, 1990; Charley and Lavenant, 1990; Cederblad and Alm, 1990; Fitzgerald-Bocarsly, 1993; Nowacki and Charley, 1993). A more comprehensive characterization of human NIPC surface markers has recently been published (Svensson *et al.*, 1996). The latter study concluded that NIPC could be immature dendritic cells. In the present study, cells isolated from foetal lymphohaematopoietic organs were analysed on the basis of

their size, internal complexity (SSC/FSC) and surface markers. PMN with a higher SSC parameter could be distinguished from mononuclear leukocytes (MOC) with lower internal complexity. The PMN/MOC ratio was found to be reduced in virus-treated foetal organs (figs. 1a, b and c). In most cases studied, PMN percentages were lower in wild-virus-treated than in dm49-4-treated foetuses (fig. 2). Because dm49-4 did not induce IFN α *in vivo*, it is likely that TGEV-induced reduction of PMN percentages is IFN α -mediated. Previous studies also showed that immunization of pig foetuses with corpuscular antigen resulted in depletion of myeloid cells from foetal lymphohaematopoietic organs (Šterzl and Kovářů, 1977).

No significant differences in leukocyte surface marker expression between stimulated and non-stimulated animals, including SLADR and CD4 previously described on adult pig blood IFN α -SCs (Nowacki and Charley, 1993), were observed in any tested cell suspensions.

In conclusion, our present results provide *in vivo* evidence of IFN α secretion, at the cell level, in pig foetal lymphohaematopoietic organs, at early stages of lymphohaematopoietic development. IFN α detected in foetal infected tissues (Lebon *et al.*, 1985) may indeed originate from NIPCs localized in foetal lymphohaematopoietic tissues. In addition, our experimental model of *in utero* induction of IFN α in pig foetuses will prove to be useful for future studies on prenatal cytokine network development.

Acknowledgements

We gratefully thank Lydia Besnardeau for preparing viruses, Christiane de Vaureix for anti-IFN α mAb, Květoslava Rybinová and Marie Zahradníčková for their excellent technical assistance during surgery, and Marta Smolová and Eva Červinková for preparing cell suspensions. We thank Christine Young of the Translation Service, INRA, for revising the English.

This work was partly supported by INRA (DRI) and grants no. 508/94/0153 and 524/97/0533 of the Grant Agency of the Czech Republic.

L'injection *in utero* de coronavirus VGET induit la sécrétion d'interféron alpha dans les organes lymphohématopoïétiques de fœtus de porc

Nous avons montré précédemment que l'incubation *in vitro* de cellules mononucléées sanguines avec du coronavirus de la gastroentérite transmissible (VGET) inactivé par irradiation UV, induisait la sécrétion d'interféron alpha (IFN α). Le travail présenté ici a consisté à réaliser sur des fœtus de porcs à différents stades de gestation une injection *in utero* (a) de VGET sauvage partiellement inactivé par les UV, ou (b) de virus sauvage ou de mutant dm49-4 totalement inactivés. La sécrétion d'IFN α par les cellules nucléées isolées, 10 ou 20 h après injection, de la rate, du foie, de la moelle osseuse et du sang fœtaux a été analysée *ex vivo* par technique ELISPOT et ELISA. L'injection de VGET a induit la présence de cellules sécrétrices d'IFN α dans les organes lymphohématopoïétiques fœtaux à mi-gestation. Une sécrétion d'IFN α plus faible, voire absente, a été observée à la suite de l'injection du mutant viral dm49-4 caractérisé par une mutation ponctuelle dans la séquence de la glycoprotéine M. L'étude en cytométrie de flux n'a pas permis de montrer de différences d'expression des marqueurs de surface leucocytaires entre les cellules de fœtus traités par le VGET ou par du milieu, ou traités par le virus sauvage ou le mutant dm49-4, à l'exception d'une réduction du pourcentage de polynucléaires, après injection par le VGET, probablement due à l'IN α sécrété.

Ces résultats montrent, *in vivo*, la sécrétion au niveau unicellulaire d'IFN α dans les tissus lymphohématopoïétiques fœtaux. De telles cellules sécrétrices d'IFN α localisées dans les tissus pourraient constituer la source de l'IFN α qui est détecté au cours d'infections fœtales.

Mots-clés: Coronavirus, Virus de la gastroentérite transmissible, IFN α ; ELISA, ELISPOT, Fœtus, Porc.

References

- Artursson, K. (1993), Studies on the interferon- α/β system of pigs. Ph.D. thesis, Uppsala.
- Capobianchi, M.R., Facchini, J., Di Marco, P., Antonelli, G. & Dianzani, F. (1985). Induction of alpha interferon by membrane interaction between viral surface and peripheral blood mononuclear cells. *Proc. Soc. Exp. Biol. Med.*, 178, 551-556.
- Cederblad, B. & Alm, G. (1990), Infrequent but efficient interferon- α -producing human mononuclear leuko-

- cytes induced by herpes simplex virus *in vitro* studied by immunoplaque and limiting dilution assays. *J. Interferon Res.*, 10, 65-73.
- Charley, B. & Laude, H. (1988), Induction of alpha interferon by transmissible gastroenteritis coronavirus: role of transmembrane glycoprotein E1. *J. Virol.*, 62, 8-11.
- Charley, B. & Lavenant, L. (1990), Characterization of blood mononuclear cells producing IFN α following induction by coronavirus-infected cells (porcine transmissible gastroenteritis virus). *Res. Immunol.*, 141, 141-151.
- Charley, B., Nowacki, W. & Vaiman, M. (1995), Frequency of interferon- α -secreting leukocytes in irradiated and bone-marrow-grafted pigs. *Vet. Res.*, 26, 292-299.
- Cukrowska, B., Šinkora, J., Řeháková, Z., Šinkora, M., Šplíchal, I., Tučková, L., Avrameas, S., Saalmüller, A., Barot-Ciorbaru, R. & Tlaskalová-Hogenová, H. (1996), Isotype and antibody specificity of spontaneously formed immunoglobulins in pig fetuses and germ-free piglets: production by CD5⁻ B cells. *Immunology*, 88, 611-617.
- De Arce, H.D., Artursson, K., L'Haridon, R., Perers, A., La Bonnardière, C. & Alm, G.V. (1992), A sensitive immunoassay for porcine interferon- α . *Vet. Immunol. Immunopathol.*, 30, 319-327.
- Dudley, D.J., Hunter, C., Mitchell, M.D. & Varner, M.W. (1996), Elevations of amniotic fluid macrophage inflammatory protein-1 alpha concentrations in women during term and preterm labor. *Obstet. Gynecol.*, 87, 94-98.
- Eloranta, M.L., Sandberg, K. & Alm, G.V. (1996), The interferon- α/β responses of mice to herpes simplex virus studied at the blood and tissue level *in vitro* and *in vivo*. *Scand. J. Immunol.*, 43, 355-360.
- Fitzgerald-Bocarsly, P. (1993), Human natural interferon- α -producing cells. *Pharmacol. Ther.*, 60, 39-62.
- Gravett, M.G., Witkin, S.S., Haluska, G.J., Edwards, J.L., Cook, M.J. & Novy, M.M.J. (1994), An experimental model for intraamniotic infection and preterm labor in rhesus monkeys. *Am. J. Obstet. Gynecol.*, 171, 1660-1667.
- Kovářů, F., Stožický, V., Kruml, J., Dlabáč, V., Donát, J. & Novotná, J. (1971), Experimental surgery in the foetal period of mammals. *Acta. Vet. Brno*, 53, 1-68.
- La Bonnardière, C. & Laude, H. (1981), High interferon titer in newborn pig intestine during experimentally induced viral enteritis. *Infect. Immun.*, 32, 28-31.
- Laude, H., Gelfi, J., Lavenant, L. & Charley, B. (1992), Single amino acid changes in the viral glycoprotein M affect induction of alpha interferon by the coronavirus transmissible gastroenteritis virus. *J. Virol.*, 66, 743-749.
- Lebon, P. (1985), Inhibition of herpes simplex virus type 1-induced interferon synthesis by monoclonal antibodies against viral glycoprotein D and by lysosomotropic drugs. *J. Gen. Virol.*, 6, 2781-2785.
- Lebon, P., Comroy-Chevalier, M.J., Robert-Galliot, B. & Chany, C. (1982a), Different mechanisms for α and β interferon induction. *Virology*, 119, 504-507.
- Lebon, P., Daffos, F., Checoursy, A., Grangeot-Keros, L., Forestier, F. & Toublanc, J.E. (1985), Presence of an acid-labile alpha-interferon in sera from fetuses and children with congenital rubella. *J. Clin. Microbiol.*, 21, 775-778.
- Lebon, P., Girard, S., Thépot, F. & Chany, Ch. (1982b), The presence of α -interferon in human amniotic fluid. *J. Gen. Virol.*, 59, 393-396.
- Nowacki, W. & Charley, B. (1993), Enrichment of coronavirus-induced interferon-producing blood leukocytes increases the interferon yield per cell: a study with pig leukocytes. *Res. Immunol.*, 144, 111-120.
- Riffault, S., Eloranta, M.-L., Carrat, Ch., Sandberg, K., Charley, B. & Alm, G. (1996), Herpes simplex virus induces appearance of interferon- α/β producing cells and partially interferon- α/β dependent accumulation of leukocytes in murine regional lymph nodes. *J. Interferon Cytok. Res.*, 16, 1007-1014.
- Roberts, N.J., Douglas, R.G., Simons, R.M. & Diamond, M.E. (1979), Virus induced interferon production by human macrophages. *J. Immunol.*, 123, 365-369.
- Romero, R., Gomez, R., Galasso, M., Munoz, H., Acosta, L., Yoon, B.H., Svinarich, D. & Cotton, D.B. (1994), Macrophage inflammatory protein-1 alpha in term and preterm parturition: effect of microbial invasion of the amniotic cavity. *Am. J. Reprod. Immunol.*, 32, 108-113.
- Saksela, E., Virtanen, I., Hovi, T., Secher, D.S. & Cantell, K. (1984), Monocyte is the main producer of human alpha interferons following Sendai virus induction. *Prog. Med. Virol.*, 30, 78-86.
- Sandberg, K., Matsson, P. & Alm, G.V. (1990), A distinct population of nonphagocytic and CD4⁺ null lymphocytes produce interferon- α after stimulation by herpes simplex virus-infected cells. *J. Immunol.*, 145, 1015-1020.
- Šplíchal, I., Bonneau, M. & Charley, B. (1994), Ontogeny of interferon alpha secreting cells in the porcine fetal hematopoietic organs. *Immunol. Lett.*, 43, 203-208.
- Šterzl, J. & Kovářů, F. (1977), Development of lymphatic tissue and immunocompetency in pig foetus and germ-free piglets. *Acta. Vet. Brno*, 46, suppl. 3, 13-53.
- Šterzl, J., Rejnek, J. & Trávníček, J. (1966), Impermeability of pig placenta for antibodies. *Folia Microbiol.*, 11, 7-10.
- Svensson, H., Johannisson, A., Nikkila, T., Alm, G.V. & Cederblad, B. (1996), The cell surface phenotype of human natural interferon- α producing cells as determined by flow cytometry. *Scand. J. Immunol.*, 44, 164-172.