

The orthopaedic experience of COVID-19: A literature review

Journal of Perioperative Practice
2021, Vol. 31(3) 102–107

© The Author(s) 2021



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1750458920971506
journals.sagepub.com/home/ppj



Ali Al-kulabi¹ , Mohamed A Mansour¹  and Azeem Thahir²

Abstract

This literature review aims to provide an account of the changes to orthopaedics in the era of COVID-19. Herein, the authors explored the use of telemedicine in orthopaedics as well as changes in surgical protocols, screening methods, work priorities and orthopaedic education. There was increased utilisation of telemedicine in orthopaedic training and outpatient cases as a means to provide continuity in education and care. The need to implement social distancing measures, coupled with the reduced availability of staff, has dictated that the practice of orthopaedics shifts to focus on acute care whilst redistributing resources to front-line specialities. This was facilitated by the cancellation of electives and the reduction of outpatient clinics. Thus, it is demonstrated that major changes have been implemented in many aspects of orthopaedic practice in order to address the challenges of the COVID-19 pandemic.

Keywords

Orthopaedics / COVID-19 / Coronavirus / Surgical protocols / Telemedicine / Orthopaedic management

Provenance and Peer review: Unsolicited contribution; Peer reviewed; Accepted for publication 12 October 2020.

Introduction

Coronavirus disease-19 (COVID-19) is an infectious disease caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pathogen which originated in the city of Wuhan, China (Li et al 2020). Fever, cough and fatigue were the most common symptoms but severe complications, such as acute respiratory distress syndrome, often resulted in hospitalisation (Fu et al 2020). Over half a million deaths worldwide have been confirmed by the World Health Organization (WHO). Realising the alarming transmissibility and virulence of SARS-CoV-2, it was characterised as a pandemic on 11 March 2020 (WHO 2020). Knowledge of SARS-CoV-2 is limited but growing, although a vaccine has not yet been developed.

Healthcare systems have undergone substantial reorganisation to release capacity, including the reallocation of staff, increasing intensive care unit (ICU) availability and prioritising acute services (Leclerc et al 2020). Healthcare providers have been under immense pressure to protect their staff as well as care for patients, presenting significant fiscal and supply challenges across the world (Rowan & Laffey 2020).

The US Surgeon General has instructed the cessation of elective surgery (Stahel 2020); as have other health authorities. Thus, despite not being considered on the 'front-line', orthopaedic practice has been impacted. COVID-19 has had ramifications for surgical protocols, theatre workflow, patient and staff safety, training and education (Al-Jabir et al 2020, De Simone et al 2020).

In this literature review, we aimed to summarise the impact of COVID-19 on the practice of orthopaedics rather than provide a systematic review or critical analysis. Some aspects such as leadership and communication were critical during the pandemic in all specialties; however, this review focussed on the changes to orthopaedic surgery.

Our findings showed significant changes in orthopaedic practice and adoption of technology.

¹School of Clinical Medicine, University of Cambridge, Cambridge, UK

²Department of Trauma and Orthopaedic Surgery, Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

Corresponding author:

Ali Al-kulabi, School of Clinical Medicine, University of Cambridge, Hills Road, Cambridge CB2 0SP, UK.
Email: afia2@cam.ac.uk

Selection criteria

PubMed, Embase, Google Scholar, Medline and Web of Science were searched up to 1 June 2020. Key words included 'Orthopaedic', 'Coronavirus', 'COVID-19' and 'Surgery'. The initial screening of the studies was done by two authors independently, for inclusion, with any discrepancies resolved by a third reviewer. All articles written in English were included if they concerned the management of orthopaedic patients and departments during the COVID-19 pandemic. Case series, research-focused, project reports and descriptive articles were included with the aim of providing a global overview. Forty-five articles were selected.

Patient screening for COVID-19

Three main stages of screening for SARS-CoV-2 infection were described in literature. Epidemiological history and symptoms were investigated at the point of admission as a method of quickly identifying suspected COVID-19 patients (Neradi et al 2020). These patients were sometimes isolated in buffer wards for further testing (Meng et al 2020, Wu et al 2020).

Computed tomography (CT) imaging and blood tests were used to further investigate suspected COVID-19 patients and patients requiring surgery (Keny et al 2020). CT was believed to have high sensitivity for COVID-19 and can be conducted quickly (Ai et al 2020). Few articles expanded on the blood tests required but COVID-19 positive patients had statistically significant differences in white blood cells, C-reactive protein, aspartate transaminase, alanine aminotransferase and lactate dehydrogenase readings (Ferrari et al 2020).

Reverse transcription polymerase chain reaction was considered the diagnostic reference standard and was used as the final confirmatory step; however, the slow turn-around in results made CT more useful in emergency situations (Meng et al 2020). Unscreened patients were strictly managed in isolated rooms with COVID-19 precautions (Wu et al 2020).

Preoperative screening for COVID-19 increased surgery waiting times, which may have increased the risk of complications (Orosz et al 2004, Pincus et al 2017). A Chinese study determined that average waiting times from injury to surgery increased by 4.1 days, and from admission to surgery by 2.0 days, compared with 2019. Patients also waited longer before presenting to hospital (Meng et al 2020).

Elective surgery

The cessation of elective surgery shifted orthopaedic focus towards emergency care (Zagra et al 2020). The cancellations reduced viral spread, preserved bed capacity and personal protective equipment (PPE) and

allowed staff to be redeployed to confront the crisis (Iannuzzi et al 2020). More urgent electives were prioritised in the weeks prior to the gradual cessation of most electives (Tadros et al 2020). Furthermore, one paper noted a difficulty in guaranteeing a safe and effective rehabilitation time for patients (Zagra et al 2020). In highly affected regions, such as Lombardy, orthopaedic electives were stopped more abruptly (Giacomo et al 2020).

Most papers excluded urgent electives (such as periprosthetic joint infections and periprosthetic fractures) and orthopaedic oncology cases from cancellation (Giacomo et al 2020, Kenanidis & Tsidiris 2020). In contrast to other articles, one paper from Singapore continued day surgical cases, deeming that the resource-burden and infection risk were reduced by expedient discharge (Liang et al 2020).

The backlog of elective cases will eventually need to be faced (Petroni et al 2020), but reduced resources in some countries may compromise this (Jain & Vaishya 2020).

Emergency surgery

Emergency, urgent elective and oncological orthopaedic services were typically kept operational throughout the crisis (Giacomo et al 2020, Giuntoli et al 2020). Changes in emergency surgery workload varied across the literature reviewed. This was partly due to the fact that some centres were designated as regional trauma centres during the crisis (Meng et al 2020, Zagra et al 2020). Generally, reductions in trauma admissions were seen, with several studies reporting a decrease in trauma workload (Giuntoli et al 2020, Luengo-Alonso et al 2020), likely related to cases of minor trauma. Factors contributing to the reduction are travel restrictions and reduced mobility encouraged by the lock-down measures (Nuñez et al 2020), and a greater fear of coming to hospital during the coronavirus pandemic (Luengo-Alonso et al 2020, Nuñez et al 2020). Some private hospitals in Singapore and the UK offered their services to help public hospitals manage trauma patients (Tadros et al 2020, Tay et al 2020).

Surgical protocol changes

The risk of COVID-19 infection has necessitated the worldwide reduction in the size of surgical teams in the operating room (Lancaster et al 2020, Meraghni et al 2020). This facilitated distancing during surgeries, reduced the likelihood of transmission and allowed scarce PPE to be used elsewhere. The redeployment of staff to the 'front line' also necessitated this reduction. To further reduce infection risk, efforts were made to reduce the duration of surgeries (Neradi et al 2020). However, the additional COVID-19 precautions (discussed below)

have also led to an increase in theatre utilisation per case in some practices (Mathai et al 2020).

The use of PPE during surgeries (such as N95 respirators) was aspired to where possible, in compliance with Centers for Disease Control and Prevention guidelines (CDC 2020, Day et al 2020, Keny et al 2020). During aerosol-generating procedures, FFP3 masks were used alongside eye protection (Tadros et al 2020).

The pandemic necessitated reorganisations of operating theatres and surrounding spaces. Some theatres were dedicated to COVID-19 patients in line with wider segregation efforts in hospitals. Donning and doffing areas were improvised, along with sterile passages between them and theatres (Keny et al 2020). Clean rooms were introduced to facilitate the exchange of materials before surgeries (Mathai et al 2020).

Particulate aerosolisation during surgery was a significant infection risk, particularly during intubation and extubation. Many protocols were employed to address this, including asking the surgical team to step out during intubation and extubation, although this was practiced in many departments before the pandemic (Liu et al 2020, Neradi et al 2020). The use of power tools during surgeries was minimised, and the use of pulsatile jet lavage was avoided (Keny et al 2020). A period of 30 minutes was left before removing patients from the operating room after a surgery to allow aerosolised material to settle before cleaning (Keny et al 2020). The use of negative pressure in operating rooms was adopted by some but not all departments owing to cost and technical challenges. One UK paper stated that their orthopaedic department continued to use positive pressure, citing that little evidence exists for negative pressure in reducing COVID-19 infection risk (Mathai et al 2020). Due to constraints in anaesthetic resources, one department in Singapore converted selected surgeries usually performed under general anaesthesia to local anaesthesia (Hwee et al 2020).

Outpatients

Outpatient clinics have implemented measures to reduce staff and patient contact amidst the pandemic. A total of 23,580 outpatient appointments (93.8%) were cancelled in one Milan centre (Zagra et al 2020). Almost all other papers noted a similar trend. A reduction in referrals was also noted (Askari et al 2020). The duration between non-urgent follow-up appointments was intentionally prolonged, thereby reducing the demand on outpatient services (Liang et al 2020).

All publications discussed the utilisation of telemedicine to conduct clinic appointments, with further details discussed below. However, in-person clinic appointments were still provided to nondeferrable

outpatient cases such as plaster removals and some wound examinations (Tadros et al 2020). As a result, rearrangement of the outpatient department was necessary to maintain social distancing and isolation protocols (Mathai et al 2020). Orthopaedic team segregation reduced cross-infection between outpatient and inpatient services. Outpatient appointments were used to review some postponed elective cases, focusing instead on addressing debilitating pain in the early stages of the crisis (Liang et al 2020).

Telemedicine

In this review, 'telemedicine' describes the use of technology to deliver medical services when distance separates the participants. Prior to COVID-19, its adoption had been slow due to cost, staff and patient technology literacy, the inability to physically examine and the reliability of technology. Some barriers to implementation such as lack of awareness (Ayatollahi et al 2015) and regulatory complications were reduced amidst the pandemic; however, others persist such as poor connectivity (Makhni et al 2020). Patient attitudes to telemedicine adoption seemed to be positive, with one report stating that 90% of patients were satisfied with virtual clinics implemented at a UK hospital in line with the NHS Long Term Plan (Gilbert et al 2020). Another paper noted that telemedicine was used in 90% of outpatient clinical evaluations (Luengo-Alonso et al 2020).

Telephone consultations were cheaper than video conferences but failed to provide a solution to the loss of physical examination (Gilbert et al 2020, Meraghni et al 2020). This can present a significant challenge in a speciality like orthopaedics where observing the patient can be critical in diagnosis. Tanaka et al (2020) describes some solutions implemented via video conferencing, such as utilising on-screen measurement tools and objects of known weight to determine length and strength, respectively.

Initial costs included costs of hardware, software, installation and staff training (Makhni et al 2020). Articles noted that long-term cost savings stemming from reduced administration and travel costs, outweighed this initial cost (Ohinmaa et al 2002).

Education

Orthopaedic trainees were redirected to COVID-19 facing areas including ICU and emergency departments, reducing their orthopaedic exposure. Conferences usually attended by orthopaedic departments were preserved by migrating online. Additional online conferences held by industry sponsors, as well as online subspecialty lectures were made available to orthopaedic trainees. The pandemic has also introduced an additional need to train staff in the use of PPE which

was delivered online and in person. Hands-on surgical experience was supplemented with virtual reality simulators used from home in one institution (Schwarzkopf et al 2020). Most institutions implemented simpler virtual teaching solutions like the use of video conferencing apps to facilitate the continuation of 'normal' teaching. Weekly journal clubs allowed for interaction with trainees (Pelt et al 2020).

National guidance in the UK also dictated that surgical exams were either deferred or cancelled (HEE 2020). A paper on the challenges in Singapore and Malaysia (Tay et al 2020) also reported that practical internal exams were postponed due to difficulty in finding patient volunteers during the pandemic.

Staff health

The COVID-19 pandemic has taken its toll on the physical and mental wellbeing of healthcare staff due to frequently changing protocols, shortages in PPE, higher patient mortality and the risk of COVID-19 infection (Walton et al 2020). Caring for infected colleagues compounded the mental and physical health burden. Furthermore, the increased relative risk of infecting family has contributed to the stress and worry of healthcare staff (Guo et al 2020). Measures adopted to curtail COVID-19 spread included changing work rotas and work patterns (Mathai et al 2020) in addition to the strategies outlined previously. Despite this, a small number of healthcare professionals, including orthopaedic staff, have contracted the virus (Guo et al 2020).

An uncertain future and disruption of social support appears to have adversely affected younger staff disproportionately. A survey completed by 611 orthopaedic surgeons from India found a significantly higher level of stress in younger staff (Sahu et al 2020).

Regular praise and acknowledgement of staff, as well as drop-in sessions with psychologists/psychiatrists were recommended based on evidence from previous outbreaks (Walton et al 2020). Conversely, many orthopaedic staff were either mildly stressed or not stressed at all as a result of the marked reduction in elective workload and a reversed work-life balance (Sahu et al 2020).

The future in a COVID-19 world

Expectations of the 'new normal' in a COVID-19 world resemble the changes discussed above. A return to elective surgery is expected, starting with simpler operations and gradually adding complex cases (North et al 2020). The literature indicates an emphasis on precaution and diligence to accompany the resumption of elective surgery, with assessment of COVID-19 patient symptom status on the day of surgery likely to become

common. Preoperative patient testing for COVID-19, as well as routine testing of surgical staff has now been implemented due to the risk of asymptomatic transmission. This relies on the availability of fast and cost-effective tests in the future (Meneghini 2020).

Patient fear may remain a hurdle; the risk of COVID-19 infection in hospital discourages patients from electing for surgery, even when medically indicated. This uncertainty extends to life-saving operations (Lancaster et al 2020). An increasing focus on same-day discharge for operations such as hip and knee arthroplasties may help to reduce this risk (Meneghini 2020).

Elective surgery deferrals are likely to greatly impact patients with non-urgent but disabling conditions, reducing their ability to work (COVID Surg Collaborative 2020). This can summate with the already high costs of treatment, leading to impoverishment, particularly in low and middle income countries (Shrime et al 2015). The resulting deterioration in population health during the struggle to 'catch up' with cancelled electives will lead to further societal costs (COVID Surg Collaborative 2020). The world has long to go before resolving the issues associated with COVID-19.

Conclusion

The COVID-19 outbreak has presented a significant challenge to global health systems and the practice of orthopaedics worldwide. To adapt, the orthopaedic community has shifted its treatment priorities to focus on acute care in concert with the efforts of other healthcare specialities. New orthopaedic surgery protocols have reflected the need to both preserve resources and limit infection risks. There has been a significant move towards virtualising both orthopaedic care and training. It remains to be seen how and if these changes will be incorporated into orthopaedic practice beyond the pandemic.

Key Phrases

1. An increased utilisation of telemedicine in the outpatient setting was observed.
2. A shift to focus on acute surgical cases rather than non-urgent surgical procedures was key during the COVID-19 pandemic.
3. A reduction in outpatient and elective cases was intentionally planned as a measure to reduce transmission and divert staff to combat COVID-19.
4. Orthopaedic surgical theatres often reduced staff as part of the response to COVID-19, including students.
5. Orthopaedic training continues in the form of virtual engagements, and attempts were made to maintain practical skills training through technology.

No competing interests declared.

ORCID iDs

Ali Al-kulabi  <https://orcid.org/0000-0001-9884-8218>

Mohamed A Mansour  <https://orcid.org/0000-0002-4558-5634>

References

- Ai T, Yang Z, Hou H et al 2020 Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: A report of 1014 cases *Radiology* 296 E32–E40
- Al-Jabir A, Kerwan A, Nicola M et al 2020 Impact of the Coronavirus (COVID-19) pandemic on surgical practice – Part 1 *International Journal of Surgery* 79 168–179
- Askari A, Arasteh P, Jabalameli M et al 2020 COVID-19 and orthopaedic surgery: Experiences from Iran *The Journal of Bone and Joint Surgery* 102 1126–1128
- Ayatollahi H, Sarabi FZP, Langarizadeh M 2015 Clinicians' Knowledge and Perception of Telemedicine Technology. *Perspectives in health information management* 12(Fall). American Health Information Management Association. Available at: <http://perspectives.ahima.org/clinicians-knowledge-and-perception-of-telemedicine-techn>. (accessed 22 July 2020).
- Centre for Disease Control and Prevention 2020 Personal Protective Equipment: Questions and Answers. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirator-use-faq.html> (accessed 29 August 2020).
- COVID Surg Collaborative 2020 Elective surgery cancellations due to the COVID-19 pandemic: Global predictive modelling to inform surgical recovery plans *British Journal of Surgery* Epub ahead of print 12 May 2020. DOI: 10.1002./bjs.11746.
- Day J, MacMahon A, Roberts MM et al 2020 Perspectives from the foot and ankle department at an academic orthopedic hospital during the surge phase of the COVID-19 pandemic in New York City *Foot & Ankle International* 41 881–884
- De Simone B, Chouillard E, Di Saverio S et al 2020 Emergency surgery during the COVID-19 pandemic: What you need to know for practice *Annals of the Royal College of Surgeons of England* 102 323–332
- Ferrari D, Ferrari D, Motta A et al 2020 Routine blood tests as a potential diagnostic tool for COVID-19 *Clinical Chemistry and Laboratory Medicine* 58 (7) 1095–1099
- Fu L, Wang B, Yuan T et al 2020 Clinical characteristics of coronavirus disease 2019 (COVID-19) in China: A systematic review and meta-analysis *Journal of Infection* 80 (6) 656–665
- Giacomo P, Damiano S, Elena D et al 2020 CoViD-19 and ortho and trauma surgery: The Italian experience *Injury* 51 1403–1405
- Gilbert AW, Billany JCT, Adam R et al 2020 Rapid implementation of virtual clinics due to COVID-19: Report and early evaluation of a quality improvement initiative *BMJ Open Quality* 9 (2) e000985.
- Giuntoli M, Bonicoli E, Bugelli G et al 2020 Lessons learnt from COVID 19: An Italian multicentric epidemiological study of orthopaedic and trauma services *Journal of Clinical Orthopaedics and Trauma* 11 721–727
- Guo X, Wang J, Hu D et al 2020 Survey of COVID-19 disease among orthopaedic surgeons in Wuhan, People's Republic of China *The Journal of Bone and Joint Surgery* 102 (10) 847–854
- Health Education England 2020 Coronavirus (Covid-19) Information for trainees. Available at: <https://www.hee.nhs.uk/coronavirus-information-trainees> (accessed 27 June 2020).
- Hwee J, Chiew J, Sechachalam S 2020 The impact of coronavirus disease 2019 (COVID-19) on the practice of hand surgery in Singapore *The Journal of Hand Surgery* 45 536–541
- Iannuzzi NP, Lack WD, Gee AO et al 2020 An orthopaedic department's response to the COVID-19 health-care crisis indirect and direct actions with thoughts for the future *Journal of Bone and Joint Surgery* 102 e65
- Jain VK and Vaishya R 2020 COVID-19 and orthopaedic surgeons: The Indian scenario *Tropical Doctor* 50 (2) 108–110.
- Kenanidis E, Tsidis E 2020 "Flattening the Curve" of COVID-19 pandemic in orthopaedics and trauma: The Greek perspective *Injury* 51 1681–1682
- Keny S, Bagaria V, Chaudhary K et al 2020 Emergency and urgent orthopaedic surgeries in non-covid patients during the COVID 19 pandemic: Perspective from India *Journal of Orthopaedics* 20 275–279
- Lancaster EM, Sosa JA, Sammann A et al 2020 Rapid response of an academic surgical department to the COVID-19 pandemic: Implications for patients, surgeons, and the community *Journal of the American College of Surgeons* 230 (6) 1064–1073
- Leclerc T, Donat N, Donat A et al 2020 Prioritisation of ICU treatments for critically ill patients in a COVID-19 pandemic with scarce resources *Anaesthesia Critical Care and Pain Medicine* 39 (3) 333–339
- Li H, Liu SM, Yu XH et al 2020 Coronavirus disease 2019 (COVID-19): Current status and future perspectives *International Journal of Antimicrobial Agents* 55 (5) 105951
- Liang ZC, Wang W, Murphy D et al 2020 Novel coronavirus and orthopaedic surgery: early experiences from Singapore *Journal of Bone and Joint Surgery* 102 745–749
- Liu J, Mi B, Hu L et al 2020 Preventive strategy for the clinical treatment of hip fractures in the elderly during the COVID-19 outbreak: Wuhan's experience *Aging* 12 (9) 7619–7625
- Luengo-Alonso G, Pérez-Tabernero FGS, Tovar-Bazaga M et al 2020 Critical adjustments in a department of orthopaedics through the COVID-19 pandemic *International Orthopaedics* 44 1557–1564
- Makhni MC, Riew GJ, Sumathipala MG 2020 Telemedicine in orthopaedic surgery: Challenges and opportunities *The Journal of Bone and Joint Surgery* 102 1109–1115
- Mathai NJ, Venkatesan AS, Key T et al 2020 COVID-19 and orthopaedic surgery: Evolving strategies and early experience *Bone & Joint Open* 1 (5) 160–166
- Meneghini RM 2020 Resource reallocation during the COVID-19 pandemic in a suburban hospital system: Implications for outpatient hip and knee arthroplasty *Journal of Arthroplasty* 35 (7) S15
- Meng Y, Leng K, Shan L et al 2020 A clinical pathway for pre-operative screening of COVID-19 and its influence on clinical outcome in patients with traumatic fractures *International Orthopaedics* 44 1549–1555
- Meraghni N, Benkaidali R, Derradji M et al 2020 Orthopedic healthcare in the time of COVID-19: Experience of the orthopedic surgery department at Mustapha Bacha Hospital, Algeria *Annals of Medicine and Surgery* 55 164–166

- Neradi D, Hooda A, Shetty A et al 2020 Management of Orthopaedic Patients During COVID-19 Pandemic in India: A guide *Indian Journal of Orthopaedics* 54 402–407
- North T, Bullock MW, Danoff JR et al 2020 Arthroplasty during the COVID-19 pandemic *Arthroplasty Today* 6 (3) 427–430
- Nuñez JH, Sallent A, Lakhani K et al 2020 Impact of the COVID-19 pandemic on an emergency traumatology service: Experience at a tertiary trauma centre in Spain *Injury* 51 (7) 1414–1418
- Ohinmaa A, Vuolio S, Haukipuro K et al 2002 A cost-minimization analysis of orthopaedic consultations using videoconferencing in comparison with conventional consulting *Journal of Telemedicine and Telecare* 8 (5) 283–289
- Orosz GM, Magaziner J, Hannan EL et al 2004 Association of timing of surgery for hip fracture and patient outcomes *Journal of the American Medical Association* 291 (14) 1738–1743
- Pelt CE, Campbell KL, Gililand JM et al 2020 The rapid response to the COVID-19 pandemic by the arthroplasty divisions at two academic referral centers *Journal of Arthroplasty* 35 S10–S14
- Petrone B, Iturriaga C, Mauri T et al 2020 COVID-19 and orthopaedics: recovery after the pandemic surge *Arthroscopy, Sports Medicine, and Rehabilitation* Epub ahead of print 23 May 2020. DOI: 10.1016/j.asmr.2020.05.009
- Pincus D, Ravi B, Wasserstein D et al 2017 Association between wait time and 30-day mortality in adults undergoing hip fracture surgery *JAMA – Journal of the American Medical Association* 318 (20) 1994–2003
- Rowan NJ, Laffey JG 2020 Challenges and solutions for addressing critical shortage of supply chain for personal and protective equipment (PPE) arising from Coronavirus disease (COVID19) pandemic – Case study from the Republic of Ireland *Science of the Total Environment* 725 138532
- Sahu D, Agrawal T, Rathod V et al 2020 Impact of COVID 19 lockdown on orthopaedic surgeons in India: A survey *Journal of Clinical Orthopaedics and Trauma* 11(Suppl 3) S283
- Schwarzkopf R, Maher NA, Slover JD et al 2020 The response of an orthopedic department and specialty hospital at the epicenter of a pandemic: The NYU Langone Health Experience *Journal of Arthroplasty* 35: S3–S5
- Shrime MG, Dare AJ, Alkire BC et al 2015 Catastrophic expenditure to pay for surgery worldwide: A modelling study *The Lancet Global Health* 3 (S2) S38–S44
- Stahel PF 2020 How to risk-stratify elective surgery during the COVID-19 pandemic? *Patient Safety in Surgery* 14 8
- Tadros BJ, Black J, Dhinsa BS 2020 COVID-19 outbreak: The early response of a UK orthopaedic department *Journal of Clinical Orthopaedics and Trauma* 11(Suppl 3) S301
- Tanaka M, Oh L, Martin S et al 2020 Telemedicine in the era of COVID-19: The virtual orthopaedic examination *The Journal of Bone and Joint Surgery* 102 e57
- Tay K, Kamal T, Woo YL et al 2020 COVID-19 in Singapore and Malaysia: Rising to the challenges of orthopaedic practice in an evolving pandemic *Malaysian Orthopaedic Journal* 14 (2) 7–15
- Walton M, Murray E, Christian MD 2020 Mental health care for medical staff and affiliated healthcare workers during the COVID-19 pandemic *European Heart Journal: Acute Cardiovascular Care* 9 (3) 241–247
- World Health Organization 2020 Timeline of WHO's response to COVID-19. Available at: <https://www.who.int/news-room/detail/29-06-2020-covidtimeline> (accessed 26 July 2020).
- Wu W, Xiong W, Kang H et al 2020 Challenges and response in the medical management of the orthopaedic department during the coronavirus disease 2019 pandemic: Strategies in Wuhan, China *International Orthopaedics* 44 1489–1495
- Zagra L, Faraldi M, Pregliasco F et al 2020 Changes of clinical activities in an orthopaedic institute in North Italy during the spread of COVID-19 pandemic: A seven-week observational analysis *International Orthopaedics* 44 1591–1598