

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# *N'*-(*E*)-2-Fluorobenzylidene]benzohydrazide

P. B. Sreeja,<sup>a</sup> M. Sithambaresan,<sup>b\*</sup> N. Aiswarya<sup>c</sup> and  
M. R. Prathapachandra Kurup<sup>c</sup>

<sup>a</sup>Department of Chemistry, Christ University, Hosur Road, Bangalore 560 029, Karnataka, India, <sup>b</sup>Department of Chemistry, Faculty of Science, Eastern University, Sri Lanka, Chenkalady, Sri Lanka, and <sup>c</sup>Department of Applied Chemistry, Cochin University of Science and Technology, Kochi 682 022, India  
Correspondence e-mail: eesanas@yahoo.com


Received 25 October 2013; accepted 21 November 2013

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.050;  $wR$  factor = 0.165; data-to-parameter ratio = 17.9.

The asymmetric unit of the title compound,  $\text{C}_{14}\text{H}_{11}\text{FN}_2\text{O}$ , contains two independent molecules, both of which adopt the *E* conformation with respect to the azomethine  $\text{C}=\text{N}$  bond. The molecules are non-planar, with dihedral angles of 26.92 (12) and 11.36 (11)° between the benzene and phenyl rings. In the crystal, molecules are linked through  $\text{N}-\text{H}\cdots\text{O}=\text{C}$  and  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bonds into chains along [101].  $\text{C}-\text{H}\cdots\text{O}$  contacts link these chains into layers parallel to (001). The three-dimensional crystal packing is stabilized by  $\pi-\pi$  interactions, the shortest separation between the centroids of benzene rings being 3.884 (1) Å.

## Related literature

For catalytic properties of hydrazones, see: Heravi *et al.* (2007). For their use as inhibitors of enzymes, see: Tamasi *et al.* (2005) and for their biological activity, see: Sreeja *et al.* (2004). For the synthesis of related compounds, see: Mangalam & Kurup (2011). For a related structure, see: Nair *et al.* (2012).



## Experimental

### Crystal data

$\text{C}_{14}\text{H}_{11}\text{FN}_2\text{O}$   
 $M_r = 242.25$   
Monoclinic,  $P2_1/n$   
 $a = 9.7010$  (6) Å  
 $b = 17.4114$  (13) Å  
 $c = 15.002$  (1) Å  
 $\beta = 104.126$  (4)°

$V = 2457.3$  (3) Å<sup>3</sup>  
 $Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 0.10$  mm<sup>-1</sup>  
 $T = 296$  K  
0.35 × 0.30 × 0.25 mm

### Data collection

Bruker Kappa APEXII CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2007)  
 $T_{\min} = 0.968$ ,  $T_{\max} = 0.977$   
18766 measured reflections  
6111 independent reflections  
3320 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$   
 $wR(F^2) = 0.165$   
 $S = 1.00$   
5972 reflections  
333 parameters  
2 restraints  
H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.21$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.17$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N}2-\text{H}2'\cdots\text{O}2^i$	0.89 (1)	2.09 (2)	2.875 (2)	147 (2)
$\text{N}2-\text{H}2'\cdots\text{N}3^i$	0.89 (1)	2.60 (2)	3.339 (2)	142 (2)
$\text{N}4-\text{H}4'\cdots\text{O}1$	0.87 (1)	2.03 (1)	2.882 (2)	171 (2)
$\text{C}7-\text{H}7'\cdots\text{O}2^i$	0.93	2.53	3.213 (2)	131
$\text{C}14-\text{H}14'\cdots\text{O}2^i$	0.93	2.49	3.402 (3)	166
$\text{C}16-\text{H}16'\cdots\text{O}2^ii$	0.93	2.55	3.450 (3)	164
$\text{C}21-\text{H}21'\cdots\text{O}1$	0.93	2.44	3.250 (2)	145
$\text{C}28-\text{H}28'\cdots\text{O}1$	0.93	2.40	3.312 (2)	166

Symmetry codes: (i)  $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (ii)  $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ .

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012) and DIAMOND (Brandenburg, 2010); software used to prepare material for publication: SHELXL97 and publCIF (Westrip, 2010).

PBS is thankful to the Center for Research, Christ University, for financial assistance. NA thanks the University Grants Commission (India) for a Junior Research Fellowship. MRPK thanks the University Grants Commission, New Delhi, for a UGC-BSR one-time grant to faculty. The authors thank the Sophisticated Analytical Instruments Facility, Cochin University of S & T, for the diffraction measurements.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YK2100).

## References

- Brandenburg, K. (2010). DIAMOND. Crystal Impact GbR, Bonn, Germany.  
Bruker (2007). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
Heravi, M. M., Ranjbar, L., Derikvand, F., Oskooie, H. A. & Bamoharram, F. F. (2007). *J. Mol. Catal. A Chem.* **265**, 186–188.  
Mangalam, N. A. & Kurup, M. R. P. (2011). *Spectrochim. Acta Part A*, **76**, 22–28.  
Nair, Y., Sithambaresan, M. & Kurup, M. R. P. (2012). *Acta Cryst.* **E68**, o2709.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Sreeja, P. B., Kurup, M. R. P., Kishore, A. & Jasmin, C. (2004). *Polyhedron*, **23**, 575–581.  
Tamasi, G., Chiasserini, L., Savini, L., Sega, A. & Cini, R. (2005). *J. Inorg. Biochem.* **99**, 1347–1359.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supplementary materials

*Acta Cryst.* (2013). E69, o1828 [doi:10.1107/S1600536813031747]

***N'*-[*E*]-2-Fluorobenzylidene]benzohydrazide**

**P. B. Sreeja, M. Sithambaresan, N. Aiswarya and M. R. Prathapachandra Kurup**

**1. Comment**

The coordination chemistry of acyl and aroyl hydrazones have been a subject of competitive research as they are multipurpose class of ligands. Apart from exhibiting physiological and biological activities (Sreeja *et al.*, 2004), they also function as catalysts (Heravi *et al.*, 2007) as well as inhibitors for many enzymes (Tamasi *et al.*, 2005).

The title compound crystallizes in monoclinic space group  $P2_1/n$ . The asymmetric unit contains two molecules, both of which adopt the *E* configuration with respect to the C=N bond (Fig. 1). They exist in the amido form with a C8=O1 bond length of 1.231 (2) Å and C22=O2 of 1.223 (2) Å, which are very close to the reported C=O bond length in closely related structure (Nair *et al.*, 2012). Both molecules adopt the *Z* conformation with respect to the amido C—N bonds, with torsion angles of  $-7.2$  (3)° and  $5.5$  (3)°.

There are eight intermolecular hydrogen bonding interactions, out of which three are classical and the rest of interactions are non-classical with D...A distances of 2.875 (2), 3.339 (2), 2.882 (2), 3.213 (2), 3.402 (3), 3.450 (3), 3.250 (2) and 3.312 (2) Å (Table 1). The hydrogen at the N2 atom forms bifurcated H bonds with the O2 and N3 atoms of the adjacent molecules (Fig. 2). Two  $\pi$ - $\pi$  interactions between the rings C15—C20 & C23—C28 (shown in blue) and between the two C23—C28 rings (shown in pink) of the adjacent molecules with the intercentroid distances of 3.8840 (14) Å and 3.9145 (14) Å, respectively, also link the molecules (Fig. 3). Fig. 4 shows the packing of the molecules by means of hydrogen bonding and  $\pi$ - $\pi$  interactions.

**2. Experimental**


The title compound was prepared by adapting a reported procedure (Mangalam & Kurup, 2011). Benzoic acid hydrazide (1 mmol, 0.136 g) was dissolved in methanol and refluxed with methanolic solution of 2-fluorobenzaldehyde (1 mmol 0.124 g), in presence of a few drops of glacial acetic acid for 6 h. On cooling the reactant media, crystals of hydrazone were separated out. The crystals were filtered and washed with minimum quantity of methanol and dried over  $P_4O_{10}$  *in vacuo*. Good quality crystals suitable for X-ray analysis were obtained from methanolic solution by slow evaporation.

**3. Refinement**


All H atoms on C were placed in calculated positions, guided by difference maps, with C—H bond distances of 0.93 Å. H atoms were assigned  $U_{iso}(H)$  values of 1.2Ueq(carrier). H atoms of N2—H2' and N4—H4' bonds were located from difference maps and the bond distances are restrained to  $0.88 \pm 0.01$  Å.

**Computing details**

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *DIAMOND* (Brandenburg, 2010); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *pubCIF* (Westrip, 2010).



**Figure 1**

ORTEP diagram of *N'*-[(*E*)-(2-fluorophenyl)methylidene]benzohydrazide with 50% probability ellipsoids.



**Figure 2**

Hydrogen-bonding interactions of the title compound.

**Figure 3** $\pi$ - $\pi$  interactions in the title compound.

**Figure 4**

Packing diagram of the title compound along the *a* axis direction.

***N'*-'[(*E*)-2-Fluorobenzylidene]benzohydrazide***Crystal data*C<sub>14</sub>H<sub>11</sub>FN<sub>2</sub>O*M<sub>r</sub>* = 242.25Monoclinic, *P*2<sub>1</sub>/*n*Hall symbol: -*P* 2<sub>1</sub>*y**n**a* = 9.7010 (6) Å*b* = 17.4114 (13) Å*c* = 15.002 (1) Å

β = 104.126 (4)°

*V* = 2457.3 (3) Å<sup>3</sup>*Z* = 8*F*(000) = 1008*D<sub>x</sub>* = 1.310 Mg m<sup>-3</sup>Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 4473 reflections

θ = 4.7–48.2°

μ = 0.10 mm<sup>-1</sup>*T* = 296 K

Block, colourless

0.35 × 0.30 × 0.25 mm

*Data collection*

Bruker Kappa APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.33 pixels mm<sup>-1</sup>

ω and φ scan

Absorption correction: multi-scan

(SADABS; Bruker, 2007)

*T<sub>min</sub>* = 0.968, *T<sub>max</sub>* = 0.977

18766 measured reflections

6111 independent reflections

3320 reflections with *I* > 2σ(*I*)*R<sub>int</sub>* = 0.026θ<sub>max</sub> = 28.3°, θ<sub>min</sub> = 2.5°*h* = -12→11*k* = -23→17*l* = -19→19*Refinement*Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.050*wR*(*F*<sup>2</sup>) = 0.165*S* = 1.00

5972 reflections

333 parameters

2 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

*w* = 1/[σ<sup>2</sup>(*F<sub>o</sub>*<sup>2</sup>) + (0.0763*P*)<sup>2</sup> + 0.4072*P*]where *P* = (*F<sub>o</sub>*<sup>2</sup> + 2*F<sub>c</sub>*<sup>2</sup>)/3(Δ/σ)<sub>max</sub> = 0.001Δρ<sub>max</sub> = 0.21 e Å<sup>-3</sup>Δρ<sub>min</sub> = -0.17 e Å<sup>-3</sup>*Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of *F*<sup>2</sup> against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on *F*<sup>2</sup>, conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F*<sup>2</sup>. The threshold expression of *F*<sup>2</sup> > σ(*F*<sup>2</sup>) is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on *F*<sup>2</sup> are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.46744 (16)	0.52151 (9)	0.31008 (10)	0.0961 (5)
F2	0.04425 (16)	0.24110 (10)	0.23115 (9)	0.0973 (5)
O1	0.42500 (14)	0.14776 (8)	0.21005 (9)	0.0613 (4)
O2	0.28675 (14)	0.15829 (9)	-0.13577 (9)	0.0630 (4)
N1	0.44609 (15)	0.29885 (10)	0.23441 (10)	0.0513 (4)
N2	0.54248 (16)	0.24863 (9)	0.28584 (10)	0.0505 (4)
N3	0.16870 (15)	0.19149 (9)	0.00186 (10)	0.0490 (4)
N4	0.30586 (16)	0.16402 (10)	0.01524 (11)	0.0512 (4)
C1	0.3744 (2)	0.50230 (12)	0.23098 (15)	0.0591 (5)
C2	0.2916 (3)	0.55868 (13)	0.18222 (19)	0.0736 (6)
H2	0.3001	0.6092	0.2028	0.088*
C3	0.1957 (3)	0.53942 (15)	0.1024 (2)	0.0862 (8)
H3	0.1382	0.5770	0.0680	0.103*
C4	0.1845 (3)	0.46473 (16)	0.07311 (19)	0.0900 (8)
H4	0.1194	0.4517	0.0187	0.108*
C5	0.2685 (2)	0.40901 (13)	0.12344 (15)	0.0706 (6)
H5	0.2595	0.3585	0.1026	0.085*
C6	0.36690 (19)	0.42641 (11)	0.20489 (13)	0.0506 (5)
C7	0.46029 (19)	0.36878 (12)	0.25810 (13)	0.0539 (5)
H7	0.5304	0.3832	0.3095	0.065*
C8	0.52872 (18)	0.17362 (11)	0.26640 (11)	0.0464 (4)
C9	0.64558 (19)	0.12175 (11)	0.31359 (12)	0.0490 (4)
C10	0.6233 (3)	0.04460 (14)	0.30166 (17)	0.0771 (7)
H10	0.5351	0.0268	0.2689	0.093*
C11	0.7297 (3)	-0.00725 (16)	0.33747 (19)	0.0953 (9)
H11	0.7136	-0.0596	0.3280	0.114*
C12	0.8586 (3)	0.01828 (16)	0.38681 (18)	0.0840 (8)
H12	0.9303	-0.0166	0.4118	0.101*
C13	0.8819 (2)	0.09439 (17)	0.39922 (19)	0.0865 (8)
H13	0.9699	0.1119	0.4327	0.104*
C14	0.7759 (2)	0.14642 (14)	0.36265 (16)	0.0711 (6)
H14	0.7932	0.1988	0.3714	0.085*
C15	-0.0531 (2)	0.24951 (13)	0.15067 (14)	0.0599 (5)
C16	-0.1862 (3)	0.27441 (13)	0.15287 (17)	0.0698 (6)
H16	-0.2083	0.2854	0.2084	0.084*
C17	-0.2853 (2)	0.28265 (13)	0.07180 (19)	0.0710 (6)
H17	-0.3766	0.2990	0.0718	0.085*
C18	-0.2513 (2)	0.26703 (13)	-0.00981 (16)	0.0674 (6)
H18	-0.3194	0.2730	-0.0650	0.081*
C19	-0.1169 (2)	0.24265 (12)	-0.01023 (14)	0.0565 (5)
H19	-0.0948	0.2323	-0.0660	0.068*
C20	-0.01365 (19)	0.23326 (10)	0.07091 (13)	0.0488 (4)
C21	0.1288 (2)	0.20552 (11)	0.07427 (13)	0.0514 (5)
H21	0.1920	0.1979	0.1311	0.062*
C22	0.35624 (18)	0.14576 (10)	-0.05743 (12)	0.0461 (4)
C23	0.49959 (18)	0.11042 (10)	-0.03891 (12)	0.0476 (4)
C24	0.5548 (2)	0.09675 (15)	-0.11316 (16)	0.0715 (6)

H24	0.5039	0.1104	-0.1719	0.086*
C25	0.6874 (3)	0.06241 (17)	-0.1000 (2)	0.0918 (8)
H25	0.7244	0.0524	-0.1505	0.110*
C26	0.7637 (3)	0.04326 (15)	-0.0146 (2)	0.0873 (8)
H26	0.8531	0.0210	-0.0067	0.105*
C27	0.7102 (2)	0.05644 (14)	0.05883 (18)	0.0747 (7)
H27	0.7626	0.0433	0.1174	0.090*
C28	0.5777 (2)	0.08945 (12)	0.04711 (14)	0.0604 (5)
H28	0.5406	0.0976	0.0979	0.073*
H4'	0.3472 (19)	0.1550 (11)	0.0722 (7)	0.060 (6)*
H2'	0.6116 (18)	0.2682 (12)	0.3300 (12)	0.077 (7)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
F1	0.0962 (10)	0.0799 (10)	0.0961 (10)	-0.0050 (8)	-0.0077 (9)	-0.0251 (8)
F2	0.0970 (10)	0.1417 (14)	0.0495 (7)	0.0320 (9)	0.0106 (7)	-0.0092 (8)
O1	0.0577 (8)	0.0644 (9)	0.0510 (7)	-0.0008 (6)	-0.0076 (6)	-0.0029 (6)
O2	0.0597 (8)	0.0844 (11)	0.0418 (7)	0.0165 (7)	0.0064 (6)	0.0050 (7)
N1	0.0475 (9)	0.0575 (11)	0.0449 (8)	0.0057 (7)	0.0036 (7)	0.0017 (7)
N2	0.0483 (9)	0.0535 (10)	0.0430 (8)	0.0049 (7)	-0.0015 (7)	-0.0018 (7)
N3	0.0430 (8)	0.0565 (10)	0.0446 (8)	0.0079 (7)	0.0054 (7)	-0.0006 (7)
N4	0.0448 (8)	0.0657 (11)	0.0392 (8)	0.0139 (7)	0.0027 (7)	0.0006 (8)
C1	0.0513 (11)	0.0597 (14)	0.0664 (13)	-0.0037 (9)	0.0146 (10)	-0.0060 (11)
C2	0.0728 (15)	0.0510 (13)	0.1003 (18)	0.0042 (11)	0.0276 (14)	0.0002 (12)
C3	0.0769 (16)	0.0693 (17)	0.102 (2)	0.0155 (13)	0.0025 (15)	0.0153 (15)
C4	0.0893 (18)	0.0762 (18)	0.0841 (17)	0.0092 (14)	-0.0184 (14)	0.0068 (14)
C5	0.0721 (14)	0.0571 (14)	0.0712 (14)	0.0045 (11)	-0.0045 (12)	0.0013 (11)
C6	0.0443 (10)	0.0538 (12)	0.0536 (11)	0.0000 (8)	0.0120 (8)	0.0014 (9)
C7	0.0508 (11)	0.0570 (13)	0.0499 (11)	0.0003 (9)	0.0047 (8)	-0.0034 (9)
C8	0.0449 (10)	0.0579 (12)	0.0339 (9)	0.0026 (8)	0.0049 (7)	-0.0002 (8)
C9	0.0527 (11)	0.0529 (12)	0.0392 (9)	0.0048 (8)	0.0066 (8)	0.0013 (8)
C10	0.0804 (16)	0.0602 (15)	0.0766 (15)	0.0064 (11)	-0.0081 (13)	-0.0036 (12)
C11	0.115 (2)	0.0582 (16)	0.097 (2)	0.0206 (14)	-0.0042 (17)	0.0005 (14)
C12	0.0837 (18)	0.0819 (19)	0.0815 (17)	0.0333 (14)	0.0107 (14)	0.0127 (14)
C13	0.0576 (13)	0.091 (2)	0.0985 (19)	0.0129 (13)	-0.0052 (13)	0.0154 (16)
C14	0.0556 (12)	0.0655 (15)	0.0806 (15)	0.0029 (10)	-0.0058 (11)	0.0068 (12)
C15	0.0647 (13)	0.0636 (13)	0.0505 (11)	0.0088 (10)	0.0124 (10)	-0.0022 (10)
C16	0.0799 (16)	0.0693 (15)	0.0690 (14)	0.0092 (12)	0.0351 (13)	-0.0027 (11)
C17	0.0558 (13)	0.0658 (15)	0.0957 (18)	0.0097 (10)	0.0266 (13)	0.0038 (13)
C18	0.0555 (12)	0.0709 (15)	0.0712 (14)	0.0073 (10)	0.0064 (11)	0.0061 (12)
C19	0.0539 (11)	0.0628 (13)	0.0524 (11)	0.0068 (9)	0.0123 (9)	0.0023 (10)
C20	0.0527 (11)	0.0445 (11)	0.0491 (10)	0.0036 (8)	0.0121 (9)	-0.0005 (8)
C21	0.0500 (10)	0.0604 (12)	0.0407 (10)	0.0085 (9)	0.0052 (8)	0.0004 (9)
C22	0.0465 (10)	0.0460 (11)	0.0436 (10)	0.0009 (8)	0.0070 (8)	0.0003 (8)
C23	0.0442 (10)	0.0446 (11)	0.0535 (11)	-0.0015 (8)	0.0108 (8)	-0.0041 (8)
C24	0.0652 (14)	0.0886 (18)	0.0640 (13)	0.0070 (12)	0.0222 (11)	-0.0062 (12)
C25	0.0751 (17)	0.112 (2)	0.101 (2)	0.0148 (15)	0.0460 (16)	-0.0117 (17)
C26	0.0539 (14)	0.0893 (19)	0.121 (2)	0.0176 (12)	0.0263 (15)	0.0003 (17)
C27	0.0528 (12)	0.0787 (16)	0.0863 (17)	0.0179 (11)	0.0046 (12)	0.0043 (13)

---

C28	0.0532 (11)	0.0659 (14)	0.0599 (12)	0.0128 (10)	0.0091 (10)	0.0017 (10)
-----	-------------	-------------	-------------	-------------	-------------	-------------

---

*Geometric parameters (Å, °)*


---

F1—C1	1.346 (2)	C11—H11	0.9300
F2—C15	1.347 (2)	C12—C13	1.350 (4)
O1—C8	1.231 (2)	C12—H12	0.9300
O2—C22	1.223 (2)	C13—C14	1.380 (3)
N1—C7	1.266 (2)	C13—H13	0.9300
N1—N2	1.372 (2)	C14—H14	0.9300
N2—C8	1.338 (2)	C15—C16	1.370 (3)
N2—H2'	0.887 (9)	C15—C20	1.372 (3)
N3—C21	1.263 (2)	C16—C17	1.362 (3)
N3—N4	1.382 (2)	C16—H16	0.9300
N4—C22	1.337 (2)	C17—C18	1.371 (3)
N4—H4'	0.865 (9)	C17—H17	0.9300
C1—C2	1.363 (3)	C18—C19	1.373 (3)
C1—C6	1.375 (3)	C18—H18	0.9300
C2—C3	1.367 (4)	C19—C20	1.384 (3)
C2—H2	0.9300	C19—H19	0.9300
C3—C4	1.368 (4)	C20—C21	1.453 (3)
C3—H3	0.9300	C21—H21	0.9300
C4—C5	1.370 (3)	C22—C23	1.484 (2)
C4—H4	0.9300	C23—C24	1.370 (3)
C5—C6	1.388 (3)	C23—C28	1.377 (3)
C5—H5	0.9300	C24—C25	1.388 (3)
C6—C7	1.453 (3)	C24—H24	0.9300
C7—H7	0.9300	C25—C26	1.356 (4)
C8—C9	1.487 (2)	C25—H25	0.9300
C9—C10	1.365 (3)	C26—C27	1.347 (4)
C9—C14	1.368 (3)	C26—H26	0.9300
C10—C11	1.378 (3)	C27—C28	1.380 (3)
C10—H10	0.9300	C27—H27	0.9300
C11—C12	1.363 (4)	C28—H28	0.9300
<hr/>			
C7—N1—N2	116.26 (15)	C14—C13—H13	119.8
C8—N2—N1	118.79 (15)	C9—C14—C13	120.6 (2)
C8—N2—H2'	123.7 (15)	C9—C14—H14	119.7
N1—N2—H2'	117.5 (15)	C13—C14—H14	119.7
C21—N3—N4	115.38 (15)	F2—C15—C16	118.13 (19)
C22—N4—N3	119.63 (14)	F2—C15—C20	118.40 (18)
C22—N4—H4'	126.0 (13)	C16—C15—C20	123.5 (2)
N3—N4—H4'	113.9 (13)	C17—C16—C15	118.4 (2)
F1—C1—C2	118.3 (2)	C17—C16—H16	120.8
F1—C1—C6	117.84 (19)	C15—C16—H16	120.8
C2—C1—C6	123.8 (2)	C16—C17—C18	120.3 (2)
C1—C2—C3	118.7 (2)	C16—C17—H17	119.8
C1—C2—H2	120.7	C18—C17—H17	119.8
C3—C2—H2	120.7	C17—C18—C19	120.1 (2)
C2—C3—C4	119.9 (2)	C17—C18—H18	119.9

---



C2—C3—H3	120.1	C19—C18—H18	119.9
C4—C3—H3	120.1	C18—C19—C20	121.1 (2)
C3—C4—C5	120.4 (2)	C18—C19—H19	119.4
C3—C4—H4	119.8	C20—C19—H19	119.4
C5—C4—H4	119.8	C15—C20—C19	116.53 (18)
C4—C5—C6	121.3 (2)	C15—C20—C21	120.25 (17)
C4—C5—H5	119.4	C19—C20—C21	123.20 (18)
C6—C5—H5	119.4	N3—C21—C20	121.54 (16)
C1—C6—C5	115.95 (18)	N3—C21—H21	119.2
C1—C6—C7	121.65 (18)	C20—C21—H21	119.2
C5—C6—C7	122.37 (19)	O2—C22—N4	121.02 (16)
N1—C7—C6	119.94 (17)	O2—C22—C23	121.70 (17)
N1—C7—H7	120.0	N4—C22—C23	117.27 (15)
C6—C7—H7	120.0	C24—C23—C28	118.78 (18)
O1—C8—N2	121.89 (16)	C24—C23—C22	117.03 (17)
O1—C8—C9	120.36 (18)	C28—C23—C22	124.17 (18)
N2—C8—C9	117.73 (15)	C23—C24—C25	119.5 (2)
C10—C9—C14	118.38 (19)	C23—C24—H24	120.3
C10—C9—C8	117.23 (18)	C25—C24—H24	120.3
C14—C9—C8	124.25 (19)	C26—C25—C24	120.9 (3)
C9—C10—C11	120.9 (2)	C26—C25—H25	119.6
C9—C10—H10	119.5	C24—C25—H25	119.6
C11—C10—H10	119.5	C27—C26—C25	120.0 (2)
C12—C11—C10	119.9 (3)	C27—C26—H26	120.0
C12—C11—H11	120.0	C25—C26—H26	120.0
C10—C11—H11	120.0	C26—C27—C28	120.0 (2)
C13—C12—C11	119.7 (2)	C26—C27—H27	120.0
C13—C12—H12	120.2	C28—C27—H27	120.0
C11—C12—H12	120.2	C23—C28—C27	120.8 (2)
C12—C13—C14	120.5 (2)	C23—C28—H28	119.6
C12—C13—H13	119.8	C27—C28—H28	119.6
C7—N1—N2—C8	177.11 (17)	C12—C13—C14—C9	0.3 (4)
C21—N3—N4—C22	178.37 (18)	F2—C15—C16—C17	-179.7 (2)
F1—C1—C2—C3	-179.3 (2)	C20—C15—C16—C17	0.8 (4)
C6—C1—C2—C3	-0.3 (4)	C15—C16—C17—C18	-0.7 (4)
C1—C2—C3—C4	0.0 (4)	C16—C17—C18—C19	0.3 (4)
C2—C3—C4—C5	0.2 (5)	C17—C18—C19—C20	0.1 (3)
C3—C4—C5—C6	-0.1 (4)	F2—C15—C20—C19	179.99 (19)
F1—C1—C6—C5	179.36 (19)	C16—C15—C20—C19	-0.5 (3)
C2—C1—C6—C5	0.4 (3)	F2—C15—C20—C21	1.8 (3)
F1—C1—C6—C7	-2.7 (3)	C16—C15—C20—C21	-178.7 (2)
C2—C1—C6—C7	178.3 (2)	C18—C19—C20—C15	0.0 (3)
C4—C5—C6—C1	-0.2 (3)	C18—C19—C20—C21	178.2 (2)
C4—C5—C6—C7	-178.1 (2)	N4—N3—C21—C20	-178.78 (17)
N2—N1—C7—C6	177.19 (16)	C15—C20—C21—N3	-178.67 (19)
C1—C6—C7—N1	176.73 (19)	C19—C20—C21—N3	3.3 (3)
C5—C6—C7—N1	-5.5 (3)	N3—N4—C22—O2	5.5 (3)
N1—N2—C8—O1	-7.2 (3)	N3—N4—C22—C23	-174.97 (16)

N1—N2—C8—C9	171.27 (15)	O2—C22—C23—C24	4.0 (3)
O1—C8—C9—C10	-9.0 (3)	N4—C22—C23—C24	-175.59 (18)
N2—C8—C9—C10	172.51 (19)	O2—C22—C23—C28	-174.60 (19)
O1—C8—C9—C14	166.5 (2)	N4—C22—C23—C28	5.8 (3)
N2—C8—C9—C14	-12.0 (3)	C28—C23—C24—C25	-0.1 (3)
C14—C9—C10—C11	-0.5 (4)	C22—C23—C24—C25	-178.8 (2)
C8—C9—C10—C11	175.3 (2)	C23—C24—C25—C26	-0.9 (4)
C9—C10—C11—C12	1.0 (4)	C24—C25—C26—C27	1.0 (4)
C10—C11—C12—C13	-0.9 (4)	C25—C26—C27—C28	0.0 (4)
C11—C12—C13—C14	0.3 (4)	C24—C23—C28—C27	1.1 (3)
C10—C9—C14—C13	-0.1 (4)	C22—C23—C28—C27	179.6 (2)
C8—C9—C14—C13	-175.6 (2)	C26—C27—C28—C23	-1.0 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2'...O2 <sup>i</sup>	0.89 (1)	2.09 (2)	2.875 (2)	147 (2)
N2—H2'...N3 <sup>i</sup>	0.89 (1)	2.60 (2)	3.339 (2)	142 (2)
N4—H4'...O1	0.87 (1)	2.03 (1)	2.882 (2)	171 (2)
C7—H7...O2 <sup>i</sup>	0.93	2.53	3.213 (2)	131
C14—H14...O2 <sup>i</sup>	0.93	2.49	3.402 (3)	166
C16—H16...O2 <sup>ii</sup>	0.93	2.55	3.450 (3)	164
C21—H21...O1	0.93	2.44	3.250 (2)	145
C28—H28...O1	0.93	2.40	3.312 (2)	166

Symmetry codes: (i)  $x+1/2, -y+1/2, z+1/2$ ; (ii)  $x-1/2, -y+1/2, z+1/2$ .