## Corrigendum

## mRNA secondary structure optimization using a correlated stem-loop prediction

Paulo Gaspar<sup>1,\*</sup>, Gabriela Moura<sup>2</sup>, Manuel A.S. Santos<sup>2</sup> and José Luís Oliveira<sup>1</sup>

<sup>1</sup>DETI/IEETA, University of Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal and <sup>2</sup>Department of Biology and CESAM, University of Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

Nucl. Acids Res. (2013) 41 (6): e73. doi: 10.1093/nar/gks1473

The authors wish to acknowledge additional funding sources. An updated Funding statement is provided below.

## **FUNDING**

European FP7 projects GEN2PHEN and MEPHITIS; FEDER funds through the Programa Operacional Factores de Competitividade – COMPETE (FCOMP-01–0124-FEDER-014284) and by National Funds through FCT - Foundation for Science and Technology in the scope of the research project ref. PTDC/BIA-GEN/110383/2009; P.G. is funded by FCT-SFRH/BD/71063/2010; Funding for open access charge: GEN2PHEN.

<sup>\*</sup>To whom correspondence should be addressed. Tel: +351 234 370 523; Fax: +351 234 370 545; Email: paulogaspar@ua.pt