

Acute mental stress and surgical performance

P. D. Grantcharov¹ , T. Boillat^{1,2}, S. Elkabany³, K. Wac^{1,4,5} and H. Rivas¹

¹Section of Bariatric and Minimally Invasive Surgery, Department of Surgery, Stanford University School of Medicine, Stanford, California, USA,

²Department of Computer Science, Lucerne University of Applied Sciences and Arts, Lucerne, Switzerland, ³International Centre for Surgical Safety, Keenan Institute for Biomedical Science, St Michael's Hospital, Toronto, Ontario, Canada, ⁴Quality of Life Technologies Laboratory, University of Geneva, Geneva, Switzerland, and University of Copenhagen, Copenhagen, Denmark, and ⁵Department of Computer Science, University of Copenhagen, Copenhagen, Denmark

Correspondence to: Mr P. D. Grantcharov, Section of Bariatric and Minimally Invasive Surgery, Department of Surgery, Stanford University School of Medicine, 291 Campus Drive, Stanford, California 94305, USA (e-mail: peter.grantcharov@gmail.com)

Background: Stress has been shown to impact adversely on multiple facets critical to optimal performance. Advancements in wearable technology can reduce barriers to observing stress during surgery. This study aimed to investigate the association between acute intraoperative mental stress and technical surgical performance.

Methods: Continuous electrocardiogram data for a single attending surgeon were captured during surgical procedures to obtain heart rate variability (HRV) measures that were used as a proxy for acute mental stress. Two different measures were used: root mean square of successive differences (RMSSD) and standard deviation of RR intervals (SDNN). Technical surgical performance was assessed on the Operating Room Black Box[®] platform using the Generic Error Rating Tool (GERT). Both HRV recording and procedure video recording were time-stamped. Surgical procedures were fragmented to non-overlapping intervals of 1, 2 and 5 min, and subjected to data analysis. An event was defined as any deviation that caused injury to the patient or posed a risk of harm.

Results: Rates of events were significantly higher (47–66 per cent higher) in the higher stress quantiles than in the lower stress quantiles for all measured interval lengths using both proxy measures for acute mental stress. The strongest association was observed using 1-min intervals with RMSSD as the HRV measure ($P < 0.001$).

Conclusion: There is an association between measures of acute mental stress and worse technical surgical performance. Further study will help delineate the interdependence of these variables and identify triggers for increased stress levels to improve surgical safety.

Funding information

No funding

Paper accepted 6 August 2018

Published online 27 September 2018 in Wiley Online Library (www.bjsopen.com). DOI: 10.1002/bjs.104

Introduction

The relationship between acute stress and performance is a well studied psychophysiological phenomenon in several high-performance industries^{1,2}. This relationship has also received some attention for surgeons and other medical professionals, but it has been focused primarily on the prevalence of burnout and other markers of chronic stress³. As acute mental stress may have a more direct relationship to surgical performance than chronic stress, a better understanding of the factors influencing the variations of surgical performance and clinical outcomes is required.

During resting conditions, there is a constant interplay between the sympathetic and parasympathetic divisions of

the autonomic nervous system (ANS). A mentally stressful event can be defined as an instance during which a balance shift in favour of the sympathetic branch (also known as the fight-or-flight response) occurs – an event often referred to as distress⁴. In the past few decades, heart rate variability (HRV) measures have been used to assess this phenomenon given their ease of measurability, the basic calculations required, and their sufficient sensitivity to observe the nearly instantaneous changes in the ANS balance.

Acute stress has been found both to facilitate and to impair performance in a variety of contexts; however, complex cognitive processing and functions, where stressors occur peripherally to the primary task, are generally

found to be impacted negatively by increased stress⁵. This is particularly relevant to the surgical environment, where stress and stressors have been shown to adversely impact functions such as decision-making^{6–12} and team performance^{12–14}. Studies have observed stress in the live operating theatre using a variety of methods^{12,15–26}, but limitations in ECG equipment and stress measurement methods have hindered research. Efforts to investigate this relationship have generally been limited to virtual environments^{27–31}. Although these studies have provided valuable insight, surgical simulations lack realism in comparison with the real surgical environment when imitating a stress scenario^{31,32}. Significant advances in wearable technology, particularly the ability reliably to measure physiological metrics unobtrusively without wired sensors, afford new opportunities to overcome these obstacles.

Standards to assess technical surgical performance^{33,34} and objective methods to estimate acute stress with ECG measurements^{35–37} have existed for decades. Despite this, no studies have employed these tools in the live surgical setting to assess the relationship between intraoperative acute mental stress levels and technical surgical performance³⁸. The aim of the present study was to investigate the association between acute intraoperative mental stress and surgical performance.

Methods

Study design

Data collection for this observational study was conducted at Stanford Hospital (Stanford, California, USA) during May and June 2017. Stress and surgical performance data were collected for one attending surgeon during this period. The study was granted institutional review board approval by Stanford University. Patient consent was not required.

Data capture

Physiological data to be used in calculations for proxy measures of acute stress were captured for the attending surgeon during each surgical procedure using the wireless Hexoskin Smart Shirt (Hexoskin, Montreal, Quebec, Canada). The Hexoskin Smart Shirt has built-in sensors that capture 256-Hz ECG data, three-dimensional 64-Hz acceleration data and 128-Hz breathing data, all of which have been validated previously³⁹. All raw data are directly downloadable from the Hexoskin online portal, with all data points time-stamped to the Coordinated Universal Time (UTC). The 256-Hz ECG capture satisfies the

recommended resolution to capture precise data to allow for HRV interpretations as set forth by the Task Force of the European Society of Cardiology and the North American Society of Pacing Electrophysiology⁴⁰.

Consecutive elective surgical procedures, both laparoscopic and endoscopic, performed between 9 May and 28 June 2017, were included in the study. Each procedure was recorded and entirely anonymized of all markers (hospital information, physician identifiers, patient data). Video recordings were time-stamped to UTC to allow for synchronization with the HRV data captured by the Hexoskin Smart Shirt. Recordings were assessed independently and blindly by a single observer for technical laparoscopic surgical performance using the Operating Room Black Box[®] platform (SST, Toronto, Ontario, Canada), with a previously validated measurement tool (Generic Error Rating Tool (GERT)⁴¹).

Stress (heart rate variability) analysis

Although it has been recommended that both time domain and frequency domain statistics be incorporated when analysing HRV, the two statistical methods correlate with one another to such a strong degree that they can be used as surrogates for each other^{42,43}. Further, time domain methods, particularly RMSSD (defined below), have been shown to have greater reliability when measuring short-term HRV than frequency domain methods⁴⁴. The following conventional time domain methods of analysing HRV were therefore selected for this study (for formulas see *Appendix S1*, supporting information): the square root of the mean sum of squares of ΔRR intervals (RMSSD) and standard deviation of RR intervals (SDNN).

As sympathetic nervous system activation begins to supersede parasympathetic nervous system activation, there becomes less interplay among these two ANS divisions, which manifests as a near-instantaneous decrease in RR-interval variation (lower HRV). As such, lower values for SDNN and RMSSD suggest greater levels of sympathetic nervous system dominance (hereafter referred to as stress).

The Hexoskin ECG processing software provides a filter for data curation of RR intervals that excludes abnormal RR intervals. These may be due to irregular heartbeats or suboptimal ECG signals.

Surgical performance

Technical laparoscopic surgical performance was evaluated with the GERT, a previously validated measurement tool.




Fig. 1 Surgical event rates during different stress levels: **a** 1-min interval, **b** 2-min interval, **c** 5-min interval. SDNN, standard deviation of RR intervals; RMSSD, root mean square of successive differences in RR intervals

The GERT allows for objective and reliable (interrater intraclass correlation coefficient for events 0.85) assessment of operative performance during laparoscopic procedures. Using this framework, both surgical errors and events are identified, with errors defined as ‘the mechanism of unintended or deviated technical task execution’ and events as ‘any deviation that causes injury to the patient or poses a risk of harm’^{41,45}. For the purposes of this study, only ‘events’ were selected as an outcome measure of surgical performance, because they occur less commonly and have a more direct relationship with clinically relevant outcomes. Examples of events include serosal tears arising from grasper slips from bowel, bleeding from needle puncturing vessel, and burns from inadvertent touching of other structures with an energy device⁴¹.

Data analysis

Software was developed by the authors to derive the HRV statistics from the ECG data files, process the surgical performance data and synchronize all data streams into the prespecified time intervals. The first interval in a procedure commenced the moment after the conclusion of the presurgery ‘time out’ checklist where video capture had begun. Data capture concluded at the end of the surgery when all surgical instruments were removed from the patient.

There is some debate in the literature regarding the optimal time intervals to ensure valid HRV recordings, with the Task Force of the European Society of Cardiology and the North American Society of Pacing Electrophysiology⁴⁰ recommending that short-term HRV

analysis be undertaken in intervals of 5 min, and others⁴⁶ suggesting that accurate HRV recordings can be taken in intervals as low as 10 s. Consideration was also given to the fact that data streams were quite distinct. For example, although physiological data were measured on a pseudo-continuous timeline (ECG readings at 256 Hz), HRV was measured in discrete intervals and surgical performance on a continuous basis. Thus, for the exploratory purposes of this study, the judgement was made to measure both data streams (HRV measurements and surgical performance) in non-overlapping intervals of 1, 2 and 5 min to best accommodate these differences.

Stress data were divided into quantiles to allow for comparison between low and high stress intervals. Five-minute intervals were presented in quintiles, rather than tertiles, owing to the reduced sample variance resulting from the increased interval length, as stress peaks may last only a fraction of the duration of an interval.

Statistical analysis

All statistical analyses were performed using Microsoft Excel[®] version 16 (Microsoft, Redmond, Washington, USA), SPSS[®] version 24 (IBM, Armonk, New York, USA) and MATLAB version 9.2 (MathWorks, Natick, Massachusetts, USA).

Stress data were assumed to be normally distributed, so comparisons between groups were performed with independent two-sample Student’s *t* tests. Surgical performance (events) was observed not to be normally distributed; therefore comparisons between groups were performed with Mann–Whitney *U* tests. $P < 0.050$ was considered statistically significant.

Table 1 Changes in heart rate variability measure between intervals with and without a surgical event

Interval	HRV measure	Mean HRV value (ms)		Difference (%)	Student's <i>t</i> test		
		0 events	≥ 1 event		<i>t</i> statistic	d.f.	<i>P</i>
1 min	SDNN	37.0	34.4	-7	2.83 (0.80, 4.44)	1452	0.005
	RMSSD	24.8	22.6	-9	2.10 (0.14, 4.30)	1452	0.036
2 min	SDNN	40.4	38.1	-6	2.47 (0.47, 4.13)	731	0.014
	RMSSD	25.3	25.5	+1	0.20 (-2.17, 2.65)	413	0.846
5 min	SDNN	43.7	41.9	-4	1.58 (-0.43, 4.09)	293	0.113
	RMSSD	27.4	25.9	-5	0.97 (-1.54, 4.49)	293	0.335

Values in parentheses are 95 per cent confidence intervals. HRV, heart rate variability; SDNN, standard deviation of RR intervals; RMSSD, root mean square of successive differences in RR intervals.

Results

Data were captured for gastric bypass (12 patients), sleeve gastrectomy (7) and peroral endoscopic myotomy (6), with a mean(s.d.) duration of 96.7(35.5) min for all procedures. As the rating framework for assessing surgical performance was tailored for laparoscopic procedures, surgical performance data were observed only for gastric bypass and sleeve gastrectomy. Stress data were observed for all procedures.

Mean(s.d.) SDNN values were 38.0(14.4) ms for 1-min intervals, 40.9(12.2) ms for 2-min intervals and 43.7(9.7) ms for 5-min intervals. Respective mean(s.d.) RMSSD values were 25.2(16.5), 26.2(14.5) and 27.1(12.5) ms. The mean(s.d.) number of events per procedure was 23.8(11.3).

Stress and surgical performance

For all time intervals there was a trend for an increasing rate of events with increasing stress levels (*Fig. 1*). For 1-, 2- and 5-min intervals, the highest stress (SDNN) quantiles had event rates that were 55.5, 62.5 and 66.1 per cent respectively greater than those of the lowest stress quantiles, all with statistically significant differences (1 min: $P = 0.002$; 2 min: $P = 0.011$; 5 min: $P = 0.028$). The same trend was observed with RMSSD as the proxy for stress, with the highest stress quantiles having 57.7, 46.8 and 54.7 per cent greater event rates than the lowest stress quantiles for each respective interval (1 min: $P < 0.001$; 2 min: $P = 0.045$; 5 min: $P = 0.046$).

When observing the inverse relationship, where mean HRV values were calculated for intervals with and without events, a similar trend was observed (*Table 1*). With the exception of RMSSD for 2-min intervals, HRV was lower (stress was greater) as measured by both SDNN and RMSSD during intervals where an event was observed *versus* intervals where an event was not observed. Levene's test for equality of variances showed equal variances for each test except for 2-min RMSSD.

Discussion

This study has demonstrated a direct relationship between acute mental stress and surgical performance in a live operating theatre. The results indicate a clear association between the occurrence of events and the level of stress experienced, with significant differences between the higher and lower stress quantiles being observed for all measured time intervals. Greater stress responses were also found during intervals where an event was observed, for all measures except the 2-min interval RMSSD.

There are several possible explanations for this finding. One may be that increased stress levels in surgeons occur in response to the occurrence of events. Although this is likely to be true, it does not preclude the possible dependency of surgical performance on stress levels. Not all events are of equal severity and are therefore not necessarily recognized by the surgeon as a deviation that causes injury or poses a risk to the patient. Further, events that are recognized by the surgeon may not manifest in stress responses if they are perceived to be innocuous and easily resolved (for example, minor bleeding). In addition, the possibility of a snowball effect cannot be precluded, whereby an event triggers a stress response that could have a negative impact on surgical performance resulting in subsequent errors and events.

Based on previous research on the effects of acute mental stress^{12,27,28,30,31} and the observations in this study, acute mental stress and surgical performance are likely to be dependent upon each other to some degree. The next step towards better understanding of this relationship would be a study of temporality. For obvious ethical reasons, this cannot be performed in the operating theatre by experimental means (that is, inducing a stress response in a surgeon and observing surgical performance), and therefore would have to be done via novel methods of interval analysis (if incorporating HRV). In this study, the intervals were predetermined from the moment that the video recording commenced, thereby introducing several limitations.

The true relationships may be more pronounced than has been found in this study. Stress responses vary greatly in length depending on the situation and, unless this stressed state fell perfectly within the bounds of an interval, the stressed state would either be diluted with more relaxed HRV data (if the stress response were shorter than the interval length) or be only partly captured (if longer than the interval length). Observing at three different interval times was an attempt to control for this, but these methods could still be improved. In addition, an event occurring near the end of an interval would be unlikely to prompt an increased stress response during that same interval. Conversely, compromised surgical skill because of a raised stress response near the end of an interval may not trigger events until the succeeding interval. Similarly, if the event occurred at the very beginning of an interval, any preceding stress increase that may have been a factor would not be grouped into the same interval.

Despite these limitations, the observed trends of the potential effect of stress on the occurrence of events should trigger further research. In particular, it would be helpful to study stress in intervals on a 'sliding window' that moves along a continuous timeline rather than fixed pre-determined intervals. The width of this window should be flexible to accommodate varying lengths of stress response. In addition, it would be interesting to study the temporality of the relationship between stress and surgical performance by comparing stress levels during the intervals preceding and succeeding events with baseline periods.

The clinical relevance of exploring variables associated with surgical events is evident. Stress can be induced in surgeons and other members of the operating team for a variety of reasons, many of which are unavoidable. It is therefore beneficial to distinguish between stress that manifests as a result of surgical events that demand increased concentration, and stress that arises as a result of external influences. The latter can be addressed by initiatives that identify and reduce such external events, whereas managing the former, which may or may not be harmful for surgical performance depending on the specific circumstance, is more complex.

It is important to explore methods, either training- or intervention-based, that diminish the occurrence, severity and potential consequences of this unavoidable phenomenon of surgical stress. Previous research^{29,47} has shown promise in this area and, given the results of the present observational study and the clear association between stress levels and surgical performance, the implementation of intervention- or training-based initiatives aimed at reducing stress in surgeons may be a viable way to reduce surgical events and adverse outcomes. Further

study is also necessary to gauge more precisely the triggers for increased stress, their individual effects on surgical performance, and the severity of this relationship.

Disclosure

The authors declare no conflict of interest.

References

- 1 Sexton JB, Thomas EJ, Helmreich RL. Error, stress, and teamwork in medicine and aviation: cross sectional surveys. *BMJ* 2000; **320**: 745–749.
- 2 Hammermeister J, Burton D. Stress, appraisal, and coping revisited: examining the antecedents of competitive state anxiety with endurance athletes. *Sport Psychol* 2001; **15**: 66–90.
- 3 Campbell DA Jr, Sonnad SS, Eckhauser FE, Campbell KK, Greenfield LJ. Burnout among American surgeons. *Surgery* 2001; **130**: 696–705.
- 4 Tomaka J, Blascovich J, Kelsey RM, Leitten CL. Subjective, physiological, and behavioral effects of threat and challenge appraisal. *J Pers Soc Psychol* 1993; **65**: 248–260.
- 5 LeBlanc VR. The effects of acute stress on performance: implications for health professions education. *Acad Med* 2009; **84**(Suppl): S25–S33.
- 6 FeldmanHall O, Raio CM, Kubota JT, Seiler MG, Phelps EA. The effects of social context and acute stress on decision making under uncertainty. *Psychol Sci* 2015; **26**: 1918–1926.
- 7 Wemm SE, Wulfert E. Effects of acute stress on decision making. *Appl Psychophysiol Biofeedback* 2017; **42**: 1–12.
- 8 Cano-López I, Cano-López B, Hidalgo V, González-Bono E. Effects of acute stress on decision making under ambiguous and risky conditions in healthy young men. *Span J Psychol* 2016; **19**: E59.
- 9 Morgado P, Sousa N, Cerqueira JJ. The impact of stress in decision making in the context of uncertainty. *J Neurosci Res* 2015; **93**: 839–847.
- 10 Keinan G, Friedland N, Ben-Porath Y. Decision making under stress: scanning of alternatives under physical threat. *Acta Psychol (Amst)* 1987; **64**: 219–228.
- 11 Cumming SR, Harris LM. The impact of anxiety on the accuracy of diagnostic decision-making. *Stress Health* 2001; **17**: 281–286.
- 12 Wetzel CM, Kneebone RL, Woloshynowych M, Nestel D, Moorthy K, Kidd J *et al*. The effects of stress on surgical performance. *Am J Surg* 2006; **191**: 5–10.
- 13 Ellis APJ. System breakdown: the role of mental models and transactive memory in the relationship between acute stress and team performance. *Acad Manage J* 2006; **49**: 576–589.
- 14 Driskell JE, Salas E, Johnston J. Does stress lead to a loss of team perspective? *Group Dyn Theory Res Pract* 1999; **3**: 291–302.
- 15 Böhm B, Rötting N, Schwenk W, Grebe S, Mansmann U. A prospective randomized trial on heart rate variability of the

- surgical team during laparoscopic and conventional sigmoid resection. *Arch Surg* 2001; **136**: 305–310.
- 16 Demirtas Y, Tulmac M, Yavuzer R, Yalcin R, Ayhan S, Latifoglu O *et al*. Plastic surgeon's life: marvelous for mind, exhausting for body. *Plast Reconstr Surg* 2004; **114**: 923–931.
 - 17 Arora S, Tierney T, Sevdalis N, Aggarwal R, Nestel D, Woloshynowych M *et al*. The imperial stress assessment tool (ISAT): a feasible, reliable and valid approach to measuring stress in the operating room. *World J Surg* 2010; **34**: 1756–1763.
 - 18 Yamamoto A, Hara T, Kikuchi K, Hara T, Fujiwara T. Intraoperative stress experienced by surgeons and assistants. *Ophthalmic Surg Lasers* 1999; **30**: 27–30.
 - 19 Kuhn EW, Choi YH, Schönherr M, Liakopoulos OJ, Rahmanian PB, Choi CY *et al*. Intraoperative stress in cardiac surgery: attendings *versus* residents. *J Surg Res* 2013; **182**: e43–e49.
 - 20 Smith WD, Berguer R, Nguyen NT. Monitor height affects surgeons' stress level and performance on minimally invasive surgery tasks. *Stud Health Technol Inform* 2005; **111**: 498–501.
 - 21 Rieger A, Stoll R, Kreuzfeld S, Behrens K, Weippert M. Heart rate and heart rate variability as indirect markers of surgeons' intraoperative stress. *Int Arch Occup Environ Health* 2014; **87**: 165–174.
 - 22 Jones KI, Amawi F, Bhalla A, Peacock O, Williams JP, Lund JN. Assessing surgeon stress when operating using heart rate variability and the State Trait Anxiety Inventory: will surgery be the death of us? *Colorectal Dis* 2015; **17**: 335–341.
 - 23 Song MH, Tokuda Y, Nakayama T, Sato M, Hattori K. Intraoperative heart rate variability of a cardiac surgeon himself in coronary artery bypass grafting surgery. *Interact Cardiovasc Thorac Surg* 2009; **8**: 639–641.
 - 24 Prichard RS, O'Neill CJ, Oucharek JJ, Holmes CY, Delbridge LW, Sywak MS. A prospective study of heart rate variability in endocrine surgery: surgical training increases consultant's mental strain. *J Surg Educ* 2012; **69**: 453–458.
 - 25 Sadiq SA, Arif A, Usmani HA. The stress of cataract surgery. *J Clin Exp Ophthalmol* 2013; **4**: 301.
 - 26 Mongin C, Dufour F, Lattanzio F, Champault G. [Evaluation of stress in surgical trainees: prospective study of heart rate during laparoscopic cholecystectomy]. *J Chir (Paris)* 2008; **145**: 138–142.
 - 27 Wetzel CM, Black SA, Hanna GB, Athanasiou T, Kneebone RL, Nestel D *et al*. The effects of stress and coping on surgical performance during simulations. *Ann Surg* 2010; **251**: 171–176.
 - 28 Hassan I, Weyers P, Maschuw K, Dick B, Gerdes B, Rothmund M *et al*. Negative stress-coping strategies among novices in surgery correlate with poor virtual laparoscopic performance. *Br J Surg* 2006; **93**: 1554–1559.
 - 29 Wetzel CM, George A, Hanna GB, Athanasiou T, Black SA, Kneebone RL *et al*. Stress management training for surgeons – a randomized, controlled, intervention study. *Ann Surg* 2011; **253**: 488–494.
 - 30 Moorthy K, Munz Y, Dosis A, Bann S, Darzi A. The effect of stress-inducing conditions on the performance of a laparoscopic task. *Surg Endosc* 2003; **17**: 1481–1484.
 - 31 Schuetz M, Gockel I, Beardi J, Hakman P, Dunschede F, Moenk S *et al*. Three different types of surgeon-specific stress reactions identified by laparoscopic simulation in a virtual scenario. *Surg Endosc* 2008; **22**: 1263–1267.
 - 32 Baker BG, Bhalla A, Doleman B, Yarnold E, Simons S, Lund JN *et al*. Simulation fails to replicate stress in trainees performing a technical procedure in the clinical environment. *Med Teach* 2017; **39**: 53–57.
 - 33 Martin JA, Regehr G, Reznick R, MacRae H, Murnaghan J, Hutchison C *et al*. Objective structured assessment of technical skill (OSATS) for surgical residents. *Br J Surg* 1997; **84**: 273–278.
 - 34 Datta V, Bann S, Mandalia M, Darzi A. The surgical efficiency score: a feasible, reliable, and valid method of skills assessment. *Am J Surg* 2006; **192**: 372–378.
 - 35 Bootsma M, Swenne CA, Van Bolhuis HH, Chang PC, Cats VM, Brusckhe AV. Heart rate and heart rate variability as indexes of sympathovagal balance. *Am J Physiol* 1994; **266**: H1565–H1571.
 - 36 Pagani M, Furlan R, Pizzinelli P, Crivellaro W, Cerutti S, Malliani A. Spectral analysis of R-R and arterial pressure variabilities to assess sympatho-vagal interaction during mental stress in humans. *J Hypertens Suppl* 1989; **7**: S14–S15.
 - 37 Hitchen M, Brodie DA, Harness JB. Cardiac responses to demanding mental load. *Ergonomics* 1980; **23**: 379–385.
 - 38 Arora S, Sevdalis N, Nestel D, Woloshynowych M, Darzi A, Kneebone R. The impact of stress on surgical performance: a systematic review of the literature. *Surgery* 2010; **147**: 318–330.e6.
 - 39 Villar R, Beltrame T, Hughson RL. Validation of the Hexoskin wearable vest during lying, sitting, standing, and walking activities. *Appl Physiol Nutr Metab* 2015; **40**: 1019–1024.
 - 40 Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. Heart rate variability: standards of measurement, physiological interpretation and clinical use. *Circulation* 1996; **93**: 1043–1065.
 - 41 Bonrath EM, Zevin B, Dedy NJ, Grantcharov TP. Error rating tool to identify and analyse technical errors and events in laparoscopic surgery. *Br J Surg* 2013; **100**: 1080–1088.
 - 42 Massin MM, Derkenne B, von Bernuth G. Correlations between indices of heart rate variability in healthy children and children with congenital heart disease. *Cardiology* 1999; **91**: 109–113.
 - 43 Bigger JT Jr, Fleiss JL, Steinman RC, Rolnitzky LM, Kleiger RE, Rottman JN. Correlations among time and frequency domain measures of heart period variability two weeks after acute myocardial infarction. *Am J Cardiol* 1992; **69**: 891–898.

- 44 Cipryan L. Within-session stability of short-term heart rate variability measurement. *J Hum Kinet* 2016; **50**: 85–92.
- 45 World Health Organization. *World Alliance for Patient Safety: WHO Draft Guidelines for Adverse Event Reporting and Learning Systems: From Information to Action*; 2005. <http://apps.who.int/iris/handle/10665/69797> [accessed 21 January 2018].
- 46 Munoz ML, van Roon A, Riese H, Thio C, Oostenbroek E, Westrik I *et al.* Validity of (ultra-)short recordings for heart rate variability measurements. *PLoS One* 2015; **10**: e0138921.
- 47 Anton NE, Howley LD, Pimentel M, Davis CK, Brown C, Stefanidis D. Effectiveness of a mental skills curriculum to reduce novices' stress. *J Surg Res* 2016; **206**: 199–205.

Supporting information

Additional supporting information can be found online in the Supporting Information section at the end of the article.