






Mitochondrial phylogenomics of pygmy grasshoppers (Orthoptera, Tetrigidae, Metrodorinae): descriptions of a new genus, two new species, and new synonyms from China

Yuemei Li^{1,2,3}, Shixiong Leng⁴, Jiasong He⁴, Weian Deng^{1,2,3,5}, Delong Guan^{5,6}

¹ Key Laboratory of Ecology of Rare and Endangered Species and Environmental Protection, Guangxi Normal University, Ministry of Education, Guilin 541006, China

² Guangxi Key Laboratory of Rare and Endangered Animal Ecology, Guangxi Normal University, Guilin 541006, China

³ College of Life Sciences, Guangxi Normal University, Guilin 541006, China

⁴ Guangxi Daguishan Crocodile Lizard National Nature Reserve, Hezhou, Guangxi 542824, China

⁵ Guangxi Key Laboratory of Sericulture ecology and Applied intelligent technology, Hechi University, Hechi, China

⁶ School of Chemistry and Bioengineering, Hechi University, Yizhou, Guangxi 546300, China

Corresponding authors: Weian Deng (dengweian5899@163.com); Delong Guan (2023660006@hcnu.edu.cn)



Academic editor: Zhu-Qing He

Received: 4 January 2025

Accepted: 7 March 2025

Published: 5 May 2025

ZooBank: <https://zoobank.org/7B675E54-601E-42D2-8B31-91C009B6A80F>

Citation: Li Y, Leng S, He J, Deng W, Guan D (2025) Mitochondrial phylogenomics of pygmy grasshoppers (Orthoptera, Tetrigidae, Metrodorinae): descriptions of a new genus, two new species, and new synonyms from China. ZooKeys 1236: 249–281. <https://doi.org/10.3897/zookeys.1236.145914>

Copyright: © Yuemei Li et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

The Chinese wingless brachypronotal pygmy grasshoppers of the subfamily Metrodorinae have often been classified within the genus *Macromotettixoides*. In this study, two undescribed species of wingless brachypronotal pygmy grasshoppers belonging to Metrodorinae were collected. To elucidate their taxonomic positions, the complete mitochondrial genomes of these two species were sequenced and analyzed. Phylogenetic analyses were conducted using 13 protein-coding genes (PCGs) from 28 tetrigid mitogenomes. Genetic distances and divergence times were estimated. Our results indicate that one species represents a new genus within Metrodorinae, while the other is a new species of *Macromotettixoides*. Consequently, a new genus and two new species of Metrodorinae from China are described and illustrated: *Edentatettix* Deng, **gen. nov.** containing *Edentatettix leyeensis* Deng, **sp. nov.**, and *Macromotettixoides yaana* Deng, **sp. nov.** Additionally, five new synonyms are proposed: *Hainantettix angustivertex* (Zha & Peng, 2021), **syn. nov.** and *Macromotettixoides angustivertex* Zha & Peng, 2021, **syn. nov.** of *Hainantettix strictivertex* Deng, 2020; *Hyboella badagongshanensis* Zheng, 2013, **syn. nov.**, *Macromotettixoides badagongshanensis* (Zheng, 2013), **syn. nov.**, and *Macromotettixoides wuyishana* Zheng, 2013, **syn. nov.** of *Macromotettixoides jiuwan-shanensis* Zheng, Wei & Jiang, 2005. For the first time, edentate ovipositors constituting an important taxonomic character within Tetrigidae is reported and discussed.

Key words: *Macromotettixoides*, Metrodoridae, mitochondrial genome, new taxa, ovipositor, phylogeny, taxonomy

Introduction

In the highly diverse orthopteran insects, tetrigids (Orthoptera: Tetrigidae) represent a relatively ancient group of orthopteran insects. Among them, Metrodorinae Bolívar, [1887] is one of the three largest subfamilies in Tetrigidae and currently includes 105 genera containing more than 648 species distributed

around the world (Deng 2016; Cigliano et al. 2024), with 16 genera (including the new genus *Edentatettix* Deng, gen. nov.) found in China. Although Metrodorinae has a wide distribution in the world, there are still many species with classification disputes due to the similarity and diversity of morphological characteristics. Since the establishment of Metrodorinae, numerous species within it have undergone transfers between genera, and the genera within the subfamily have been revised frequently (Storozhenko 2014; Tumbrinck and Telnov 2014; Tumbrinck 2019; Skejo et al. 2019; Peng et al. 2021; Kasalo 2022; Kasalo et al. 2023b).

Pygmy grasshoppers (Tetrigidae) is a taxonomic challenging group, exhibiting striking polymorphism in various morphological traits such as body color, patterns, wing lengths, and pronotum size and shape. Long et al. (2023) systematically scrutinized type specimens of pygmy grasshoppers stored in China, revealing 23 new synonyms of *Tetrix japonica* (Bolívar, 1887), indicating notable morphological diversity. Concurrently, Zhang et al. (2023) found similar variability in *Tetrix japonica* morphology throughout their life cycles. Given the complexity and variability of morphological features within Tetrigidae, relying solely on these traits often leads to recurrent classification errors, hindering the achievement of the rigorous accuracy standards required in contemporary species classification. Consequently, it is necessary to utilize molecular data as an auxiliary verification tool to ensure the accuracy of classification. In recent years, with the rapid development of high-throughput sequencing, especially the progress of whole mitochondrial genome sequencing, insect mitochondrial genomes have been widely used as a molecular marker to investigate phylogenetic relationships, biological identification, and the genetic structure of populations (Boore 1999; Cameron 2014; Song et al. 2019; Nie et al. 2021; Li et al. 2021b). Mitogenomes have greatly improved our understanding of the phylogenetics of Tetrigidae, and some scholars have begun to use these methods to answer questions about the taxonomy and phylogeny of tetrigids (Fang et al. 2010; Pavón-Gozalo et al. 2012; Lin et al. 2015; Chen et al. 2018; Adžić et al. 2020; Huang et al. 2022; Kasalo et al. 2023a).

In this study, we obtained two unknown species of wingless pygmy grasshoppers of Metrodorinae. To determine their taxonomic positions, we sequenced their mitochondrial genomes, constructed molecular phylogenetic relationships, and estimated genetic distances and divergence times. Finally, it was determined that one of them belongs to a new genus of Metrodorinae, while the other belongs to a new species of the genus *Macromotettixoides*. We establish a new genus *Edentatettix* Deng, gen. nov., characterized by the absence or degeneration of the unique saw-like teeth in the female ovipositor. *Edentatettix leyeensis* Deng, sp. nov. is described as type species. Meanwhile, *Macromotettixoides yaana* Deng, sp. nov., is described as new to science. Based on a re-examination of type specimens, we propose five critical taxonomic revisions as follows: 1) synonymization of *Hainantettix strictivertex* Deng, 2020 with *H. angustivertex* (Zha & Peng, 2021); 2) reclassification of *Macromotettixoides angustivertex* Zha & Peng, 2021 under *Hainantettix*; 3) consolidation of *Macromotettixoides jiuwanshanensis* Zheng, Wei & Jiang, 2005 with *Hyboella badagongshanensis* Zheng, 2013, *M. badagongshanensis* (Zheng, 2013), and *M. wuyishana* Zheng, 2013 based on overlapping diagnostic characters. In addition, the taxonomic significance of toothless ovipositors in Tetrigidae is discussed.

Materials and methods

Mitogenome sequencing, assembly, annotation and analysis

Total genomic DNA was extracted from muscle tissues of the hind femur using TIANamp Genomic DNA Kit (TIANGEN) and sent to Berry Genomics (Beijing, China) for genomic sequencing using Next Generation Sequencing (NGS) method. Separate 350-bp insert libraries were created from the whole genomic DNA and sequenced using the Illumina HiSeq X Ten sequencing platform. A total of 5 Gb of 150-bp paired-end (PE) reads were generated in total for each sample.

The mitochondrial genome was assembled by NOVOPlasty 4.2.1 and annotated using the MITOS2 Web Server (<http://mitos2.bioinf.uni-leipzig.de/index.py>, accessed on 17 May 2024; Donath et al. 2019). The annotated mitogenome sequences were checked in CLC Genomics Workbench 12.0.1, MEGA 11.0.13, and Geneious Prime 11.0.18. The maps of the mitogenomes were generated using the Proksee website (<https://proksee.ca>, accessed on 10 August 2024, Grant et al. 2023). The nucleotide composition of the mitogenome, PCGs, three codon positions of PCGs, tRNA genes, rRNA genes, and the control regions (CR) was computed in PhyloSuite v.1.2.3 and MEGA 11.0.13. The AT and GC skews were calculated using the formula: $AT\text{-skew} = (A - T) / (A + T)$ and $GC\text{-skew} = (G - C) / (G + C)$.

Phylogenetic analysis

To determine the phylogenetic positions of two new species in Metrodoridae, a total of 30 mitogenomes, including 28 downloaded from NCBI and two from this study were employed. *Mirhipipteryx andensis* (Ripipterygidae: Ripipteryginae) and *Ellipes minuta* (Tridactylidae: Tridactylinae) were selected as outgroups (Table 1). The analysis was performed using PhyloSuite v. 1.2.3. Redundant sequences were removed, and PCGs in the mitochondrial genome were extracted, aligned in batches, and concatenated with MAFFT (Katoh and Standley 2013). Thirteen PCGs were aligned and concatenated through the MAFFT v. 7.505 (Katoh and Standley 2013) plugin in PhyloSuite v. 1.2.3. ModelFinder v. 2.2.0 (Kalyanamoorthy et al. 2017) was used to select the best-fit model using AICc and BICc standards. The phylogenetic tree was assessed with the maximum likelihood method and the bayesian inference method. The maximum likelihood estimations with IQ-TREE v. 2.2.0 (Nguyen et al. 2015) under the Edge-linked partition model with 15,000 ultrafast bootstrap replicates (Minh et al. 2013) and Bayesian inference used MrBayes v. 3.2.7a under the GTR+I+G4 model (with two parallel runs, 2,000,000 generations). Genetic distances were calculated using the Tajima-Nei model based on the sequence of 13 PCGs with the bootstrap method of 1000 replications using MEGA 11.0.13.

Divergence time analysis

The divergence times were estimated using BEAST v. 1.8.4 (Drummond and Rambaut 2007; Drummond et al. 2012), and the calibration times estimated from previous studies were queried from the timetree.org website (<http://time-tree.org/>, accessed on 9 December 2024; Kumar et al. 2022). The nodes of *Tetrix* and *Alulatettix* (4.35~7.98 MYA) was selected for calibration (Song et al. 2015). The parameters, including time priors and prior distributions, were set as “Yule Process” and “Normal”. The range was strictly set according to the

Table 1. Accession numbers and references of the mitogenomes of Tetrigidae included in this study.

Subfamily	Species	Accession number	Reference
Batrachideinae	<i>Saussurella borneensis</i>	MZ169555	Deng et al. 2021
Tripetalocerinae	<i>Tripetaloceroides tonkinensis</i>	MW770353	Zhang et al. 2021
Scelimeninae	<i>Eucriotettix oculatus</i>	MT162546	Li et al. 2021a
	<i>Loxilobus prominenculus</i>	MT162545	Li et al. 2021b
Metrodorinae	<i>Bolivaritettix sikkinensis</i>	KY123120	Yang et al. 2017b
	<i>Bolivaritettix yuanbaoshanensis</i>	KY123121	Yang et al. 2017b
	<i>Bolivaritettix lativertex</i>	MN083173	Chang et al. 2020
	<i>Macromotettixoides maoershanensis</i>	OR030790	Luo et al. 2024
	<i>Macromotettixoides brachycorna</i>	OR003899	Luo et al. 2024
	<i>Macromotettixoides orthomargina</i>	OR030789	Luo et al. 2024
	<i>Macromotettixoides yaana</i> Deng, sp. nov.	PQ826485	This study
	<i>Edentatettix leyeensis</i> Deng, sp. nov.	PQ826484	This study
	<i>Systolederus spicupennis</i>	MH791445	Yang 2017a
	<i>Teredorus bashanensis</i> = <i>Systolederus bashanensis</i> (Devriese & Husemann, 2023)	MZ041208	Li et al. 2021c
	<i>Teredorus anhuiensis</i> = <i>Systolederus anhuiensis</i> (Devriese & Husemann, 2023)	NC_071822	Unpublished
	<i>Teredorus guangxiensis</i> = <i>Systolederus zhengi</i> (Devriese & Husemann, 2023)	NC_082935	Guan et al. 2024
	<i>Teredorus hainanensis</i> = <i>Systolederus hainanensis</i> (Devriese & Husemann, 2023)	NC_063117	Li et al. 2021c
	<i>Teredorus nigropennis</i> = <i>Systolederus nigropennis</i> (Devriese & Husemann, 2023)	MN938922	Li et al. 2020b
Tetriginae	<i>Coptotettix longtanensis</i>	OK540319	Unpublished
	<i>Coptotettix longjiangensis</i>	KY798413	Lin et al. 2017
	<i>Euparatettix tridentatus</i>	NC_082933	Guan et al. 2024
	<i>Euparatettix variabilis</i>	NC_046542	Chang et al. 2020
	<i>Euparatettix bimaculatus</i>	NC_046541	Chang et al. 2020
	<i>Exothotettix guangxiensis</i>	NC_082934	Guan et al. 2024
	<i>Formosatettix qinlingensis</i>	KY798412	Lin et al. 2017
	<i>Alulatettix yunnanensis</i>	NC_018542	Xiao et al. 2012a
	<i>Tetrix japonica</i>	NC_018543	Xiao et al. 2012b
	<i>Tetrix ruyuanensis</i> = <i>Tetrix japonica</i> (Long et al., 2023)	NC_046412	Chang et al. 2020
Outgroup	<i>Mirhipipteryx andensis</i>	NC_028065	Song et al. 2015
	<i>Ellipes minuta</i>	NC_014488	Sheffield et al. 2010

queried calibration times. The uncorrelated relaxed clock model was selected with the relaxed distribution set as lognormal, and the same partition generated from the IQ-TREE was adopted. Finally, the MCMC generations and burn-ins were set as 10 million and 10%, respectively. The generated trees were imported into the Treeannotator to yield a consensus tree. The resulting phylogenetic tree was viewed and edited in ChiPlot (v. 2.6.1) (<https://www.chiplot.online/>, accessed on 10 December 2024; Xie et al. 2023).

Taxonomy, nomenclature, terminology, and measurements

Taxonomy follows the Orthoptera Species File [OSF] (Cigliano et al. 2024), a database of Orthoptera taxonomy. Nomenclature follows the International Code of the Zoological Nomenclature (ICZN 1999). Morphological terminology and landmark-based measurement methods followed those used by Zheng (2005a), Deng et al. (2007), Tumbrinck and Telnov (2014), Tumbrinck (2019),

Tan and Artchawakom (2015), Muhammad et al. (2018), and Zha et al. (2017). Measurements are given in millimeters (mm). Grasshopper specimens were examined using a Motic-SMZ-168 stereo-microscope and photographed using a KEYENCE VHX-600 Digital Microscope (Keyence Corporation, Osaka, Japan). All images were processed with Adobe Photoshop 24.0.0. The distribution map was prepared by ArcMap 10.8.1 and edited in Adobe Photoshop 24.0.0.

The species of *E. leyeensis* Deng, sp. nov. was collected from Wutaishan Forest Park, Leye County, Guangxi Province, China (24°51'11"N, 106°32'17"E) on 23 August 2021 by Wei-An Deng. Specimens of *M. yaana* Deng, sp. nov. were collected from Longdong (Ganyanggou), Baoxing County, Yaan City, Sichuan Province, China (30°24'19"N, 102°35'45"E) on 2 August 2016 by Wei-An Deng. The collected specimens were preserved in 100% anhydrous ethanol and stored in the refrigerator at −20 °C at the College of Life Science, Guangxi Normal University, Guilin, China (CLSGNU).

Type specimen depositories

The specimens examined in this study, including all holotypes and paratypes, have been deposited in the following institutions:

CLSGNU College of Life Science, Guangxi Normal University, Guilin, China;
EMHU Entomological Museum of Hechi University, Hechi, China;
IZSNU Institute of Zoology, Shaanxi Normal University, Xi'an, China;
HNU Huaibei Normal University, Huaibei, Anhui, China.

Results

Mitogenome characteristics of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov.

Genome organization and nucleotide composition

The mitogenomes of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. are circularized, with sizes of 15,813 bp and 16,379 bp, respectively (Fig. 1). Both mitogenomes contain 13 PCGs, 22 tRNAs, 2 rRNA (*rrnS* and *rrnL*), and a control region. Most genes (9 PCGs and 14 tRNA genes) were encoded on the majority strand (J-strand), and the rest of the genes (4 PCGs, 8 tRNAs, and 2 rRNAs) were located on the minority strand (N-strand) (Table 2).

The AT-skew and nucleotide composition of the mitogenomes of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. are shown in Table 3. With asymmetric nucleotide composition (*E. leyeensis* Deng, sp. nov.: 42.4% A, 30.2% T, 17.7% C, and 9.8% G; *M. yaana* Deng, sp. nov.: 44.3% A, 30.2% T, 16.5% C, and 9.3% G) and A+T-biased (*E. leyeensis* Deng, sp. nov.: 72.6%; *M. yaana* Deng, sp. nov.: 74.2%). This nucleotide composition pattern is consistent with other species of Tetrigidae (Xiao et al 2012a; Zhang et al 2021). The AT skews of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. are 0.169 and 0.194, respectively, and the CG skews are -0.287 and -0.276, respectively. The AT-skew is positive, and the CG skew is negative. This shows that the content of bases C is higher than that of G, and A is higher than T in the whole (Table 3).

Table 2. Mitochondrial genome organization of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov.

Genes	Strand	Anticodon	Location	Length (bp)	Intergenic nucleotides	Start codon	Stop codon
<i>trnI</i>	J	GAT	1–64/1–65	64/65	0/.	–	–
<i>trnQ</i>	N	TTG	66–134/.	69/.	1/.	–	–
<i>trnM</i>	J	CAT	134–201/134–205	68/72	-1/.	–	–
<i>nad2</i>	J	–	202–1209/221–1216	1008/996	0/15	ATG/ATT	TAA/.
<i>trnW</i>	J	TCA	1208–1272/1215–1282	65/68	-2/.	–	–
<i>trnC</i>	N	GCA	1265–1326/1275–1338	62/64	-8/.	–	–
<i>trnY</i>	N	GTA	1327–1389/1339–1402	63/64	0/.	–	–
<i>cox1</i>	J	–	1387–2925/1400–2938	1539/.	-3/.	ATC/.	TAA/.
<i>trnL2</i>	J	TAA	2921–2983/2934–2997	63/64	0/-5	–	–
<i>cox2</i>	J	–	2985–3662/2999–3682	678/684	1/.	ATG/.	TAA/.
<i>trnD</i>	J	GTC	3666–3728/3682–3744	63/.	3/-1	–	–
<i>trnK</i>	J	CTT	3732–3797/3746–3811	67/66	2/1	–	–
<i>atp8</i>	J	–	3797–3958/3815–3973	162/159	-1/3	ATA/ATG	TAA/.
<i>atp6</i>	J	–	3952–4623/3967–4638	672/.	-7/.	ATG/.	TAA/.
<i>cox3</i>	J	–	4623–5411/4638–5444	789/807	-1/.	ATG/.	TAA/.
<i>trnG</i>	J	TCC	5411–5473/5428–5489	63/62	-1/-17	–	–
<i>nad3</i>	J	–	5474–5827/5490–5843	354/.	0/.	ATT/.	TAG/.
<i>trnA</i>	J	TGC	5826–5892/5842–5906	67/65	-2/.	–	–
<i>trnR</i>	J	TCG	5892–5953/5906–5965	62/60	-1/.	–	–
<i>trnN</i>	J	GTT	5954–6017/5966–6032	64/67	0/.	–	–
<i>trnS1</i>	J	GCT	6018–6081/6033–6099	64/67	0/.	–	–
<i>trnE</i>	J	TTC	6082–6145/6100–6163	64/.	0/.	–	–
<i>trnF</i>	N	GAA	6144–6207/6162–6223	64/62	-2/.	–	–
<i>nad5</i>	N	–	6208–7930/6224–7946	1723/.	0/.	ATG/.	T(AA)/.
<i>trnH</i>	N	GTG	7933–7996/7951–8015	64/65	2/4	–	–
<i>nad4</i>	N	–	7996–9315/8015–9340	1320/1326	-1/.	ATA/ATG	TAG/.
<i>nad4l</i>	N	–	9315–9602/9334–9624	288/291	-1/-7	ATT/.	TAA/.
<i>trnT</i>	J	TGT	9605–9666/9627–9688	62/.	2/.	–	–
<i>trnP</i>	N	TGG	9667–9729/9689–9755	63/67	0/.	–	–
<i>nad6</i>	J	–	9740–10225/9769–10248	486/480	10/13	ATT/ATA	TAA/.
<i>cob</i>	J	–	10225–11361/10248–11387	1137/1140	-1/.	ATG/.	TAA/.
<i>trnS2</i>	J	TGA	11370–11434/11386–11452	65/67	8/-2	–	–
<i>nad1</i>	N	–	11450–12491/11811–12749	942/945	115/358	ATT/.	TAA/.
<i>trnL1</i>	N	TAG	12492–12555/12750–12813	64/.	0/.	–	–
<i>rrnL</i>	N	–	12560–13852/12815–14099	1293/1285	4/1	–	–
<i>trnV</i>	N	TAC	13857–13924/14103–14170	68/.	4/3	–	–
<i>rrnS</i>	N	–	13923–14657/14170–14913	735/744	-2/-1	–	–
CR	–	–	14658–15813/14914–16379	1156/1466	–	–	–

Note: “H” indicates the majority strand and “L” indicates the minority strand.

Protein-coding genes and codon usage

The mitogenomes of both *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. contain 13 PCGs, with *nad5* being the longest and *atp8* being the shortest. The total length of the 13 PCGs in *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. are 11,098 bp and 11,116bp, respectively, approximately accounting for 70.18% and 67.87% of the whole mitogenome, respectively (Table 2). Nine of the 13 PCGs are encoded on the J-strand (*cox1*, *cox2*, *cox3*, *cytb*, *nad2*, *nad3*, *nad6*, *atp6*, *atp8*),




Table 3. Nucleotide composition of mitochondrial genome of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov.

Genes or partitions	A (%)	T (%)	G (%)	C (%)	A+T (%)	AT-skew	GC-skew
Whole genome	42.40/44.30	30.20/29.90	9.80/9.30	17.70/16.50	72.60/74.20	0.169/0.194	-0.287/-0.276
PCGs	31.30/34.60	39.80/38.00	13.70/12.30	15.20/.	71.10/72.60	-0.120/-0.047	-0.054/-0.106
PCGs-1 st	34.30/37.90	34.40/32.60	17.90/16.30	13.00/13.30	68.70/70.50	0/0.075	0.155/0.100
PCGs-2 nd	20.60/23.00	45.70/43.50	14.60/14.50	19.40/19.00	66.30/66.50	-0.379/-0.308	-0.149/-0.134
PCGs-3 rd	39.00/42.90	39.40/37.90	8.50/6.00	13.00/13.20	78.40/80.80	-0.006/-0.061	-0.208/-0.372
tRNA	36.70/39.50	37.20/35.10	14.70/14.00	11.40/.	73.90/74.60	-0.008/0.059	0.130/0.104
rRNA	28.90/48.40	45.20/27.70	17.10/8.00	8.80/15.9	74.10/76.10	-0.220/0.272	0.319/-0.332
CR	49.74/51.09	30.97/32.67	6.49/6.62	12.80/9.62	80.70/83.80	0.233/0.220	-0.327/-0.185

Note: AT-skew = $(A - T) / (A + T)$; GC-skew = $(G - C) / (G + C)$.

The relative synonymous codon usage (RSCU) values of the mitogenome are summarized (Fig. 2). The codon distribution analysis shows that the two codons UUA (Leu2) and UCU (Ser2) are the most frequently used in *E. leyeensis* Deng, sp. nov. The codons of UUA (Leu2) and UCA (Ser2) in *M. yaana* Deng, sp. nov. are the most frequently used. The frequency of the codons ending with A/U is much higher than with G/C, suggesting that the AU composition at the third position of codons has a positive influence on the nucleotide AT (or AU) bias of the PCGs.

The lengths of the 22 tRNA genes in *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. are 1418 bp and 1435 bp, respectively, with size ranges of 62 to 69 bp for *E. leyeensis* Deng, sp. nov. and 60 to 72 bp for *M. yaana* Deng, sp. nov. Two rRNA genes (*rrnL* and *rrnS*), separated by *trnV*, are located between *trnL1*



Figure 2. Relative synonymous codon usage (RSCU) in the mitogenomes of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov.

and A+T-rich region. The total AT-skew of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. in rRNA are -0.220 and 0.272, respectively. The total GC-skew of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. in rRNA are 0.319 and -0.332, respectively. All tRNA genes could be folded into the typical clover-leaf structure, except *trnS1* and *trnV*, which lacked the dihydrouridine (DHU) arm (Fig. 3).

Phylogenetic analysis

We constructed maximum likelihood (ML) and Bayesian inference (BI) trees based on the sequences of 13 PCGs from the mitochondrial genomes of 28 species from five subfamilies (Tetriginae, Metrodorinae, Scelimeninae, Tripetalocerinae, and Batrachideinae) of Tetrigidae and two outgroup species with the best partition schemes (Table 1). All phylogenetic analyses used the same data matrices, yet different methods yielded the same topology. Similar topologies were obtained from BI and ML trees (Fig. 4).

In this study, ML and BI trees had the same topological structures, and both phylogenies revealed the non-monophyletic relationships among species of the subfamily Metrodorinae. Tetrigidae was retrieved as monophyletic with strong support (posterior probability, PP = 1). However, only one species' datum was available for Batrachideinae, and Tripetalocerina, making it impossible to determine their monophyly.

The results of phylogeny analysis conducted by ML and BI methods are as follows: (Batrachideinae + (Tripetalocerinae + ((Scelimeninae + Metrodorinae) + (Metrodorinae + (Metrodorinae + Tetriginae))))). *Saussurella borneensis*




Figure 3. The secondary structures for 22 tRNA genes of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. Watson-Crick base pairings and mismatches are represented by dashes (–) and stars, respectively.

Hancock, 1912 (Batrachideinae) split off earliest from the other taxa, suggesting that it is the earliest species within Tetrigidae. The subfamily Tetriginae represented the most evolutionarily advanced group within Tetrigidae, while the subfamily Metrodorinae occupied an intermediate position. These findings are consistent with previous studies (Chen 2005; Lin et al. 2015; Wang 2022). Although *E. leyeensis* Deng, sp. nov. bears a striking morphological similarity to species of *Macromotettixoides*, both belonging to wingless pygmy grasshoppers of the Metrodorinae, phylogenetic evidence does not justify classifying *E. leyeensis* within *Macromotettixoides*. Instead, it constitutes a separate new genus, namely *Edentatettix* Deng, gen. nov. Furthermore, another new species *M. yaana* Deng, sp. nov. forms a sister-group relationship with *Macromotettixoides orthomargina* Wei & Deng, 2023, which is strongly supported (PP = 1), confirming that *M. yaana* Deng, sp. nov. belongs to *Macromotettixoides*.




Figure 4. Phylogenetic tree obtained from ML and BI analysis based on 13 PCGs.




Figure 5. Pairwise genetic distances of mitochondrial DNA for 28 species of Tetrigidae based on 13PCGs.

Genetic distances

In Tetrigidae, the genetic distances between different species pairs varied significantly, with a range from 0.018 to 0.612 (Fig. 5). In this study, the intraspecies genetic distance of *Tetrix japonica* (= *Tetrix ruyuanensis* Liang, 1998, syn. nov.) (Long et al. 2023) was 0.004. The genetic distances of the new species *M. yaana* Deng, sp. nov. to the three species of *Macromotettixoides* ranged from 0.083

to 0.200, which were greater than the interspecific distance (0.004), confirming that *M. yaana* Deng, sp. nov. is a separate species. Furthermore, the genetic distances between *E. leyeensis* Deng, sp. nov. and the other 27 species ranged from 0.403 to 0.582, which is significantly greater than the intergeneric genetic distances observed in *Systolederus* (0.018–0.360), *Bolivaritettix* (0.053–0.096), *Euparatettix* (0.133–0.227), and *Macromotettixoides* (0.087–0.228). Therefore, despite morphological similarities to species like *Macromotettixoides*, the genetic distances indicate that *E. leyeensis* Deng, sp. nov. warrants classification in a new genus.

Divergence time analysis

Divergence time analysis shows that Metrodorinae appeared around 120.52 Mya (Fig. 6). This analysis also identifies considerable variability in the timings of differentiation across diverse genera within this subfamily. For example, the new genus *Edentatettix* Deng, gen. nov. can be traced back to approximately 112.23 Mya, which is close to the estimated split time of Metrodorinae. The early divergence of this genus may indicate distinct adaptive shifts in its evolutionary history, leading to the development of unique ecological niches and morphological characteristics. Additionally, *Tetrix japonica*, with a divergence timescale of approximately 0.71 Mya, represents one of the most recently evolved species in Tetrigidae. It also exhibits the broadest global distribution among its relatives, reflecting its remarkable adaptability and ecological success.




Figure 6. Dated phylogenetic tree of Tetrigidae based on the PCGs dataset. A time scale is provided. Mya refers to one million years ago.

Taxonomy of the subfamily Metrodorinae Bolívar, 1887

Key to the Chinese genera of Metrodorinae Bolívar, 1887

Modified from Lu and Deng (2021), Metrodorinae currently includes 16 genera in China (including the new genus *Edentatettix* Deng, gen. nov.).

- 1 Fastigium of the vertex distinctly projecting before anterior margin of compound eyes.....**2**
 - Fastigium of the vertex not or slightly projecting before anterior margin of compound eyes**4**
- 2 Fastigium of the vertex calyptriform protruding before anterior margin of compound eyes ***Calyptraeus* Wang, 2001**
 - Fastigium of the vertex angular projecting or square projecting**3**
- 3 Fastigium of the vertex angular projecting; antennal grooves inserted between inferior margin of compound eyes ***Rhopalotettix* Hancock, 1910**
 - Fastigium of the vertex square projecting; antennal grooves inserted at lowest third of compound eyes height ***Miriatrioides* Zheng & Jiang, 2002**
- 4 Saw-like teeth of female ovipositor absent or degenerate
 - ***Edentatettix* Deng, gen. nov.**
 - Saw-like teeth of female ovipositor present**5**
- 5 With a distinct obtuse projection under each lateral carina of prozona; pronotum platy, in dorsal view, dorsum of pronotum with irregular concavities ***Concavetettix* Deng, 2021**
 - Without obtuse projection under each lateral carina of prozona; pronotum tectiform.....**6**
- 6 Vertex narrow, still narrower towards front, eyes drawn to each other in front and elevated..... ***Systolederus* Bolívar, 1887**
 - Vertex not as above.....**7**
- 7 Posterior margins of each lateral lobes of pronotum only with ventral sinus ***Macromotettixoides* Zheng, Wei & Jiang, 2005**
 - Posterior margins of each lateral lobe of pronotum with ventral sinus and tegminal (upper) sinus**8**
- 8 Humeral apex ridge and lower margin of pronotum connected in the middle or behind middle of lower margin of pronotum.....
 - ***Macromotettix* Günther, 1939**
 - Humeral apex ridge and lower margin of pronotum connected before middle of lower margin of pronotum.....**9**
- 9 Tegmina present and hind wings absent ***Paramphinotus* Zheng, 2004**
 - Tegmina and hind wings present**10**
- 10 Ventral margins of fore femora and middle femora with two big teeth.....
 - ***Orthotettixoides* Zheng, 1998**
 - Ventral margins of fore femora and middle femora without big teeth**11**
- 11 Head and eyes not exserted above pronotal surface**12**
 - Head and eyes distinctly exserted above pronotal surface.....**13**
- 12 Anterior part of pronotum strongly widened, arched and uplifted
 - ***Hyboella* Hancock, 1915**
 - Anterior part of pronotum normal and strongly widened
 - ***Bolivaritettix* Günther, 1939**

- 13 Antennal grooves inserted far below inferior margin of compound eyes **14**
 - Antennal grooves inserted at lowest third of compound eye height or between inferior margin of compound eyes **15**
- 14 The vertex horn is distinctly raised above the dorsal margin of eyes and the vertex is deeply depressed between eyes in frontal view; pronotum between the shoulders is not elevated to an obtuse gibbosity ***Xistra* Bolívar, 1887**
 - In frontal view, the vertex horn slightly raised above the dorsal margin of eyes (or not) and the vertex is slightly depressed (or not depressed) between eyes; pronotum between the shoulders generally strongly elevated to an obtuse gibbosity ***Xistrella* Bolívar, 1909**
- 15 Antennal grooves inserted at lowest third of compound eye height; median carina of pronotum with a series of projections
 - ***Cotysoides* Zheng & Jiang, 2000**
 - Antennal grooves inserted between inferior margin of compound eyes; median carina of pronotum generally straight or undulated ... ***Mazarredia* Bolívar, 1887**

Descriptions

Genus *Edentatettix* Deng, gen. nov.

<https://zoobank.org/A21EDA29-0040-4AF1-B899-87A4DB4D19DF>

Type species. *Edentatettix leyeensis* Deng, sp. nov., here designated.

Diagnosis. The new genus can be easily distinguished from other genera of Metrodorinae by the saw-like teeth of the female ovipositor absent (Fig. 8G). *Edentatettix* is allied to *Macromotettixoides* Zheng, Wei & Jiang, 2005, but differs as follows: head and eyes slightly exserted above pronotal surface (not exserted in *Macromotettixoides*), dorsal surface of pronotum low and flat and tectiform is not obvious (distinctly tectiform in *Macromotettixoides*), saw-like teeth of female ovipositor absent (present in *Macromotettixoides*). *Edentatettix* is also similar to *Concavetettix* Deng, 2021 but differs from the latter by the obtuse projection under each lateral carina of the prozona absent (present in *Concavetettix*), dorsum of pronotum flat and slightly depressed in the middle part between the shoulders (dorsum of pronotum with irregular concavities in *Concavetettix*), saw-like teeth of female ovipositor absent (present in *Concavetettix*).

Description. General characters and coloration. Size small, brachypronotal. Coloration uniformly brown, antennae and face dark brown, middle of the dorsal surface of pronotum with a dark spot. Body surface is interspersed with sparse carinae and notches.

Head. Head and eyes slightly exserted above pronotal surface. Fastigium of vertex short; in dorsal view, width of vertex between eyes 1.5–1.6× width of compound eye. In lateral view, frontal ridge and vertex forming a rounded-angle shape and slightly projected inferior margin of the compound eye, frontal costa distinctly concave between lateral ocelli. In frontal view, frontal costa bifurcated above lateral ocelli, the bifurcation of the frontal costa in the middle of the compound eye height; width of longitudinal furrow of frontal ridge narrower than antennal groove diameter. Antennae short, filiform, antennal grooves inserted below inferior margin of compound eyes. Eyes globose, lateral (paired) ocelli located lowest third of compound eye height.

Thorax. Dorsal surface of pronotum low and flat and tectiform is not obvious; pronotal surface interspersed with sparse carinae and notches between shoulders and behind the middle, slightly depressed in the middle part between the shoulders. Pronotum with truncate anterior margin, median carina entire and nearly straight in profile; lateral carinae of prozona parallel; humeral angle obtuse; with interhumeral carina; hind pronotal process narrow, nearly reaching apex of hind femur and its apex narrowly rounded. Posterior angles of lateral lobes produced outwards, end of posterior angles truncate, posterior margins of lateral lobes of pronotum only with ventral sinus. Tegmina and hind wings invisible.

Legs. Fore and middle femora slightly compressed, margins finely serrated and carinate and ventral margins slightly undulated. Hind femora robust and short, 2.8× as long as wide; with carinated. Length of first segment of posterior tarsi equal to third.

Abdomen. Female ovipositor narrow and long, dorsal margins of upper valvulae and ventral margins of lower valvulae without saw-like tooth or saw-like teeth indistinct (Fig. 8G).

Etymology. The generic epithet is derived from *edent*, referring to the absent saw-like teeth of female ovipositor (Fig. 8G).

***Edentatettix leyeensis* Deng, sp. nov.**

<https://zoobank.org/8DCD114B-F272-41D3-9C83-0426CD226421>

Figs 7–9

Type material. *Holotype* • ♀, CHINA, Guangxi Prov., Leye county (Wutaishan Forest Park), 24°51'11"N, 106°32'17"E, 1200 m alt., 23 August 2021, collected by Wei-An Deng, CLSGNU. *Paratypes* • 2♂, 5♀, same data, CLSGNU • 3♂, 6♀, same data, 18 August 2022, collected by Wei-An Deng and Yue-Mei Li, CLSGNU.

Diagnosis. As there is only one species in the genus, see the generic diagnosis.

Description. Female. Small size, short, body surface interspersed with sparse carinae and notches.

Head. Head and eyes slightly exserted above pronotal surface. Fastigium of vertex short; in dorsal view, width of vertex between eyes 1.5–1.6× width of compound eye; anterior margin of fastigium arched and not surpassing anterior margin of eye; median carina visible; lateral margins turned backward; vertex uneven with paired fossulae. In lateral view, frontal ridge and vertex forming a rounded-angle shape and slightly projected inferior margin of the compound eye, frontal costa distinctly concave between lateral ocelli, protruding anteriorly and broadly rounded between antennal grooves. In frontal view, frontal costa bifurcated above lateral ocelli, the bifurcation of the frontal costa in the middle of the compound eye height; longitudinal furrow widely divergent between antennae, width of longitudinal furrow of frontal ridge narrower than antennal groove diameter. Antennae short, filiform, antennal grooves inserted below inferior margin of compound eyes, 15-segmented; the 10th and 11th segment are the longest, ~ 7.0–8.0× longer than its width. Eyes globose, lateral (paired) ocelli located lowest third of compound eye height.

Thorax. Brachypronotal. Dorsal surface of pronotum low and flat and tectiform is not obvious; pronotal surface interspersed with sparse carinae and notches between shoulders and behind the middle, slightly depressed in the middle part between the shoulders. Pronotum with truncate anterior margin,




Figure 7. *E. leyeensis* Deng, sp. nov., holotype female **A** body, lateral view **B** the same, dorsal view **C** head and anterior part of pronotum, dorsal view **D** the same, lateral view.




Figure 8. *E. leyeensis* Deng, sp. nov., holotype female **A** head, frontal view **B** left fore femur, lateral view **C** left mid femur, lateral view **D** left hind femur, lateral view **E** left hind tibia, lateral view **F** left posterior tarsus, lateral view **G** ovipositor of female, lateral view **H** subgenital plate of female, ventral view.




Figure 9. *Edentatettix leyeensis* Deng, sp. nov., paratype male **A** body in dorsal view **B** body in lateral view.

median carina entire and nearly straight in profile; lateral carinae of prozona parallel; humeral angle obtuse; with interhumeral carina; hind pronotal process narrow, nearly reaching apex of hind femur and its apex narrowly rounded. Lower margin of hind process nearly straight, lateral carinae of metazona slightly curved, width of infrascapular area is 0.7 mm. Posterior angles of lateral lobes produced outwards, end of posterior angles truncate, posterior margins of lateral lobes of pronotum only with ventral sinus. Tegmina and hind wings invisible.

Legs. Fore and middle femora slightly compressed, margins finely serrated and carinate and ventral margins slightly undulated. Hind femora robust and short, 2.8× as long as wide; with carinated, dorsal margins smooth and ventral margins finely serrated; antegenicular denticles and genicular denticles acute. Outer and inner side of hind tibia with five or six spines. Length of first segment of posterior tarsi equal to third, three pulvilli of first segment of posterior tarsi: first small, second and third large; apices of all pulvilli obtuse.

Abdomen. Ovipositor narrow and long, length of upper valvulae 3.8× its width, dorsal margins of upper valvulae and ventral margins of lower valvulae without saw-like tooth or saw-like teeth indistinct (Fig. 8G). Length of subgeni-

tal plate slightly longer than its width, middle of posterior margin of subgenital plate triangular projecting.

Coloration. Body brown; antennae and face dark brown. Middle of dorsal surface of pronotum with a dark spot. Fore and middle femora and tibia brown, with two dark brown transverse spots. Hind femora dark brown, outer side with two pale stripes. Hind tibia black, with two pale rings in the middle.

Male. Similar to female, but smaller and narrower. Width of vertex between eyes 1.3–1.5× width of compound eye. Ventral margins of middle femora undulated. Subgenital plate short, cone-shaped, apex bifurcated.

Measurements (mm). Length of body: ♂ 9.0–9.5, ♀ 11.5–12.0; length of pronotum: ♂ 8.0–8.5, ♀ 9.5–10.0; length of hind femur: ♂ 5.5–5.8, ♀ 6.5–7.0.

Etymology. The specific name refers to the type locality: Leye, Guangxi, China; adjective.

Distribution. P. R. China.

***Macromotettixoides yaana* Deng, sp. nov.**

<https://zoobank.org/63C2CA79-9127-44CB-A783-4E2F135833CB>

Figs 11–13

Type material. *Holotype* • ♀, CHINA, Sichuan Prov., Yaan City, Baoxing county, Longdong (Ganyanggou), 30°24'19"N, 102°35'45"E, 1400 m alt., 02 August 2016, collected by Wei-An Deng, CLSGNU. *Paratypes* • 8♂, 7♀, same data, CLSGNU.

Diagnosis. This new species is similar to *Macromotettixoides convexa* Deng, 2020, from which it differs in that the pronotal surface is without a tuberculiform convex between shoulders (pronotal surface with a distinctly tuberculiform convexity between shoulders in *M. convexa*); lower margin of hind pronotal process straight (lower margin of hind pronotal process curved in *M. convexa*); median carina of pronotum undulated in profile (median carina of pronotum distinctly arch-like before shoulders and undulated behind shoulders in profile in *M. convexa*); lower outer carina of hind femora without projections (postmedian of lower outer carina of hind femora with two inconspicuous projections in *M. convexa*).

Description. Female. Small size, short, body surface interspersed with coarse protuberances.

Head. Head and eyes not exserted above pronotal surface. Fastigium of vertex short; in dorsal view, width of vertex between eyes 1.4–1.6× width of compound eye; anterior margin of fastigium slightly concave in the middle, slightly surpassing anterior margin of eye; median carina visible anteriorly; lateral margins turned backward; vertex uneven with paired fossulae. In lateral view, frontal ridge and vertex forming a rounded-angle shape, frontal costa distinctly concave between eyes, protruding anteriorly and broadly rounded between antennal grooves. In frontal view, frontal costa bifurcated above lateral ocelli, the bifurcation of the frontal costa in the middle of the compound eye height; longitudinal furrow widely divergent between antennae, width of longitudinal furrow of frontal ridge narrower than antennal groove diameter. Antennae short, filiform, antennal grooves inserted below inferior margin of compound eyes, 14-segmented, the 10th and 11th segment are the longest, ~ 2.5–3.0× longer than its width. Eyes globose, lateral (paired) ocelli located lowest third of compound eye height.

Thorax. Brachypronotal. Pronotum with distinctly tectiform, pronotal surface interspersed with dense protuberances of variable sizes and notches. Pronotum with truncate anterior margin, median carina slightly lamellar and entire and undulated in profile; lateral carinae of prozona slightly lamellar and parallel; humeral angle obtuse; hind pronotal process narrow, nearly reaching apex of hind femur and its apex narrowly rounded. Lower margin of hind process straight, lateral carinae of metazona curved, width of infrascapular area is 0.9 mm. Posterior angles of lateral lobes slightly produced outwards, end of posterior angles truncate, posterior margins of lateral lobes of pronotum with distinctly ventral sinus and very weakly tegminal sinus. Tegmina and hind wings strongly reduced and covered by infrascapular area and invisible or slightly visible.

Legs. Fore and middle femora slightly compressed, margins finely serrated and carinate and ventral margins with two inconspicuous projections and undulated. Hind femora robust and short, 2.8× as long as wide; with carinated and margins finely serrated; antegenicular denticles and genicular denticles acute. Outer side of hind tibia with 6–8 spines, inner side with six or seven spines. Length of first segment of posterior tarsi slightly longer than third, three pulvilli of first segment of posterior tarsi: first and second small and apices acute, third large and apex a right angle.

Abdomen. Ovipositor narrow and long, length of upper valvulae 3.5× its width, upper and lower valvulae with slender saw-like teeth. Length of subgenital plate slightly longer than its width, middle of posterior margin of subgenital plate slightly triangular projecting.




Figure 10. Distribution map of *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov.




Figure 11. *Macromotettixoides yaana* Deng, sp. nov., holotype female **A** body, lateral view **B** the same, dorsal view **C** head and anterior part of pronotum, dorsal view **D** the same, lateral view **E** head, frontal view.




Figure 12. *Macromotettixoides yaana* Deng, sp. nov., holotype female **A** left fore femur, lateral view **B** left mid femur, lateral view **C** left hind femur, lateral view **D** left hind tibia, lateral view **E** left posterior tarsus, lateral view **F** subgenital plate of female, lateral view **G** subgenital plate of female, ventral view.

Coloration. Body dark brown or brown; antennae dark brown. Hind tibia dark brown, with two pale rings in the middle.

Male. Similar to female, but smaller and narrower. Width of vertex between eyes 1.4–1.5× width of compound eye. Ventral margins of fore and middle femora slightly undulated. Subgenital plate short, cone-shaped, apex bifurcated.

Measurements (mm). Length of body: ♂ 6.5–7.0, ♀ 8.5–9.0; length of pronotum: ♂ 5.2–5.6, ♀ 7.0–7.5; length of hind femur: ♂ 4.0–4.5, ♀ 5.0–5.5.

Etymology. The specific name refers to the type locality: Yaan, Jinxiu, Sichuan, China; adjective.

Distribution. P. R. China: Sichuan (Fig. 10).




Figure 13. *Macromotettixoides yaana* Deng, sp. nov., paratype male **A** body in dorsal view **B** body in lateral view.

New synonyms

Hainantettix strictivertex Deng, 2020

Fig. 14

Hainantettix strictivertex Deng in Zhang, Zhao, Wu & Deng, 2020: 552 [description] (holotype – ♀, China: Hainan Prov., Qiongzong, Limushan; paratypes – 4♀, China: Hainan Prov., Wuzhishan; in EMHU; examined).

Macromotettixoides angustivertex Zha & Peng in Peng, Shi, Ding & Zha, 2021: 48 [description] (holotype – ♀, China: Hainan Prov., Wuzhishan, in HNU, not examined).

Hainantettix angustivertex (Zha & Peng): Subedi 2022: 40; syn. nov.




Figure 14. **A–C** Holotype of *Hainantettix strictivertex* Deng, 2020 **A** body in dorsal view **B** body in lateral view **C** labels **D**, **E** holotype of *Macromotettixoides angustivertex* Zha & Peng, 2021, syn. nov. **D** body in dorsal view **E** body in lateral view (photographs from Peng et al.).

Remarks. *Macromotettixoides angustivertex* Zha & Peng was described by Peng et al. (2021) and was later transferred to the genus *Hainantettix* by Subedi (2022). We examined the type specimen of *Hainantettix strictivertex* Deng, 2020. Although we did not examine the type material of *M. angustivertex* from Hainan, according to the original description and photographs of the type specimens in Peng et al. (2021), we found that the two species share identical morphological structures, type locality, and coloration. These two taxa are conspecific and characterized by the vertex very strongly narrowed towards the front drawing the eyes together; antennal grooves inserted below inferior margin of compound eyes; rami strongly divergent, width of longitudinal furrow of frontal ridge is distinctly wider than antennal groove diameter.

***Macromotettixoides jiuwanshanensis* Zheng, Wei & Jiang, 2005**

Figs 15, 16

Macromotettixoides jiuwanshanensis Zheng, Wei & Jiang, 2005: 366 [description] (holotype – ♀, China: Guangxi Prov., Luocheng (jiuwanshan), in IZSNU, examined). Zheng et al. 2005b: 176; Zheng et al. 2006: 604; Deng et al. 2007: 160; Zheng and Shi 2009: 572; Deng 2011: 544; Zheng 2013a: 242; Deng et al. 2014: 548; Deng et al. 2015: 165; Deng 2016: 156; Zha et al. 2017: 16; Han et al. 2020: 563; Li et al. 2020a: 110; Fan et al. 2023: 128.

Hyboella badagongshanensis Zheng, 2013b: 11 (holotype – ♀, China: Hunan Prov., Sangzhi (Badagongshan), in IZSNU, examined); Deng 2016: 150; syn. nov.

Macromotettixoides badagongshanensis (Zheng, 2013b): Zha, Yu, Boonmee, Eungwanichayapant, Wen, 2017: 23; Han et al. 2020: 564; Li et al. 2020a: 112; Fan et al. 2023: 129; syn. nov.

Macromotettixoides wuyishana Zheng, 2013: 242 [description] (holotype – ♀, China: Fujian Prov., Wuyishan, in IZSNU, examined); Deng 2016: 156; Zha et al. 2017: 15; Han et al. 2020: 563; Li et al. 2020a: 111; Fan et al. 2023: 128; syn. nov.

Remarks. *Hyboella badagongshanensis* was described by Zheng (2013b) but was later transferred to the genus *Macromotettixoides* by Zha et al. (2017). We examined the type specimens of *M. jiuwanshanensis*, *M. badagongshanensis*, and *M. wuyishana* and found that the structures and coloration of the body are the same in these three taxa. Therefore, we consider *M. badagongshanensis*, and *M. wuyishana* as synonyms of *M. jiuwanshanensis*. These three taxa are conspecific and characterized by the width of the vertex between the eyes being 2.1–2.2× the width of the compound eye; anterior margin of fastigium arched and surpassing anterior margin of eye; the anterior margin of pronotum is slightly obtuse protruding; and the lower carinae of fore and mid femora is straight.

Discussion

In this study, we describe two new pygmy grasshoppers of Metrodorinae found in China, *E. leyeensis* Deng, sp. nov. and *M. yaana* Deng, sp. nov. Through detailed morphological description and mitochondrial genome sequencing, we conducted a comprehensive analysis of the two new species. We have discovered




Figure 15. A–C Holotype of *Macromotettixoides jiuwanshanensis* Zheng, Wei & Jiang, 2005 **A** body in dorsal view **B** body in lateral view **C** labels **D–F** holotype of *Hyboella badagongshanensis* Zheng, 2013, syn. nov. **D** body in dorsal view **E** body in lateral view **F** labels.




Figure 16. *Macromotettixoides jiuwanshanensis* Zheng, Wei & Jiang, 2005. Holotype of *Macromotettixoides wuyishana* Zheng, 2013, syn. nov. **A** body in dorsal view **B** lateral view **C** body in labels.




Figure 17. ovipositor of female, lateral view **A** *E. leyeensis* Deng, sp. nov. **B** *M. yaana* Deng, sp. nov. **C** *M. orthomargina* **D** *M. brachycorna* **E** *M. maoershanensis*.

something quite intriguing. Although *E. leyeensis* Deng, sp. nov. morphologically resembles the genus *Macromotettixoides*, its uniqueness lies in the absence of serrated teeth on the female ovipositor (Fig. 17), which is extremely rare among Tetrigidae. Based on morphological characteristics, phylogeny, and divergence time analysis, *E. leyeensis* Deng, sp. nov. is set apart in classification, leading us to classify it under the new genus *Edentatettix* Deng, gen. nov.

The existence of an ovipositor serves as a shared derived characteristic (synapomorphy) among Insecta (Barbosa and Fianco 2024). Across this expansive class, a diverse array of morphological and functional characteristics can be observed in the ovipositors among different orders, and the varied structures of ovipositors in Orthoptera are no exception, vividly mirroring their unique adaptations to diverse ecological niches (Austin and Browning 1981; Kluge 2016). For instance, the sword-like or spear-shaped ovipositors of katydids and crickets exemplify their high adaptability for laying eggs in plant tissues or soil, ensuring both egg safety and enhanced larval survival rates (Kluge 2016; Chen et al. 2021). Conversely, the ovipositors of locusts and pygmy grasshoppers (Tetrigidae) are geared more towards excavating functionality, with the former having

a conical shape facilitating efficient soil digging (Kalogianni 1995; Zhao et al. 2022). The latter, while also adept at digging, may further enhance oviposition precision and efficiency by having fine teeth along the dorsal and ventral edges of their ovipositors (Kluge 2016). These varied oviposition strategies typify Orthopteran responses to environmental pressures, embodying a co-evolutionary mechanism between form and function.

The emergence of *E. leyeensis* Deng, sp. nov. with a toothless ovipositor provides not only morphological evidence for early Tetrigidae diversification but also insights into the ancient origins of complex reproductive structure transformations. The toothless ovipositor may represent a unique adaptive strategy, potentially associated with specific oviposition habitats or behaviors, meriting further investigation.

Importantly, we note for the first time that toothless ovipositors are an important taxonomic feature. This distinctive feature intimates the likelihood of diverse morphological variations in reproductive organs that may have emerged throughout the evolutionary journey of this group. Such variations are plausibly influenced by a multitude of factors, encompassing genetics, environmental conditions, and reproductive strategies (Hornig et al. 2018; Chen et al. 2021; Barbosa and Fianco 2024). Moreover, the timing of the emergence of this toothless characteristic (*E. leyeensis* Deng, sp. nov.), paralleling the divergence period of Tetrigidae, implies that it is an ancient character. This finding reinforces the hypothesis that Tetrigidae insects have undergone intricate morphological transformations, implicating their reproductive structures, over the course of their evolutionary progression.

Fossil records from the Early Miocene and Middle Eocene epochs affirm that the ovipositors of female Tetrigidae were adorned with saw-like teeth (Heads et al. 2014; Skejo et al. 2024), congruent with observations in contemporary species. The ancient divergence time of *E. leyeensis* Deng, sp. nov., given the constraints posed by the fossil record, accentuates that our current paleontological evidence likely constitutes merely a fraction of the extensive and diverse history of life. Some species, especially those inhabiting environments unfavorable to fossil preservation, may have irretrievably lost their historical footprint. Consequently, integrating insights from molecular biology, morphology, and paleontology becomes indispensable for elucidating the profound evolutionary mechanisms underlying global biodiversity.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by the Science & Technology Fundamental Resources Investigation Program of China (2023FY100200), the National Natural Science Foundation of China (32360124), and the Guangxi Natural Science Foundation under Grant No. 2023GXNSFDA026037.

Author contributions

Conceptualization: Wei-An Deng. Investigation: Wei-An Deng, Shi-Xiong Leng, Jia-Song He. Species identification: Wei-An Deng. Software: Yue-Mei Li, De-Long Guan. Analysed the data: Yue-Mei Li, De-Long Guan. Original draft writing: Yue-Mei Li, Wei-An Deng. Review and editing: Wei-An Deng. Funding acquisition: Wei-An Deng.

Author ORCIDs

Yuemei Li  <https://orcid.org/0009-0001-7327-5334>

Shixiong Leng  <https://orcid.org/0009-0002-3422-3862>

Jiasong He  <https://orcid.org/0009-0002-7224-594X>

Weian Deng  <https://orcid.org/0000-0002-8023-2498>

Delong Guan  <https://orcid.org/0000-0001-6866-0943>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Adžić K, Deranja M, Franjević D, Skejo J (2020) Are Scelimeninae (Orthoptera: Tetrigidae) monophyletic and why it remains a question? *Entomological News* 129(2): 128–146. <https://doi.org/10.3157/021.129.0202>
- Austin AD, Browning TO (1981) A mechanism for movement of eggs along insect ovipositors. *International Journal of Insect Morphology and Embryology* 10(2): 93–108. [https://doi.org/10.1016/S0020-7322\(81\)80015-3](https://doi.org/10.1016/S0020-7322(81)80015-3)
- Barbosa DN, Fianco M (2024) A Tale's blade: Understanding evolutionary features of oviposition behavior based on Tettigoniidae (Insecta, Orthoptera, Ensifera) ovipositor morphology. *Arthropod Structure and Development* 79: 101332. <https://doi.org/10.1016/j.asd.2024.101332>
- Bolívar I (1887) Essai sur les Acridiens de la tribu des Tettigidae. *Annales de la Societe Entomologique de Belgique* 31: 173–313.
- Bolívar I (1909) Nouvelles espèces d'Acridiens du Musée de Genève. *Boletín de la Real Sociedad Española de Historia Natural* 9: 399–400.
- Boore JL (1999) Animal mitochondrial genomes. *Nucleic Acids Research* 27(8): 1767–1780. <https://doi.org/10.1093/nar/27.8.1767>
- Cameron SL (2014) Insect mitochondrial genomics: implications for evolution and phylogeny. *Annual Review of Entomology* 59(1): 95–117. <https://doi.org/10.1146/annurev-ento-011613-162007>
- Chang HH, Nie YM, Zhang N, Zhang X, Sun HM, Mao Y, Qiu ZY, Huang Y (2020) MtOrt: an empirical mitochondrial amino acid substitution model for evolutionary studies of Orthoptera insects. *BMC Evolutionary Biology* 20: 1–5. <https://doi.org/10.1186/s12862-020-01623-6>
- Chen AH (2005) Phylogenetic Relationship Research of Tetrigoidea in China. Nanjing Normal University, M.Sc. thesis, 52 pp.
- Chen YZ, Deng WA, Wang JM, Lin LL, Zhou SY (2018) Phylogenetic relationships of Scelimeninae genera (Orthoptera: Tetrigoidea) based on COI, 16S rRNA and 18S rRNA gene sequences. *Zootaxa* 4482(2): 392–400. <https://doi.org/10.11646/zootaxa.4482.2.11>

- Chen L, Gu JJ, Yang Q, Ren D, Blanke A, Béthoux O (2021) Ovipositor and mouthparts in a fossil insect support a novel ecological role for early orthopterans in 300 million years old forests. *Elife* 10: e71006. <https://doi.org/10.7554/eLife.71006>
- Cigliano MM, Braun H, Eades DC, Otte D (2024) Orthoptera Species File Online. Version 5.0/5.0. 2024 <http://Orthoptera.SpeciesFile.org> [Accessed 13 December 2023]
- Deng WA (2011) A taxonomic study on the genus *Macromotettixoides* Zheng (Orthoptera, Tetrigoidea, Metrodorinae). *Acta Zootaxonomica Sinica* 36(3): 543–546.
- Deng WA (2016) Taxonomic Study of Tetrigoidea from China. Huazhong Agricultural University, Ph.D. Dissertation, 341 pp.
- Deng WA, Zheng ZM, Wei SZ (2007) Fauna of Tetrigoidea from Yunnan and Guangxi. Science and Technology Press, Nanning, 458 pp.
- Deng WA, Lei CL, Zheng ZM, Li XD, Lin LL, Lin MP (2014) Description of a new species of the genus *Macromotettixoides* Zheng (Orthoptera: Tetrigoidea: Metrodorinae) from China. *Neotropical Entomology* 43: 547–554. <https://doi.org/10.1007/s13744-014-0237-6>
- Deng WA, Zheng ZM, Li XD, Lin MP, Wei SZ, Yuan BD, Lin LL (2015) The groundhopper fauna (Orthoptera: Tetrigidae) of Shiwanshan (Guangxi, China) with description of three new species. *Zootaxa* 3925(2): 151–178. <https://doi.org/10.11646/zootaxa.3925.2.1>
- Deng WA, Zhang RJ, Li XD, Xin L (2021) The complete chloroplast genome of *Saussurella borneensis* (Orthoptera: Tetrigoidea) from China and its phylogenetic analysis. *Mitochondrial DNA Part B* 6(9): 2739–2740. <https://doi.org/10.1080/23802359.2021.1966336>
- Devriese H, Husemann M (2023) *Afrosystolederus garmsi* (Orthoptera, Tetrigidae), a new genus and species from Mount Gibi (Liberia) with remarks on *Systolederus*, *Pseudosystolederus* and *Teredorus*. *Zootaxa* 5258(3): 331–341. <https://doi.org/10.11646/zootaxa.5258.3.6>
- Donath A, Jühling F, Al-Arab M, Bernhart SH, Reinhardt F, Stadler PF, Middendorf M, Berrnt M (2019) Improved annotation of protein-coding genes boundaries in metazoan mitochondrial genomes. *Nucleic Acids Research* 47(20): 10543–10552. <https://doi.org/10.1093/nar/gkz833>
- Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* 7: 1–8. <https://doi.org/10.1186/1471-2148-7-214>
- Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* 29(8): 1969–1973. <https://doi.org/10.1093/molbev/mss075>
- Fan CM, Li M, Mao BY (2023) Descriptions of two new species of the genus *Macromotettixoides* (Orthoptera: Tetrigidae: Metrodorinae) from China. *Zootaxa* 5306(1): 127–134. <https://doi.org/10.11646/zootaxa.5306.1.6>
- Fang N, Xuan WJ, Zhang YY, Jiang GF (2010) Molecular phylogeny of Chinese Tetrigoids (Orthoptera, Tetrigoidea) based on the mitochondrial cytochrome C oxidase I gene. *Acta Zootaxonomica Sinica* 35: 696–702.
- Grant JR, Enns E, Marinier E, Mandal A, Herman EK, Chen C, Graham M, Van Domselaar G, Stothard P (2023) Proksee: In-depth characterization and visualization of bacterial genomes. *Nucleic Acids Research* 51(W1): W484–492. <https://doi.org/10.1093/nar/gkad326>
- Guan DL, Huang CM, Deng WA (2024) Reassessment of the phylogenetics of two pygmy grasshopper generic groups *Tetrix* and *Systolederus* through mitochondrial phylogenomics using four new mitochondrial genome assemblies. *Insects* 15(3): 174. <https://doi.org/10.3390/insects15030174>

- Günther K (1939) Revision der Acrydiinae (Orthoptera), III. Sectio Amorphopi (Metrodorae Bol. 1887 auct.). Abhandlungen und Berichte der Museum für Tierkunde und Volkerkunde zu Dresden (Ser. A: Zool.) (N.F.) 20: 16–335.
- Han YA, Li M, Mao BY (2020) A taxonomic study of the genus *Macromotettixoides* (Tetrigidae, Metrodorinae) with descriptions of two new species and two newly discovered males. Zootaxa 4718(4): 562–572. <https://doi.org/10.11646/zootaxa.4718.4.9>
- Hancock JL (1910) Third paper on the Tetriginae (Orthoptera) in the Oxford University Museum. Transactions of the Entomological Society of London 58(3): 346–365. <https://doi.org/10.1111/j.1365-2311.1910.tb01173.x>
- Hancock JL (1915) Indian Tetriginae (Acrydiinae). Records of the Indian Museum 11(1): 55–137. <https://doi.org/10.26515/rzsi/v11/i1/1915/163610>
- Hedges SW, Thomas MJ, Wang Y (2014) A remarkable new pygmy grasshopper (Orthoptera, Tetrigidae) in Miocene amber from the Dominican Republic. ZooKeys 30(429): 87–100. <https://doi.org/10.3897/zookeys.429.8020>
- Hornig MK, Haug C, Schneider JW, Haug JT (2018) Evolution of reproductive strategies in dictyopteran insects—clues from ovipositor morphology of extinct roachoids. Acta Palaeontologica Polonica 63(1): 1–24. <https://doi.org/10.4202/app.00324.2016>
- Huang CM, Deng WA, Zhang RJ, He CX (2022) The molecular phylogenetic analysis of some species of Tetriginae based on *COI*, 16S rRNA & 18S rRNA genes. Genomics and Applied Biology 41(5): 970–980.
- Kalogianni E (1995) Physiological properties of wind-sensitive and tactile trichoid sensilla on the ovipositor and their role during oviposition in the locust. Journal of Experimental Biology 198(6): 1359–1369. <https://doi.org/10.1242/jeb.198.6.1359>
- Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermiin LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Kasalo N (2022) *Metamazarredia* is *Rosacris* is *Mazarredia* (Orthoptera: Tetrigidae) – a long-awaited sequel to a taxonomic dilemma. Zootaxa 5200(2): 495–500. <https://doi.org/10.11646/zootaxa.5200.5.7>
- Kasalo N, Skejo J, Husemann M (2023a) DNA barcoding of pygmy hoppers – the first comprehensive overview of the BOLD Systems’ Data Shows Promise for Species Identification. Diversity 15(6): 696. <https://doi.org/10.3390/d15060696>
- Kasalo N, Yong S, Rebrina F, Skejo J (2023b) Definition of the tribe Metrodorini (Orthoptera: Tetrigidae) with notes on biogeography and evolution of Metrodorinae and Cladonotinae. Acta Entomologica Musei Nationalis Pragae 63(1): 187–193. <https://doi.org/10.37520/aemnp.2023.010>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular Biology and Evolution 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kluge NJ (2016) Structure of ovipositors and Cladoendosis of Saltatoria, or Orchesopia. Entomological Review 96: 1015–1040. <https://doi.org/10.1134/S0013873816080078>
- Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G, Hedges SB (2022) TimeTree 5: an expanded resource for species divergence times. Molecular Biology and Evolution 39(8): msac174. <https://doi.org/10.1093/molbev/msac174>
- Li M, Deng Y, Yin FX, Mao BY (2020a) A new species of *Macromotettixoides* (Orthoptera: Tetrigidae: Metrodorinae) from Yunnan, China. Entomotaxonomia 42(2): 110–115.
- Li XD, Wang YQ, Deng WA, Rong WT, Li R (2020b) First record of mitochondrial genome of *Teredorus nigropennis* (Orthoptera: Tetrigidae) and phylogenetic analysis. Mitochondrial DNA B Resour 5(2): 1145–1146. <https://doi.org/10.1080/23802359.2020.1730259>

- Li RR, Li M, Yan J, Bai M, Zhang HF (2021a) Five mitochondrial genomes of the genus *Eysarcoris* Hahn, 1834 with phylogenetic implications for the Pentatominae (Hemiptera: Pentatomidae). *Insects* 12(7): 597. <https://doi.org/10.3390/insects12070597>
- Li R, Ying XL, Deng WA, Rong WT, Li XD (2021b) Mitochondrial genomes of eight Scelimeninae species (Orthoptera) and their phylogenetic implications within Tetrigoidea. *PeerJ* 9: e10523. <https://doi.org/10.7717/peerj.10523>
- Li XJ, Liu YX, Lin LL (2021c) Comparative mitogenomes and phylogenetic analysis reveal taxonomic relationship of genera *Teredorus* and *Systolederus* (Orthoptera, Tetrigoidea). *Zootaxa* 5027(1): 127–135. <https://doi.org/10.11646/zootaxa.5027.1.7>
- Lin LL (2014) Mitochondrial Genome Analysis and Phylogenetic Relationship of Three Species of Tetrigoidea. Shaanxi Normal University, Ph.D. Dissertation, 140 pp.
- Lin MP, Li XD, Wei SZ, Deng WA (2015) Mitochondrial *COI* gene partial sequences and phylogenetic relationship of some groups of Tetrigoidea. *Journal of Huazhong Agricultural University* 34(6): 40–48.
- Lin LL, Li XJ, Zhang HL, Zheng ZM (2017) Mitochondrial genomes of three Tetrigoidea species and phylogeny of Tetrigoidea. *PeerJ* 5: e4002. <https://doi.org/10.7717/peerj.4002>
- Long Y, Teng CL, Huang CM, Zhang RJ, Deng WA, Lin LL (2023) Twenty-three new synonyms of the Eastern common groundhopper, *Tetrix japonica* (Bolívar, 1887) (Orthoptera, Tetrigidae). *ZooKeys* 1187: 135–167. <https://doi.org/10.3897/zookeys.1187.110067>
- Lu XY, Deng WA (2021) New genus and new species of the subfamily Metrodorinae from China (Orthoptera: Tetrigidae). *Zootaxa* 4964(2): 345–362. <https://doi.org/10.11646/zootaxa.4964.2.6>
- Luo JL, Zhang RJ, Deng WA (2024) First mitogenomic characterization of *Macromotettixoides* (Orthoptera, Tetrigidae), with the descriptions of two new species. *ZooKeys* 1195: 95–120. <https://doi.org/10.3897/zookeys.1195.112623>
- Minh BQ, Nguyen MA, Von Haeseler A (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>.
- Muhammad AA, Tan MK, Abdullah NA, Azirun MS, Bhaskar D, Skejo J (2018) An annotated catalogue of the pygmy grasshoppers of the tribe Scelimenini Bolívar, 1887 (Orthoptera: Tetrigidae) with two new *Scelimena* species from the Malay Peninsula and Sumatra. *Zootaxa* 4485(1): 1–70. <https://doi.org/10.11646/zootaxa.4485.1.1>
- Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Nie R, Vogler AP, Yang XK, Lin M (2021) Higher-level phylogeny of longhorn beetles (Coleoptera: Chrysomeloidea) inferred from mitochondrial genomes. *Systematic Entomology* 46(1): 56–70. <https://doi.org/10.1111/syen.12447>
- Pavón-Gozalo P, Manzanilla J, García-Paris M (2012) Taxonomy and morphological characterization of *Allotettix simony* (Bolívar, 1890) and implications for the systematics of Metrodorinae (Orthoptera: Tetrigidae). *Zoological Journal of the Linnean Society* 164(1): 52–70. <https://doi.org/10.1111/j.1096-3642.2011.00764.x>
- Peng T, Shi TZ, Ding JH, Zha LS (2021) Two *Macromotettixoides* species from Hainan Island, PR China, with taxonomic notes for the genus (Metrodorinae, Tetrigidae). *Entomological News* 130(1): 47–60. <https://doi.org/10.3157/021.130.0104>
- Sheffield NC, Hiatt KD, Valentine MC, Song H (2010) Mitochondrial genomics in Orthoptera using MOSAS. *Mitochondrial DNA* 21(3–4): 87–104. <https://doi.org/10.3109/19401736.2010.500812>

- Skejo J, Gupta SK, Chandra K, Panhwar WA, Franjevic D (2019) Oriental macropterous leaf-mimic pygmy grasshoppers – genera *Oxyphyllum* and *Paraphyllum* (Orthoptera: Tetrigidae) and their taxonomic assignment. *Zootaxa* 4590(5): 546–560. <https://doi.org/10.11646/zootaxa.4590.5.3>
- Skejo J, Kasalo N, Thomas MJ, Heads SW (2024) A new long-winged pygmy grasshopper in Eocene Baltic amber raises questions about the evolution of reduced tegmenula in Tetrigidae (Orthoptera). *Journal of Orthoptera Research* 33(1): 1–26. <https://doi.org/10.3897/jor.33.105144>
- Song H, Amédégnato C, Cigliano MM, Desutter-Grandcolas L, Heads SW, Huang Y, Otte D (2015) 300 million years of diversification: elucidating the patterns of orthopteran evolution based on comprehensive taxon and gene sampling. *Cladistics* 31(6): 621–651. <https://doi.org/10.1111/cla.12116>
- Song N, Li XX, Yin XM, Li X, Yin J, Pan P (2019) The mitochondrial genomes of palaeopteran insects and insights into the early insect relationships. *Scientific Reports* 9(1): 17765. <https://doi.org/10.1038/s41598-019-54391-9>
- Storozhenko SY (2014) To synonymy of the genus *Xistrella* Bolívar, 1909 (Orthoptera: Tetrigidae, Metrodorinae). *Far Eastern Entomologist* 277: 47–48.
- Subedi M (2022) A new genus and a new groundhopper species from Nepal (Orthoptera: Tetriginae: *Skejotettix netrajyoti* gen. et sp. nov.). *Zootaxa* 5205(1): 35–54. <https://doi.org/10.11646/zootaxa.5205.1.3>
- Tan MK, Artchawakom T (2015) A new species from the genus *Gorochovitettix* (Tetrigidae: Metrodorinae) from Thailand. *Zootaxa* 3990(3): 444–450. <https://doi.org/10.11646/zootaxa.3990.3.9>
- Tumbrinck J, Telnov D (2014) Taxonomic revision of the Cladonotinae (Orthoptera: Tetrigidae) from the islands of South-East Asia and from Australia, with general remarks to the classification and morphology of the Tetrigidae and descriptions of new genera and species from New Guinea and New Caledonia. *Biodiversity, Biogeography and Nature Conservation in Wallacea and New Guinea*. 2: 345–396.
- Tumbrinck J (2019) Taxonomic and biogeographic revision of the genus *Lamellitettigodes* (Orthoptera: Tetrigidae) with description of two new species and additional notes on *Lamellitettix*, *Probolotettix*, and *Scelimena*. *Journal of Orthoptera Research* 28(2): 167–180. <https://doi.org/10.3897/jor.28.34605>
- Wang YZ (2001) A new genus and new species of Metrodoridae (Orthoptera: Tetrigidae) recorded in Yunnan, China. *Journal of Southwest Agricultural University* 23: 147–148.
- Wang MQ (2022) Phylogenetic Analyses of Metrodorinae (Orthoptera: Tetrigidae) based on the Mitochondrial Genome. Dali University, M.Sc. thesis, 116 pp.
- Xiao B, Chen W, Hu CC, Jiang GF (2012a) Complete mitochondrial genome of the groundhopper *Alulatettix yunnanensis* (Insecta: Orthoptera: Tetrigoidea). *Mitochondrial DNA* 23(4): 286–287. <https://doi.org/10.3109/19401736.2012.674122>
- Xiao B, Feng X, Miao WJ, Jiang GF (2012b) The complete mitochondrial genome of grouse locust *Tetrix japonica* (Insecta: Orthoptera: Tetrigoidea). *Mitochondrial DNA* 23(4): 288–289. <https://doi.org/10.3109/19401736.2012.674123>
- Xie JM, Chen YR, Cai GJ, Cai RL, Hu Z, Wang H (2023) Tree Visualization By One Table (tvBOT): a web application for visualizing, modifying and annotating phylogenetic trees. *Nucleic Acids Research* 51(W1): W587–592. <https://doi.org/10.1093/nar/gkad359>
- Yang J (2017a) Mitochondrial genome sequencing of three Tetrigoidea and comparative analysis of mitochondrial genome of Acrididae. Shaanxi Normal University, Master thesis, 87 pp.

- Yang J, Lu C, Zhang ZB, Huang Y, Lin LL (2017b) Mitochondrial genomes of two pygmy grasshoppers (Orthoptera: Tetrigoidea) and a comparative analysis of Caelifera mitogenomes. *Zoological Science* 34(4): 287–294. <https://doi.org/10.2108/zs160217>
- Zha LS, Yu FM, Boonmee S, Eungwanichayapant PD, Wen TC (2017) Taxonomy of *Macromotettixoides* with the description of a new species (Tetrigidae, Metrodorinae). *ZooKeys* (645): 13–25. <https://doi.org/10.3897/zookeys.645.9055>
- Zhang RJ, Zhao CL, Wu FP, Deng WA (2020) Molecular data provide new insights into the phylogeny of Cladonotinae (Orthoptera: Tetrigoidea) from China with the description of a new genus and species. *Zootaxa* 4809(3): 547–559. <https://doi.org/10.11646/zootaxa.4809.3.8>
- Zhang RJ, Xin L, Deng WA (2021) Characterization of the complete mitochondrial genome of *Tripetaloceroides tonkinensis* (Orthoptera: Tetrigoidea) from China and its phylogenetic analysis. *Mitochondrial DNA Part B* 6(7): 1990–1991. <https://doi.org/10.1080/23802359.2021.1938724>
- Zhang RJ, Wang JG, Huang DL, Deng WA (2023) Life cycle and other biological characteristics of *Tetrix japonica*. *Journal of Environmental Entomology* 45(4): 965–973.
- Zhao XM, Su YZ, Shao T, Fan ZY, Cao LL, Liu WM, Zhang JZ (2022) Cuticle protein gene *LmCP8* is involved in the structural development of the ovipositor in the migratory locust *Locusta migratoria*. *Insect Molecular Biology* 31(6): 747–759. <https://doi.org/10.1111/imb.12801>
- Zheng ZM (1998) A study of Tetrigoidea from Xishuangbanna (Orthoptera). *Acta Zootaxonomica Sinica* 23(2): 161–180.
- Zheng ZM (2005a) Fauna of the Tetrigoidea from Western China. Science Press, Beijing, 501 pp.
- Zheng ZM (2013a) A new species of the genus *Macromotettixoides* Zheng (Orthoptera: Metrodorida) from Fujian Province. *Entomotaxonomia* 35(4): 241–244.
- Zheng ZM (2013b) Key to the species of *Systolederus*, *Hyboella*, *Bolivaritettix* (Orthoptera: Tetrigoidea: Metrodoridae) from China with descriptions of three new species. *Journal of Shangqiu Normal University* 29(12): 1–13.
- Zheng ZM, Jiang GF (2000) New genus and new species of Metrodoridae from Guangxi (Orthoptera: Tetrigoidea). *Acta Zootaxonomica Sinica* 25(4): 402–405.
- Zheng ZM, Jiang GF (2002) One new genus and seven new species of Tetrigoidea from southern region of Guangxi. *Zoological Research* 23(5): 409–416.
- Zheng ZM, Ou XH (2004) Three new species of the genus *Formosatettix* Tinkham from Hengduan Mountain region of western Yunnan (Orthoptera, Tetrigidae). *Acta Zootaxonomica Sinica* 29(1): 107–109.
- Zheng ZM, Shi FM (2009) Five new species of Tetrigoidea from Jiangxi Province of China (Orthoptera). *Acta Zootaxonomica Sinica* 34(3): 572–577.
- Zheng ZM, Wei ZM, Jiang GF (2005b) A new genus and a new species of Metrodoridae (Orthoptera) from China. *Acta Zootaxonomica Sinica* 30(2): 366–367.
- Zheng ZM, Li P, Wan B (2006) A revision of *Macromotettixoides* Zheng from China (Orthoptera: Metrodoridae). *Journal of Huazhong Agricultural University* 25(6): 603–605.