

Synthesis of Tetrahydroazepines through Silyl Aza-Prins Cyclization Mediated by Iron(III) Salts

Victoria Sinka,* Israel Fernández, and Juan I. Padrón*



Cite This: *J. Org. Chem.* 2022, 87, 11735–11742



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: A new methodology for the synthesis of seven-membered unsaturated azacycles (tetrahydroazepines) was developed. It is based on the powerful aza-Prins cyclization in combination with the Peterson-type elimination reaction. In a single reaction step, a C–N, C–C bond and an endocyclic double bond are formed. Under mild reaction conditions and using iron(III) salts as sustainable catalysts, tetrahydroazepines with different degrees of substitution are obtained directly and efficiently. DFT calculations supported the proposed mechanism.



INTRODUCTION

Unsaturated seven-membered ring azacycles (tetrahydroazepines) are found in numerous natural and non-natural products with remarkable pharmaceutical activity.^{1–3} Also, they serve as substructures of more complex molecules or as precursors of hydroazepines with different biological properties.^{4,5} Most of these natural products come from terrestrial sources such as balanol (1), an unusual metabolite isolated from fungus *Verticillium balanoides*, which is a potent inhibitor of PKC¹ or (–)-galanthamine (2), commercially known as Reminyl and used for the symptomatic treatment of Alzheimer's (Figure 1).⁶ The unnatural azepane 3 proved to be an excellent agent against lung cancer, where the heterocyclic nitrogenous core is essential for its bioactivity (IC₅₀ of 4.18 nM) (Figure 1).²

Classical methods to synthesize this type of heterocycles include Brønsted or Lewis acid-mediated cyclizations,^{7,8} atom-transfer radical cyclization (ATRC),⁹ cycloadditions,^{10,11} conjugate addition cyclizations,^{12,13} ring expansions (cyclopropanes,¹⁴ aziridines,¹⁵ azetidines,^{16–18} 2-cyano-6-oxazolopiperidine¹⁹), and ring-closing metathesis.^{20–25} Among the different types of acid-mediated cyclizations, the aza-Prins cyclization is a powerful tool for obtaining nitrogenated heterocycles. It has been widely used for the synthesis of piperidines and pyrrolidines.^{26–33} However, there are few examples of synthesis of tetrahydroazepines through this methodology.^{7,34} In 2016, Barbero and co-workers achieved the synthesis of azepane rings, with an exocyclic double bond, through a diastereoselective silyl aza-Prins cyclization mediated by InCl₃ (Scheme 1).⁷ Nevertheless, this reaction, inspired by the work of the Dobbs' group, is a relatively high-energy demanding reaction.³⁰ Therefore, new methods for the synthesis of seven-membered azacycles via Prins cyclizations remain challenging and highly desirable.

RESULTS AND DISCUSSION

Based on our previous work, which allowed the diastereoselective synthesis of oxepenes through the Prins-Peterson cyclization (PPC) strategy,³⁵ we decided to approach the nitrogen version and thus synthesize different tetrahydroazepines. If successful, it would constitute a direct and straightforward method that, using a sustainable iron(III) catalyst, builds a C–N, C–C bond and an endocyclic double bond in a single reaction step (Scheme 1).

We started with the simplest tetrahydroazepines, that is, bearing a single substituent that comes from the corresponding aldehyde. 1-Amino-3-triphenylsilyl-4-pentenenes **6a–b**, precursors of the silyl aza-Prins cyclization (SAPC), could be accessed in three reaction steps (see Supporting Information). Next, the SAPC assays were performed with different sustainable metal catalysts, based on our previous experience for oxepene synthesis, affording the desired tetrahydroazepine contaminated with variable amounts of the corresponding pyrrolidine. The best results were obtained using substoichiometric amounts (0.1 equiv) of FeBr₃ and FeCl₃, the yield and ratio being slightly higher when the former was used (see Supporting Information).

Once FeBr₃ was set as the best catalyst, we focused on optimizing the reaction conditions to avoid the formation of pyrrolidine **8** resulting from the intramolecular hydroamination side reaction, whose relative stereochemistry was determined by ¹H-GOESY NMR experiments (Table 1).³⁶

Received: June 13, 2022

Published: August 17, 2022







Figure 1. Representative bioactive hydroazepines/tetrahydroazepines.

Scheme 1. Previous Reports on Silyl Aza-Prins Cyclization and Our Work to Synthesize Unsaturated Seven-Membered Ring Azacycles

a) Previous work (Barbero, 2016)



b) This work





Table 1. Optimization of Reaction Conditions for SAPC Reaction with FeBr₃^a





entry	FeBr ₃ (equiv)	T (°C)	time (min)	yield 7a:8 (%)
1	0.20	r.t.	15	58:35
2	0.10	r.t.	35	67:10
3	0.05	r.t.	35	82:13
4	0.20	0	30	83:13
5	0.10	0	35	90:0
6	0.05	0	300	89:7
7	0.20	-20	60	85:0
8	0.10	-20	300	85:0
9	0.05	-20	120	52:35

^aReaction conditions: **6a** (0.20 mmol), isovaleraldehyde (0.24 mmol), FeBr₃ (0.01–0.04 mmol), dry DCM (0.1 M). Isolated yield. The conversion in all cases was 100%.


The increase in catalyst loading at room temperature, with respect to that initially tested, favored the undesired azacycle **8** (Table 1, entry 1), whereas the use of 0.05 equiv of FeBr₃ increased the yield of tetrahydroazepine **7a** (Table 1, entries 2 and 3). We observed that by decreasing the temperature, the formation of the pyrrolidine was reduced (Table 1, entries 4 and 7). However, lower catalyst loading at low temperatures led to long reaction times, which did not avoid the presence of azacycle **8** (Table 1, entries 6 and 9). The best result involved a compromise between 0 °C and 0.10 equiv of FeBr₃, suppressing the formation of azacycle **8** and increasing the yield of tetrahydroazepine **7a** up to 90% (Table 1, entry 5). With the optimized reaction conditions in hand, we set out to explore the scope of this SAPC. The reaction conditions could be applied to tosylated and mesylated amines **6a–b**, respectively, and to a variety of aldehydes (Scheme 2).

Scheme 2. Scope of Monosubstituted Tetrahydroazepines 7 Synthesis From SAPC^a




^aReaction conditions: **6a–6b** (0.20 mmol), aldehyde (0.24–0.40 mmol), FeBr₃ (0.02, mmol), dry DCM (0.1 M). Isolated yield. The conversion in all cases was 100%. The pyrrolidinic byproduct *rac-8* was not observed in any example made. ^b2 g scale of **6a** (4.0 mmol) afforded 60% yield of **7a** (741 mg, 2.41 mmol).

Systems with aliphatic chains such as isovaleraldehyde and heptanal, with both tosylated and mesylated 1-amino-3-triphenylsilyl-4-pentenes **6a–b**, respectively, gave yields of 92, 80, 73, and 81% (tetrahydroazepines **7a**, **7b**, **7e**, and **7f**) (Scheme 2). Likewise, good results were observed with 4-pentenal, which led to tetrahydroazepines **7c** and **7g** with an

Table 2. Optimization of Reaction Conditions for SAPC Reaction with FeCl₃^a

entry	FeCl ₃ (equiv)	DCM (M)	T (°C)	time (min)	yield ^{b,g} 10a:11 (%)
1 ^c	0.1	0.1	10	360	13:8
2	0.3	0.1	10	140	45:18
3	0.5	0.1	10	120	42:15
4 ^d	0.3	0.1	0	360	28:18
5 ^e	0.3	0.1	0 → r.t.	300	40:24
6	0.3	0.1	r.t.	240	40:22
7	0.6	0.2	r.t.	100	53:18
8	0.3	0.3	r.t.	120	47:22
9 ^f	1.0 + 0.3	0.1	-20 → r.t.	220	72:4

^aReaction conditions: **9a** (0.098 mmol), isovaleraldehyde (0.15 mmol), FeCl₃ (0.098–0.13 mmol), dry DCM. ^bIsolated yield. ^cConversion of 63%. ^dConversion of 54%. ^eThe order of addition was changed: DCM, aldehyde, FeCl₃, and amine **9a** in portions. ^fInitial FeCl₃ load was 1.0 equiv at -20 °C. After 3 h of reaction, another 0.3 equiv of catalyst was added at -20 °C, and after 10 min the bath was removed, leaving the reaction at room temperature. ^g**10a** is obtained as an inseparable mixture of *cis/trans* diastereomers by flash chromatography. The *trans* isomer was identified as the major one through NMR studies see [Supporting Information](#).

Scheme 3. Scope of Disubstituted Tetrahydroazepines Synthesis from Silyl Aza-Prins Cyclization^a


^aReaction conditions: **9a–b** (0.098 mmol), aldehyde (0.15 mmol), FeCl₃ (0.13, mmol), dry DCM (0.1 M). Isolated yield. The conversion in all cases was 100%. The pyrrolidinic byproduct **rac-11** was observed in small amounts (4% yield) in each example.

endocyclic and an exocyclic olefin. Reaction with phenylacetaldehyde and hydrocinnamaldehyde allowed a phenyl substituent to be incorporated into the heterocycles **7d**, **7h**, and **7i** in 65, 63, and 60% yield, respectively. Our SAPC protocol also worked when using an aldehyde with a benzoate carrier chain, so future derivatizations of tetrahydroazepine **7k** are feasible. However, there was no reactivity with aromatic aldehydes.

Subsequent efforts were devoted to extend this methodology to the synthesis of disubstituted tetrahydroazepine **10**. Once we synthesized the precursor amines **9** substituted at the β position with methyl and allyl groups (see [Supporting Information](#)), we made the preliminary SAPC assays. On this occasion, FeCl₃, which also gave good yields in the synthesis of monosubstituted tetrahydroazepines, was the catalyst that provided the best yields for the corresponding disubstituted tetrahydroazepines (see [Supporting Information](#)). The next optimization step consisted of adjusting the amount of FeCl₃, the concentration of the solvent, and the temperature ([Table 2](#)).

Because the 1-amino-3-triphenylsilyl-4-pentene **9a** showed a different reactivity than **6a–b** (see above), we decided to adjust the SAPC conditions again ([Table 2](#)). The amount of FeCl₃ was varied between 0.1 and 0.5 equiv, maintaining the temperature at 10 °C, the best result being 30 mol % of the catalyst ([Table 2](#), entries 1–3). Dropping to 0 °C prevented complete consumption of starting material, and lower yield of tetrahydroazepine **10a** was observed ([Table 2](#), entries 4 and 5). In addition, at room temperature, the concentration was increased to favor the intermolecular reaction to generate the desired disubstituted tetrahydroazepine **10a**. However, seven-membered azacycle **10a** and pyrrolidine byproduct **11** were obtained in 50 and 20% yield on average, respectively ([Table 2](#), entries 6–8). Therefore, we decided to increase the catalyst loading up to 1.0 equiv at low temperatures, to avoid the formation of pyrrolidine **11**. The starting material was completely consumed, adding an extra 0.3 equiv of FeCl₃ at -20 °C and allowing the reaction to reach the room temperature. Thus, a conversion of 100% and the predominant

Scheme 4. Mechanistic Proposal for SAPC Supported by DFT Calculations



formation (95%) of tetrahydroazepine **10a** was observed with 72% yield.

The scope of the SAPC reaction with amines **9a–b** was then tested (Scheme 3). On the one hand, the 1-amino-3-triphenylsilyl-4-pentene **9a** bearing a methyl unit was reactive with aliphatic aldehydes such as isovaleraldehyde and heptanal, giving tetrahydroazepines **10a** and **10b** in 72 and 61% yield, respectively. Likewise, it was possible to obtain tetrahydroazepine **10c** through SAPC with 4-pentenal (Scheme 3). On the other hand, the precursor amine **9b** bearing an allyl moiety at β position showed lower reactivity than the amine **9a**. Only SAPC with isovaleraldehyde was possible generating tetrahydroazepine **10d** in 60% yield (Scheme 3).

The mechanistic proposal for this transformation is based on that reported for the analogous process forming oxepenes³³ (Scheme 4). First, a condensation occurs between the 1-amino-3-triphenylsilyl-4-pentene **6a** and the Lewis acid-activated aldehyde. The zwitterionic species **12** leads to the amino-alcohol **13** and evolves to the iminium ion **14**. This last species is intramolecularly trapped by the double bond, generating the carbocation **15**, stabilized by the presence of the silyl group (β effect). Intermediate **15** undergoes a Peterson-type elimination, leading to tetrahydroazepines **7** and triphenylsilanol as a byproduct (Scheme 4).

Density functional theory (DFT) calculations were carried out [PCM(CH₂Cl₂)–B3LYP-D3/def2-TZVPP//PCM(CH₂Cl₂)–B3LYP-D3/def2-SVP level] to gain more insights into the proposed reaction mechanism. In particular, we focused on the key intramolecular cyclization step involving **13**, where the tosyl group was replaced by a mesyl group and R¹ = Me. Our calculations indicate that the release of the anion FeBr₃OH[−], thus forming **14**, which is endergonic ($\Delta G_R = 13.1$ kcal/mol), and the subsequent nucleophilic cyclization possesses a barrier of only 9.5 kcal/mol. Therefore, from **13**, this S_N1-type reaction is compatible with the reaction

conditions used in the experiments. Alternatively, **13** can evolve into an intimate ion pair **14'**, where the FeBr₃OH[−] fragment is tightly bonded to the cation **14** in a less endergonic reaction ($\Delta G_R = 8.7$ kcal/mol). Then, a similar nucleophilic cyclization occurs with a rather similar, low activation barrier of 9.0 kcal/mol. The initial slightly endergonic C–O(H)FeBr₃ bond rupture in **15** is efficiently compensated by the final Peterson elimination, which is strongly exergonic ($\Delta G_R > -45$ kcal/mol) therefore driving the complete catalytic cycle forward (see computed profiles in the Supporting Information).

CONCLUSIONS

In conclusion, we have developed an efficient and straightforward approach for the formation of mono- and disubstituted seven-membered Δ^4 -unsaturated azacycles. A variety of mono- and disubstituted tetrahydroazepines can be accessed through SAPC, and in a single reaction step, C–C, C–N, and endocyclic C=C bonds are formed. In addition, it has been shown that the functionalization of the precursor amines determines their reactivity. Increasing the substitution of tetrahydroazepine up to three substituents is currently ongoing in our laboratory.

EXPERIMENTAL SECTION

General Information. Reagents were obtained from commercial sources (Sigma-Aldrich, Merck, Alfa Aesar), without further purification. Solvents (DCM, Et₂O, THF, and DMF) were used from the PureSolv system. The dispensing system allows easy access to the anhydrous solvents. EtOH was purified by distillation and dried following the procedure in the literature.³⁵ Chemical reactions and the separation of the crudes were monitored by thin-layer chromatography (TLC). TLC was performed on aluminum foil sheets 60 F254 manufactured by MERCK. The solvent or solvent mixture was *n*-hexane/ethyl acetate (EtOAc) in different ratios. Flash column chromatography was performed using silica gel (0.015–0.04 mm) and

n-hexane/EtOAc solvent systems. Automated flash column chromatography was performed using the Biotage Isolera System (Isolera Prime). It includes simultaneous UV detection on all wavelengths and baseline correction, which enable detection of poor UV absorbing compounds. NMR spectra were recorded on Bruker Avance instruments. ¹H NMR spectra were recorded at 400, 500, and 600 MHz, and ¹³C NMR spectra were recorded at 100, 125, and 150 MHz, VTU 298.0 K. The residual solvent peak was used as an internal reference (CDCl₃: δ_H 7.26, δ_C 77.0).³⁶ High-resolution mass spectra were recorded on an LCT Premier XE mass spectrometer. It is provided with different ionization sources: electrospray (ESI), an atmospheric pressure chemical ionization (APCI) source, and an orthogonal acceleration time of flight (oa-TOF) analyzer that provides high sensitivity, resolution, and accurate mass measurement.

General Procedure (1) for Silyl Aza-Prins Cyclization of 1-Amino-3-triphenylsilyl-4-pentenes 6a–b. To a solution of amines 6a–b (0.30–0.12 mmol, 1.0 equiv) in dry DCM (3.0–1.2 mL, 0.1 M) at 0 °C were added the aldehyde (0.45–0.18 mmol, 1.5 equiv) and the catalyst (0.030–0.012 mmol, 0.1 equiv). Once the reaction was complete, checked by TLC, it was quenched with water. The layers were separated, and the aqueous phase was extracted three times with DCM. The combined organic layers were dried over anhydrous MgSO₄, filtrated, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/EtOAc solvent system).

General Procedure (2) for Silyl Aza-Prins Cyclization of 1-Amino-3-triphenylsilyl-4-pentene 9a–b. To a solution of amines 9a–b (0.098–0.074 mmol, 1.0 equiv) in DCM (1.0–0.7 mL, 0.1 M) at –20 °C under inert atmosphere were added the aldehyde (0.15–0.11 mmol, 1.5 equiv) and FeCl₃ (0.098–0.074 mmol, 1.0 equiv). After stirring the mixture for 2 h at –20 °C, an extra amount of FeCl₃ (0.029–0.022 mmol, 0.3 equiv) was added, and then, the bath was removed. The reaction mixture was stirred at room temperature for 30 min. Once the reaction was completed, it was quenched with water. The phases were separated, and the aqueous layer was extracted with 3× DCM. The combined organic phases were dried over anhydrous MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/DCM/MeOH 69:29:2 solvent system).

2-Isobutyl-1-tosyl-2,3,6,7-tetrahydro-1H-azepine (7a). Following the general procedure (1), to a solution of amine 6a (80 mg, 0.16 mmol, 1.0 equiv) in 1.6 mL of dry DCM (0.1 M) at 0 °C were added isovaleraldehyde (21 μL, 0.19 mmol, 1.2 equiv) and FeBr₃ (4.8 mg, 0.019 mmol, 0.1 equiv) to obtain 45 mg of tetrahydroazepine 7a as a pale yellow oil (0.147 mmol, 92%). *R*_f = 0.63 (*n*-hexane/EtOAc 80:20); ¹H NMR (CDCl₃, 400 MHz): δ = 7.69 (d, *J* = 8.1 Hz, 2H), 7.26 (d, *J* = 7.3 Hz, 2H), 5.70 (m, 1H), 5.55 (m, 1H), 4.13 (m, 1H), 3.66 (brddd, *J* = 14.5, 4.8 & 3.4 Hz, 1H), 3.14 (brddd, *J* = 14.0, 11.0 & 2.2 Hz, 1H), 2.40 (s, 3H), 2.38–2.27 (m, 2H), 2.15 (m, 2H), 1.37 (m, 3H), 0.82 (d, *J* = 6.4 Hz, 3H), 0.80 (d, *J* = 6.3 Hz, 3H); ¹³C{¹H}-NMR (CDCl₃, 125 MHz): δ = 142.8 (C), 138.6 (C), 130.7 (CH), 129.5 (2× CH), 127.2 (CH), 127.1 (2× CH), 53.4 (CH), 41.2 (CH₂), 40.2 (CH₂), 32.5 (CH₂), 30.3 (CH₂), 24.6 (CH), 22.9 (CH₃), 22.3 (CH₃), 21.5 (CH₃); HRMS (ESI⁺): *m/z* [M + Na]⁺ calcd for C₁₇H₂₅NO₂NaS, 330.1505; found, 330.1504.

2,3-Trans-2-methyl-1-tosyl-3-(triphenylsilyl) pyrrolidine (rac-8). A solution of 60 mg of amine 6a (0.12 mmol, 1.0 equiv) in 1.2 mL of dry DCM (0.1 M) at room temperature in the presence of 59 mg of FeBr₃ (0.024 mmol, 0.20 equiv) gave 21 mg of pyrrolidine rac-8 (0.042 mmol, 35% yield) as a white amorphous solid. *R*_f = 0.52 (*n*-hexane/EtOAc 80:20); ¹H NMR (CDCl₃, 400 MHz): δ = 7.50 (d, *J* = 8.1 Hz, 2H), 7.47–7.39 (m, 9H), 7.37–7.30 (m, 6H), 7.14 (d, *J* = 8.1 Hz, 2H), 3.77 (m, 1H), 3.40 (m, 1H), 3.29 (m, 1H), 2.42 (s, 3H), 1.95 (m, 2H), 1.29 (m, 4H); ¹³C{¹H}-NMR (CDCl₃, 100 MHz): δ = 142.7 (C), 135.9 (6× CH), 135.7 (C), 135.4 (C), 133.0 (2× C), 129.7 (3× CH), 129.5 (2× CH), 128.0 (6× CH), 127.3 (2× CH), 59.4 (CH), 49.3 (CH₂), 33.5 (CH), 28.8 (CH₂), 24.0 (CH₃), 21.5 (CH₃); HRMS (ESI⁺): *m/z* [M + Na]⁺ calcd for C₃₀H₃₁NO₂NaSi, 520.1742; found, 520.1745.

2-Hexyl-1-tosyl-2,3,6,7-tetrahydro-1H-azepine (7b). Following the general procedure (1), to a solution of amine 6a (0.150 mg, 0.30 mmol, 1.0 equiv) in 3.0 mL of dry DCM (0.1 M) at 0 °C were added heptanal (51 μL, 0.36 mmol, 1.2 equiv) and FeBr₃ (9.0 mg, 0.030 mmol, 0.1 equiv) to obtain 81 mg of tetrahydroazepine 7b as a pale yellow oil (0.24 mmol, 80%). *R*_f = 0.57 (*n*-hexane/EtOAc 80:20); ¹H NMR (CDCl₃, 400 MHz): δ = 7.70 (d, *J* = 8.2 Hz, 2H), 7.26 (d, *J* = 8.3 Hz, 2H), 5.69 (m, 1H), 5.56 (m, 1H), 4.03 (m, 1H), 3.68 (brddd, *J* = 14.5, 5.1 & 3.1 Hz, 1H), 3.14 (brddd, *J* = 13.3, 10.9 & 2.1 Hz, 1H), 2.41 (s, 3H), 2.38–2.28 (m, 2H), 2.19 (m, 2H), 1.46 (m, 2H), 1.29–1.11 (m, 6H), 1.10–0.98 (m, 2H), 0.85 (t, *J* = 7.0 Hz, 3H); ¹³C{¹H}-NMR (CDCl₃, 100 MHz): δ = 142.7 (C), 138.5 (C), 130.6 (CH), 129.4 (2× CH), 127.0 (CH), 126.9 (2× CH), 55.4 (CH), 41.3 (CH₂), 32.3 (CH₂), 31.6 (CH₂), 31.2 (CH₂), 30.5 (CH₂), 29.0 (CH₂), 26.1 (CH₂), 22.5 (CH₂), 21.4 (CH₃), 14.0 (CH₃); HRMS (ESI⁺): *m/z* [M + Na]⁺ calcd for C₁₉H₂₉NO₂NaS, 358.1817; found, 358.1821.

2-(But-3-en-1-yl)-1-tosyl-2,3,6,7-tetrahydro-1H-azepine (7c). Following the general procedure (1), to a solution of amine 6a (0.150 mg, 0.30 mmol, 1.0 equiv) in 3.0 mL of dry DCM (0.1 M) at 0 °C were added 4-pentenal (61 μL, 0.60 mmol, 2.0 equiv) and FeBr₃ (9.0 mg, 0.030 mmol, 0.1 equiv) to obtain 61 mg of tetrahydroazepine 7c as a pale yellow oil (0.20 mmol, 67%). *R*_f = 0.54 (*n*-hexane/EtOAc 80:20); ¹H NMR (CDCl₃, 400 MHz): δ = 7.68 (d, *J* = 8.4 Hz, 2H), 7.26 (d, *J* = 7.8 Hz, 2H), 5.71 (m, 2H), 5.54 (m, 1H), 4.94 (m, 2H), 4.06 (m, 1H), 3.65 (ddd, *J* = 14.5, 5.4 & 3.3 Hz, 1H), 3.18 (ddd, *J* = 14.6, 10.6 & 2.4 Hz, 1H), 2.40 (s, 3H), 2.37–2.26 (m, 2H), 2.18 (m, 2H), 1.90 (m, 2H), 1.59 (m, 2H); ¹³C{¹H}-NMR (CDCl₃, 100 MHz): δ = 142.8 (C), 138.3 (C), 137.8 (CH), 130.8 (CH), 129.5 (2× CH), 127.0 (2× CH), 126.7 (CH), 114.8 (CH₂), 55.2 (CH), 41.5 (CH₂), 31.8 (CH₂), 30.7 (CH₂), 30.3 (CH₂), 30.2 (CH₂), 21.4 (CH₃); HRMS (ESI⁺): *m/z* [M + Na]⁺ calcd for C₁₇H₂₃NO₂NaS, 328.1347; found, 328.1349.

2-Benzyl-1-tosyl-2,3,6,7-tetrahydro-1H-azepine (7d). Following the general procedure (1), to a solution of amine 6a (0.150 g, 0.30 mmol, 1.0 equiv) in 3.0 mL of dry DCM (0.1 M) at 0 °C were added phenylacetaldehyde (43 μL, 0.36 mmol, 1.2 equiv) and FeBr₃ (9.0 mg, 0.030 mmol, 0.1 equiv) to obtain 67 mg of tetrahydroazepine 7d as a pale yellow oil (0.195 mmol, 65%). *R*_f = 0.46 (*n*-hexane/EtOAc 80:20); ¹H NMR (CDCl₃, 400 MHz): δ = 7.62 (m, 2H), 7.23 (m, 5H), 7.12 (m, 2H), 5.73 (m, 1H), 5.55 (m, 1H), 4.30 (m, 1H), 3.64 (ddd, *J* = 14.4, 5.7 & 3.4 Hz, 1H), 3.33 (ddd, *J* = 14.4, 10.4 & 2.6 Hz, 1H), 2.84 (m, 2H), 2.47–2.34 (m, 4H), 2.26 (m, 2H), 2.09 (m, 1H); ¹³C{¹H}-NMR (CDCl₃, 100 MHz): δ = 142.8 (C), 138.5 (C), 137.9 (C), 130.8 (CH), 129.5 (2× CH), 129.2 (2× CH), 128.4 (2× CH), 127.0 (2× CH), 126.5 (CH), 126.4 (CH), 58.1 (CH), 41.9 (CH₂), 38.8 (CH₂), 30.6 (CH₂), 30.5 (CH₂), 21.4 (CH₃); HRMS (ESI⁺): *m/z* [M + Na]⁺ calcd for C₂₀H₂₃NO₂NaS, 364.1347; found, 364.1349.

2-Isobutyl-1-(methylsulfonyl)-2,3,6,7-tetrahydro-1H-azepine (7e). Following the general procedure (1), to a solution of amine 6b (0.500 g, 1.19 mmol, 1.0 equiv) in 12 mL of dry DCM (0.1 M) at 0 °C were added isovaleraldehyde (0.19 mL, 1.79 mmol, 1.5 equiv) and FeBr₃ (35 mg, 0.12 mmol, 0.1 equiv) to obtain 0.201 g of tetrahydroazepine 7e as a pale yellow oil (0.87 mmol, 73%). *R*_f = 0.49 (*n*-hexane/EtOAc 70:30); ¹H NMR (CDCl₃, 400 MHz): δ = 5.78 (m, 1H), 5.68 (m, 1H), 4.07 (dq, *J* = 6.9 & 4.0 Hz, 1H), 3.66 (dt, *J* = 14.9 & 4.0 Hz, 1H), 3.21 (ddd, *J* = 14.6, 11.5 & 2.9 Hz, 1H), 2.87 (s, 3H), 2.48 (m, 2H), 2.28 (m, 2H), 1.53 (m, 2H), 1.38 (m, 1H), 0.93 (brd, *J* = 1.6 Hz, 3H), 0.91 (brd, *J* = 1.6 Hz, 3H); ¹³C{¹H}-NMR (CDCl₃, 100 MHz): δ = 130.5 (CH), 127.1 (CH), 54.1 (CH), 41.3 (CH₂), 40.6 (CH₃), 40.5 (CH₂), 33.1 (CH₂), 30.7 (CH₂), 24.7 (CH), 22.9 (CH₃), 22.3 (CH₃); HRMS (ESI⁺): *m/z* [M + Na]⁺ calcd for C₁₁H₂₁NO₂NaS, 254.1191; found, 254.1191.

2-Hexyl-1-(methylsulfonyl)-2,3,6,7-tetrahydro-1H-azepine (7f). Following the general procedure (1), to a solution of amine 6b (50 mg, 0.12 mmol, 1.0 equiv) in 1.2 mL of dry DCM (0.1 M) at 0 °C were added heptanal (25 μL, 0.18 mmol, 1.5 equiv) and FeBr₃ (3.6 mg, 0.012 mmol, 0.1 equiv) to obtain 25 mg of tetrahydroazepine 7f as a pale yellow oil (0.097 mmol, 81%). *R*_f = 0.62 (*n*-hexane/EtOAc 70:30); ¹H NMR (CDCl₃, 400 MHz): δ = 5.76 (m, 1H), 5.68 (m,

0.98 (d, $J = 7.2$ Hz, 1.5H), 0.85 (dt, $J = 7.1$ & 2.3 Hz, 4.5H); $^{13}\text{C}\{^1\text{H}\}$ -NMR (CDCl_3 , 100 MHz) (diastereomeric mixture 1:0.5): $\delta = 142.8$ (C), 142.7 (C), 139.0 (CH), 138.0 (C), 137.4 (CH), 136.4 (CH), 129.5 (2 \times CH), 129.4 (2 \times CH), 127.0 (4 \times CH), 125.5 (CH), 124.1 (CH), 56.4 (CH), 54.7 (CH), 48.9 (CH₂), 47.1 (CH₂), 35.9 (CH), 35.2 (CH), 33.1 (CH₂), 32.6 (CH₂), 31.7 (2 \times CH₂), 30.6 (CH₂), 29.8 (CH₂), 29.1 (CH₂), 29.0 (CH₂), 26.4 (CH₂), 26.0 (CH₂), 22.5 (2 \times CH₂), 21.4 (2 \times CH₃), 19.1 (CH₃), 18.9 (CH₃), 14.0 (2 \times CH₃); HRMS (ESI⁺): m/z [$M + \text{Na}$]⁺ calcd for $\text{C}_{20}\text{H}_{31}\text{NO}_2\text{NaS}$: 372.1973, found, 372.1974.

2-(But-3-en-1-yl)-6-methyl-1-tosyl-2,3,6,7-tetrahydro-1H-azepine (10c). Following the general procedure (2), to a solution of amine **9a** (50 mg, 0.098 mmol, 1.0 equiv) in 1.0 mL of dry DCM (0.1 M) were added 4-pentenal (21 μL , 0.15 mmol, 1.5 equiv) and FeCl_3 (21 mg, 0.13 mmol, 1.3 equiv) to afford 17 mg of tetrahydroazepine **10c** as a pale yellow oil (0.052 mmol, 53%). $R_f = 0.70$ (*n*-hexane/EtOAc 80:20); ^1H NMR (CDCl_3 , 500 MHz) (*trans/cis* diastereomeric mixture 1:0.7): $\delta = 7.69$ (dd, $J = 13.0$ & 8.2 Hz, 3H), 7.31–7.26 (m, 3H), 5.70 (m, 1.7H), 5.58–5.53 (dt, $J = 7.7$ & 3.5 Hz, 0.7H), 5.52–5.47 (m, 1H), 5.44–5.36 (m, 1.6H), 5.00–4.90 (m, 3H), 4.06 (m, 1H), 3.94 (m, 0.7H), 3.63 (dd, $J = 14.9$ & 3.5 Hz, 1H), 3.45 (dd, $J = 13.2$ & 2.4 Hz, 0.7H), 3.23 (dd, $J = 13.3$ & 7.9 Hz, 0.7H), 2.84 (dd, $J = 14.9$ & 10.8 Hz, 1H), 2.53 (m, 1H), 2.45–2.37 (m, 5H), 2.24–2.17 (m, 1H), 2.07 (ddd, $J = 16.9$, 8.5 & 4.6 Hz, 1H), 1.95–1.84 (m, 3H), 1.70–1.59 (m, 2H), 1.58–1.54 (m, 2H), 1.27 (m, 3H), 1.05 (d, $J = 7.4$ Hz, 2H), 0.98 (d, $J = 7.4$ Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ -NMR (CDCl_3 , 125 MHz) (diastereomeric mixture 1:0.7): $\delta = 142.9$ (C), 142.8 (C), 138.9 (C), 138.1 (CH), 137.9 (CH), 137.8 (CH), 137.0 (C), 136.6 (CH), 129.6 (2 \times CH), 129.4 (2 \times CH), 127.0 (4 \times CH), 125.2 (CH), 123.8 (CH), 114.9 (CH₂), 114.8 (CH₂), 56.0 (CH), 54.3 (CH), 49.4 (CH₂), 47.1 (CH₂), 35.7 (CH), 35.1 (CH), 32.6 (CH₂), 32.1 (CH₂), 30.6 (CH₂), 30.1 (CH₂), 29.9 (CH₂), 29.1 (CH₂), 21.5 (2 \times CH₃), 19.2 (CH₃), 19.1 (CH₃); HRMS (ESI⁺): m/z [$M + \text{Na}$]⁺ calcd for $\text{C}_{18}\text{H}_{25}\text{NO}_2\text{NaS}$, 342.1504; found, 342.1510.

6-Allyl-2-isobutyl-1-tosyl-2,3,6,7-tetrahydro-1H-azepine (10d). Following the general procedure 3.6, to a solution of amine **9b** (40 mg, 0.074 mmol, 1.0 equiv) in 0.7 mL of dry DCM (0.1 M) were added isovaleraldehyde (12 μL , 0.11 mmol, 1.5 equiv) and FeCl_3 (16 mg, 0.096 mmol, 1.3 equiv) to afford 15 mg of tetrahydroazepine **10d** as a pale yellow oil (0.044 mmol, 60%). $R_f = 0.58$ (*n*-hexane/EtOAc 90:10); ^1H NMR (CDCl_3 , 500 MHz) (*trans/cis* diastereomeric mixture 1:0.6): $\delta = 7.68$ (dd, $J = 11.7$ & 8.2 Hz, 3H), 7.27 (m, 3H), 5.75 (m, 1.6H), 5.62–5.43 (m, 3H), 5.05 (m, 3H), 4.14 (m, 1H), 4.00 (m, 0.6H), 3.66 (dd, $J = 14.8$ & 3.2 Hz, 1H), 3.47 (dd, $J = 13.4$ & 2.7 Hz, 0.6H), 3.33 (dd, $J = 13.5$ & 7.4 Hz, 0.6H), 2.83 (dd, $J = 14.9$ & 10.8 Hz, 1H), 2.48 (m, 1.6H), 2.42 (m, 5H), 2.21 (m, 1H), 2.18–2.10 (m, 1H), 2.10–2.07 (m, 2H), 1.41–1.20 (m, 9H), 0.81 (brt, $J = 6.6$ Hz, 9H); $^{13}\text{C}\{^1\text{H}\}$ -NMR (CDCl_3 , 100 MHz) (diastereomeric mixture 1:0.6): $\delta = 142.8$ (C), 138.9 (C), 137.3 (C), 136.2 (C), 136.0 (CH), 135.7 (2 \times CH), 134.6 (CH), 129.5 (2 \times CH), 129.4 (2 \times CH), 127.2 (2 \times CH), 127.1 (2 \times CH), 126.3 (CH), 125.2 (CH), 117.1 (CH₂), 116.7 (CH₂), 54.2 (CH), 52.5 (CH), 47.3 (CH₂), 45.0 (CH₂), 41.5 (CH₂), 40.0 (CH), 39.9 (CH), 39.6 (CH₂), 37.9 (CH₂), 37.7 (CH₂), 33.0 (CH₂), 29.7 (CH₂), 24.8 (CH), 24.5 (CH), 23.6 (CH₃), 22.8 (2 \times CH₃), 22.3 (CH₃), 21.6 (CH₃), 21.5 (CH₃); HRMS (ESI⁺): m/z [$M + \text{Na}$]⁺ calcd for $\text{C}_{20}\text{H}_{29}\text{NO}_2\text{NaS}$, 370.1817; found, 370.1818.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.2c01396>.

Experimental details and procedures, compound characterization data, and NMR spectra (PDF)

AUTHOR INFORMATION

Corresponding Authors

Victoria Sinka – Instituto de Productos Naturales y Agrobiología (IPNA), CSIC, 38206 La Laguna, Tenerife (Spain); Email: vsinka@ipna.csic.es

Juan I. Padrón – Instituto de Productos Naturales y Agrobiología (IPNA), CSIC, 38206 La Laguna, Tenerife (Spain); orcid.org/0000-0002-0745-2259; Email: jipadron@ipna.csic.es

Author

Israel Fernández – Departamento de Química Orgánica and Centro de Innovación en Química Avanzada (ORFEO-CINQA), Facultad de Ciencias Químicas, Universidad Complutense de Madrid, 28040 Madrid, Spain; orcid.org/0000-0002-0186-9774

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.joc.2c01396>

Author Contributions

J.I.P. conceived and designed this work. V.S. performed experiments and analyzed the data. Both authors wrote the manuscript. I.F. performed DFT calculations and wrote the corresponding part.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Grants (PGC2018-094503-B-C22 to J.I.P. and PID2019-106184GB-I00 and RED2018-102387-T to I.F.) funded by MCIN/AEI/10.13039/501100011033 and, as appropriate, by “ERDF A way of making Europe”, by the “European Union” or by the “European Union NextGeneration EU/PRTR”. V.S. thanks the Spanish MCIU for an FPU fellowship. The manuscript was edited by Guido Jones, also funded by the Cabildo de Tenerife under the same TFinnova support.

REFERENCES

- (1) Koide, K.; Bunnage, M. E.; Gomez Paloma, L.; Kanter, J. R.; Taylor, S. S.; Brunton, L. L.; Nicolaou, K. C. Molecular Design and Biological Activity of Potent and Selective Protein Kinase Inhibitors Related to Balanol. *Chem. Biol.* **1995**, *2*, 601–608.
- (2) Lelais, G.; Epple, R.; Marsilje, T. H.; Long, Y. O.; McNeill, M.; Chen, B.; Lu, W.; Anumolu, J.; Badiger, S.; Bursulaya, B.; DiDonato, M.; Fong, R.; Juarez, J.; Li, J.; Manuia, M.; Mason, D. E.; Gordon, P.; Groessl, T.; Johnson, K.; Jia, Y.; Kasibhatla, S.; Li, C.; Isbell, J.; Spraggon, G.; Bender, S.; Michellys, P.-Y. Discovery of (R,E)-N-(7-Chloro-1-(1-[4-(Dimethylamino)but-2-Enoyl]Azepan-3-Yl)-1H-Benzo[d]Imidazole-2-Yl)-2-Methylisonicotinamide (EGF816), a Novel, Potent, and WT Sparing Covalent Inhibitor of Oncogenic (L858R, Ex19del) and Resistant (T790M) EGFR Mutants. *J. Med. Chem.* **2016**, *59*, 6671–6689.
- (3) Maddili, S. K.; Chowrasia, R.; Kannekanti, V. K.; Gandham, H. 1,3-Di-*n*-Butylimidazolium Tribromide [BBim]Br₃: An Efficient Recyclable Catalyst Mediated Synthesis of *N*-Substituted Azepines and Their Biological Evaluation-Interaction Study with Human Serum Albumin. *J. Photochem. Photobiol., B* **2018**, *178*, 101–107.
- (4) Greger, H. Structural Relationships, Distribution and Biological Activities of Stemona Alkaloids. *Planta Med.* **2006**, *72*, 99–113.
- (5) Greger, H. Structural Classification and Biological Activities of Stemona Alkaloids. *Phytochem. Rev.* **2019**, *18*, 463–493.
- (6) Marco, L.; Carreiras, M. d. C. Galanthamine, a Natural Product for the Treatment of Alzheimer's Disease. *Recent Pat. CNS Drug Discov.* **2006**, *1*, 105–111.

- (7) Barbero, A.; Diez-Varga, A.; Pulido, F. J.; González-Ortega, A. Synthesis of Azepane Derivatives by Silyl-Aza-Prins Cyclization of Allylsilyl Amines: Influence of the Catalyst in the Outcome of the Reaction. *Org. Lett.* **2016**, *18*, 1972–1975.
- (8) Kumar Boominathan, S. S.; Reddy, M. M.; Hou, R.-J.; Chen, H.-F.; Wang, J.-J. A Simple and Efficient Method for Constructing Azepino[4,5-b]Indole Derivatives via Acid Catalysis. *Org. Biomol. Chem.* **2017**, *15*, 1872–1875.
- (9) Hou, L.; Zhou, Z.; Wang, D.; Zhang, Y.; Chen, X.; Zhou, L.; Hong, Y.; Liu, W.; Hou, Y.; Tong, X. DPPF-Catalyzed Atom-Transfer Radical Cyclization via Allylic Radical. *Org. Lett.* **2017**, *19*, 6328–6331.
- (10) Zhu, C.-Z.; Feng, J.-J.; Zhang, J. Rhodium(I)-Catalyzed Intermolecular Aza-[4+3] Cycloaddition of Vinyl Aziridines and Dienes: Atom-Economical Synthesis of Enantiomerically Enriched Functionalized Azepines. *Angew. Chem., Int. Ed.* **2017**, *56*, 1351–1355.
- (11) Zhou, M.-B.; Song, R.-J.; Li, J.-H. Hexafluoroantimonic Acid Catalysis: Formal [3+2+2] Cycloaddition of Aziridines with Two Alkynes. *Angew. Chem., Int. Ed.* **2014**, *53*, 4196–4199.
- (12) Guevara-Pulido, J. O.; Andrés, J. M.; Ávila, D. P.; Pedrosa, R. Enantioselective Synthesis of Seven-Membered Carbo- and Heterocycles by Organocatalyzed Intramolecular Michael Addition. *RSC Adv.* **2016**, *6*, 30166–30169.
- (13) Zhang, Y.-H.; Yuan, Y.-H.; Zhang, S.-Y.; Tu, Y.-Q.; Tian, J.-M. Asymmetric Intramolecular Friedel–Crafts Reaction Catalyzed by a Spiropyrrrolidine Organocatalyst: Enantioselective Construction of Indolizine and Azepine Frameworks. *Tetrahedron Lett.* **2018**, *59*, 4015–4018.
- (14) Chen, C.; Kattanguru, P.; Tomashenko, O. A.; Karpowicz, R.; Siemiaszko, G.; Bhattacharya, A.; Calasans, V.; Six, Y. Synthesis of Functionalised Azepanes and Piperidines from Bicyclic Halogenated Aminocyclopropane Derivatives. *Org. Biomol. Chem.* **2017**, *15*, 5364–5372.
- (15) Zhou, J.; Yeung, Y.-Y. N-Bromosuccinimide-Induced Aminocyclization–Aziridine Ring-Expansion Cascade: An Asymmetric and Highly Stereoselective Approach toward the Synthesis of Azepane. *Org. Lett.* **2014**, *16*, 2134–2137.
- (16) Masson, G.; Gomez Pardo, D.; Cossy, J. Azetidiniums: Ring-Expansion to Pyrrolidines, Piperidines, Azepanes, and Azocanes. *Eur. J. Org. Chem.* **2020**, *2020*, 7103–7118.
- (17) Drouillat, B.; Dorogan, I. V.; Kletsii, M.; Burov, O. N.; Couty, F. Competitive Ring Expansion of Azetidines into Pyrrolidines and/or Azepanes. *J. Org. Chem.* **2016**, *81*, 6677–6685.
- (18) Sivaprakasam, M.; Couty, F.; David, O.; Marrot, J.; Sridhar, R.; Srinivas, B.; Rama Rao, K. A Straightforward Synthesis of Enantiopure Bicyclic Azetidines. *Eur. J. Org. Chem.* **2007**, *2007*, 5734–5739.
- (19) Cutri, S.; Bonin, M.; Micouin, L.; Husson, H.-P.; Chiaroni, A. Practical Asymmetric Synthesis of 1,2-Diamines in the 3-Aminoazepane Series. *J. Org. Chem.* **2003**, *68*, 2645–2651.
- (20) Zakaszewska, A.; Najda-Mocarska, E.; Makowiec, S. Preparation of Bicyclic β -Lactam and Bicyclic 1,3-Oxazinone Scaffolds Using Combined Cycloaddition and Metathesis Processes. *Synth. Commun.* **2018**, *48*, 1793–1804.
- (21) Varela, A.; Garve, L. K. B.; Leonori, D.; Aggarwal, V. K. Stereocontrolled Total Synthesis of (–)-Stemaphylline. *Angew. Chem., Int. Ed.* **2017**, *56*, 2127–2131.
- (22) Wakamatsu, H.; Yoshimura, Y.; Sasaki, M.; Kawahata, K.; Yamaguchi, Y. Synthesis of Various Heterocycles Having a Dienamide Moiety by Ring-Closing Metathesis of Ene-Ynamides. *Synthesis* **2018**, *50*, 3467–3486.
- (23) Majumdar, K. C.; Samanta, S.; Chattopadhyay, B.; Nandi, R. K. Synthesis of Azepine- and Azocine-Annulated Heterocycles by Aromatic Aza-Claisen Rearrangement and Ring-Closing Metathesis. *Synthesis* **2010**, *2010*, 863–869.
- (24) Majumdar, K. C.; Mondal, S.; Ghosh, D. Concise Access to Pyrimidine-Annulated Azepine and Azocine Derivatives by Ruthenium-Catalyzed Ring-Closing Metathesis. *Synthesis* **2010**, *2010*, 1080–1176.
- (25) Qadir, M.; Cobb, J.; Sheldrake, P. W.; Whittall, N.; White, A. J. P.; Hii, K. K.; Horton, P. N.; Hursthouse, M. B. Conformation Analyses, Dynamic Behavior and Amide Bond Distortions of Medium-Sized Heterocycles. 1. Partially and Fully Reduced 1-Benzazepines. *J. Org. Chem.* **2005**, *70*, 1545–1551.
- (26) Carballo, R. M.; Ramírez, M. A.; Rodríguez, M. L.; Martín, V. S.; Padrón, J. I. Iron(III)-Promoted Aza-Prins-Cyclization: Direct Synthesis of Six-Membered Azacycles. *Org. Lett.* **2006**, *8*, 3837–3840.
- (27) Miranda, P. O.; Carballo, R. M.; Martín, V. S.; Padrón, J. I. A New Catalytic Prins Cyclization Leading to Oxa- and Azacycles. *Org. Lett.* **2009**, *11*, 357–360.
- (28) Carballo, R. M.; Valdomir, G.; Purino, M.; Martín, V. S.; Padrón, J. I. Broadening the Synthetic Scope of the Iron(III)-Catalyzed Aza-Prins Cyclization. *Eur. J. Org. Chem.* **2010**, *2010*, 2304–2313.
- (29) Mohr, P.; Imhoff, M.-P.; Reggiani, F. Stereoselective Synthesis of Pyrrolidines from B'-Amino-Allylsilanes. *Tetrahedron Lett.* **2015**, *56*, 2262–2264.
- (30) Dobbs, A. P.; Guesné, S. J. J.; Martinović, S.; Coles, S. J.; Hursthouse, M. B. A Versatile Indium Trichloride Mediated Prins-Type Reaction to Unsaturated Heterocycles. *J. Org. Chem.* **2003**, *68*, 7880–7883.
- (31) Kobayashi, N.; Kaneko, K.; Amemiya, S.; Noguchi, K.; Yamanaka, M.; Saito, A. Alkyne Aza-Prins Cyclization of N-(Hexa-3,5-Diynyl)Tosylamides with Aldehydes Using Triflic Acid and a Binuclear Aluminum Complex. *Chem. Commun.* **2019**, *55*, 8619–8622.
- (32) Sawant, R. T.; Stevens, M. Y.; Odell, L. R. Acetic Acid-Promoted Cascade N-Acyliminium Ion/Aza-Prins Cyclization: Stereoselective Synthesis of Functionalized Fused Tricyclic Piperidines. *Chem. Commun.* **2017**, *53*, 2110–2113.
- (33) Hanessian, S.; Tremblay, M. Tandem Functionalization of Nonactivated Alkenes and Alkynes in Intramolecular N-Acyloxyiminium Ion Carbocyclization. Synthesis of 6-Substituted Hydroindole 2-Carboxylic Acids. *Org. Lett.* **2004**, *6*, 4683–4686.
- (34) Guyot, B.; Pernet, J.; Miginiac, L. Synthèse de N-Alkyl Tétrahydro-2,3,6,7 Azépines et de N-Alkyl Hexahydro-1,2,3,4,7,8 Azocines à Partir d' ω -Monoalkylaminoallyltriméthylsilanes Par Processus d'aminométhylation-Désilylation Intramoléculaire. *J. Organomet. Chem.* **1990**, *386*, 19–28.
- (35) Cruz, D. A.; Sinka, V.; Martín, V. S.; Padrón, J. I. Iron-Catalyzed Prins–Peterson Reaction for the Direct Synthesis of Δ^4 -2,7-Disubstituted Oxepenes. *J. Org. Chem.* **2018**, *83*, 12632–12647.
- (36) Pérez, S. J.; Purino, M. A.; Cruz, D. A.; López-Soria, J. M.; Carballo, R. M.; Ramírez, M. A.; Fernández, I.; Martín, V. S.; Padrón, J. I. Enantiodivergent Synthesis of (+)- and (–)-Pyrrolidine 197B: Synthesis of Trans-2,5-Disubstituted Pyrrolidines by Intramolecular Hydroamination. *Chem.—Eur. J.* **2016**, *22*, 15529–15535.