

Draft Genome Sequence of *Terasakiispira papahanaumokuakeensis* PH27A^T, a Spiral Bacterium from the Northwestern Hawaiian Islands

Xuehua Wan,^a Luis Dasilveira,^b Shaobin Hou,^a Jennifer A. Saito,^a Stuart P. Donachie^{a,b}

Advanced Studies in Genomics, Proteomics and Bioinformatics, University of Hawai'i at Mānoa, Honolulu, Hawai'i, USA^a; Department of Microbiology, University of Hawai'i at Mānoa, Honolulu, Hawai'i, USA^b

The genus *Terasakiispira* hosts only *Terasakiispira papahanaumokuakeensis* PH27A^T, cultivated from an anchialine pond on Pearl and Hermes Atoll, Northwestern Hawaiian Islands. The strain's genome sequence may provide insights into the evolution of free-living *Oceanospirillaceae*.

Received 25 August 2016 Accepted 26 August 2016 Published 20 October 2016

Citation Wan X, Dasilveira L, Hou S, Saito JA, Donachie SP. 2016. Draft genome sequence of *Terasakiispira papahanaumokuakeensis* PH27A^T, a spiral bacterium from the Northwestern Hawaiian Islands. *Genome Announc* 4(5):e01166-16. doi:10.1128/genomeA.01166-16.

Copyright © 2016 Wan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stuart P. Donachie, donachie@hawaii.edu.

The genus *Terasakiispira* was established to accommodate strain PH27A^T, the sole strain of *Terasakiispira papahanaumokuakeensis* (1, 2). PH27A^T cells are spiral and motile by bipolar tufts of flagella. The strain shares 93.3 and 92.1% nucleotide identity in the 16S rRNA gene with its nearest-described neighbors, *Marinospirillum celere* v1c_Sn-red^T and *M. alkaliphilum* Z4^T, respectively. PH27A^T was distinguished from *Marinospirillum* species by its hydrolysis of gelatin and the abundances of ubiquinone Q-9 and cadaverine (1, 2). Cells of PH27A^T accumulate poly-β-hydroxybutyrate (PHB), a biodegradable polymer that has potential applications in medicine and the food industry. We present here the draft genome sequence of *T. papahanaumokuakeensis* PH27A^T.

Genomic DNA was isolated from a 48-h culture of PH27A^T in marine broth 2216E (Difco) using the Wizard genomic DNA purification kit (Promega, USA), with cetyltrimethylammonium bromide (CTAB) and phenol extraction steps. We generated 123.6 Mb of shotgun sequences and 84.3 Mb of 8-kb-span paired-end sequences using the Roche 454 GS FLX+ Titanium pyrosequencing platform, providing ~50× genome coverage. Newbler 2.8 built two scaffolds totaling 4,048,029 bp (scaffold N₅₀, 4,042,489 bp), comprising 22 contigs (contig N₅₀, 404.2 kb); 99.99% of the assembled bases have a base accuracy quality value above the Phred quality score of Q40. The genome's G+C content is 51.3%.

The genome was annotated in the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) and Rapid Annotation using Subsystem Technology (RAST) (3–5). PGAP identified 3,404 protein-coding genes, 62 tRNA coding regions, and 5 rRNA coding regions. RAST identified 3,647 protein-coding open reading frames, 62 tRNA coding regions, and 480 subsystems. RAST identified the flagellar assembly system and chemotaxis system in PH27A^T, consistent with its formal description (1). Thirty-seven metalloproteases were predicted by BLASTP against the MEROPS database 10.0 (6). Homologs of three enzymes (acetyl-coenzyme A [acetyl-CoA] acetyltransferase, acetoacetyl-CoA reductase, and poly-β-

hydroxybutyrate polymerase) in the PHB biosynthetic pathway were predicted in the PH27A^T proteome with BLASTP, as were three enzymes listed in the MetaCyc database ($E < 1e^{-10}$) (7). Further data mining of the PH27A^T genome will shed light on the physiological characteristics and evolutionary features that distinguish PH27A^T and this genus from neighboring *Oceanospirillaceae* and other *Gammaproteobacteria*.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under accession number [MDTQ00000000](https://www.ncbi.nlm.nih.gov/nuclink/MDTQ00000000). The version described in this paper is the first version, MDTQ01000000.

ACKNOWLEDGMENTS

This research was supported by Advanced Studies in Genomics, Proteomics and Bioinformatics at the University of Hawai'i at Mānoa, Hawai'i.

REFERENCES

- Zepeda VK, Busse HJ, Golke J, Saw JH, Alam M, Donachie SP. 2015. *Terasakiispira papahanaumokuakeensis* gen. nov., sp. nov., a gammaproteobacterium from Pearl and Hermes Atoll, Northwestern Hawaiian Islands. *Int J Syst Evol Microbiol* 65:3609–3617. <http://dx.doi.org/10.1099/ijsem.0.000438>.
- Donachie SP, Hou S, Lee K-S, Riley CW, Pikina A, Belisle C, Kempe S, Gregory TS, Bossuyt A, Boerema J, Liu J, Freitas TA, Malahoff A, Alam M. 2004. The Hawaiian archipelago: a microbial diversity hotspot. *Microb Ecol* 48:509–520. <http://dx.doi.org/10.1007/s00248-004-0217-1>.
- Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta) genomic annotation. *Omic* 12:137–141. <http://dx.doi.org/10.1089/omi.2008.0017>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F,

- Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
6. Rawlings ND, Morton FR. 2008. The Merops batch blast: a tool to detect peptidases and their non-peptidase homologues in a genome. *Biochimie* 90:243–259. <http://dx.doi.org/10.1016/j.biochi.2007.09.014>.
7. Caspi R, Billington R, Ferrer L, Foerster H, Fulcher CA, Keseler IM, Kothari A, Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley S, Subhraveti P, Weaver DS, Karp PD. 2016. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res* 44:D471–D480. <http://dx.doi.org/10.1093/nar/gkv1164.26527732>.