


RESEARCH ARTICLE

Open Access

Factors associated with the use of maternal health services by mothers in a post-conflict area of western Côte d'Ivoire in 2016



Mamadou Samba¹, Akissi Régine Attia-Konan^{2*} , Abou Dramane Sangaré¹, Gotré Jules Youan³, Luc Philippe Kouadio² and Ramata Bakayoko-Ly¹

Abstract

Background: In Côte d'Ivoire, maternal health service utilization indicators remain low despite improvements in health coverage and the availability of free health care for pregnant women. The objective of the study was to identify the determinants associated with the use of maternal health services in the department of Bloléquin, in western Côte d'Ivoire.

Methods: We conducted a cross-sectional study with an analytical focus. The study sample size was 400 women. Study participants were selected through a two-stage cluster survey. The data were collected using a standardized questionnaire whose items concerned socio-demographic data, the different uses of maternal health services, namely childbirth assisted by a health professional, use of family planning, prenatal consultation and postnatal consultation. Logistic regression was used to investigate factors associated with the use of maternal health services. The significance of the statistical tests was set at 5%. The odds ratios and 95% confidence intervals were calculated and interpreted.

Results: The results showed that women made less use of family planning services (OR = 0.4), prenatal consultation (OR = 0.2) and assisted childbirth (OR = 0.2) when they provided the funding for care themselves. Women with monthly incomes above \$26.8 used family planning services 4 times more than those with lower incomes. Married women used prenatal consultations 3 times more often than unmarried women ($CI_{95\%} = 1.4-7.3$). Desiring pregnancy increased the use of post-natal consultations by 3 times ($CI_{95\%} = 1.5-6.1$).

Conclusion: Improving the use of maternal health services in western Côte d'Ivoire requires taking into account women's socio-cultural and economic challenges. In initiatives related to the financial empowerment of women, efforts must be made at the level of emotional considerations related to pregnancy.

Keywords: Accessibility, Maternal health services, Determinants, Côte d'Ivoire

* Correspondence: yaoregci@yahoo.fr

²Department of Public Health, Hydrology and Toxicology UFR of Pharmaceutical and Biological Sciences / Unversity Félix Houphouet-Boigny, Abidjan, Côte d'Ivoire

Full list of author information is available at the end of the article



Background

Access to maternal health services is a challenge for health systems in developing countries. In fact, the use of these services makes it possible to reduce obstetric complications, thus maternal mortality. Although the number of maternal deaths decreased from 532,000 to 303,000 between 1990 and 2015 worldwide, maternal mortality remains the highest in sub-Saharan Africa with a rate of 546 deaths per 100,000 live births [1]. To reduce this rate, most states in Africa have introduced an exemption from the cost of care for pregnant women [2]. However, maternal health services are underutilized, especially in low- and middle-income countries.

In Côte d'Ivoire, indicators of maternal health service utilization are low despite the existence of a national policy of "free care" for pregnant women. The coverage rate in ANC4 was 38.23% in 2015, the modern contraceptive prevalence rate was 17%, and the assisted childbirth rate was 55.10% [3]. The maternal mortality ratio was estimated at 645 per 100,000 live births, well above the regional average [1]. The western part of the country, which was particularly affected by the military-political crisis between 2002 and 2011, is experiencing alarming rates. There is significant underutilization of maternal health services with a 25% ANC 4 rate compared to 35% for the national average, with high ANC drop-out rates of up to 67% [3]. To improve access to care, the Government has introduced a policy of free care for pregnant women, taking into account prenatal consultations (ANC), childbirth and caesarean section throughout the country. In addition, there has been an increase in the coverage of health facilities. However, maternal health services remain underutilized and the factors associated with this underutilization by women are not well known, although the literature describes some of them, including economic, social, cultural and organizational parameters related to the health system [4]. A good knowledge of these determinants will certainly make it possible to better address the issue of the use of maternal health services. This is why it seemed wise to understand the social, cultural and economic logic of the use of maternal care of people living in this part of Côte d'Ivoire. The objective of the study was to identify the determinants associated with the use of maternal health services in the department of Bolequin, in western Côte d'Ivoire.

Methods

Study area

The transversal analytical study took place in the department of Bolequin located in the West of Côte d'Ivoire. It has one general hospital, two urban health centers and five rural health centers at the public health level. In 2016, the number of health workers was estimated at 36, including 7 general practitioners, 19 nurses and 10

midwives, for an estimated population of 103,189 inhabitants and 21,324 women of childbearing age [3].

Sampling and setting

The study population consisted of women with a child aged up to 3 years old at the time and living in the Bolequin department at the time of the survey. Were included in the study sample, women meeting these inclusion criteria and giving their informed consent. The sample size was calculated from Schwartz's formula:

$$n = p(1-p) \times \frac{z^2}{d^2}$$

with $p = 38\%$ (corresponding to the national level of ANC 4 rate according to the 2013 RASS); $z = 1.96$; $d = 0.05$; $n = 362$. By increasing this number by 10%, the sample size was 400 women.

For the recruitment of women to be interviewed, two-stage cluster sampling was used. The eight sanitary areas of the department formed the clusters. In the first degree, six areas were drawn at random. Then, in the second degree, a random draw of two secondary roads in each sanitary area was carried out.

Data collection

The data was collected by 4 investigators trained under the supervision of a lead investigator. Once in the chosen locality, the direction to be taken by the investigator was determined by the bottle method. Thus, once in the street drawn at random, the direction was indicated by the direction taken by the tip of the bottle following a throw. Then a random draw for a concession from a random number table was used to find the first household to survey. The survey was carried out from one person to the next until the number of women required per area was reached. The number of women required per area was allocated in proportion to the number of women of childbearing age in each area. The data were collected using a standardized questionnaire developed for this study whose items related socio-demographic, economic, maternal health practices and constraints related to geographical and financial accessibility to health facilities (Supplementary file 2). Mothers with at least one 3-year-old child were interviewed only once about their use of care.

Data analysis

A descriptive analysis of the variables was carried out using frequencies and percentages. A univariate analysis tested the association between the variables of interest and each of the explanatory variables. This step identified the variables to be included in the multivariate analysis if p value is < 0.20 as recommended. Logistic regression was used to investigate factors associated with the use of maternal health services.

There were four variables of interest related to service utilization: childbirth attended by a health professional, family planning (FP), prenatal consultation (PN), and post-natal consultation (PNC). The independent variables included age, marital status, household type, place of residence, gestational age, desire for pregnancy, estimated income, source of funding and distance to a health facility. The final model of the logistic regression was obtained by the top-down step-by-step method. The odds ratios were calculated to measure the strength of association between the variables studied. The significance of the statistical tests was set at 5%. The data collected were analyzed by the STATA 12.1 software. The authorizations of the administrative and health authorities of the department of Bolequin were obtained as well as the verbal informed consent of the women surveyed. The confidentiality of the data was preserved throughout the study.

Ethical considerations

The study protocol was approved by the National Research Ethics Committee of the Ministry of Health and Public Hygiene of Côte d'Ivoire.

Results

Sociodemographic and economic characteristics

Women of reproductive age were predominantly young and lived in couples (85.5%). Nearly half of the women were between the ages of 22 and 30 years old. They were mostly housewives (47%) and had no regular income (36.5%). They lived in cities less than 5 km from a health facility (75%). Women lived 75% in a monogamous home. The spouse provided care financing for 80% of the women (Table 1).

Frequency of use of maternal health services

Family planning services are the least (40%), followed by ANC (58.8%). Assisted childbirth and PNCs were the most commonly used, with respectively 78.5 and 78.25%, or about 8 out of 10 women (Table 2).

Determinants of maternal health service utilization

Table 3 presents the results of the univariate analysis and the variables to be included in the logistic regression model ($p < 0,2$).

Family planning services

Women with monthly incomes of at least \$ 26.8 use more family planning services than women with no income (OR = 4.0 and OR = 3.9). When women self-financed care, they used family planning services less (OR = 0.4) (Table 4).

Prenatal consultations service (ANC services)

Marital status and source of funding significantly influence the use of ANC services. Married women used

Table 1 Socio-demographic and economic characteristics of the sample ($N = 400$)

Variables	Frequency	Percentage (%)
Age (year)		
[16–22[99	24.7
[22–31[178	44.5
[31–49]	123	30.8
Marital status		
Never married	32	8.0
Married	342	85.5
Widowed/Divorced	26	6.5
Professional activities		
Housewife	188	47.0
Farmer	76	19.0
tradeswoman	52	13.0
Craftsman	48	12.0
State agent	24	6.0
Other	16	4.0
Monthly income (US\$)		
No income	146	36.5
< 26,8	78	19.5
26,8 - 53,5	85	21.3
> 53,5	91	22.8
Type of relationship		
Polygamous	101	25.3
Monogamous	299	74.7
Place of residence		
Urban	234	58.5
Rural	166	41.5
Distance to health facility		
≤ 5 km	300	75.0
> 5 km	100	25.0
Funding source		
Spouse	322	80.5
Personal income	45	11.3
Parents	33	8.2
Desired pregnancy		
Yes	354	88.5
No	46	11.5
Gestated		
1	80	20.0
> 1	320	80.0

ANC services more than unmarried women (OR = 3.2). Those who financed their own health expenses used ANC services less (OR = 0.2) (Table 5).

Table 2 Frequency of use of maternal health services (N = 400)

Type of service	Frequency	Percentage (%)
Family planning (FP)	160	40.0
Prenatal consultations (ANC)	235	58.8
Assisted childbirth (PNC)	314	78.5
CPON	313	78.3

Delivery assisted by qualified personnel

Women in couples, as well as widows and divorced women, knew that assisted childbirth was almost 4 times more common than unmarried women (OR = 3.6; OR =

3.9). Women who self-supported their health expenses used less assisted childbirth (OR = 0.2) (Table 6).

Postnatal consultations services (PNC services)

Desiring pregnancy increased the use of CPONs by 3 times (CI_{95%} = 1.5–6.1). The fact of financing her own care leads women to consult less after childbirth (OR = 0.3; IC_{95%} = 0.1–0.5) (Table 7).

Discussion

The use of maternal health services in the Bolequin department is low and far from the national average [3].

Table 3 Results of univariate analysis (N = 400)

Variables	FP		ANC		Professional delivery care		PNC	
	OR	p	OR	p	OR	p	OR	p
Age (year)								
[16–22[1		1		1		1	
[22–30[2.3	0.008	1.4	0.346	2.0	0.053	1.8	0.114
[31–49]	1.6		1.4		1.3		1.6	
Marital status								
Never married	1		1		1		1	
Married	2.6	0.039	1.4	0.346	7.2	0.000	2.0	0.001
Widowed/Divorced	1.6		1.4		2.8		0.5	
Residence area								
Urban	1		1		1		1	
Rural	1.2	0.341	0.9	0.753	1.7	0.096	0.6	0.016
Monthly income								
No income	1		1		1		1	
< 26,8\$	1.6	0.000	1.1	0.486	0.9	0.172	1.6	0.431
26,8\$ – 53,5\$	3.8		1.2		1.3		1.4	
> 53,5\$	3.7		1.5		1.9		0.9	
Type of relationship								
Polygamous	1		1		1		1	
Monogamous	0.7	0.297	0.7	0.192	1.5	0.376	1.1	0.760
Distance to health facility								
≤ 5 km	1		1		1		1	
> 5 km	0.9	0.637	1.6	0.051	1.6	0.113	1.8	0.052
Funding source								
Spouse	1		1		1		1	
Personal income	0.5	0.012	0.4	0.000	0.3	0.000	0.5	0.000
Parents	0.4		0.2		0.2		0.2	
Desired pregnancy								
No	1		1		1		1	
Yes	2.3	0.012	2.7	0.002	2.7	0.004	4.1	0.000
Gestated								
1	1		1		1		1	
> 1	1.9	0.020	1.5	0.130	1.7	0.086	1.8	0.052

Table 4 Determinants of family planning service utilization in Bolequin department: logistic regression (N = 400)

Variables	N	n (%)	OR	CI _{95%}
Age (year)				
[16–22[99	28 (28.3)	–	–
[22–30[178	84 (47.2)	1.8	0.9–3.2
[31–49]	123	48 (39.0)	1.02	0.5–2.1
Marital status				
Never married	32	7 (21.3)	–	–
Married	342	145 (42.4)	1.2	0.4–3.3
Widowed/Divorced	26	8 (30.8)	1.7	0.5–6.4
Monthly income				
No income	146	36 (24.7)	–	–
< 26,8\$	78	27 (34.6)	1.7	0.9–3.2
26,8\$ – 53,5\$	85	47 (55.3)	4.0	2.2–7.2
> 53,5\$	91	50 (55.0)	3.9	2.2–7.1
Funding source				
Spouse	322	140 (435)	–	–
Personal income	45	11 (24.4)	0.4	0.1–0.8
Parents	33	9 (27.3)	0.8	0.3–2.0
Desired pregnancy				
No	46	11 (23.9)	–	–
Yes	354	149 (42.1)	1.7	0.8–3.7
Gestated				
1	80	23 (28.8)	–	–
> 1	320	137 (42.8)	1.2	0.6–2.3

Assisted childbirth and postnatal consultations were the most commonly used services with similar rates. Po's study showed a positive association between the three

Table 5 Determinants of the use of ANC services (N = 400)

Variables	N	n (%)	OR	CI _{95%}
Marital status				
Never married	32	10 (31.3)	–	–
Married	342	214 (62.6)	3,2	1.4–7.3
Widowed/Divorced	26	11 (42.3)	1,5	0.5–4.7
Distance to health facility				
≤ 5 km	300	168 (56)	–	–
> 5 km	100	67 (67)	1.6	0.9–2.6
Funding source				
Spouse	322	189 (80.4)	–	–
Personal income	45	27 (60)	0.2	0.1–0.5
Parents	33	19 (57.6)	0.5	0.2–1.2
Desired pregnancy				
No	46	17 (37.0)	–	–
Yes	354	218 (61.6)	1.9	0.9–3.8

Table 6 Determinants of the use of assisted childbirth (N = 400)

Variables	N	n (%)	OR	CI _{95%}
Age (year)				
[16–22[99	71 (71.7)	–	–
[22–30[178	149 (83.7)	1.4	0.7–2.8
[31–49]	123	94 (76.4)	0.9	0.4–1.7
Marital status				
Never married	32	13 (40.6)	–	–
Married	342	284 (83.0)	3.6	1.5–8.7
Widowed / Divorced	26	17 (65.4)	3.9	1.1–13.3
Monthly income				
No income	146	111 (76.0)	–	–
< 26,8	78	57 (73.1)	0.7	0.3–1.4
26,8 – 53,5	85	68 (80.0)	0.9	0.4–2.0
> 53,5	91	78 (85.7)	1.2	0.4–2.9
Residence area				
Urban	166	137 (82.5)	–	–
Rural	234	177 (65.4)	2.0	1.0–4.0
Funding source				
Spouse	322	271 (84.2)	–	–
Personal income	45	24 (53.3)	0.2	0.1–0.5
Parents	33	19 (57.6)	0.5	0.2–1.2
Desired pregnancy				
Non	46	28 (80.7)	–	–
Oui	354	286 (60.9)	1.5	0.7–3.3
Gestated				
1	80	57 (71.3)	–	–
> 1	320	257 (80.3)	1.1	0.5–2.4

stages of obstetric care, ANC, childbirth and postnatal consultations (PNC) [5]. Hence the importance of setting up a monitoring reference system to ensure the continuity of services and thus increase the use of maternal health services.

Our study confirms the existence of a financial barrier to the use of maternal health services. Although maternal care is free in Côte d'Ivoire, there is still underutilization of maternal health services. A study conducted in Kenya, showed an increase in the use of maternal health services in cases of exemption measures in regions where there were public sector obstetric services available [6]. Hence the importance of increasing geographical accessibility to health centers. However, the financial barrier remains and women sometimes give up maternal care. When pregnant women funded care themselves, they used maternal health services less often. This is why the majority of them have proposed free care to improve its use. Indeed, most of the women in our sample were housewives and did not have a regular income. Several women indexed the quality of

Table 7 Determinants of the use of a post-natal consultation service (N = 400)

Variables	N	n (%)	OR	CI _{95%}
Age (year)				
[16–22[99	70 (70.7)	–	–
[22–30[178	145 (81.5)	1.5	0.8–3.0
[31–49]	123	98 (79.7)	1.4	0.7–3.1
Marital status				
Never married	32	22 (68.8)	–	–
Married	342	278 (81.3)	0.8	0.3–2.2
Widowed/Divorced	26	13 (50.0)	0.4	0.1–1.2
Residence area				
Urban	166	120 (72.3)	–	–
Rural	234	193 (82.5)	0.6	0.4–1.0
Distance to health facility				
≤ 5 km	300	228 (76.0)	–	–
> 5 km	100	58 (85.0)	1.1	0.5–2.2
Funding source				
Spouse	322	266 (82.6)	–	–
Personal income	45	24 (53.3)	0.3	0.1–0.5
Parents	33	23 (69.7)	0.7	0.3–1.7
Desired pregnancy				
No	46	24 (52.2)	–	–
Oui	354	289 (81.6)	3.0	1.5–6.1
Gestated				
1	80	56 (70.0)	–	–
> 1	320	257 (80.3)	1.1	0.6–2.3

care, including reception and transportation costs, to a health facility. The provision of lower quality care has a negative impact on the use of care [7]. Transport-related expenses, which are not often taken into account in health expenditure, sometimes constitute an obstacle to the use of health services [8] even in the case of exemption from fees as observed in the in charge of tuberculosis in Burkina Faso [9]. Therefore, in addition to the obstacle of studies, they mentioned other obstacles to the use of maternal health services. These include mothers' level of education, perception of care, lack of knowledge of obstetric complications and living in rural areas [10, 11]. In addition, situations of political instability and armed conflict in a country compound the lack of access to maternal health services due to the deterioration of the health care system [12]. According to Gaber, the various socio-political crises that Côte d'Ivoire has experienced have had a negative impact on the supply of care [13].

Family Planning (FP) services were the least used among maternal health services despite the steps taken by the government to make FP products accessible with the help of development partners. FP services were more

frequented by women with incomes. This suggests that having financial autonomy leads women to make a decision in favour of their health by avoiding multiple pregnancies. This result seems to contradict the fact that women used their income less to make prenatal consultations (ANC). He suggests that women pay more attention to family planning so they are willing to contribute financially. Nevertheless, it seems advisable to develop income-generating activities for these women in order to increase their financial autonomy. This action requires the popularization of literacy, although little cited by women as a means of improving the use of care (3 out of 400 women), as access to maternal care increases with the level of education [14]. However, women recognize that good awareness will promote better access, which can be increased through literacy. In fact, the use of maternal health services is increased when women have a better understanding of the number of recommended NPCs [15].

Marital status also influenced the use of maternal health services. Married women made greater use of ANC services and more use of assisted childbirth. Married women often benefit from the financial support of their spouses [15]. Single women who are pregnant use care less because they have fewer financial resources and do not always want to become pregnant [15]. They often leave the community to escape the accusatory gaze of society [16]. This phenomenon is magnified in girls although our study did not reveal age as an obstacle to the use of obstetric care. In fact, the latter often have a low rate of access to antenatal care [15, 17].

The level of acceptability of pregnancy impacts the use of PNC services. When women did not want to become pregnant, they used these services less. Beninguisse's 2005 study identified this relationship but with ANC [18]. Thus, the opportunity of pregnancy appears to be a determinant of maternal care use [19]. Therefore, the social dimension of the disease, especially when it is chronic or stigmatized, involves taking into account the psycho-emotional state of the patient to promote this use of care. This is what is observed in the management of HIV / AIDS. Although pregnancy is not a disease, it creates emotions for women that can influence their use of care. Improving well-being during pregnancy is therefore a concern for achieving the Sustainable Development Goals [1].

The parity and geographical access often associated with the use of obstetric care were not highlighted in our study [15]. The majority of women lived within 5 km of a health facility.

Our study presents the limitations of all observational studies regarding possible desirability information bias. Women may overestimate the frequency of use of maternal health services. Thus, we have higher reported

rates than those calculated by the Ministry of Health, which collects data from health facility registers. This uniformly distributed information bias throughout the questionnaire was minimized by the administration of the questions by health workers with a good knowledge of the field.

Conclusion

The underutilization of maternal health services is one of the causes of the high mortality rates observed in developing countries. The results of this study showed a positive association between the use of maternal health services, the marital status of women and the source of funding for care. Improving the use of maternal health services in the department of Blolequin involves taking into account these socio-cultural and economic challenges of women. Investing in the economic empowerment of women is a challenge in this area of Côte d'Ivoire that has been the scene of armed conflict. This autonomy will reduce the woman's dependence and lead her to make informed choices about health. In addition to economic challenges, emotional factors of pregnancy need to be taken into account to improve this access. The social dimension of pregnancy should be developed in maternal health services, where staff often only consider health aspects. Taking into account these different factors will contribute to achieving the objectives for sustainable development in Côte d'Ivoire.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12913-020-4976-2>.

Additional file 1. Questionnaire to pregnant women or women who have been pregnant for 3 years prior to survey

Acknowledgements

Not Applicable.

Authors' contributions

SM oversaw the design of the study, development of the sample and conduct of data collection. AKAR and YGJ conducted quantitative analyses, searched the literature, and contributed in manuscript writing. SM, AKAR, SAD contributed to interpretation of data and led the writing of the manuscript. KLP, B-LR contributed to study design. All authors read and approved the manuscript.

Funding

No funding sources.

Availability of data and materials

The datasets used and analysed during the current study are freely available from the corresponding author upon reasonable request.

Ethics approval and consent to participate

The study protocol was approved by the National Research Ethics Committee of the Ministry of Health and Public Hygiene of Côte d'Ivoire. The informed consent of the respondents was obtained in written prior to their participation in the study. Each study participant was briefed on the study objective and confidentiality was assured for any information provided.

Consent for publication

Not applicable.

Competing interests

The authors do not declare any conflict of interest.

Author details

¹Public Health and Medicinal Plants Laboratory, UFR Odonto-Stomatology University Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire. ²Department of Public Health, Hydrology and Toxicology UFR of Pharmaceutical and Biological Sciences / University Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire. ³NGO Amepouh Bloléquin, Abidjan, Côte d'Ivoire.

Received: 23 July 2019 Accepted: 11 February 2020

Published online: 22 February 2020

References

- World Health Organization. Trends in maternal mortality: 1990 to 2015. Estimated by WHO, UNICEF, UNFPA, World Bank Group, and United Nations population division. Geneva: World Health Organization; 2015.
- Ridde V, Girard J-E. Douze ans après l'initiative de Bamako: Constats et implications politiques pour l'équité d'accès aux services de santé des indigents africains. *Santé Publique*. 2004;16:37–51.
- Ministère de la santé et de la lutte contre le sida Côte d'Ivoire. Rapport Annuel sur la Situation sanitaire (RASS) année 2015. Available on <https://dipe.info/index.php/fr/component/jdownloads/send/6-rapport-annuel-sur-la-situation-sanitaire/50-rapport-annuel-sur-la-situation-sanitaire-rass?Itemid=0>
- Gelato A, Chojenta C, Musa A, Loxton D. Barriers to access and utilization of emergency obstetric care at health facilities in sub-Saharan Africa: a systematic review of literature. *Systematic Rev*. 2018;7:183.
- Owili PO, Muga MA, Chou YJ, Hsu YE, Huang N, Chien LY. Relationship between women's characteristics and continuum of care for maternal health in Kenya: complex survey analysis using structural equation modeling. *Women Health*. 2017;57(8):942–61.
- Njuguna J, Kamau N, Muruka C. Impact of free delivery policy on utilization of maternal health services in county referral hospitals in Kenya. *BMC Health Serv Res*. 2017;17(1):429.
- Doumbouya ML. Pauvreté et accessibilité aux services de santé. Le cas de la Guinée. *Economie et Solidarités*. 2007;38(2):137–53.
- Barasa EW, Marina T, Ravishankar N. Assessing the impoverishing effects, and factors associated with the incidence of catastrophic health care payments in Kenya. *Int J for Equity Health*. 2017;16:31.
- Laokri S, Weil O, Drabo KM, Dembelé SM, Kafando B, Dujardin B. Removal of user fees no guarantee of universal health coverage: observations from Burkina Faso. *Bull World Health Organ*. 2013;91(4):277–82.
- Adeyanju O, Tubeuf S, Ensor T. Socio-economic inequalities in access to maternal and child healthcare in Nigeria: changes over time and decomposition analysis. *Health Policy Plan*. 2017;32(8):1111–9.
- Ponna SN, Upadrashta VP, Babu Geddam JJ, Dudala SR, Sadasivuni R, Bathina H. Regional variation in utilization of antenatal care services in the state of Andhra Pradesh. *J Family Med Prim Care*. 2017;6:231–9.
- Kabakian-Khasholian T, Shayboub R, El-Kak F. Seeking maternal Care at Times of conflict: the case of Lebanon. *Health Care Women Int*. 2013;34(5):352–62.
- Gaber S, Patel P. Tracing health system challenges in post-conflict Côte d'Ivoire from 1893 to 2013. *Glob Public Health*. 2013;8(6):698–712.
- Wilunda C, Quaglio G, Putoto G, Takahashi R, Calia F. Determinants of utilisation of antenatal care and skilled birth attendant at delivery in south west Shoa zone, Ethiopia: a cross sectional study. *Reprod Health*. 2015;12:74.
- Messi E, Yaye W. Contraintes à l'accès aux soins de santé maternelle dans la ville de Maroua. *Int J Eng Sci*. 2017;6(1):13–21.
- Légrand TK, Mbacké SM. Teenage pregnancy and child health in the urban Sahel. *Stud Family Plan*. 1993;24(3):137–49.
- Agunwa CC, Obi EI, Ndu AC, Omotowo IB, Idoko AK, et al. Determinants of patterns of maternal and child health service utilization in a rural community in south eastern Nigeria. *BMC Health Serv Res*. 2017;17:715.
- Beninguisse G. La déperdition des soins prénatals au Tchad. Mémoire en ligne. 2005. Available on <https://www.memoireonline.com/05/09/2090/La-deperdition-des-soins-prenatals-au-Tchad.html>.

19. Beninguisse G, Nikiéma B. La discontinuité des soins obstétricaux en Afrique Centrale et de l'Ouest. Niveaux, Schémas et facteurs associés. Montréal: Projet ARC6AUF; 2005. p. 7.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

