




Proteome of a *Moraxella catarrhalis* Strain under Iron-Restricted Conditions

Luke V. Blakeway,^a Aimee Tan,^{a*} Ian R. Peak,^{a,b}  John M. Attack,^a  Kate L. Seib^a

^aInstitute for Glycomics, Griffith University, Gold Coast, Queensland, Australia

^bSchool of Medical Science, Griffith University, Gold Coast, Queensland, Australia

ABSTRACT *Moraxella catarrhalis* is a leading cause of otitis media and exacerbations of chronic obstructive pulmonary disease; however, its response to iron starvation during infection is not completely understood. Here, we announce a sequential window acquisition of all theoretical fragment ion spectra mass spectrometry (SWATH-MS) data set describing the differential expression of the *M. catarrhalis* CCRI-195ME proteome under iron-restricted versus iron-replete conditions.

Moraxella catarrhalis is a human-restricted bacterial pathogen that causes otitis media in children and exacerbations of chronic obstructive pulmonary disease in adults. Iron is a cofactor required by proteins involved in respiration and DNA replication and thus is essential to life for most bacterial species (1). As iron is sequestered by host tissues, all host-adapted pathogens encounter periods of iron starvation. Previous studies have defined the repertoire of genes required for the growth of *M. catarrhalis* under iron-restricted conditions (2) and described differential expression at the transcriptional level (3). However, global changes in protein expression in response to iron starvation have not been investigated.

M. catarrhalis CCRI-195ME grown in brain heart infusion (BHI) broth with the iron chelator desferal (30 μ M) was compared with bacteria grown in BHI only. Cells were grown at 37°C with six replicates per condition. Cell lysates were equalized to 1 μ g/ μ l protein in 50 mM triethylammonium hydrogen carbonate buffer (TEAB). A total of 100 μ g of each sample was reduced with 5 mM dithiothreitol at 65°C for 30 mins, alkylated with 10 mM iodoacetamide at 25°C for 30 mins, and digested with 1:25 trypsin at 37°C for 16 hours. Samples were acidified with 1 μ l 100% formic acid, pelleted, and resuspended in 100 μ l of loading buffer (2% acetonitrile and 0.1% formic acid). Information-dependent acquisition (IDA) and sequential window acquisition of all theoretical fragment ion spectra mass spectrometry (SWATH-MS) experiments were performed by the Australian Proteome Analysis Facility (Sydney, NSW, Australia) using a 5600 TripleTOF mass spectrometer (AB Sciex) with an Ultra nano-liquid chromatography (nanoLC) system (Eksigent) as previously described (4). For 1D IDA, replicates were pooled and 10 μ l was injected. For 2D IDA, all samples were pooled and fractionated with high-pH reversed-phase high-performance liquid chromatography (HPLC). A total of 13 fractions were collected, dried, and resuspended in 20 μ l of loading buffer, and 10 μ l was injected. Protein identification was performed with ProteinPilot (v4.2) in “thorough mode” using the database *M. catarrhalis* CCRI-195ME coding DNA sequence (CDS) (2,136 proteins) ASM208012v1. SWATH library construction was performed by merging 1D-IDA and 2D-IDA runs using SwathXtend (5) (<https://www.bioconductor.org/packages/release/bioc/html/SwathXtend.html>). For SWATH-MS, each sample was diluted 1:1 in loading buffer, and 10 μ l was injected in random order with one blank run between samples. SWATH data were extracted using PeakView (v2.1) with the following parameters: the 6 most intense fragments of each

Citation Blakeway LV, Tan A, Peak IR, Attack JM, Seib KL. 2020. Proteome of a *Moraxella catarrhalis* strain under iron-restricted conditions. *Microbiol Resour Announc* 9:e00064-20. <https://doi.org/10.1128/MRA.00064-20>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2020 Blakeway et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kate L. Seib, k.seib@griffith.edu.au.

* Present address: Aimee Tan, Department of Microbiology and Immunology, The Peter Doherty Institute for Infection and Immunity, The University of Melbourne, Melbourne, Victoria, Australia.

Received 24 January 2020

Accepted 25 February 2020

Published 19 March 2020

peptide were extracted (75-ppm mass tolerance and 10-min retention time window), shared and modified peptides were excluded, and peptides with a confidence level of $\geq 99\%$ and false discovery rate (FDR) of $\leq 1\%$ were used for quantitation. Peptide peak areas were normalized to the total peak area for each run, and statistical analysis was performed as previously described (4). Proteins are considered differentially abundant if the quantitation ratios were ≥ 1.5 or ≤ 0.67 and respective *P* values were ≤ 0.05 .

SWATH-MS identified 695 proteins, of which 94 were differentially abundant under iron-restricted versus iron-replete conditions. Under iron restriction, 39 proteins were more abundant, including the lactoferrin- and transferrin-binding proteins LbpA, LbpB, and TbpB (6, 7), while 55 proteins were less abundant, including the superoxide dismutase SodA (8) and the nitrate reductase subunit NarJ. This data set will act as the primary proteomic resource for studying the response to iron restriction in *M. catarrhalis* and facilitate study of this important human otopathogen.

Data availability. The SWATH mass spectrometry proteomics data and processed differential abundance analysis have been deposited in the ProteomeXchange Consortium via the PRIDE (9) partner repository with the data set identifier [PXD017206](https://doi.org/10.1093/bioinformatics/btad006).

ACKNOWLEDGMENTS

This work was supported by the Australian National Health and Medical Research Council (NHMRC) (career development fellowship to K.L.S.; project grant 1099279 to K.L.S. and J.M.A.), the Garnett Passe and Rodney Williams Memorial Foundation (research training fellowship to A.T. and grant-in-aid supplementation to K.L.S. and J.M.A.), and an Australian Government research training program scholarship to L.V.B.

REFERENCES

- Andrews S, Norton I, Salunkhe AS, Goodluck H, Aly WSM, Mourad-Agha H, Cornelis P. 2013. Control of iron metabolism in bacteria. *Met Ions Life Sci* 12:203–239. https://doi.org/10.1007/978-94-007-5561-1_7.
- de Vries SPW, Burghout P, Langereis JD, Zomer A, Hermans PWM, Bootsma HJ. 2013. Genetic requirements for *Moraxella catarrhalis* growth under iron-limiting conditions. *Mol Microbiol* 87:14–29. <https://doi.org/10.1111/mmi.12081>.
- Wang W, Reitzer L, Rasko DA, Pearson MM, Blick RJ, Laurence C, Hansen EJ. 2007. Metabolic analysis of *Moraxella catarrhalis* and the effect of selected in vitro growth conditions on global gene expression. *Infect Immun* 75:4959–4971. <https://doi.org/10.1128/IAI.00073-07>.
- Bjelosevic S, Pascovici D, Ping H, Karlaftis V, Zaw T, Song X, Molloy MP, Monagle P, Ignjatovic V. 2017. Quantitative age-specific variability of plasma proteins in healthy neonates, children and adults. *Mol Cell Proteomics* 16:924–935. <https://doi.org/10.1074/mcp.M116.066720>.
- Wu JX, Song X, Pascovici D, Zaw T, Care N, Krisp C, Molloy MP. 2016. SWATH mass spectrometry performance using extended peptide MS/MS assay libraries. *Mol Cell Proteomics* 15:2501–2514. <https://doi.org/10.1074/mcp.M115.055558>.
- Bonnah RA, Wong H, Loosmore SM, Schryvers AB. 1999. Characterization of *Moraxella (Branhamella) catarrhalis* *lbpB*, *lbpA*, and lactoferrin receptor *orf3* isogenic mutants. *Infect Immun* 67:1517–1520. <https://doi.org/10.1128/IAI.67.3.1517-1520.1999>.
- Luke NR, Campagnari AA. 1999. Construction and characterization of *Moraxella catarrhalis* mutants defective in expression of transferrin receptors. *Infect Immun* 67:5815–5819. <https://doi.org/10.1128/IAI.67.11.5815-5819.1999>.
- Luke NR, Karalus RJ, Campagnari AA. 2002. Inactivation of the *Moraxella catarrhalis* superoxide dismutase SodA induces constitutive expression of iron-repressible outer membrane proteins. *Infect Immun* 70:1889–1895. <https://doi.org/10.1128/iai.70.4.1889-1895.2002>.
- Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, Inuganti A, Griss J, Mayer G, Eisenacher M, Pérez E, Uszkoreit J, Pfeuffer J, Sachsenberg T, Yilmaz S, Tiwary S, Cox J, Audain E, Walzer M, Jarnuczak AF, Ternent T, Brazma A, Vizcaino JA. 2019. The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* 47:D442–D450. <https://doi.org/10.1093/nar/gky1106>.