

Continuous Flow Sodiation of Substituted Acrylonitriles, Alkenyl Sulfides and Acrylates

Johannes H. Harenberg⁺, Niels Weidmann⁺, Konstantin Karaghiosoff, and Paul Knochel^{*}


Abstract: The sodiation of substituted acrylonitriles and alkenyl sulfides in a continuous flow set-up using NaDA (sodium diisopropylamide) in EtNMe₂ or NaTMP (sodium 2,2,6,6-tetramethylpiperidide)·TMEDA in *n*-hexane provides sodiated acrylonitriles and alkenyl sulfides, which are subsequently trapped in batch with various electrophiles such as aldehydes, ketones, disulfides and allylic bromides affording functionalized acrylonitriles and alkenyl sulfides. This flow-procedure was successfully extended to other acrylates by using Barbier-type conditions.

The metalation of unsaturated nitriles and sulfides is an important synthetic procedure.^[1] After quenching with various electrophiles, highly functionalized unsaturated products are obtained, which may be useful building blocks for biologically active heterocycles and natural products.^[2] The batch-metalation of alkenyl nitriles or sulfides with lithium bases is often complicated due to competitive allylic lithiations.^[3] The use of stronger, more polar bases like sodium or potassium amides may avoid such limitations. However, the sodiation of such unsaturated compounds is much less explored.^[4] Moreover, the use of sodium organometallics is of high interest due to the low price, high abundance and low toxicity of sodium salts.^[5] Recently, arylsodium compounds have been prepared by Collum using NaDA (sodium diisopropylamide) as deprotonating agent^[4e,6] and by Asako and Takai, who have investigated the utility of arylsodiums in catalytic cross-couplings.^[7] Yoshida, Ley, Organ and others have demonstrated a high functional group tolerance performing challenging metalations in a continuous flow set-up.^[8] Based on these studies, we have extended the Collum procedure to the preparation of sodiated aryl and heteroaryl derivatives which are difficult to generate otherwise and decompose upon batch-sodiation.^[9] KDA·TMEDA (potassium diisopropylamide·*N,N,N',N'*-tetramethylethylenediamine) in *n*-hexane was used in continuous flow for similar

metalations.^[10] Herein, we wish to report that NaDA and NaTMP (TMPH = 2,2,6,6-tetramethylpiperidine) were efficient bases for the regioselective flow-metalation of various substituted acrylonitriles and alkenyl sulfides.^[11] In first experiments, we have optimized the sodiation of cinnamionitrile (**1a**) and have found that metalation with NaDA (0.24 M in DMEA (dimethylethylamine), 1.2 equiv) at -78°C using a combined flow-rate of 10 mL min^{-1} and a 0.02 mL reactor proceeded best with a residence time of 0.12 s affording organosodium **2a**. Subsequent trapping with electrophiles of type **3** such as aldehydes, ketones, disulfides and allylic bromides afforded 2-substituted cinnamionitriles of type **4** with usually high *E/Z* ratios (Table 1, entries 1–10). Thus, for a quenching with aromatic aldehydes, we obtained the *Z*-product of type **4** as major product, whereas for more sterically hindered ketones the *E*-product was formed.

The diastereoselectivity of products of type **4** obtained after the addition to a carbonyl electrophile was tentatively explained by assuming that the sodiated nitrile **2a** reacted fast with an aldehyde (RCHO) according to pathway A leading to the allylic alcohol **Z-4**. In contrast, by using ketones, an equilibration to the cummulene form **2a'** may occur and the cyclic transition state **A** would be disfavoured due to steric hindrance. *E/Z* isomerization of the cummulene structure **2a'** occurred affording the *E-4* product via transition state **B** (Scheme 1).

We have then extended this flow procedure to various functionalized arylacrylonitriles of type **5**. Electron-rich cinnamionitrile derivatives (**5a–5d**) were selectively metalated in 2-position using NaDA in a continuous flow set-up within 0.12 s at -78°C . The resulting organosodiums (**6a–d**) were trapped in batch with various carbonyl electrophiles, such as *m*-anisaldehyde (**3k**), cyclohexanecarboxaldehyde (**3l**) or cyclohexanone (**3m**), and with 3-bromocyclohexene



Scheme 1. Tentative mechanism for the stereoselective addition of sodiated phenylacrylonitrile **2a** or **2a'** to aldehydes or ketones.

[*] J. H. Harenberg,^[+] N. Weidmann,^[+] Prof. Dr. K. Karaghiosoff,

Prof. Dr. P. Knochel

Department Chemie; Ludwig-Maximilians-Universität München

Butenandtstrasse 5–13, Haus F, 81377 München (Germany)

E-mail: paul.knochel@cup.uni-muenchen.de

[+] These authors contributed equally.

Supporting information and the ORCID identification number(s) for

the author(s) of this article can be found under:

<https://doi.org/10.1002/anie.202012085>.

© 2020 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Table 1: Sodiation of cinnamitrile (**1a**) using a microflow reactor and subsequent batch quench of the intermediate sodium organometallic **2a** with various electrophiles of type **3** leading to functionalized cinnamitriles of type **4**.

flow		batch			
#	electrophile	product ^[a]	#	electrophile	product ^[a]
1		 4aa: 95%, Z/E > 99/1 ^[b]	6		 4af: 93%, Z/E = 54/46
2		 4ab: 92%, Z/E > 99/1 ^[b]	7		 4ag: 82%, E/Z > 99/1 ^[c]
3		 4ac: 74%, Z/E = 89/11 ^[b]	8		 4ah: 82%, E/Z > 99/1 d.r. > 99/1 ^[c]
4		 4ad: 93%, Z/E > 99/1 ^[c]	9		 4ai: 78%, E/Z > 99/1 ^[b]
5		 4ae: 93%, E/Z = 9/1 ^[b]	10		 4aj: 87%, E/Z > 99/1 ^[b]

[a] Yield of analytically pure product. [b] The *E*- or *Z*-diastereoselectivity was assigned in analogy to related products, for which X-ray data were obtained. [c] The diastereoselectivity was determined by crystal structure analyses, see Supporting Information. [d] 10 mol% CuCN·2LiCl.

(**3e**) using 10 mol% CuCN·2LiCl as catalyst, affording the desired alcohols (**7ak**, **7bk**, **7cl** and **7dm**) and an allylated cinnamitrile derivative (**7de**) in 57–97% yield with diastereomeric ratios up to > 99/1 (Table 2, entries 1–5). Similarly, regioselective sodiation of electron-deficient 3-(4-(trifluoromethyl)phenyl)acrylonitrile (**5e**) followed by copper-catalyzed allylation with 3-bromocyclohexene (**3e**) led to the functionalized phenylacrylonitrile (**7ee**) in 66% yield with an *E/Z* ratio > 99/1. Furthermore, an extension to methoxy- and ethoxyacrylonitriles **5f** and **5g** was possible resulting in secondary alcohols (**7fn**, **7fd**, **7gc** and **7gl**) after batch-quench with aromatic aldehydes (**3c**, **3d** and **3n**), and aliphatic aldehyde (**3l**) in 91–98% and *Z/E* ratios > 99/1 (entries 7–10). An alkenyl sulfide such as phenyl-


(styryl)sulfane (**5h**) provided the sodium derivative (**6h**) upon metalation with NaDA, which after trapping with sterically demanding ketones such as adamantanone (**3o**) and benzophenone (**3g**) gave tertiary alcohols (**7ho** and **7hg**) in 85–95% yield and comparable *E/Z* ratios to the starting material **5h** (entries 11–12).

Extension to alkyl-substituted acrylonitriles such as geranyl nitrile (**8a**, *E/Z* = 50/50) and the related nitrile **8b** (*E/Z* = 65/35) was possible under the standard sodiation conditions providing after electrophilic quench the desired functionalized nitriles (**10ap**, **10al**, **10aq**, **10br**) in 60–98% yield as *E/Z* mixtures (Table 3, entries 1–4). Interestingly, starting from the diastereomerically pure acrylonitrile **8c** (*E/Z* > 99/1) the desired product **10cq** was obtained in 67% yield (*Z/E* = 58/42) after quench with α -tetralone (**3q**) (entry 5) showing the prevalence of the cumulene structure of the sodiated nitriles (see **2a'** in Table 1). However, the methoxy-substituted acrylonitrile **8d** (*E/Z* = 80/20) afforded after continuous flow sodiation and quenching with *o*-anisaldehyde (**3k**) the allylic alcohol **10dk** as single diastereoisomer in 58% yield (*Z/E* > 99/1) showing the importance of the methoxy group for controlling the stereochemistry of the intermediate sodiated nitrile (entry 6). Also, the dienyl nitrile **8e** was sodiated in flow and trapping with an allylic bromide (**3e**) or an aldehyde (**3k**) furnished the functionalized dienyl nitriles (**10ee** and **10ek**) in 74–82% yield (entries 7–8).

Recently, Takai and Asako published a straightforward synthesis of lithium-free sodium 2,2,6,6-tetramethylpiperidide (NaTMP) in *n*-hexane by using sodium dispersion, TMPH, TMEDA and isoprene.^[12] This method would allow us to avoid the use of the amine DMEA as solvent and therefore making our method more practical. Using the Takai procedure, we have prepared hexane-soluble NaTMP·TMEDA^[13] and have performed an efficient continuous flow sodiation of cinnamitrile (**1a**) selectively in 2-position within 0.12 s at -78°C . A subsequent batch trapping of **2a'** with various ketones of type **3** afforded the desired tertiary alcohols of type **4** in 58–83% yield as single regioisomers (Scheme 2). Similarly, ethoxyacrylonitrile **5g** gave, after batch quench with *o*-anisaldehyde (**3k**) and benzophenone (**3g**), the allylic alcohols (**7gk** and **7gg**) in 65–78% yield (*Z/E* > 99/1). Further, geranyl nitrile (**8a**) provided the organosodium **9a** upon metalation with NaTMP·TMEDA, which after a copper-catalyzed allylation using 3-bromocyclohexene (**3e**) led to the desired product (**10ae**) in 54% yield with a *E/Z* ratio of 52/48.

However, the sodiation of other acrylates still remained challenging. Applying our standard sodiation method to ethyl cinnamate (**11a**) afforded solely the condensation product **13a** showing that the sodiation of **11a** was possible, but difficult to control. Thus, the intermediate organosodium **12a** reacted instantaneously with another molecule of **11a** before the desired electrophile quench proceeded (Scheme 3a). To prevent this self-condensation reaction, sterically hindered *tert*-butyl cinnamate (**11b**) was used affording organosodium **12b** after continuous flow sodiation. A copper-catalyzed batch allylation with 3-bromocyclohexene (**3e**) gave the desired product **13be** in 61% yield with an *E/Z* ratio > 99/1 (Scheme 3b). To overcome the need of sterically hindered esters, we envisioned a Barbier-type in situ trapping^[14] of the

Table 2: Sodiation of substituted acrylonitriles and alkenyl sulfides of type **5** using a microflow reactor and subsequent batch quench of the intermediate sodium organometallics of type **6** with various electrophiles of type **3** leading to functionalized phenylacrylonitriles and alkenyl sulfides of type **7**.




#	SM	product ^[a]	#	SM	product ^[a]
1	5a $E/Z = 76/24$	7ak : 97% ^[b] $Z/E = 89/11$	7	5f $E/Z = 83/17$	7fn : 93% ^[c] $Z/E > 99/1$
2	5b $E/Z = 79/21$	7bk : 84% ^[c] $Z/E = 90/10$	8	5f $E/Z = 83/17$	7fd : 98% ^[c] $Z/E > 99/1$
3	5c $E/Z = 83/17$	7cl : 74% ^[b] $Z/E > 99/1$	9	5g $E/Z = 68/32$	7gc : 95% ^[c] $Z/E > 99/1$
4	5d $E/Z = 79/21$	7dm : 67% ^[c] $E/Z > 99/1$	10	5g $E/Z = 68/32$	7gl : 91% ^[b] $Z/E > 99/1$
5	5d ^[d] $E/Z = 79/21$	7de : 57% ^[b] $E/Z > 99/1$	11	5h $E/Z = 71/29$	7ho : 95% ^[c] $E/Z = 77/23$

Table 2: (Continued)

#	SM	product ^[a]	#	SM	product ^[a]
6	5e ^[d] $E/Z = 78/22$	7ee : 66% ^[c] $E/Z > 99/1$	12	5h $E/Z = 71/29$	7hg : 85% ^[b] $E/Z = 68/32$

[a] Yield of analytically pure product. [b] The *E*- or *Z*-diastereoselectivity was assigned in analogy to related products, for which X-ray data were obtained. [c] The diastereoselectivity was determined by crystal structure analyses, see Supporting Information. [d] 10 mol% CuCN·2LiCl.



Scheme 2. General set-up for the sodiation of functionalized acrylonitriles with NaTMP-TMEDA in a microflow reactor and subsequent batch quench of the intermediate sodium organometallics with various electrophiles leading to functionalized acrylonitriles. [a] The diastereoselectivity was determined by crystal structure analyses, see Supporting Information. [b] The *E*- or *Z*-diastereoselectivity was assigned in analogy to related products, for which X-ray data were obtained. [c] 10 mol% CuCN·2LiCl.

highly reactive organosodiums of type **12**. Interestingly, ethyl cinnamate (**11a**), which underwent self-condensation side reactions applying our standard flow conditions (Scheme 3a), was sodiated at -78 °C under Barbier-conditions and afforded organosodium **12a**, which was instantaneously trapped by adamantanone (**3o**), outcompeting self-condensation and resulting in the tertiary alcohol **13ao** in 66% yield ($E/Z > 99/1$). Similarly, methyl-3-methoxyacrylate (**11c**) was sodiated in 3-position in the presence of adamantanone (**3o**) using NaDA (1.2 equiv) affording the spiroactone **13co** in 58% yield (Scheme 3c).

In summary, we have reported the sodiation of substituted acrylonitriles and alkenyl sulfides in a continuous flow set-up using NaDA (sodium diisopropylamide) in EtNMe₂ (DMEA) and NaTMP (sodium 2,2,6,6-tetramethylpiperidide)·TMEDA


Table 3: Sodiation of alkyl- and alkenyl-substituted acrylonitriles of type **8** using a microflow reactor and subsequent batch quench of the intermediate sodium organometallics of type **9** with various electrophiles of type **3** leading to functionalized alkyl- and alkenyl-substituted acrylonitriles of type **10**.

#	substrate	electrophile	product ^[a]
1		I ₂	10ap : 75%, Z/E = 68/32
2			10al : 60%, Z/E = 64/36
3			10aq : 98%, Z/E = 53/47
4			10br : 85%, Z/E = 55/45
5			10cq : 67%, Z/E = 58/42
6			10dk : 58%, Z/E > 99/1
7			10ee : 74%, 2E/2Z = 77/23
8			10ek : 82%, 2Z/2E = 76/24


[a] Yield of analytically pure product. [b] 10 mol% CuCN·2LiCl.

in *n*-hexane. The resulting sodiated acrylonitriles and alkenyl sulfides were subsequently trapped in batch with various electrophiles such as aldehydes, ketones, disulfides and allylic bromides affording functionalized acrylonitriles and alkenyl sulfides. This flow-procedure was successfully extended to other acrylates by using Barbier-type conditions.


a) Sodiation of ethyl cinnamate **11a** leading to self-condensation side product:



b) Sodiation of sterically hindered acrylates and subsequent batch trapping:



c) Sodiation of acrylates using Barbier-type trapping conditions:



Scheme 3. Sodiation of substituted acrylates of type **11** using a microflow reactor under standard-flow conditions and Barbier conditions. *In situ* quench of the intermediate sodium organometallics of type **12** with adamantanone (**3o**) afforded functionalized acrylates of type **13**. [a] 0.20 M in THF, 1.0 equiv [b] 0.30 M in THF, 1.5 equiv.

Acknowledgements

N. Weidmann thanks the German Academic Scholarship Foundation for a fellowship. We thank the DFG and LMU for financial support. We further thank BASF (Ludwigshafen) and Albemarle (Frankfurt) for the generous gift of chemicals and Uniqsis for technical support. Open access funding enabled and organized by Projekt DEAL.

Conflict of interest

The authors declare no conflict of interest.

Keywords: acrylonitriles · alkenyl sulfides · Barbier-type reactions · flow chemistry · sodiation

- [1] a) F. F. Fleming, Q. Wang, Z. Zhang, O. W. Steward, *J. Org. Chem.* **2002**, *67*, 5953; b) J. Doroszuk, M. Musiejuk, L. Poniakowski, D. Witt, *Eur. J. Org. Chem.* **2018**, 6333; c) O. De Lucchi, L. Pasquato, *Tetrahedron* **1988**, *44*, 6755; d) B. M. Trost, A. C. Lavoie, *J. Am. Chem. Soc.* **1983**, *105*, 5075; e) B. Bartels, R. Hunter, C. D. Simon, G. D. Tomlinson, *Tetrahedron Lett.* **1987**, *28*, 2985; f) A. B. Flynn, W. W. Ogilvie, *Chem. Rev.* **2007**, *107*, 4698; g) F. F. Fleming, Q. Wang, *Chem. Rev.* **2003**, *103*, 2035;

- h) G. Dagousset, C. François, T. León, R. Blanc, E. Sansiaume-Dagousset, P. Knochel, *Synthesis* **2014**, 46, 3133.
- [2] a) S. Sengupta, V. Snieckus, *J. Org. Chem.* **1990**, 55, 5680; b) M. A. Reed, M. T. Chang, V. Snieckus, *Org. Lett.* **2004**, 6, 2297; c) E. Block, S. Ahmad, J. L. Catalfamo, M. K. Jain, R. Apitz-Castro, *J. Am. Chem. Soc.* **1986**, 108, 7045; d) G. Brooks, K. Coleman, J. S. Davies, P. A. Hunter, *J. Antibiot.* **1988**, 41, 892; e) T. H. Morris, E. H. Smith, R. Walsh, *J. Chem. Soc. Chem. Commun.* **1987**, 964.
- [3] a) F. F. Fleming, S. Gudipati, J. A. Aitken, *J. Org. Chem.* **2007**, 72, 6961; b) F. F. Fleming, V. Gudipati, O. W. Steward, *Tetrahedron* **2003**, 59, 5585; c) B. A. Feit, U. Melamed, R. R. Schmidt, H. Speer, *Tetrahedron Lett.* **1981**, 37, 2143; d) D. C. Harrowven, H. S. Poon, *Tetrahedron Lett.* **1994**, 35, 9101; e) D. C. Harrowven, H. S. Poon, *Tetrahedron* **1996**, 52, 1389; f) R. R. Schmidt, J. Talbiersky, P. Russegger, *Tetrahedron Lett.* **1979**, 20, 4273; g) R. R. Schmidt, R. Hirsenkorn, *Tetrahedron* **1983**, 39, 2043; h) R. Knorr, E. Lattek, *Chem. Ber.* **1981**, 114, 2116; i) M. A. Ganiek, M. R. Becker, M. Ketels, P. Knochel, *Org. Lett.* **2016**, 18, 828.
- [4] a) A. A. Morton, F. D. Marsh, R. D. Coombs, A. L. Lyons, S. E. Penner, H. E. Ramsden, V. B. Baker, E. L. Little, R. L. Letsinger, *J. Am. Chem. Soc.* **1950**, 72, 3785; b) A. A. Morton, E. J. Lanpher, *J. Org. Chem.* **1955**, 20, 839; c) R. A. Benkeser, D. J. Foster, D. M. Sauve, J. F. Nobis, *Chem. Rev.* **1957**, 57, 867; d) R. A. Woltornist, Y. Ma, R. F. Algera, Y. Zhou, Z. Zhang, D. B. Collum, *Synthesis* **2020**, 52, 1478; e) R. F. Algera, Y. Ma, D. B. Collum, *J. Am. Chem. Soc.* **2017**, 139, 11544; f) Y. Huang, G. H. Chan, S. Chiba, *Angew. Chem. Int. Ed.* **2017**, 56, 6544; *Angew. Chem.* **2017**, 129, 6644.
- [5] a) D. Seyferth, *Organometallics* **2006**, 25, 2; b) D. Seyferth, *Organometallics* **2009**, 28, 2; c) N. Wiberg, *Lehrbuch der Anorganischen Chemie*, 102. ed., De Gruyter Verlag, Berlin, **2007**, pp. 1259–1299.
- [6] a) R. F. Algera, Y. Ma, D. B. Collum, *J. Am. Chem. Soc.* **2017**, 139, 15197; b) R. F. Algera, Y. Ma, D. B. Collum, *J. Am. Chem. Soc.* **2017**, 139, 7921; c) R. F. Algera, Y. Ma, D. B. Collum, *J. Am. Chem. Soc.* **2017**, 139, 11544; d) Y. Ma, R. F. Algera, D. B. Collum, *J. Org. Chem.* **2016**, 81, 11312.
- [7] S. Asako, H. Nakajima, K. Takai, *Nat. Catal.* **2019**, 2, 297.
- [8] a) M. B. Plutschack, B. Pieber, K. Gilmore, P. H. Seeberger, *Chem. Rev.* **2017**, 117, 11796; b) J. Britton, T. F. Jamison, *Nat. Protoc.* **2017**, 12, 2423; c) T. Brodmann, P. Koos, A. Metzger, P. Knochel, S. V. Ley, *Org. Process Res. Dev.* **2012**, 16, 1102; d) M. Colella, A. Nagako, R. Luisi, *Chem. Eur. J.* **2020**, 26, 19; e) B. Gutmann, C. O. Kappe, *J. Flow Chem.* **2017**, 7, 65; f) M. Teci, M. Tilley, M. A. McGuire, M. G. Organ, *Org. Process Res. Dev.* **2016**, 20, 1967; g) D. A. Thaisrivongs, J. R. Naber, N. J. Rogus, G. Spencer, *Org. Process Res. Dev.* **2018**, 22, 403; h) F. Ullah, T. Samarakoon, A. Rolfe, R. D. Kurtz, P. R. Hanson, M. G. Organ, *Chem. Eur. J.* **2010**, 16, 10959; i) J. Y. F. Wong, J. M. Tobin, F. Vilela, G. Barker, *Chem. Eur. J.* **2019**, 25, 12439; j) H. Kim, H.-J. Lee, D.-P. Kim, *Angew. Chem. Int. Ed.* **2015**, 54, 1877; *Angew. Chem.* **2015**, 127, 1897; k) H. Kim, A. Nagaki, J.-i. Yoshida, *Nat. Commun.* **2011**, 2, 264.
- [9] N. Weidmann, M. Ketels, P. Knochel, *Angew. Chem. Int. Ed.* **2018**, 57, 10748; *Angew. Chem.* **2018**, 130, 10908.
- [10] J. H. Harenberg, N. Weidmann, P. Knochel, *Angew. Chem. Int. Ed.* **2020**, 59, 12321; *Angew. Chem.* **2020**, 132, 12419.
- [11] Commercially available equipment from *Uniqsis* was used. For a detailed optimization, see Supporting Information.
- [12] a) S. Asako, M. Koderia, H. Nakajima, K. Takai, *Adv. Synth. Catal.* **2019**, 361, 3120; b) D. R. Armstrong, A. R. Kennedy, R. E. Mulvey, S. D. Robertson, *Chem. Eur. J.* **2011**, 17, 8820; c) R. E. Mulvey, S. D. Robertson, *Angew. Chem. Int. Ed.* **2013**, 52, 11470; *Angew. Chem.* **2013**, 125, 11682; d) R. McLellan, M. Uzelac, L. J. Bole, J. M. Gil-Negrete, D. R. Armstrong, A. R. Kennedy, R. E. Mulvey, E. Hevia, *Synthesis* **2019**, 51, 1207; e) B. Gehrus, P. H. Hitchcock, A. R. Kennedy, M. F. Lappert, R. E. Mulvey, P. J. A. Rodger, *J. Organomet. Chem.* **1999**, 587, 88.
- [13] For the synthesis of NaTMP·TMEDA, see Supporting Information.
- [14] a) M. G. Ganiek, M. V. Ivanova, B. Martin, P. Knochel, *Angew. Chem. Int. Ed.* **2018**, 57, 17249; *Angew. Chem.* **2018**, 130, 17496; b) M. A. Ganiek, M. R. Becker, G. Berionni, H. Zipse, P. Knochel, *Chem. Eur. J.* **2017**, 23, 10280; c) N. Weidmann, J. H. Harenberg, P. Knochel, *Org. Lett.* **2020**, 22, 5895.

Manuscript received: September 4, 2020

Accepted manuscript online: October 7, 2020

Version of record online: November 3, 2020