

Predictors of disability in patients with chronic low back pain

Elena Sirbu¹, Roxana Ramona Onofrei², Simona Szasz³, Monica Susan⁴

¹Department of Physical Therapy and Special Motility, West University of Timișoara, Timișoara, Romania

²Department of Rehabilitation, Physical Medicine and Rheumatology, “Victor Babes” University of Medicine and Pharmacy Timisoara, Romania

³Department of Rheumatology, The University of Medicine and Pharmacy Tg Mures, Romania

⁴Department of Internal Medicine I, “Victor Babes” University of Medicine and Pharmacy Timisoara, Romania

Submitted: 10 October 2019; **Accepted:** 12 December 2019

Online publication: 8 July 2020

Arch Med Sci 2023; 19 (1): 94–100

DOI: <https://doi.org/10.5114/aoms.2020.97057>

Copyright © 2020 Termedia & Banach

Corresponding author:

Roxana Ramona Onofrei
Department of Rehabilitation,
Physical Medicine and
Rheumatology
“Victor Babes” University
of Medicine and Pharmacy
Timisoara
2 Eftimie Murgu Square,
300041 Timisoara, Romania
E-mail:
onofrei.roxana@umft.ro

Abstract

Introduction: Chronic low back pain (CLBP) is a common and disabling health problem. In this study, we aimed to assess the relationship between pain intensity, the components of catastrophizing, depression and disability in patients with chronic low back pain.

Material and methods: Seventy-six patients diagnosed with CLBP (age range 25–77 years; 73.7% female) participated in the study. Participants' socio-demographic data were collected: age, gender, height, weight, and work status (employed or retired). All participants were asked to complete the Pain Catastrophizing Scale (PCS), the visual analogue scale (VAS), the Oswestry Disability Questionnaire (ODQ), and the Beck Depression Inventory (BDI).

Results: The mean group scores revealed moderate CLBP complaints (VAS – 4 [3–6]), mild depression (BDI – 10 [5–16]), a moderate level of catastrophizing (PCS total score 20.5 [10–34]) and moderate disability (Oswestry Disability Index [ODI] – 31 [14–38]). Positive significant correlations were found between ODI and age, residence, work status, VAS, PCS-rumination, PCS-magnification, PCS-helplessness and BDI, and also between PCS subscales and VAS. Our multivariate linear regression analysis showed that age, pain intensity, PCS-helplessness and depression can predict disability in patients with CLBP, explaining 84% of the variance of disability ($R^2 = 0.851$, adjusted $R^2 = 0.843$).

Conclusions: A multidisciplinary approach is needed for patients with CLBP and should include physical, mental and social evaluation in order to offer an optimal treatment.

Key words: chronic low back pain, catastrophizing, depression, disability.

Introduction

Chronic low back pain (CLBP) is defined as low back pain lasting more than 12 weeks, affecting more than 50% of the general population. It is estimated that more than 70% of the population experience at least one episode of lower lumbar pain, in a certain moment of life. Prevalence is higher among young people, being the second cause of absence from the workplace and one of the main reasons for consulting a doctor. Back

pain typically occurs between the ages of 30 and 50, due to the ageing processes but also secondary to a sedentary lifestyle. Approximately 10–40% of low back pain complaints are chronic, of which 85% are nonspecific [1].

Data from the European Health surveys show a wide variation in the prevalence of self-reported low back pain. This ranges from less than 12% of respondents reporting ever having doctor-diagnosed low back pain in France to nearly 33% in Austria reporting ever having had this condition. Among the European countries, the highest proportion of people reporting ever having had low back pain (including not diagnosed by a doctor) is in Slovenia (40.7%) [2]. In Romania the prevalence of low back pain in adults is high (62%), second only to headaches (79%) in the ranking of painful disorders [3].

This condition may negatively impact the patient's quality of life, and may influence activity for long periods of time, leading to absenteeism in the workplace. Apart from pain and disability the patients can present depression, due to chronic pain, and increased prescription drugs consumption [4]. Patients with chronic pain are more likely to develop depression and even higher levels of pain and worse health-related quality of life [5–12].

Increased attention has recently been given to the concept of pain catastrophizing, often encountered in patients with chronic conditions. This term was introduced to describe an inappropriately cognitive style used by patients with anxiety and depressive disorders. Catastrophic thinking with regard to pain involves a set of exaggerated and negative cognitive and emotional feelings related to actual or anticipated painful stimulation [13]. Some patients define catastrophizing as a tendency to magnify or exaggerate the value or seriousness of pain sensations, while others emphasized pain-related worry and fear, associated with the inability to divert the attention away from pain [14]. According to Sullivan *et al.*, the catastrophic thinking incorporates magnification of pain-related symptoms, rumination about pain, helplessness, and pessimism about the underlying disease [15]. Previous studies demonstrated the association of catastrophizing and sensitization with increased clinical pain among patients with chronic low back pain and myofascial pain [16–18]. Central sensitization seems to mediate the relation between psychological factors (anxiety, depression, catastrophic thinking) and pain intensity [19, 20].

In this study, we aimed to assess the relationship between pain intensity, the components of catastrophizing, depression and disability in patients with chronic low back pain.

Material and methods

Subjects

A total of 100 consecutive patients who were seeking help for their low back pain in a rehabilitation medicine clinic in the period from December 2018 to February 2019 were selected.

Inclusion criteria were chronic low back pain, defined as pain in the back, located between the last rib and the gluteal fold, with mechanical characteristics lasting more than 3 months. Subjects with sciatica, history of back surgery, spinal tumor, spinal fracture, spinal stenosis or radiculopathy, fibromyalgia, inflammatory and infectious spinal diseases were excluded. Chronic pain relief drug users were also excluded from this study.

This study conformed to the principles outlined in the Helsinki Declaration and was approved by the local Ethics Committee (approval no. 198/2018). Participation in the study was voluntary. All subjects who met the inclusion criteria read and signed the informed consent.

Assessment

Participants' socio-demographic data were collected – age, gender, height, weight, and work status (employed or retired). The main anthropometric parameters measured were weight, height and body mass index of the participants. All participants were asked to complete the Pain Catastrophizing Scale (PCS), the visual analogue scale (VAS), the Oswestry Disability Questionnaire (ODQ), and the Beck Depression Inventory (BDI).

The PCS was used to measure the degree of catastrophic thinking about pain. The scale has 13 items and three dimensions of pain catastrophizing: rumination, magnification, and helplessness. All items are scored through a 5-point Likert scale ranging from 0 (not at all) to 4 (all the time), relating to the past painful experience. Separate subscores for the three dimensions can be calculated. For example, rumination is the sum of items 8, 9, 10 and 11; magnification is the sum of items 6, 7 and 13; and helplessness is the sum of items 1, 2, 3, 4, 5 and 12. A total score ranging from 0 to 52 points can be calculated for the PCS. Higher scores denote a higher degree of catastrophizing. Patients were categorized as high or low catastrophizers based on a median split of PCS scores [15].

Pain intensity was self-completed by each patient on a single-item scale (Visual Analog Scale). This scale is most commonly anchored by “no pain” (score of 0) and “pain as bad as it could be” or “worst imaginable pain” (score of 10). Respondents were asked to report “current” pain intensity or pain intensity “in the last 24 hours”.

The ODQ scale was used to measure the limitation in everyday life activities. It is based on

Table I. Patients socio-demographic data

Characteristics	Value
Sex, n (%):	
Male	20 (26.3)
Female	56 (73.7)
Age [years], mean \pm SD	53.79 \pm 13.82
Weight [kg], mean \pm SD	76.11 \pm 12.5
Height [cm], mean \pm SD	167.35 \pm 9.17
BMI [kg/m ²], mean \pm SD	27.19 \pm 3.75
Residence, n (%):	
Rural	20 (26.3)
Urban	56 (73.7)
Work status, n (%):	
Employed	51 (67.1)
Retired	25 (32.9)

Table II. Summary of scores

Variables	Median [IQR]
Pain Catastrophizing Scale:	
PCS total	20.5 [10–34]
Rumination	7 [4–12]
Magnification	4.5 [3–7]
Helplessness	9 [3–17]
VAS	4 [3–6]
Oswestry Disability Index	31 [14–38]
Beck Depression Index	10 [5–16]

10 sections with six levels each, assessing the limitation of various activities of daily living [21, 22]. The values range from 0 (the best health state) to 100 (the worst health state). For each section of the questionnaire, the total possible score is 5. The first statement was scored 0, and consecutive statements were scored from 1 to 5. The total score was then divided by the total possible score and expressed as a percentage to produce the Oswestry Disability Index (ODI). The ODI is interpreted as follows: 0–20%, minimal disability; 21–40%, moderate disability; 41–60%, severe disability; 61–80%, crippled; 81–100%, patients are either bed-bound or exaggerate their symptoms [21, 22].

The Beck Depression Inventory (BDI) is a 21-item, self-report rating inventory that measures the intensity of depression [23]. Each question had a set of four possible responses, ranging in intensity, and a score of 0 to 3 is assigned for each answer. The sum of the 21 scored questions gives the total score. A score of 0 to 9 indicates minimal depression; 10 to 18 indicates mild depression; 19 to 29 indicates moderate depression; 30 to 63 indicates

severe depression. Higher total scores indicate more severe depressive symptoms.

Statistical analysis

The statistical analysis was performed using MedCalc Statistical Software version 19 (MedCalc Software bvba, Ostend, Belgium). All data were tested for normality with Shapiro-Wilk's test. Descriptive statistics were calculated for patients' characteristics (mean and standard deviations) and for VAS, PCS, ODI and BDI (median and interquartile range). Correlation (Spearman's rank correlation coefficient) and multivariate linear regression with a stepwise procedure were conducted to examine the relationship between ODI and the other variables (gender, age, BMI, work status, pain intensity (VAS), rumination, magnification, helplessness and BDI). The criterion for entry into the regression model was $p < 0.05$ and for removal from the regression was $p > 0.1$. The significance level was set at $p < 0.05$ for all tests.

In order to calculate the required sample size for a multiple regression analysis, the G*Power 3.1 software was used. Using up to 8 variables in the regression analysis, at a large effect size of 0.35, at an alpha level of 0.05 and a power of 0.8, a minimum sample of 52 participants was required [24].

Results

Of the 100 patients recruited for this study, only 76 (aged ranged 25–77 years; 73.7% female) agreed to participate, signed the informed consent and answered the questionnaires. Table I outlines the patients' socio-demographic data.

The responses of the VAS, PCS scores, ODI and BDI are presented in Table II. None of the variables showed a normal distribution. The mean group scores revealed moderate CLBP complaints (VAS – 4 [3–6]), mild depression (BDI – 10 [5–16]), a moderate level of catastrophizing (PCS total score 20.5 [10–34]) and moderate disability (ODI – 31 [14–38]). The majority of the patients (55.26%) were low catastrophizers (PCS total $<$ 24), with a VAS of 4 [2–4], a BDI of 7 [2–11] and an ODI of 18 [12–28]. The high catastrophizers (44.74%) had significantly higher scores (VAS – 7 [5–7]), BDI – 16 [11–30] and ODI – 38 [34–52]). Twenty-four patients (31.58%) were scored as having minimal disability, 36 patients (47.37%) moderate disability, and 16 patients (21.05%) severe disability.

There were positive weak significant correlations between socio-demographic data (age, residence, work status) and pain, PCS subscales, depression and disability. Positive significant correlations were found between ODI and age, residence, work status, VAS, PCS total, PCS-rumination, PCS-magnification, PCS-helplessness and

Table III. Correlation among measured variables

	Sex	Age	Residence	Work status	BMI	VAS	ODI	PCS total	PCS-R	PCS-M	PCS-H	BDI
Sex												
Age	0.30*											
Residence	-0.19	-0.06										
Work status	0.22	0.80*	0.15									
BMI	0.03	0.05	0.05	0.02								
VAS	0.08	0.29*	0.30*	0.46*	-0.15							
ODI	0.15	0.63*	0.31*	0.61*	0.10	0.72*						
PCS total	0.11	0.56*	0.32*	0.57*	0.06	0.75*	0.85*					
PCS-R	0.08	0.44*	0.30*	0.48*	0.07	0.67*	0.74*	0.94*				
PCS-M	0.0	0.48*	0.29*	0.51*	0.06	0.71*	0.80*	0.95*	0.91*			
PCS-H	0.18	0.58*	0.28*	0.56*	0.10*	0.74*	0.87*	0.96*	0.85*	0.88*		
BDI	0.06	0.44*	0.26*	0.47*	0.08	0.66*	0.77*	0.76*	0.69*	0.74*	0.75*	

Spearman rank correlation coefficient. * $P < 0.05$. BMI – body mass index, VAS – visual analogue scale for pain, ODI – Oswestry Disability Index, PCS-R – Pain Catastrophizing Scale-Rumination, PCS – Pain catastrophizing Scale-Magnification, PCS-H – Pain catastrophizing Scale-Helplessness, BDI – Beck Depression Index.

Table IV. Summary of stepwise regression analysis of disability (ODI)

Predictors	B	β	t	P-value	R ²	Adjusted R ²
Age	0.216	0.193	3.320	0.001	0.851	0.843
VAS	1.779	0.223	3.164	0.002		
PCS-helplessness	0.651	0.302	3.235	0.002		
BDI	0.511	0.365	4.808	< 0.0001		

BDI, and also between PCS total score, PCS subscales and VAS (Table III).

A multivariate linear regression with a stepwise procedure was conducted to examine the relationship between ODI and the other variables. Only those variables that were significant in the correlation analysis were entered into the regression. A significant model emerged ($F(4,71) = 101.38$, $p < 0.0001$), explaining 84% of the variance of disability ($R^2 = 0.851$, adjusted $R^2 = 0.843$). Age, VAS, PCS-helplessness and BDI contributed significantly in this model. The excluded variables from this regression model were residence, work status, PCS-rumination and PCS-magnification. The regression coefficients of the predictors are shown in Table IV.

Discussion

This study investigated whether socio-demographic data, catastrophizing, pain intensity and depression would predict disability in patients with chronic low back pain. In our regression model, age, pain intensity, helplessness and depression explained the variance of disability in patients with nonspecific CLBP.

Socio-demographic (e.g. age, sex, work status, BMI), pain characteristics (intensity, duration, localization, causes) and psychological factors (fear-avoidance beliefs, catastrophizing, anxiety, depression) are reported as predictors of disability not only in patients with chronic pain, but also in patients with CLBP [25–29].

Our results showed that age, residence and work status were significantly correlated with the disability, but only age contributed significantly in the prediction of disability in CLBP patients, similar to previous studies [30–32].

Consistent with previous research, we found that pain intensity was correlated with disability and was one of the independent variables which significantly influenced the disability in our CLBP patients. Significant correlations between CLBP intensity and disability have also been reported in the literature [33–36]. Pain intensity was proved to be a strong predictor of disability in populations with non-specific low back pain in previous studies [37–39].

Catastrophizing has been defined as exaggerated negative emotions regarding pain experience and stimuli [15, 40]. It was reported that catastrophizing could predict pain intensity, disability and psychological distress, and that it has a detrimen-

tal effect on general health perception, functional status, medical consumption and quality of life [41–43]. There are a lot of studies in the literature regarding the role of pain catastrophizing in the prognosis and disability of patients with low back pain. Catastrophizing was strongly correlated with disability and the impact on daily living in populations with low back pain [33, 34, 38, 44, 45]. In contrast, there are contradicting studies which report that catastrophizing does not explain the unique variance in disability when assessed alongside factors such as pain-related fear or coping styles [35, 46, 47].

Our study revealed that helplessness was the only component of catastrophizing that significantly predicted disability in patients with NCLBP. One explanation for this situation could be the strong correlations observed between the three PCS subscales. It is possible for some patients to identify so much with pain and to pay more attention to it so that they develop in time helplessness phenomena and implicitly pain-related disability. Similar results were reported by Rosenstiel and Keefe, who found that helplessness (assessed with the Coping Strategy Questionnaire) was associated with greater disability [48]. Sullivan *et al.* found that helplessness was the strongest predictor of pain-related disability in the group of patients off work for more than 4 years [49]. In the same study they observed that for patients who had been off work for 2–4 years rumination was the only component that predicted disability. Vienneau *et al.* observed in their study that helplessness was the best predictor of disability in patients with CLBP [50]. Our results are not in accordance with other studies. Sullivan *et al.* reported that rumination was the component of catastrophizing that was strongly associated with disability in patients with soft-tissue injuries [51]. Ogunlana *et al.* found that the magnification and rumination components of pain catastrophizing explained 22.6% of the variance of disability in patients with NLBP [34].

It is known that depression is the most frequent emotional condition in individuals with chronic pain or chronic conditions [52, 53]. In our study the correlation between disability and depression was positive and significant, suggesting that the greater the disability related to chronic back pain is, the greater is the possibility of depression-related symptoms. Depression contributed significantly in our model for predicting CLBP-related disability. It is common for depressed patients to avoid activities they used to do, such as housework, sports, social interaction, etc., which leads to disability-related symptoms. Our findings are in accordance with previous research. Hung *et al.* reported that depression was the most powerful factor associated with disability in their studied populations with CLBP [54].

Another interesting study focused on mechanisms that might be involved in the development of depression and disability. The authors suggest that a potential mechanism is the individual's health locus of control (HLoC). In psychology, health locus of control refers to people's beliefs that they have control over different events in their lives (internal locus of control) or that their own actions are a result of external factors beyond their control (external locus of control). The primary outcome of this study is that peoples with low control over their health (internal HLoC) report high levels of depression resulting from pain and pain interference, but moderate levels of disability [55].

There are a number of limitations of our study that should be considered. We did not take into account the stage of chronicity of low back pain or the use of medication. We also did not evaluate the fear avoidance beliefs and the coping strategies that could interfere with catastrophizing and depression in our study sample.

Our findings can contribute to a better understanding of the predictive factors of disability in patients with CLBP, such as age, pain intensity, catastrophizing pain and depression. The need to assess the abovementioned variables is an essential element in the complex approach of the patient with CLBP. Moreover, the knowledge of these factors can provide support for health professionals to choose more effective therapeutic approaches.

Acknowledgments

Elena Sirbu and Simona Szasz have contributed equally to the study.

Conflict of interest

The authors declare no conflict of interest.

References

1. Key J. The problem of back pain. In: Key J, editor. Back Pain – A Movement Problem: A clinical approach incorporating relevant research and practice. Edinburgh: Churchill Livingstone; 2010; 3-10.
2. eumusc.net. Musculoskeletal Health in Europe Report v5.0. Available at: <http://www.eumusc.net/myUploadData/files/Musculoskeletal Health in Europe Report v5.pdf>.
3. Busse R, Schreyögg J, Smith PC. Variability in healthcare treatment costs amongst nine EU countries – results from theHealthBASKET project. Health Econ 2008; 17: S1-8.
4. Sathya P, Ramakrishnan KS, Shah PP. Prevalence of depression, anxiety and stress in patients with mechanical low back pain. Int J Ther Rehabil Res 2015; 4: 67-72.
5. Hiyama A, Watanabe M, Katoh H, Sato M, Sakai D, Mochida J. Effect of depression and neuropathic pain using questionnaires on quality of life in patients with low back pain; cross-sectional retrospective study. Eur Spine J 2016; 25: 2750-60.
6. Manchikanti L, Fellows B, Pampati V, Beyer C, Damron K, Barnhill RC. Comparison of the psychological status of

- chronic pain patients and the general population. *Pain Physician* 2002; 5: 40-8.
7. Tsuji T, Matsudaira K, Sato H, Vietri J. The impact of depression among chronic low back pain patients in Japan. *BMC Musculoskelet Disord* 2016; 17: 447.
 8. López-López D, Vilar-Fernández JM, Calvo-Lobo C, Losa-Iglesias ME, Rodríguez-Sanz D, Becerro-de-Bengoa-Vallejo R. Evaluation of depression in subacute low back pain: a case control study. *Pain Physician* 2017; 20: E499-E506.
 9. López-López D, Becerro-de-Bengoa-Vallejo R, Losa-Iglesias ME, Rodríguez-Sanz D, Palomo-López P, Calvo-Lobo C. Relationship between depression scores and degree of skin perspiration: a novel cross-sectional study. *Int Wound J* 2019; 16: 139-43.
 10. Calvo-Lobo CC, Vilar-Fernández JM, Losa-Iglesias ME, et al. Depression symptoms among older adults with and without subacute low back pain. *Rehabil Nurs* 2019; 44: 47-51.
 11. Baskan E, Yagci N, Cavlak U. Factors affecting body awareness in older adults with chronic musculoskeletal pain. *Arch Med Sci* 2021; 17: 934-9.
 12. Kuzstal M, Trafidło E, Madziarska K, et al. Depressive symptoms but not chronic pain have an impact on the survival of patients undergoing maintenance hemodialysis. *Arch Med Sci* 2018; 14: 265-275.
 13. Leung L. Pain catastrophizing: an updated review. *Indian J Psychol Med* 2012; 34: 204-17.
 14. Craner JR, Sperry JA, Evans MM. The relationship between pain catastrophizing and outcomes of a 3-week comprehensive pain rehabilitation program. *Pain Med* 2016; 17: 2026-35.
 15. Sullivan MJL, Bishop SR, Pivik J. The pain catastrophizing scale: development and validation. *Psychol Assess* 1996; 7: 524-32.
 16. Meints SM, Mawla I, Napadow V, et al. The relationship between catastrophizing and altered pain sensitivity in patients with chronic low-back pain. *Pain* 2019; 160: 833-43.
 17. San-Antolín M, Rodríguez-Sanz D, Becerro-de-Bengoa-Vallejo R, et al. Central sensitization and catastrophism symptoms are associated with chronic myofascial pain in the gastrocnemius of athletes. *Pain Med* 2020; 21: 1616-25.
 18. Huysmans E, Ickmans K, Van Dyck D, et al. Association between symptoms of central sensitization and cognitive behavioral factors in people with chronic nonspecific low back pain: a cross-sectional study. *J Manipulative Physiol Ther* 2018; 41: 92-101.
 19. Shigetoh H, Tanaka Y, Koga M, Osumi M, Morioka S. The mediating effect of central sensitization on the relation between pain intensity and psychological factors: a cross-sectional study with mediation analysis. *Pain Res Manag* 2019; 2019: 3916135.
 20. Aoyagi K, He J, Nicol AL, et al. A subgroup of chronic low back pain patients with central sensitization. *Clin J Pain* 2019; 35: 869-79.
 21. Davidson M, Keating JL. A Comparison of Five Low Back Disability Questionnaires: reliability and responsiveness. *Phys Ther* 2002; 82: 8-24.
 22. Copay AG, Glassman SD, Subach BR, Berven S, Schuler TC, Carreon LY. Minimum clinically important difference in lumbar spine surgery patients: a choice of methods using the Oswestry Disability Index, Medical Outcomes Study Questionnaire Short Form 36, and Pain Scales. *Spine J* 2008; 8: 968-74.
 23. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961; 4: 561-71.
 24. Faul F, Erdfelder ER, Lang AG, Buchner A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007; 39: 175-91.
 25. Heapy A, LaChappelle K, Higgins D, et al. Demographic, psychological, and clinical predictors of Roland Morris Disability Questionnaire score among patients with chronic low back pain. *J Pain* 2016; 17: S4-5.
 26. Raftery MN, Sarma K, Murphy AW, De la Harpe D, Normand C, McGuire BE. Chronic pain in the Republic of Ireland – community prevalence, psychosocial profile and predictors of pain-related disability: results from the Prevalence, Impact and Cost of Chronic Pain (PRIME) study, Part 1. *Pain* 2011; 152: 1096-103.
 27. McIntyre A, Alibrahim F, Mehta S, et al. Predictors of pain disability among individuals who use opioids attending a chronic pain clinic. *Arch Phys Med Rehabil* 2018; 99: e64.
 28. Crowell A, Younger J, Lobato R, et al. Demographic and psychosocial predictors of disability due to pain. *J Pain* 2007; 8: S76.
 29. Lötters F, Franche RL, Hogg-Johnson S, Burdorf A, Pole JD. The prognostic value of depressive symptoms, fear-avoidance, and self-efficacy for duration of lost-time benefits in workers with musculoskeletal disorders. *Occup Environ Med* 2006; 63: 794-801.
 30. Häuser W, Schmutzer G, Brähler E, Schiltenswolf M, Hilbert A. The impact of body weight and depression on low back pain in a representative population sample. *Pain Med* 2014; 15: 1316-27.
 31. Schmidt CO, Raspe H, Pflingsten M, et al. Back pain in the German adult population: prevalence, severity, and sociodemographic correlates in a multiregional survey. *Spine* 2007; 32: 2005-11.
 32. Webb R, Brammah T, Lunt M, Urwin M, Allison T, Symons D. Prevalence and predictors of intense, chronic, and disabling neck and back pain in the UK general population. *Spine* 2003; 28: 1195-202.
 33. Thomas EN, Pers YM, Mercier G, et al. The importance of fear, beliefs, catastrophizing and kinesiophobia in chronic low back pain rehabilitation. *Ann Phys Rehabil Med* 2010; 53: 3-14.
 34. Ogunlana MO, Odole AC, Adejumo A, Odunaiya N. Catastrophizing, pain, and disability in patients with nonspecific low back pain. *Hong Kong Physiother J* 2015; 33: 73-9.
 35. Woby SR, Watson PJ, Roach NK, Urmston M. Are changes in fear-avoidance beliefs, catastrophizing, and appraisals of control, predictive of changes in chronic low back pain and disability? *Eur J Pain* 2004; 8: 201-10.
 36. Jensen MP, Romano JM, Turner JA, Good AB, Wald LH. Patient beliefs predict patient functioning: further support for a cognitive-behavioural model of chronic pain. *Pain* 1999; 81: 95-104.
 37. Peters ML, Vlaeyen JWS, Weber WEJ. The joint contribution of physical pathology, pain-related fear and catastrophizing to chronic back pain disability. *Pain* 2005; 113: 45-50.
 38. van den Hout JHC, van den Vlaeyen JWS, Heuts PHTG, Sillen WJT. Functional disability in nonspecific low back pain: the role of pain-related fear and problem-solving skills. *Int J Behav Med* 2001; 8: 134-48.
 39. Mannion AF, Junge A, Taimela S, Müntener M, Lorenzo K, Dvorak J. Active therapy for chronic low back pain: part 3. Factors influencing self-rated disability and its change following therapy. *Spine* 2001; 26: 920-9.
 40. Chaves JF, Brown JM. Spontaneous cognitive strategies for the control of clinical pain and stress. *J Behav Med* 1987; 10: 263-76.

41. de Boer MJ, Struys MMRĲ, Versteegen GJ. Pain-related catastrophizing in pain patients and people with pain in the general population. *Eur J Pain* 2012; 16: 1044-52.
42. Severeijns R, Vlaeyen JWS, van den Hout MA, Weber WEJ. Pain catastrophizing predicts pain intensity, disability, and psychological distress independent of the level of physical impairment. *Clin J Pain* 2001; 17: 165-72.
43. Severeijns R, van den Hout MA, Vlaeyen JWS, Picavet HSJ. Pain catastrophizing and general health status in a large Dutch community sample. *Pain* 2002; 99: 367-76.
44. Ramírez-Maestre C, Esteve R, Ruiz-Párraga G, Gómez-Pérez L, López-Martínez AE. The key role of pain catastrophizing in the disability of patients with acute back pain. *Int J Behav Med* 2017; 24: 239-48.
45. Picavet HSJ, Vlaeyen JWS, Schouten JSAG. Pain catastrophizing and kinesiophobia: predictors of chronic low back pain. *Am J Epidemiol* 2002; 156: 1028-34.
46. Woby SR, Urmston M, Watson PJ. Self-efficacy mediates the relation between pain-related fear and outcome in chronic low back pain patients. *Eur J Pain* 2007; 11: 711-8.
47. Turner JA, Jensen MP, Romano JM. Do beliefs, coping, and catastrophizing independently predict functioning in patients with chronic pain? *Pain* 2000; 85: 115-25.
48. Rosenstiel AK, Keefe FJ. The use of coping strategies in chronic low back pain patients: relationship to patient characteristics and current adjustment. *Pain* 1983; 17: 33-44.
49. Sullivan MJL, Sullivan ME, Adams HM. Stage of chronicity and cognitive correlates of pain-related disability. *Cogn Behav Ther* 2002; 31: 111-8.
50. Vienneau TL, Clark AJ, Lynch ME, Sullivan MJL. Catastrophizing, functional disability and pain reports in adults with chronic low back pain. *Pain Res Manag* 1999; 4: 93-6.
51. Sullivan MJL, Stanish W, Waite H, Sullivan M, Tripp DA. Catastrophizing, pain, and disability in patients with soft-tissue injuries. *Pain* 1998; 77: 253-60.
52. Amaricai E, Poenaru DV. The post-stroke depression and its impact on functioning in young and adult stroke patients of a rehabilitation unit. *J Ment Health* 2016; 25: 137-41.
53. Ericsson M, Poston WSC, Linder J, Taylor JE, Haddock CK, Foreyt JP. Depression predicts disability in long-term chronic pain patients. *Disabil Rehabil* 2002; 24: 334-40.
54. Hung CI, Liu CY, Fu TS. Depression: an important factor associated with disability among patients with chronic low back pain. *Int J Psychiatry Med* 2015; 49: 187-98.
55. Campbell P, Hope K, Dunn KM. The pain, depression, disability pathway in those with low back pain: a moderation analysis of health locus of control. *J Pain Res* 2017; 10: 2331-9.