

Apoptosis Induced by 13-S-hydroxyoctadecadienoic acid in the Breast Cancer Cell Lines, MCF-7 and MDA-MB-231

Masoumeh Tavakoli-Yaraki¹, Fatemeh Karami-Tehrani^{1*}

¹ Department of Clinical Biochemistry, Cancer Research Laboratory, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

ARTICLE INFO

Article type:
Original article

Article history:
Received: Jul 10, 2012
Accepted: Sep 10, 2012

Keywords:
Apoptosis
MCF-7
MDA-MB-231
PPAR- δ
13-S-hydroxyoctadecadienoic acid

ABSTRACT

Objective(s): The 15-Lipoxygenase-1(15-LOX-1) pathway has become of considerable interest as a promising molecular approach for the modulation of cancer cell growth. 13-S-hydroxyoctadecadienoic acid (13(S)-HODE) is a main metabolite of 15-LOX-1 which is proposed to influence the cancer cell's growth. This study aims to investigate the role of 13(S)-HODE in the regulation of cell growth and apoptosis in the breast cancer cell lines.

Materials and Methods: MTT assay was used to examine the cytotoxic effect of 13(S)-HODE in the breast cancer cells, MCF-7 and MDA-MB-231. Annexin-V-FITC staining and cell cycle analysis were performed using flow cytometry. The effect of 13(S)-HODE on the expression level of Peroxisome proliferator-activated receptors- δ (PPAR- δ) was also evaluated.

Results: The results demonstrated that 13(S)-HODE inhibited cell growth in a dose and time dependant manner in MCF-7 and MDA-MB-231 cell lines. The reduction of cell growth was associated with the induction of cell cycle arrest and apoptosis in the breast cancer cell lines. Moreover, PPAR- δ was down-regulated in response to 13(S)-HODE administration.

Conclusion: This study conducted evidences in to the stimulatory effect of 13(S)-HODE on the inhibition of cell growth and induction of apoptosis in the breast cancer cell lines.

► Please cite this paper as:

Tavakoli-Yaraki M, Karami-Tehrani F. Apoptosis Induced by 13-S-hydroxyoctadecadienoic acid in the Breast Cancer Cell Lines, MCF-7 and MDA-MB-231. Iran J Basic Med Sci: 2013;16: 653-9.

Introduction

Breast cancer is considered as the leading cause of cancer-related mortality among women in the world (1). Many scientific efforts have been focused on finding better therapeutic strategies and new molecular approaches to reduce the mortality.

The homeostasis of cancer cells depends on the balance between apoptosis and cell proliferation(2); therefore, identification of the critical mechanisms associated with the loss of apoptosis and increased cell proliferation is important for molecular targeting. 15-lipoxygenase-1(15-LOX-1) belongs to the non-heme iron-containing di-oxygenases family which is expressed mainly in the reticulocytes, eosinophils, macrophages, and airway

epithelium(3). 15-LOX-1 catalyzes the dioxygenation of linoleic acid at the 13th carbon and produces 13-S-hydroxyoctadecadienoic acid (13(S)-HODE) (4). Linoleic acid is an essential poly unsaturated fatty acid in the human diet and changes in the level of 13(S)-HODE is associated with the pathological conditions (5). It has been shown that down-regulation of 15-LOX-1 in the lung(6), colorectal (7)esophageal (8), pancreatic(9) and breast(10) cancer, associated with the reduction of 13(S)-HODE, leads to the cancer progression (5,6,9,11). Moreover, 13(S)-HODE is implicated to the cell differentiation and apoptosis in colorectal cancer (12-15) and 13(S)-HODE acts as a modulator of tumor invasion and metastasis (16,17).

Peroxisome proliferator -activated receptors (PPARs)

* Corresponding author: Fatemeh Karami-Tehrani, Department of Clinical Biochemistry, Cancer Research Laboratory, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran. Tel: +98- 21- 82883567; Fax: +98-21- 82884555; E-mail: karamitf@modares.ac.ir

© 2013 mums.ac.ir All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

are ligand-activated transcription factors belong to the nuclear receptor super family (18). Three isoforms of PPARs have been recognized including PPAR α , PPAR δ and PPAR γ . The involvement of PPARs in lipid homeostasis and cellular differentiation, tumorigenesis and cancer development have been well-established so far (19,20). Polyunsaturated fatty acids such as 13(S)-HODE and their related metabolites modulate PPAR- δ expression and function (19). It has been shown that 13(S)-HODE significantly reduces PPAR- δ expression and activity in colorectal cancer (21). 15-LOX-1 expression is down-regulated in the breast cancer. However, the mechanistic contribution in the breast tumorigenesis remains to be elucidated(10). On the other hand, the involvement of 13(S)-HODE in the regulation of breast cancer cell growth is not clearly determined. Accordingly, the current study is designed to evaluate the effect of 13(S)-HODE on the inhibition of cell growth, induction of apoptosis and cell cycle distribution in human breast cancer cell lines, MDA-MB-231 and MCF-7. The expression level of PPAR- δ as an endogenous receptor for 13(S)-HODE has been assessed in both cell lines. The results demonstrate that 13(S)-HODE suppresses the growth of MDA-MB-231 and MCF-7 cells which is associated with the cell cycle arrest and induction of apoptosis. The effect of 13(S)-HODE is also accompanied by the reduction of PPAR- δ expression level.

Materials and Methods

Chemical materials and Cell culture

RPMI 1640, trypsin/EDTA, NaCl/Pi, penicillin and streptomycin were obtained from Gibco (Rockville, USA). 13-S-hydroxyoctadecadienoic acid (13(S)-HODE) was provided by Cayman Chemicals (Michigan, USA). Annexin-V-FITC apoptosis detection kit, Propidium Iodid (PI) ,MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide], dimethylsulfoxide were purchased from Sigma-Aldrich (Munich, Germany).

Human breast cancer cell lines, MCF-7 and MB-MDA-231 were purchased from Pasteur Institute of Iran. Cells were cultured in RPMI 1640 containing 10% (v/v) fetal bovine serum, 100 U/ml of penicillin and 100 μ g/ml of streptomycin and were maintained at 37 °C with an atmosphere of 5% CO₂ and 100% humidity. Cells were used freshly and collected by trypsinization at the confluence of 70-100%. The results of 13(S)-HODE on the breast cancer cell growth were always compared to the cells treated with medium alone and cells treated with Ethanol as a solvent of 13(S)-HODE, in order to validate the effect of 13(S)-HODE. Untreated cells and cells treated with different concentrations of Ethanol equal to 13(S)-HODE had no effect on MCF-7 and MDA-MB-231 viability, apoptosis and cell cycle distribution (data not shown).

Cell viability assay: MTT assay

MTT assay was used to determine the viability of treated and untreated cells. MCF-7 and MB-MDA-231 cell lines were cultured in 96-well plates at 5×10^3 cell/well in RPMI 1640 (supplemented with 10% fetal bovine serum, 100 U/ml of penicillin and 100 μ g/ml of streptomycin) in 5% CO₂ at 37 °C until nearly confluent. 13(S)-HODE was dissolved in ethanol and cells were treated with the indicated concentrations of 13(S)-HODE and incubated in RPMI 1640 (supplemented with 2% FBS) for 24, 48 and 72 hr. Cells treated with the Ethanol (as a vehicle) were considered as a control. 20 μ l of MTT (5mg/ml in PBS) was added to each well and cells were incubated for 4 hr at 37 °C. The supernatant of each well was removed and formazan crystals were dissolved in 200 μ l of dimethylsulfoxide, respectively. The absorbance was read by microplate reader at 570 nm (Bio-Rad, Hercules, CA, USA). Treated groups were compared to vehicle controls and the percentages of viable cells were reported, respectively. The Data represent at least three separate repeated experiments.

Flow cytometric analysis of apoptosis

To quantify the apoptotic cell death, Annexin-V-FITC kit was applied and the results were analyzed using flow cytometry according to the manufacturer's specifications. Briefly, cells were harvested at the density of 1×10^6 cells/ml following treatment with various concentrations of 13(S)-HODE (Cells treated with the Ethanol (as a vehicle) were considered as a control.) and washed twice with PBS. Cell pellets were resuspended in 500 μ l of 1 \times binding buffer, following gentle vortex, 5 μ l of annexin-V-FITC and 5 μ l of PI were added to each sample. Cells were incubated 10 min at room temperature in the dark and further analyzed by FACSCalibur flow cytometer (Becton Dickinson, San Jose, USA) using the supplied software in the instrument (BD CellQuest software).

Analysis of Peroxisome proliferator -activated receptors- δ (PPAR- δ) mRNA expression

Trizol (Invitrogen, Grand Island, USA) was used to extract total RNA from cells according to the manufacturer's instructions. To evaluate RNA integrity agarose gel electrophoresis was applied. Then 1 μ g of total RNA was reverse-transcribed to synthesize single-stranded cDNA using High Capacity RNA-to-cDNA Kit according to the manufacturer's specifications (Invitrogen, Grand Island, USA). Specific cDNA was used as a template and quantitative real-time RT-PCR analysis of PPAR- δ was performed using the SYBR Green kit (Qiagen) in an ABI 7500 Sequence Detection System (Applied Biosystems) according to the instructions and the relative gene expression levels were normalized to the expression level of GAPDH (housekeeping gene). The relative expression level was evaluated us-

ing $2^{-\Delta\Delta Ct}$ analysis method. RT-PCR analysis was also performed and the products were analyzed by agarose gel electrophoresis 2% (w/v).

The following primer sets were applied:

PPAR- δ (NM-006238): Forward, 5'- agcagccttcctcaacgag-3', and Reverse, 5'- ggtctcggttcggctcttctgat-3',
glyceraldehydes-3-phosphatedehydrogenase(GAPDH) (NM-002046):

Forward: 5'- aatgacccttcattgacctc-3', Reverse: 5'-agttgtcatggatgaccttg-3'

Cell cycle analysis

PI staining was used to measure the DNA content and cell distribution. The MCF-7 and MB-MDA-231 cells were treated with different concentrations of 13(S)-HODE for 48 hr (Cells treated with the Ethanol (as a vehicle) were considered as a control). Then cells were harvested and fixed with 70% (v/v) ice-cold ethanol and washed twice with ice-cold PBS and stained with 0.1mg/ml of RNase A (Sigma) and 0.05 mg/ml of PI (Sigma) for 1 hr at 37 °C in the dark.

Cell cycle distribution and the percentage of cells in various cell cycle phases were determined using FACS-Calibur flow cytometer (Becton Dickinson, San Jose, USA) and inbuilt software (BD CellQuest software). The sub-G1 population was considered as apoptotic cells.

Statistical analysis

Non-parametric one-way analysis of variance (ANOVA) with *post hoc* Dennet's test was conducted to analyze and compare data using software GraphPad Prism. Data was presented as mean \pm SD and all experiments were repeated at least three times. Differences were taken significant for $P < 0.05$ and determined by asterisk in corresponding Figure.

Results

The effect of 13(S)-HODE on the cell growth inhibition in the breast cancer cell lines

The effect of 13(S)-HODE was examined on the growth of MCF-7 and MDA-MB-231 breast cancer cell lines. Cells were treated with different concentrations of 13(S)-HODE (5-100 μ M) and incubated in RPMI 1640 9 supplemented with 2% FBS for 24, 48 and 72 hr and the cell viability was measured by MTT assay. As indicated in Figure 1A, a significant reduction of MCF-7 cell viability was observed in a time and dose dependent manner. Treatment of MCF-7 cells with 100 μ M of 13(S)-HODE reduced the cell viability about 53.25% and 62.5% after 48 and 72 hr, respectively. Furthermore, treatment of MDA-MB-231 cells with 13(S)-HODE resulted in a significant decrease in the percent-

age of viable cells in a time and dose dependent manner. Treatment of MDA-MB-231 cells with 100 μ M of 13(S)-HODE reduced the cell viability about 57.75% and 62.25% after 48 and 72 hr, respectively (Figure 1.B). The IC50 value (the effective dose of 13(S)-HODE that causes 50% growth inhibition), for MCF-7 and MDA-MB-231 after 48 hr were 76.3 μ M and 80.23 μ M, respectively.




Figure 1. 13(S)-HODE inhibited cell growth in breast cancer cell lines. MCF-7(A) and MDA-MB-231(B) cells were treated with different concentrations of 13(S)-HODE (5-100 μ M) for 24 h, 48 h and 72 hr. The proportion of viable cells was determined by MTT assay. Cell-growth inhibition occurred in a dose and time-dependent manner in both cell lines. The data represents Mean \pm SD of three separate experiments. Statistical differences between 13(S)-HODE-treated and untreated control groups were analyzed by ANOVA (*= $P < 0.05$, **= $P < 0.01$, ***= $P < 0.001$).

The Effect of 13(S)-HODE on the cell cycle progression

The effect of 13(S)-HODE on the cell cycle progression was studied in the breast cancer cell lines. MCF-7 and MDA-MB-231 cells were treated with 20, 50, 100 μ M of 13(S)-HODE and incubated in RPMI 1640 (supplemented with 2% FBS) for 48 hr and cell cycle analysis were performed using flow cytometry. A significant increase in the percentage of Sub-G1 population in MCF-7 ($P < 0.001$) and MDA-MB-231 ($P < 0.001$) was observed (Figure 2). The elevation in the sub-G1 population occurred in a dose dependent manner in both cell lines. The results demonstrated that the treatment of the cells with 13(S)-HODE were associated with the disturbance in the cell cycle progression, raising the cell cycle arrest in the breast cancer cell lines.




Figure 2. The effect of 13(S)-HODE on cell cycle distribution in breast cancer cell lines. MCF-7(A) and MDA-MB-231(B) cells were treated with 20, 50 and 100 μ M of 13(S)-HODE for 48 hr and pattern of cell cycle distribution were analyzed by flow cytometry using PI staining as described in Materials and methods. 13(S)-HODE significantly increased the proportion of sub-G1 population in a dose-dependant manner in both cell lines. Data represent Mean \pm SD of three separate experiments. Statistical differences between 13(S)-HODE-treated and untreated control groups were analyzed by ANOVA (*= $P < 0.05$, **= $P < 0.01$, ***= $P < 0.001$).

The effect of 13(S)-HODE on the induction of apoptosis in MCF-7 and MDA-MB-231 cell lines

To examine the effect of 13(S)-HODE on the induction of apoptosis, annexin-V and PI double staining method was employed and analyzed using flow cytometry with BD CellQuest software. According to the instruction, annexin-V positive, PI negative cells are recognized as early apoptotic and annexin-V positive, PI positive are considered as late apoptotic cells. MCF-7 and MDA-MB-231 cells were treated with 20, 50 and 100 μ M of 13(S)-HODE and incubated in RPMI 1640 (supplemented with 2% FBS) for 48 hr, and then analyzed by flow cytometry. Annexin-negative and PI-negative cells were considered as controls. A significant increase was observed in the percentages of early ($P < 0.001$) and late ($P < 0.01$) apoptotic cells for MCF-7 cells (Figure 3A). A significant elevation in the level of early ($P < 0.001$) and late ($P < 0.01$) apoptotic cells was also detected in MDA-MB-231 cells (Figure 3B). The treatment of




Figure 3. 13(S)-HODE induced apoptosis in breast cancer cell lines. MCF-7(A) and MDA-MB-231(B) cells were treated with 20, 50 and 100 μ M of 13(S)-HODE for 48 hr. Apoptosis was detected by flow cytometric analysis using annexin-V and PI staining as described in Materials and Methods. 13(S)-HODE enhanced the percentage of early apoptotic cells (Annexin-V positive cells) and late apoptotic cells (Annexin-V and PI positive cells) in a dose-dependant manner in both cell lines. The data represents Mean \pm SD of three separate experiments. Statistical differences between 13(S)-HODE-treated and untreated control groups were analyzed by ANOVA (*= $P < 0.05$, **= $P < 0.01$, ***= $P < 0.001$).

the cells through increasing the concentrations of 13(S)-HODE resulted in a notable shift from viable to apoptotic cells in a dose dependant manner in both cell lines. The rate of apoptotic cells was ranged from 3.5% to 43.28% for MCF-7 and 2.99% to 40.2 % for MDA-MB-231.

The effect of 13(S)-HODE on the PPAR- δ gene expression in the breast cancer cell lines

The expression level of PPAR- δ was studied following the treatment by 13(S)-HODE to determine whether PPAR- δ expression is affected by 13(S)-HODE. Cells were treated in different concentrations of 13(S)-HODE (50 and 100 μ M) for 48 hr and then analyzed by real-time RT-PCR experiments. The results demonstrated that PPAR- δ expression level reduced in response to 13(S)-HODE in MCF-7 cell line, while 13(S)-HODE had no significant effect on the PPAR- δ expression level in MDA-MB-231 (Figure 4A), the result of RT-PCR analysis also supported the above data (Figure 4B).





Figure 4. The effect of 13(S)-HODE on the expression level of PPAR- δ in breast cancer cells.

Cells were treated with 50 and 100 μ M of 13(S)-HODE for 48 hr. Relative PPAR- δ gene expression level was evaluated. The expression level of PPAR- δ was decreased in response to 13(S)-HODE in MCF-7 while 13(S)-HODE revealed no effect on the expression of PPAR- δ in MDA-MB-231(A). RT-PCR analysis of mRNA expression in both cell lines (PPAR- δ , 437bp and GAPDH, 495bp) (B). The data represents Mean \pm SD of three separate experiments. Statistical differences between 13(S)-HODE -treated and untreated control groups were analyzed by ANOVA (*= $P < 0.05$)

Discussion

Lilonic acid is an essential poly-unsaturated fatty acid which produces 13(S)-HODE via 15-lipoxygenase (15-LOX-1) pathway (22). 13(S)-HODE mediates important molecular functions in mammalian cells including cellular adhesion(16) and proliferation(12). Aberrant expression of enzymes involved in the oxygenation of fatty acids is correlated with the modulation of cancer cell-growth. The subsequent effect depends on the expression of enzyme, type of formed metabolite and type of cancer involved. It has been well-established that 15-LOX-1 mediates anti-tumorigenic effect in various cancers, however, the correlated mechanism is not unraveled yet (10,17,23-25). Recent studies have shown that 13(S)-HODE mediates inhibition of cell-growth and induction of apoptosis in malignant cells(23). The pro-apoptotic role of 13(S)-HODE has been shown in colorectal cancer (5,7). It has been revealed that the reduction of 15-LOX-1 expression and 13(S)-HODE for-

mation are associated with the progression of breast cancer (10) These findings have emphasized on the possible effect of 13(S)-HODE on the inhibition of different cancer progression. The emergence of 13(S)-HODE as a novel molecular approach to trigger cell death pathways and lack of solid evidences on the precise role of 13(S)-HODE in the regulation of breast cancer cell growth, motivated us to study the effect of 13(S)-HODE in the regulation of cell growth and apoptosis in breast cancer which is the most leading cause of death in women. Treatment of MCF-7 and MDA-MB-231 cells with increasing concentrations of 13(S)-HODE inhibited cell growth in a dose and time-dependent manner in both cell lines. Our data is consistent with the inhibitory effect of exogenous 13(S)-HODE on the growth of colorectal (7) and Leukemia cell lines (23). The results revealed that the effect of 13(S)-HODE on the breast cancer cell growth is associated with an increase in the percentage of sub-G1 population which is consistent with the effective role of 13(S)-HODE in the regulation of cell-growth in human colon cancer cells (14) Our findings showed that the growth inhibition mediated by 13(S)-HODE was accompanied by a significant induction of apoptosis in a dose dependent manner in MCF-7 and MDA-MB-231 cell lines. In addition, it has been reported that the activation of PPAR- δ is correlated with cancer development in liver (26), prostate (27), breast (20,28,29) cancers. Ligand activation of PPAR- δ increases breast cancer cell proliferation and promotes mammary gland cancer development (27-29); polyunsaturated fatty acids and their related metabolites modulate PPAR- δ expression and function (19); therefore, the possible association of 13(S)-HODE with PPAR- δ in the breast cancer cells has been evaluated in this study and it has been demonstrated that 13(S)-HODE significantly reduces PPAR- δ expression and activity in colorectal cancer (21). Our findings indicate that PPAR- δ is down-regulated in response to 13(S)-HODE in MCF-7, however, no significant differences have been observed in MDA-MB-231. This is in line with the previous report which indicated that the growth of estrogen receptor negative breast cancer cells were not influenced by PPAR- δ (27). Our findings have demonstrated the effective role of 13(S)-HODE on the expression of this receptor, which is in consistency with the down-regulation of PPAR beta/delta by 13(S)-HODE in human colorectal cancer(21). However, reports about the effect of PPAR- δ on the regulation of breast cancer cell growth are in conflict. It has been reported that the ligand activation of PPAR- δ inhibits mouse mammary gland cells' growth (20), while the activation of PPAR- δ has been involved in the proliferation of mammary epithelial cells(29). This conflicting data can be related to the differences in the type of ligands, analysis procedures and involvement of other molecular pathways. However, it might be postulated that PPAR- δ may be involved in the apoptosis induced by 13(S)-HODE in breast cancer.

To characterize whether the 13(S)-HODE mediated growth inhibition is dependent on the estrogen activity, estrogen receptor positive (MCF-7) and negative (MDA-MB-231) breast cancer cell lines are used in this study. Our results demonstrate that no significant differences are observed in the sensitivity of the cells to the effect of 13(S)-HODE. Reports on the stimulatory or inhibitory effect of 13(S)-HODE on the cancer cell growth are discussed controversially. It has been reported that 13(S)-HODE in combination with EGF (epidermal growth factor) exhibit mitogenic effect on the SHE (Syrian hamster embryo) (30) rat hepatoma cells(31). Additionally, the elevation of 15-LOX-1 expression level and 13(S)-HODE formation have been reported in the tumors of prostate (32). Exogenous 13(S)-HODE has also increased the level of cell proliferation in prostate cancer cell lines (33). The results of other studies as well as our data support the anti-tumorigenic effect of 13(S)-HODE in cancer cells(23). This conflicting data on the significant role of 13(S)-HODE in the regulation of cell-growth can be explained by the tissue specificity of 13(S)-HODE and other possible molecular mechanisms that might be involved in the 15-LOX-1 signaling pathway. It is clear that further studies are required to characterize the precise molecular mechanism by which 13(S)-HODE affects cancer cell-growth.

Conclusion

Taken together, the data presented here demonstrates that 13(S)-HODE induces apoptosis and cell cycle arrest in estrogen receptor positive and negative breast cancer cell lines. The result of present study underlines the mediatory role of 13(S)-HODE for the induction of apoptosis in breast cancer cells and provides evidence on the possible relevance of PPAR- δ in the regulation of cell growth mediated by 13(S)-HODE; however, further studies are needed to clarify the specific role of PPAR family. Based on the data presented here and other investigations, 13(S)-HODE might emerge as a promising molecular approach for the induction of apoptosis in breast cancer.

Acknowledgement

This study was partly supported by a PhD grant (Masoumeh Tavakoli-Yaraki) by Tarbiat Modares University.

References

- Lambrechts S, Decloedt J, Neven P. Breast cancer prevention: Life-style changes and chemoprevention. *Acta Clin Belg* 2011; 66:283-292.
- Kasibhatla S, Tseng B. Why target apoptosis in cancer treatment? *Mol Cancer Ther* 2003; 2:573-580.
- Brash AR. Lipoxygenases: Occurrence, functions, catalysis, and acquisition of substrate. *J Biol Chem* 1999; 274:23679-23682.
- Brash AR, Jisaka M, Boeglin WE, Chang MS, Keeney DS, Nanney LB, et al. Investigation of a second 15s-lipoxygenase in humans and its expression in epithelial tissues. *Adv Exp Med Biol* 1999; 469:83-89.
- Shureiqi I, Wojno KJ, Poore JA, Reddy RG, Moussalli MJ, Spindler SA, et al. Decreased 13-s-hydroxyoctadecadienoic acid levels and 15-lipoxygenase-1 expression in human colon cancers. *Carcinogenesis* 1999; 20:1985-1995.
- Yuan H, Li MY, Ma LT, Hsin MK, Mok TS, Underwood MJ, Chen GG. 15-lipoxygenases and its metabolites 15(s)-hete and 13(s)-hode in the development of non-small cell lung cancer. *Thorax* 2010; 65:321-326.
- Nixon JB, Kim KS, Lamb PW, Bottone FG, Eling TE. 15-lipoxygenase-1 has anti-tumorigenic effects in colorectal cancer. *Prostaglandins Leukot Essent Fatty Acids* 2004; 70:7-15.
- Xu XC, Shappell SB, Liang Z, Song S, Menter D, Subbarayan V, et al. . Reduced 15s-lipoxygenase-2 expression in esophageal cancer specimens and cells and upregulation *in vitro* by the cyclooxygenase-2 inhibitor, ns398. *Neoplasia* 2003; 5:121-127.
- Hennig R, Kehl T, Noor S, Ding XZ, Rao SM, Bergmann F, et al. TE. 15-lipoxygenase-1 production is lost in pancreatic cancer and overexpression of the gene inhibits tumor cell growth. *Neoplasia* 2007; 9:917-926.
- Jiang WG, Watkins G, Douglas-Jones A, Mansel RE. Reduction of isoforms of 15-lipoxygenase (15-lox)-1 and 15-lox-2 in human breast cancer. *Prostaglandins Leukot Essent Fatty Acids* 2006; 74:235-245.
- Shureiqi I, Chen D, Lee JJ, Yang P, Newman RA, Brenner DE, et al. 15-lox-1: A novel molecular target of nonsteroidal anti-inflammatory drug-induced apoptosis in colorectal cancer cells. *J Natl Cancer Inst* 2000; 92:1136-1142.
- Hsi LC, Xi X, Lotan R, Shureiqi I, Lippman SM. The histone deacetylase inhibitor suberoylanilide hydroxamic acid induces apoptosis via induction of 15-lipoxygenase-1 in colorectal cancer cells. *Cancer Res* 2004; 64:8778-8781.
- Kumar KA, Arunasree KM, Roy KR, Reddy NP, Aparna A, Reddy GV, Reddanna P. Effects of (15s)-hydroperoxyeicosatetraenoic acid and (15s)-hydroxyeicosatetraenoic acid on the acute- lymphoblastic-leukaemia cell line jurkat: Activation of the fas-mediated death pathway. *Biotechnol Appl Biochem* 2009; 52:121-133.
- Shureiqi I, Chen D, Lotan R, Yang P, Newman RA, Fischer SM, et al. 15-lipoxygenase-1 mediates nonsteroidal anti-inflammatory drug-induced apoptosis independently of cyclooxygenase-2 in colon cancer cells. *Cancer Res* 2000; 60:6846-6850.
- Shureiqi I, Wu Y, Chen D, Yang XL, Guan B, Morris JS, et al. The critical role of 15-lipoxygenase-1 in colorectal epithelial cell terminal differentiation and tumorigenesis. *Cancer Res* 2005; 65:11486-11492.
- Pasqualini ME, Heyd VL, Manzo P, Eynard AR. Association between e-cadherin expression by human colon, bladder and breast cancer cells and the 13-hode:15-hete ratio. A possible role of their metastatic potential. *Prostaglandins Leukot Essent Fatty Acids* 2003; 68:9-16.
- Cimen I, Tuncay S, Banerjee S. 15-lipoxygenase-1 expression suppresses the invasive properties of colorectal carcinoma cell lines hct-116 and ht-29. *Cancer Sci* 2009; 100:2283-2291.
- Youssef J, Badr M. Peroxisome proliferator-activated receptors and cancer: Challenges and opportunities. *Br J Pharmacol* 2011;164:68-82.
- Chawla A, Repa JJ, Evans RM, Mangelsdorf DJ. Nuclear receptors and lipid physiology: Opening the x-files. *Science* 2001;294:1866-1870.
- Foreman JE, Sharma AK, Amin S, Gonzalez FJ, Peters JM. Ligand activation of peroxisome proliferator-activated receptor-beta/delta (pparbeta/delta) inhibits cell growth in a mouse mammary gland cancer cell line. *Cancer Lett* 2010; 288:219-225.
- Zuo X, Wu Y, Morris JS, Stimmel JB, Leesnitzer LM, Fischer SM, et al. Oxidative metabolism of linoleic acid modulates ppar-beta/delta suppression of ppar-gamma activity. *Oncogene* 2006; 25:1225-1241.
- Sigal E, Grunberger D, Highland E, Gross C, Dixon RA, Craik CS. Expression of cloned human reticulocyte 15-lipoxygenase and immunological evidence that 15-lipoxygenases of different cell

- types are related. *J Biol Chem* 1990; 265:5113-5120.
23. Mahipal SV, Subhashini J, Reddy MC, Reddy MM, Anilkumar K, Roy KR, et al. Effect of 15-lipoxygenase metabolites, 15-(s)-hpete and 15-(s)-hete on chronic myelogenous leukemia cell line k-562: Reactive oxygen species (ros) mediate caspase-dependent apoptosis. *Biochem Pharmacol* 2007; 74:202-214.
 24. Kim JH, Chang JH, Yoon JH, Lee JG, Bae JH, Kim KS. 15-lipoxygenase-1 induced by interleukin-4 mediates apoptosis in oral cavity cancer cells. *Oral Oncol* 2006; 42:825-830.
 25. Viita H, Pacholska A, Ahmad F, Tietavainen J, Naarala J, Hyvarinen A, et al. 15-lipoxygenase-1 induces lipid peroxidation and apoptosis, and improves survival in rat malignant glioma. *In Vivo* 2012; 26:1-8.
 26. Glinghammar B, Skogsberg J, Hamsten A, Ehrenborg E. Ppardelta activation induces cox-2 gene expression and cell proliferation in human hepatocellular carcinoma cells. *Biochem Biophys Res Commun* 2003; 308:361-368.
 27. Stephen RL, Gustafsson MC, Jarvis M, Tatoud R, Marshall BR, Knight D, et al. Activation of peroxisome proliferator-activated receptor delta stimulates the proliferation of human breast and prostate cancer cell lines. *Cancer Res* 2004; 64:3162-3170.
 28. Yin Y, Russell RG, Dettin LE, Bai R, Wei ZL, Kozikowski AP, et al. Peroxisome proliferator-activated receptor delta and gamma agonists differentially alter tumor differentiation and progression during mammary carcinogenesis. *Cancer Res* 2005; 65:3950-3957.
 29. Ghosh M, Ai Y, Narko K, Wang Z, Peters JM, Hla T. Ppardelta is pro-tumorigenic in a mouse model of cox-2-induced mammary cancer. *Prostaglandins Other Lipid Mediat* 2009; 88:97-100.
 30. Glasgow WC, Hui R, Everhart AL, Jayawickreme SP, Angerman-Stewart J, Han BB, et al. The linoleic acid metabolite, (13s)-hydroperoxyoctadecadienoic acid, augments the epidermal growth factor receptor signaling pathway by attenuation of receptor dephosphorylation. Differential response in syrian hamster embryo tumor suppressor phenotypes. *J Biol Chem* 1997; 272:19269-19276.
 31. Sauer LA, Dauchy RT, Blask DE, Armstrong BJ, Scalici S. 13-hydroxyoctadecadienoic acid is the mitogenic signal for linoleic acid-dependent growth in rat hepatoma 7288ctc *in vivo*. *Cancer Res* 1999; 59:4688-4692.
 32. Kelavkar UP, Cohen C, Kamitani H, Eling TE, Badr KF. Concordant induction of 15-lipoxygenase-1 and mutant p53 expression in human prostate adenocarcinoma: Correlation with gleason staging. *Carcinogenesis* 2000; 21:1777-1787.
 33. Kelavkar UP, Nixon JB, Cohen C, Dillehay D, Eling TE, Badr KF. Over-expression of 15-lipoxygenase-1 in pc-3 human prostate cancer cells increases tumorigenesis. *Carcinogenesis* 2001; 22:1765-1773.