

# Original Research





Received: Apr 22, 2021 Revised: Aug 6, 2021 Accepted: Aug 30, 2021 Published online: Nov 8, 2021

#### §Corresponding Authors:

#### Min-Jeong Shin

School of Biosystems and Biomedical Sciences, College of Health Science, Korea University, 145 Anam-ro, Seongbuk-gu, Seoul 02841, Korea.

Tel. +82-2-3290-5643 Fax. +82-2-940-2849 Email. mjshin@korea.ac.kr

#### Clara Yongjoo Park

Department of Food and Nutrition, Chonnam National University, 77 Yongbong-ro, Buk-gu, Gwangju 61186, Korea.

Tel. +82-62-530-1354 Fax. +82-62-530-1339 Email. parkcy@jnu.ac.kr

©2022 The Korean Nutrition Society and the Korean Society of Community Nutrition
This is an Open Access article distributed under the terms of the Creative Commons
Attribution Non-Commercial License (https://creativecommons.org/licenses/by-nc/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

#### **ORCID iDs**

Dahyun Park 🕞

https://orcid.org/0000-0002-9200-6754 Mi-Kyung Choi

https://orcid.org/0000-0001-6026-1374

# Higher food literacy scores are associated with healthier diet quality in children and adolescents: the development and validation of a two-dimensional food literacy measurement tool for children and adolescents

Dahyun Park (1) 1, Mi-Kyung Choi (1) 2, Yoo Kyoung Park (1) 3,4, Clara Yongjoo Park (1) 58, and Min-Jeong Shin (1) 1,68

Interdisciplinary Program in Precision Public Health, Graduate School, Korea University, Seoul 02841, Korea <sup>2</sup>Faculty of Food and Health Sciences, College of Natural Sciences, Keimyung University, Daegu 42601, Korea <sup>3</sup>Department of Medical Nutrition, Graduate School of East-West Medical Science, Kyung Hee University, Yongin 17104, Korea

<sup>4</sup>Department of Medical Nutrition, AgeTech-Service Convergence Major, Graduate School of East-West Medical Science, Kyung Hee University, Yongin 17104, Korea

<sup>5</sup>Department of Food and Nutrition, Chonnam National University, Gwangju 61186, Korea

<sup>6</sup>School of Biosystems and Biomedical Sciences, College of Health Science, Korea University, Seoul 02841, Korea

# **ABSTRACT**

**BACKGROUND/OBJECTIVES:** Most child and adolescent food literacy measurement tools focus on nutrition and food safety. However, the importance of aspects related to the food system such as food distribution and food waste and their effects on environmental sustainability is growing. We therefore developed and validated a two-dimensional tool for children (8–12 years old) and adolescents (13–18 years old) that can comprehensively measure food literacy. The association of food literacy with diet quality and self-reported health was assessed.

**SUBJECTS/METHODS:** First, we developed a food literacy conceptual framework that contains food system and literacy dimensions through a literature review, focus group interviews, and expert review. After a face validity study, we conducted the main survey (n = 200) to validate the questionnaire. Construct validity and reliability were assessed using exploratory factor analysis (EFA), confirmatory factor analysis (CFA), and Cronbach's alpha. **RESULTS:** As a result of the Delphi study, content validity was confirmed for the remaining 30 items after two items were excluded (content validity ratio = 0.86). Eleven items were excluded from the EFA results, while the CFA results indicated appropriate fit indices for the proposed model (comparative fit index = 0.904, root mean square error of approximation = 0.068). The final food literacy questionnaire consisted of 19 questions and comprised 5 factors: production, distribution, selection, preparation and cooking, and intake. Food literacy was positively associated with diet quality, as assessed by the Nutrition Quotient score, in both children and adolescents and with self-reported health in adolescents.



Yoo Kyoung Park https://orcid.org/0000-0002-8536-0835 Clara Yongjoo Park https://orcid.org/0000-0001-5868-7130

Min-Jeong Shin (D)

https://orcid.org/0000-0002-8952-4008

#### Funding

This work was carried out with the support of the Cooperative Research Program for Agriculture Science and Technology Development (Project No. PJ014709022021), Rural Development Administration, Republic of Korea. This work was also supported by the National Research Foundation of Korea (NRF) grant funded by the Ministry of Science and ICT (NRF-2020R1A2C2005580).

#### **Conflicts of Interest**

The authors declare no potential conflicts of interests.

#### **Author Contributions**

Conceptualization: Shin MJ; Data curation: Shin MJ, Choi MK, Park D, Park YK, Park CY; Formal analysis: Choi MK, Park D; Writing original draft: Park D, Park CY, Shin MJ. **CONCLUSIONS:** Based on this integrated conceptual framework, a two-dimensional 19-item food literacy measurement tool was developed and verified for practical use to improve the diet quality and food-related environmental sustainability awareness of children and adolescents.

**Keywords:** Literacy; health literacy; questionnaire; eating behavior; sustainable development; diet, food, and nutrition

# INTRODUCTION

Childhood and adolescence are periods of rapid physiological, sexual, neurological, and behavioral changes [1]. During these life stages, proper nutrition is important to achieve one's full growth potential [2]. However, in modern society, the high accessibility of energy-dense foods and reduced physical activity, due to increased screen time and sedentary activity, has prompted energy imbalances in children [3]. This has led to the 47% increase in overweight and obese children and adolescents during the past few decades worldwide [4]; the obesity rates of Korean children and adolescents show a similar pattern [5].

As children and adolescents gradually become more independent in lifestyle decision-making, including food selection and intake, their food literacy during this crucial period may impact lifelong eating patterns and behaviors. Food literacy is defined as "the scaffolding that empowers individuals, households, communities, or nations to protect diet quality through change and strengthen dietary resilience over time. It is composed of a collection of inter-related knowledge, skills and behaviors required to plan, manage, select, prepare, and eat food to meet needs and determine intake" [6]. Thus, enhancing food literacy during child development may benefit dietary behaviors by increasing the knowledge of healthy and sustainable diets and choice of appropriate foods. In addition, children and adolescents may be strongly affected by peers, parental modeling, mass media, and body image when selecting and consuming foods [7]. Therefore, the ability, skills, and behaviors that enable one to search for, comprehend, and apply food knowledge is essential. In adults, a positive association between food literacy and dietary behavior has been demonstrated [8], but this association has not been investigated in children and adolescents.

In addition to personal nutrition, emphasis on the impact of food selection and intake on the food system and environment is increasing. More than half of the 17 United Nations Sustainable Development Goals are related to food, including health, environment, food production and consumption patterns, and food security [9]. In line with this trend, recent research on food literacy also covers a broad spectrum—from individual health to the food system and its interactions with the society and environment [10,11]. Therefore, the term "food literacy" should incorporate complex decision-making concepts related to food [12]. The food system encompasses all processes related to food, including production, processing, distribution, consumption, and disposal, and operates within a social, political, economic, and natural environment [13]. By integrating the food system into the conceptual framework of food literacy, we may adequately address issues such as food safety and sustainable agriculture, which were limited in previous approaches [14].

Unfortunately, children and adolescents have few opportunities to be educated on the effects of food selection and intake on the society and environment as most school curricula do not focus on this aspect [15,16]. Moreover, most food literacy interventions in children and



adolescents have focused on food preparation and nutrition, but not on the food system [17]. Recently, we developed and validated a food literacy measurement tool for adults based on a new two-dimensional conceptual framework which integrates a food system dimension with the existing literacy measurement tools [18]. Using this 25-item measurement tool, we found that food literacy was positively associated with food knowledge and diet quality in adults [18]. However, no food literacy measurement tool that covers these comprehensive concepts exists for children and adolescents [10].

In the current study, we aimed to develop and validate a food literacy measurement tool based on a comprehensive conceptual framework applicable to children and adolescents. Using this measurement tool, we explored the associations of food literacy with diet quality and subjective health in Korean children and adolescents.

# SUBJECTS AND METHODS

The current questionnaire was developed and validated following the methods previously described [18]. Briefly, the conceptual framework, based on a literature review of food literacy in adults [6,12,19-22], was adapted to children and adolescents through the generation of new questions, a focus group interview, Delphi study, and face validity study (**Fig. 1**). A main survey was then performed, and the results were validated against Nutrition Quotient (NQ) scores and subjective health. All of the study processes were approved in advance by the Institutional Review Board of Korea University (KUIRB-2019-0306-01). All study participants voluntarily assented, and their legal guardians consented to their participation in the study.

# Development of a food literacy conceptual framework and items for children and adolescents

A detailed description of the development of the conceptual framework can be found elsewhere [18]. Briefly, the conceptual framework was developed based on a literature review on food literacy measurement tools with the addition of a food system dimension [18]. This two-dimensional framework consisted of a literacy dimension and a food system dimension. Initially, the literacy dimension had three domains (functional, interactive, and critical literacy) and the food system dimension had eight domains (production, processing, distribution, planning and management, selection, preparation and cooking, intake, and disposal) [18]. The framework was revised through focus group interviews that included 12 older adults (≥ 65 years of age), 12 younger adults (19–64 years of age), and 12 children and adolescents (8-18 years of age) who reflected a balance between urban and rural area residents. An item (question) pool (total of 562 items) was generated from the 547 items determined by the literature review and the 15 newly composed life-stage specific items collected from the results of the focus group interviews. For instance, for this study in children and adolescents, items such as "I search recipes through various media," "I usually willingly participate in preparing or cooking meals," and "I try different foods that I am not used to eating every day" were included. Questions were then excluded if the meaning of one question overlapped with another question's, if the question measured characteristics limited to a narrow age range or specific gender, or if multiple questions measured the same domain. In the latter case, the question easiest to understand and that measured the broadest range of content was selected. A 5-point Likert scale was used to measure food literacy (strongly disagree=1 point, disagree=2 points, neutral=3 points, agree=4 points, and strongly agree=5 points).






Fig. 1. The study design for the development and validation of a food literacy measurement tool.

# Questionnaire validation

We confirmed the content validity of questionnaire with a Delphi survey of 15 food and nutrition experts. Each item was evaluated on a four-point Likert scale to calculate the content validity ratio (CVR) [23,24]; a question was determined to be appropriate if its CVR was 0.49 or higher [23]. The questions evaluated as inappropriate were corrected according to the experts' suggestions, and the survey was repeatedly reviewed until all questions were valid.

To evaluate the face validity of the questionnaire, a pilot survey was conducted in ten children and adolescents aged 8–18 years. Participants were recruited through posters and online recruitment notices, i.e., convenience sampling. Interviews were performed to assess the clarity of the questions.

The main survey was conducted (online or offline) on 200 Korean children and adolescents aged 8-18 years old in August 2020. Participants were selected through convenience sampling. The appropriate sample size was determined according to statistical criteria that considered factor loading and the number of questions [25,26]. An exploratory factor analysis (EFA) using a maximum likelihood analysis with a direct oblimin rotation pattern matrix was conducted



to verify construct validity, thus determining the number and nature of the questionnaire's underlying factors. We measured sample appropriateness via the Kaiser–Meyer–Olkin (KMO) method and evaluated the factor analysis using Bartlett's test of sphericity and total variance explained. Using the EFA, each item was allocated to the factor with the highest factor loading. Items were deleted if they did not meet the inclusion criteria (i.e., factor loading < 0.4, communalities extraction < 0.4). Internal consistency and reliability for the total questionnaire and each factor were confirmed through Cronbach's alpha value (> 0.7) [27]. A confirmatory factor analysis (CFA) was conducted to test whether the data fit the hypothesized measurement model that was extracted by the EFA. The goodness of fit of the model was evaluated using various indices: the root mean square error of approximation (RMSEA), which tests the fit of the model to the covariance matrix, was considered to show an acceptable fit when the value was 0.05–0.08 and a good fit when < 0.05; the comparative fit index (CFI) was considered to demonstrate a good fit with a value of  $\geq$  0.9; and the relative  $\chi^2$  ( $\chi^2$ /degrees of freedom) with a value of 3 or less was considered as acceptable [28-32].

#### NQ score and assessment of subjective health

The survey included questions for NQ score and subjective health to evaluate dietary quality and overall health, respectively. The NQ score assesses overall diet quality and food behaviors [33,34]; higher NQ scores denote healthier diet quality and food behavior. Subjective health was measured with the question "What do you usually think about your health?" Participants answered on a 5-point scale: 1 = very poor, 2 = poor, 3 = fair, 4 = good, and 5 = very good.

Linear regression analyses were used to assess the association of food literacy scores with NQ scores and subjective health. Crude analyses (Model 1) and an adjustment for age and gender (Model 2) were performed. All analyses were conducted using the Statistical Package for Social Sciences for Windows, version 24.0 (IBM Corp, New York, NY, USA).

# **RESULTS**

## Questionnaire development

Of the 562 items in the item pool, 530 items were deleted according to the exclusion criteria, resulting in 32 candidate items. Based on the focus group interviews, the items in the planning and management domain were reclassified and the planning and management domain was deleted.

#### **Questionnaire validation**

The Delphi study was conducted three times to reach a consensus. Two items were deleted during this process, resulting in a total of 30 items. In addition, items in the processing domain were reassigned to the selection domain during the Delphi survey. Therefore, the remaining 30 items were classified into 18 areas (three food literacy domains × six food system domains). In the literacy dimension, there were seventeen, seven, and six items corresponding to the functional, interactive, and critical literacy domains, respectively. In the food system dimension, there were five, three, nine, five, five, and three items in the production, distribution, selection, preparation and cooking, intake, and disposal domains, respectively. The questionnaire's content validity was confirmed by CVR (mean CVR: 0.86).

The pilot study to assess the questionnaire's face validity found that most items were comprehensible for the general public and could adequately measure food literacy. Some



questions were rewritten to clarify their meaning, seven questions were edited with simpler or more specific vocabulary, and examples were added to four questions to increase understanding.

**Table 1** shows the characteristics of the 200 children and adolescents that participated in the main survey. The participants were equally organized by gender and age group. The validity of the questionnaire was confirmed through the EFA, using the maximum likelihood method (**Table 2**). Eleven items were excluded due to communalities or factor loading values less than 0.4. Five factors were derived from the 19 items included in the final tool (Factor 1: selection, Factor 2: production, Factor 3: distribution, Factor 4: preparation and cooking, and Factor 5:

Table 1. Main survey participants' characteristics

Variables	Total (n = 200)	Boys (n = 104)	Girls (n = 96)	<i>P</i> -value
Age (yrs)				0.20
8-12	23.5 (47)	14.0 (28)	9.5 (19)	
13-15	34.0 (68)	17.0 (34)	17.0 (34)	
16-18	47.5 (85)	21.0 (42)	21.5 (43)	
Residential area				0.37
Urban	95.5 (191)	94.2 (98)	96.9 (93)	
Rural	4.5 (9)	5.8 (6)	3.1 (3)	
Subjective health				0.28
Very poor	1.0 (2)	0.0 (0)	2.1 (2)	
Poor	10.0 (20)	8.7 (9)	11.5 (11)	
Fair	36.0 (72)	33.7 (35)	38.5 (37)	
Good	36.0 (72)	36.5 (38)	35.4 (34)	
Very good	17.0 (34)	21.2 (22)	12.5 (12)	
Nutrition quotient-score	$52.16 \pm 8.69$	$53.54 \pm 8.68$	$50.67 \pm 8.49$	0.02

Values are presented as mean  $\pm$  SD for continuous variables or numbers and percentages for categorical variables, % (No.).

 $\textbf{Table 2.} \ \ \textbf{Validity results from the exploratory factor analysis of the main survey (n=200)}$ 

Factor	ltem	Communalities	Factor loading				
			1	2	3	4	5
Selection	I can find or ask about the correct methods to recycle food packaging and food waste. 66	0.57	0.74	-0.08	0.05	0.08	0.04
	I can determine how well food hygiene is practiced by watching the meal preparation and cooking process. <sup>54</sup>	0.57	0.65	0.05	0.02	-0.19	0.06
	I can explain the pros and cons of my usual diet. <sup>cs</sup>	0.55	0.61	-0.03	-0.07	-0.07	-0.04
	I can determine whether a food is necessary for me by watching/reading food advertisements. <sup>bs</sup>	0.59	0.60	0.14	-0.17	-0.08	0.07
	I can talk about the pros and cons of Korean food culture. $^{\mbox{\tiny CS}}$	0.48	0.48	-0.08	-0.10	-0.09	0.09
	I can find or ask about various ways to judge the quality (taste, freshness, etc.) of food. c3	0.78	0.45	-0.12	-0.21	0.01	0.06
	I am aware of the environmental impact of food waste and take care when disposing of it. b6	0.42	0.41	-0.06	-0.12	-0.12	0.20
Production	I usually check for the food's country of origin. <sup>a1</sup>	0.44	-0.01	-0.78	0.03	0.07	0.02
	I usually check for the genetically modified organism label on food products. <sup>a1</sup>	0.43	0.08	-0.70	0.02	-0.10	0.00
	I usually check for agricultural food certification (organic, pesticide free, etc.) labels. <sup>a1</sup>	0.52	-0.12	-0.57	-0.24	-0.13	0.03
Distribution	I am aware of how different food production methods impact the environment and society. $^{\mbox{\tiny cl}}$	0.50	-0.04	0.02	-0.90	0.02	0.06
	I can find information on various food distribution methods, such as local food. $^{\text{b2}}$	0.55	0.18	-0.07	-0.65	-0.03	-0.11
	I am aware of how different food transportation methods impact the environment and society. $^{\!\!\!^{\rm C2}}$	0.49	0.10	-0.24	-0.41	-0.03	0.11
Preparation	I usually willingly participate in preparing or cooking meals.°4	0.56	-0.10	-0.05	-0.02	-0.66	0.15
and cooking	I search for recipes through various media. <sup>b4</sup>	0.55	0.07	-0.03	-0.04	-0.63	0.04
	I can cook according to the recipes shown on processed food packages. <sup>a4</sup>	0.46	0.30	-0.03	0.03	-0.60	-0.15
Intake	I usually try to eat a variety of food groups, including grains, fish, meat, vegetables, fruits, dairy, etc. <sup>a5</sup>	0.46	0.06	-0.06	-0.01	0.04	0.63
	I try different foods that I am not used to eating every day. <sup>a3</sup>	0.52	-0.06	0.03	-0.01	-0.13	0.63
	I try to get accurate information about food and health. a4	0.46	0.31	-0.08	-0.01	0.06	0.56
Eigenvalue		-	6.89	1.82	1.34	1.25	0.95
Cumulative %	% of variance	-	33.50	40.63	45.06	49.08	52.15

a,b,cDomain of literacy dimension (a: functional literacy, b: interactive literacy, and c: critical literacy); 1-6 Domain of food system dimension (l: production,

<sup>2:</sup> distribution, 3: selection, 4: preparation and cooking, 5: intake, and 6: disposal). Some questions may overlap with questions developed for adults [18].



Table 3. Regression analysis of food literacy score with NQ score and subjective health (n = 200)

	Model 1		Model 2		
	Coef. (95% CI)	<i>P</i> -value	Coef. (95% CI)	P-value	
NQ score					
Children (8-12 years)	0.428 (0.211-0.645)	< 0.001	0.431 (0.204-0.658)	< 0.001	
Adolescents (13-18 years)	0.200 (0.089-0.310)	< 0.001	0.230 (0.120-0.340)	< 0.001	
Adolescents (13-15 years)	0.191 (0.047-0.334)	0.010	0.206 (0.067-0.345)	0.004	
Adolescents (16-18 years)	0.229 (0.045-0.414)	0.016	0.280 (0.102-0.459)	0.003	
Subjective health					
Children (8-12 years)	0.014 (-0.009-0.037)	0.220	0.014 (-0.011-0.038)	0.261	
Adolescents (13-18 years)	0.022 (0.010-0.034)	< 0.001	0.024 (0.012-0.037)	< 0.001	
Adolescents (13-15 years)	0.023 (0.008-0.037)	0.002	0.023 (0.008-0.037)	0.002	
Adolescents (16-18 years)	0.024 (0.001-0.046)	0.044	0.028 (0.006-0.051)	0.014	

NQ, Nutrition Quotient; Coef., coefficient; CI, confidential interval. Model 1: crude model, Model 2: adjusted for age and gender.

intake). We confirmed that the KMO measure of sampling adequacy was appropriate (0.886) and that the Bartlett's test results were statistically significant ( $X^2 = 1527.630$ , df = 171, and P-value < 0.001). The Cronbach's alpha for each of the five factors were all appropriate, i.e., above 0.7: Factor 1 = 0.848, Factor 2 = 0.765, Factor 3 = 0.790, Factor 4 = 0.728, and Factor 5 = 0.707. The CFA results indicated a desirable fit of the proposed models, with the CFI = 0.904, RMSEA = 0.068, TLI = 0.888, AIC = 9945.865, and BIC = 10150.360. The mean ( $\pm$  SD) food literacy score of the final 19 items was 60.0 ( $\pm$  11.9), ranging from 19 to 88.

## Association of food literacy with NQ scores and subjective health

Food literacy scores were positively associated with diet quality as measured by NQ scores (**Table 3**). When food literacy scores increased by one unit, NQ scores increased by 0.428 and 0.200 in children and adolescents, respectively. That relationship remained positive after adjusting for age and gender. Subjective health was also positively associated with food literacy in adolescents, regardless of adjustment.

# **DISCUSSION**

We developed a food literacy measurement tool for children and adolescents based on a two-dimensional conceptual framework consisting of five factors and nineteen questions. This tool includes the concept of the food system and expands its scope to encompass sustainability of the environment and society. We also confirmed that food literacy is positively associated with NQ scores in both children and adolescents and with adolescents' subjective health status.

The current food literacy tool for children and adolescents integrated a food system domain which may promote food-related sustainability in addition to food safety and nutrition [35]. As food-related choices and intake manifested during youth can last a lifetime [36], enhancing food literacy from a young age should be a focus of effective food and nutrition education. Food literacy with respect to not only nutrition and food safety but also the society and environment is essential even for younger individuals. Previous studies show that environmental behavior [37], as well as its antecedents [38] such as ecological awareness [39,40] and environmental attitudes [41], begin to form and develop in early childhood. However, previous food literacy measuring instruments developed for children and adolescents did not incorporate domains related to the food system [42-45]. In Australian adolescents, four of the five top important aspects of food literacy chosen were related to food safety and nutrition [46], possibly due to



the limited focus on food-related environmental sustainability in the classroom [15]. When we assessed Korean children and adolescents' awareness of environmental sustainability during the development of the current food literacy measurement tool, we found that their awareness seemed to be higher than that of Korean adults. As a result, items related to disposal were included in the selection factor, and the separate disposal factor was omitted. In contrast, the disposal factor was retained in adults [18]. This indicates that Korean children and adolescents consider food disposal when selecting which foods to consume. In addition, the distribution factor was included in the tool for children and adolescents, while it was omitted in the one for adults. This unique factor in the tool for children and adolescents indicates that more variation may exist among the younger generation in items related to food distribution, which may impact food literacy scores. Moreover, the item "I am aware of how different food production methods impact the environment and society" in the distribution factor was excluded from the tool for adults due to the EFA results, but was retained in the tool for children and adolescents. The current food literacy measurement tool may be used to identify gaps in and convey the importance of food knowledge, beliefs, and behavior that impact the society and environment to children and adolescents.

Many items in the present tool for children and adolescents were newly developed or assigned to different factors from those in the adult tool, reflecting the unique circumstances of younger individuals. For instance, several questions that belonged to the preparation and cooking factor in adults corresponded to the selection factor in children and adolescents [18]. This reflects that the preparation and cooking of foods are mostly adult tasks, while children and adolescents mostly select from already prepared foods. This may be owing to not only children's limited cooking capabilities but also to limited opportunities and time constraints for preparing and cooking food due to school-related activities and/or a lack of confidence in food preparation [46]. Thus, the three questions pertaining to the preparation and cooking factor in the current tool differ from those in the tool for adults. These questions were included based on the results of the focus group interviews and Delphi study to reflect the characteristics of children and adolescents concerning cooking-related capacity and attitudes. On the other hand, items that pertained to the production and selection factors in adults comprised an independent factor (distribution) in children and adolescents. Finally, the child and adolescent measurement tool included the question "I try different foods that I am not used to eating every day" in the intake factor. This question highlights the willingness of children and adolescents to actively rather than passively consume a variety of foods. This amenability to trying unfamiliar foods can also influence the selection of foods when the environment is at stake. The present tool was tailored to children and adolescents to capture their unique life stage and to incorporate the food system.

Although cross-sectional, the positive relationship between NQ scores and food literacy scores demonstrates that an increase in food literacy may increase nutrition-related diet quality in children and adolescents. The NQ scores and self-reported health measures are valid measures of diet quality [33,34] and physical and emotional well-being [47], respectively, in children and adolescents. According to prior research on adults, improving food literacy is key to supporting healthy food selection and preparation and may lead to improvements in dietary quality and ultimately health outcomes [8]. Similarly, a few, but not all, observational studies have shown a positive relationship between food literacy and dietary intake in adolescents [48]. However, these studies did not include the concept of food system when assessing food literacy. In addition, only a few food literacy interventions have been implemented in children or adolescents, mostly targeting the cooking aspect of



food literacy [17]. Most of these interventions lacked a theoretical basis, which may be the reason for their low success rate in influencing positive dietary behaviors [17]. Childhood and especially adolescence can present unique challenges for developing and implementing interventions due to the time crunch created by increased school workloads and the greater influence of peer pressure at those ages [7]. On the other hand, self-reported health status indicators show greater stability during early and middle adolescence [49], which may explain why the reported positive association between subjective health and food literacy in adolescents was not found in children. In line with the positive association between food literacy and diet quality, increased food literacy may heighten environment-related behaviors in children and adolescents. Well-designed interventions to increase food literacy, with additional emphasis on the food system, in children and adolescents are required.

Despite its brevity, the current food literacy measurement tool, which incorporates East Asian culture, was validated across a wide age range of growing individuals. Previous food literacy assessment tools for children or adolescents were developed for narrow age ranges. For example, the preschool-Food Literacy Assessment Tool (FLAT) was developed for 3- to 6-year-old children in Italy [42], while the Tool for Food Literacy Assessment in Children (TFLAC) [43] and Food and Nutrition Literacy (FNLIT) [44] were developed for those aged 9-11 years in the U.S. and 10-12 years in Iran, respectively. Another tool was recently developed for Danish adolescents 12-14 years of age [45]. Because the present tool has been developed and validated for subjects within a wide age range (8-18 years old), it is possible to assess food literacy across childhood development and to compare differences by age groups. The positive association between food literacy status and NO score, which was developed for children and adolescents separately, indicates that this measurement tool can accurately convey food literacy for both life stages despite the differences between them. In addition, compared to the existing tools, which on average consist of 31 items, the present 19-item measurement tool is shorter, enabling a simple measurement of food literacy. In addition, food literacy may differ by geographical and cultural backgrounds while food systems may differ by country and region [42,44], yet no measurement tool has been developed for Korean or East Asian children or adolescents [48]. Thus, a strength of the current validated tool is that it may more easily capture food literacy status in both children and adolescents in Korea.

This study is not without limitations. The survey participants were restricted to Koreans. However, to broaden the applicability of this tool, we included individuals of both genders and various ages and regions in the study. Although the questionnaire was validated in subjects aged 8–18, we were not able to validate the questionnaire in more detailed age groups due to the small sample size. For wider use, the tool should be translated and validated in children and adolescents of other countries. However, this tool is meaningful since it is the first to set a framework to assess food literacy that includes the food system in children and adolescents. It should also be noted that this study used convenience sampling when recruiting survey participants. This food literacy measurement tool can be employed to conduct future research on practical education and policy to increase food literacy and their influence on food-related behaviors.

We developed a food literacy measurement tool for children and adolescents based on a systematic conceptual framework that includes the food system. Food literacy and NQ scores were positively associated in both children and adolescents. This tool provides an understanding of the food literacy status of both children and adolescents that can serve as



a basis for establishing effective strategies to solve various food-related problems and for achieving environmental sustainability associated with food-related behaviors.

## **ACKNOWLEDGMENTS**

We would like to thank the experts for their input and feedback in generating the items for the self-perceived food literacy scale.

#### REFERENCES

- Das JK, Salam RA, Thornburg KL, Prentice AM, Campisi S, Lassi ZS, Koletzko B, Bhutta ZA. Nutrition in adolescents: physiology, metabolism, and nutritional needs. Ann N Y Acad Sci 2017;1393:21-33.
   PUBMED | CROSSREF
- 2. Story M. Nutritional requirements during adolescence. In: McAnarney ER, Kreipe RE, Orr DE, Comerci GD, editors. Textbook of Adolescent Medicine. Philadelphia (PA): W.B. Saunders; 1992. p.75-84.
- Kremers SP, de Bruijn GJ, Visscher TL, van Mechelen W, de Vries NK, Brug J. Environmental influences on energy balance-related behaviors: a dual-process view. Int J Behav Nutr Phys Act 2006;3:9.
   PUBMED I CROSSREF
- 4. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, Mullany EC, Biryukov S, Abbafati C, Abera SF, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet 2014;384:766-81.

#### PUBMED | CROSSREF

- 5. Jung E. Korea Health Statistics 2019: Korea National Health and Nutrition Examination Survey (KNHANES VIII-1). Cheongju: Korea Disease Control and Prevention Agency; 2020.
- Vidgen HA, Gallegos D. Defining food literacy and its components. Appetite 2014;76:50-9.
   PUBMED | CROSSREF
- 7. Story M, Stang J. Understanding adolescent eating behaviors. In: Story M, Stang J, editors. Guidelines for Adolescent Nutrition Services. Minneapolis (MN): Center for Leadership, Education and Training in Maternal and Child Nutrition; 2005. p. 9-19.
- 8. Begley A, Paynter E, Butcher LM, Dhaliwal SS. Effectiveness of an adult food literacy program. Nutrients 2019;11:1-15.

#### PUBMED | CROSSREF

- 9. United Nations. THE 17 GOALS | Sustainable Development [Internet]. New York (NY): United Nations; 2021 [cited 2021 March 16]. Available from: https://sdgs.un.org/goals.
- Yuen EY, Thomson M, Gardiner H. Measuring nutrition and food literacy in adults: a systematic review and appraisal of existing measurement tools. Health Lit Res Pract 2018;2:e134-60.

  PUBMED | CROSSREF
- Widener P, Karides M. Food system literacy: empowering citizens and consumers beyond farm-to-fork pathways. Food Cult Soc 2014;17:665-87.
- 12. Cullen T, Hatch J, Martin W, Higgins JW, Sheppard R. Food literacy: definition and framework for action. Can J Diet Pract Res 2015;76:140-5.

#### PURMED I CROSSREE

- Farmland Information Center. A primer on community food systems: linking food, nutrition and agriculture – FIC. Northampton (MA): Farmland Information Center; 2019 [cited 2021 July 27]. Available from: https://farmlandinfo.org/publications/a-primer-on-community-food-systems-linking-foodnutrition-and-agriculture/.
- Hanh N. Sustainable food systems concept and framework. Rome: Food and Agriculture Organization of the United Nations; 2018.
- Ronto R, Ball L, Pendergast D, Harris ND. Food literacy at secondary schools in Australia. J Sch Health 2016;86:823-31.

PUBMED | CROSSREF



 Bailey CJ, Drummond MJ, Ward PR. Food literacy programmes in secondary schools: a systematic literature review and narrative synthesis of quantitative and qualitative evidence. Public Health Nutr 2019;22:2891-913.

#### PUBMED | CROSSREF

 Brooks N, Begley A. Adolescent food literacy programmes: a review of the literature. Nutr Diet 2014;71:158-71.

#### CROSSREF

- Park D, Park YK, Park CY, Choi MK, Shin MJ. Development of a comprehensive food literacy measurement tool integrating the food system and sustainability. Nutrients 2020;12:1-13.
- 19. Azevedo Perry E, Thomas H, Samra HR, Edmonstone S, Davidson L, Faulkner A, Petermann L, Manafò E, Kirkpatrick SI. Identifying attributes of food literacy: a scoping review. Public Health Nutr 2017;20:2406-15.
- Krause C, Sommerhalder K, Beer-Borst S, Abel T. Just a subtle difference? Findings from a systematic review on definitions of nutrition literacy and food literacy. Health Promot Int 2018;33:378-89.
   PUBMED | CROSSREF
- 21. Slater J, Falkenberg T, Rutherford J, Colatruglio S. Food literacy competencies: a conceptual framework for youth transitioning to adulthood. Int J Consum Stud 2018;42:547-56.
- Thomas H, Azevedo Perry E, Slack J, Samra HR, Manowiec E, Petermann L, Manafò E, Kirkpatrick SI. Complexities in Conceptualizing and measuring food literacy. J Acad Nutr Diet 2019;119:563-73.
   PUBMED | CROSSREF
- 23. Lawshe CH. A quantitative approach to content validity. Person Psychol 1975;28:563-75.
- 24. Lynn MR. Determination and quantification of content validity. Nurs Res 1986;35:382-5. PUBMED | CROSSREF
- Kyriazos TA. Applied psychometrics: sample size and sample power considerations in factor analysis (EFA, CFA) and SEM in general. Psychology (Irvine) 2018;9:2207-30.
- Guadagnoli E, Velicer WF. Relation of sample size to the stability of component patterns. Psychol Bull 1988;103:265-75.

#### PUBMED | CROSSREF

- 27. Nunally JC, Bernstein IH. Psychometric Theory. New York (NY): McGraw-Hill; 1978.
- Hu LT, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. Struct Equ Modeling 1999;6:1-55.

  CROSSREF
- Tabachnick BG, Fidell LS. Using Multivariate Statistics Title: Using Multivariate Statistics. London: Pearson: 2019.
- 30. Kline RB. Principles and Practice of Structural Equation Modeling. New York (NY): The Guilford Press; 2011.
- 31. Marsh HW, Hau KT, Wen Z. In search of golden rules: Comment on hypothesis-testing approaches to setting cutoff values for fit indexes and dangers in overgeneralizing Hu and Bentler's (1999) findings. Struct Equ Modeling 2004;11:320-41.

#### CROSSREF

 Bentler PM, Bonett DG. Significance tests and goodness of fit in the analysis of covariance structures. Psychol Bull 1980;88:588-606.

#### CROSSREF

33. Kim HY, Lee JS, Hwang JY, Kwon S, Chung HR, Kwak TK, Kang MH, Choi YS. Development of NQ-A, nutrition quotient for Korean adolescents, to assess dietary quality and food behavior. J Nutr Health 2017;50:142-57.

#### CROSSREF

34. Kim HY, Kwon S, Lee JS, Choi YS, Chung HR, Kwak TK, Park J, Kang MH. Development of a Nutrition Quotient (NQ) equation modeling for children and the evaluation of its construct validity. Korean J Nutr 2012;45:390.

#### CROSSREF

- Food and Agriculture Organization of the United Nations. Sustainable Food Systems. Concept and Framework. Rome: Food and Agriculture Organization of the United Nations; 2018. p.1-8.
- Cunningham SJ. Predicting obesity in young adulthood from childhood and parental obesity. Emerg Off Pediatr 1997;10:204.



37. Evans GW, Otto S, Kaiser FG. Childhood origins of young adult environmental behavior. Psychol Sci 2018;29:679-87.

#### PUBMED | CROSSREF

38. Corraliza JA, Collado S. Ecological awareness and children's environmental experience. Pap Psicol 2019;40:190-6.

#### CROSSREE

39. Ewert A, Place G, Sibthorp J. Early-life outdoor experiences and an individual's environmental attitudes. Leis Sci 2005;27:225-39.

#### CROSSREF

40. Otto S, Evans GW, Moon MJ, Kaiser FG. The development of children's environmental attitude and behavior. Glob Environ Change 2019;58:101947.

#### CDOSSDE

41. Jørgensen KA. Bringing the jellyfish home: environmental consciousness and 'sense of wonder' in young children's encounters with natural landscapes and places. Environ Educ Res 2016;22:1139-57.

42. Tabacchi G, Battaglia G, Messina G, Paoli A, Palma A, Bellafiore M. Validity and internal consistency of the preschool-flat, a new tool for the assessment of food literacy in young children from the training-to-health project. Int J Environ Res Public Health 2020;17:1-15.

PUBMED | CROSSREF

43. Amin SA, Lehnerd M, Cash SB, Economos CD, Sacheck JM. Development of a tool for food literacy assessment in children (TFLAC). J Nutr Educ Behav 2019;51:364-9.

#### PUBMED | CROSSREF

44. Doustmohammadian A, Omidvar N, Keshavarz-Mohammadi N, Abdollahi M, Amini M, Eini-Zinab H. Developing and validating a scale to measure Food and Nutrition Literacy (FNLIT) in elementary school children in Iran. PLoS One 2017;12:e0179196.

#### PUBMED | CROSSREF

45. Stjernqvist NW, Elsborg P, Ljungmann CK, Benn J, Bonde AH. Development and validation of a food literacy instrument for school children in a Danish context. Appetite 2021;156:104848.

 Ronto R, Ball L, Pendergast D, Harris N. Adolescents' perspectives on food literacy and its impact on their dietary behaviours. Appetite 2016;107:549-57.

#### PUBMED | CROSSREF

47. Vingilis ER, Wade TJ, Seeley JS. Predictors of adolescent self-rated health. Analysis of the National Population Health Survey. Can J Public Health 2002;93:193-7.

#### PUBMED | CROSSREF

48. Vaitkeviciute R, Ball LE, Harris N. The relationship between food literacy and dietary intake in adolescents: a systematic review. Public Health Nutr 2015;18:649-58.

#### PUBMED | CROSSREF

49. Fosse NE, Haas SA. Validity and stability of self-reported health among adolescents in a longitudinal, nationally representative survey. Pediatrics 2009;123:e496-501.

PUBMED | CROSSREF