



Predictors of Health Behaviors Among Undergraduate Students During the COVID-19 Pandemic: A Cross-Sectional Predictive Study

This article was published in the following Dove Press journal:
Journal of Multidisciplinary Healthcare

Wanich Suksatan ^{1,2}
Bovornpot Choompunuch³
Apinya Koontalay⁴
Vachira Posai⁵
Ali H Abusafia ⁶

¹Faculty of Nursing, HRH Princess Chulabhorn College of Medical Science, Chulabhorn Royal Academy, Bangkok, Thailand; ²Trudy Busch-Valentine School of Nursing, Saint Louis University, St. Louis, MO, USA; ³Faculty of Education, Maharakham University, Maharakham, Thailand; ⁴Faculty of Nursing, Mahidol University, Bangkok, Thailand; ⁵Nursing Department, Sunpasithiprasong Hospital, Ubon Ratchathani, Thailand; ⁶Nursing Program, USM Health Campus, Kelantan, Malaysia

Background: The COVID-19 pandemic has affected the health behaviors of university students. Thus, factors influencing students' health behaviors during the COVID-19 outbreak should be examined.

Purpose: To our knowledge, little is known about stress, adversity quotient, and health behaviors of university students in Thailand. Therefore, the aims of this study were to determine the relationships among socioeconomic factors, stress, and adversity quotient influencing university students' health behavior during the COVID-19 crisis.

Patients and Methods: We conducted a cross-sectional predictive study; it was included 416 undergraduate students ≥ 18 years of age at a university in Thailand. We used descriptive statistics, chi-squared test, and stepwise multiple linear regression for data analysis.

Results: The results indicated that most participants were women (71.90%), and the highest proportion of students were enrolled in College of Politics and Governance (25.24%). Most of the students had a high stress level ($M = 3.54$, $SD = 0.53$), high adversity quotient ($M = 3.77$, $SD = 0.63$), and good health behavior ($M = 3.06$, $SD = 0.53$). Adversity quotient, stress, and current faculty/college were significant predictors of health behaviors of undergrad students (total variance: 37.2%).

Conclusion: Total adversity quotient was the strongest predictor of health behavior, followed by stress and current faculty/colleges. On the basis of our results, we suggested that faculty members, caretakers, and interdisciplinary care teams should consider adversity quotient and stress in developing activities to encourage and promote students' physical and mental health behaviors, particularly during the COVID-19.

Keywords: COVID-19 pandemic, Thai students, stress, adversity quotient, health behaviors

Introduction

In December 2019, the first cases of coronavirus disease 2019 (COVID-19) were found in the city of Wuhan, Hubei Province, China.¹ The pneumonia-causing virus was identified as the new coronavirus named as severe acute respiratory syndrome (SARS-CoV) 2,² and the disease was named by the World Health Organization as coronavirus disease 2019 (COVID-19).³ The virus causes the same symptoms as those with respiratory infections, ranging from mild to severe stuffy nose, sore throat, cough, and fever.⁴ Some people have pneumonia or difficulty breathing symptoms to diseases that cause serious illnesses, eg, Middle East Respiratory Syndrome (MERS-CoV) and SARS-CoV, which causes respiratory illnesses in people as well.⁵ If the symptoms are very severe, death may result.⁶ In clinical

Correspondence: Bovornpot Choompunuch
Faculty of Education, Maharakham University, Maharakham, Thailand
Tel +66 915575640
Email bovrnpot.c@msu.ac.th

epidemiology, laboratory characteristics, and radiographic characteristics, cells likely to be destruction of cells in the lower respiratory tract, resulting in respiratory failure.⁷

As of February 8, 2021, there were 106,965,292 cases of COVID-19 and 2,334,952 deaths related to it.⁸ In Thailand, 23,557 cases of COVID-19 have been confirmed, with 17,410 cured and 79 deaths.⁸ Seventy-nine confirmed deaths between 1 month and 97 years old. In terms of gender, male to female ratio 1.13:1, Thai nationality were 4,597, unknown nationals (597 cases), Myanmar (125 cases), India (77 cases), Cambodia (73 cases), and others (53 cases) (Emergency Operation Center Department of Disease Control, 2020). Since December 25, 2020, COVID-19 re-emerged in Thailand, and the tendency of patients to spread in each province is increasing in all age groups, including university students.⁹

In many countries, including Thailand, policies have been established to prevent the spread, control, and resolve COVID-19 outbreaks, such as strict city-wide lockdown or closure of areas that are home to large numbers of people, which prevents epidemics from spreading,¹⁰ and ban on foreign tourists from entering the country. These measures have considerably influenced people's lives, including changing their health habits, lifestyle, and slowing down the economy.¹⁰ It also affects the teaching and learning of students who must adapt to government policies, including wearing masks; staying away from people with respiratory symptoms (eg, cold symptoms, cough, and sneezing); washing their hands thoroughly with soap and water or alcohol hand sanitizer; avoiding unnecessary contact with eyes, nose, and mouth; avoiding sharing belongings with others; eating hot cooked food; and avoiding crowded places.⁹ If this is inevitable, students must protect themselves by wearing a surgical mask.

The university has also adopted new classroom teaching (eg, online learning) that suits the learners' context and the circumstances that arise during this time.¹¹ University students specifically report more significant mental health problems, including stress levels, compared with individuals who are not studying.¹² Because of the closure of universities or schools, teachers, professors, and students have moved to online distance learning, which may cultivate sensations of separation or disconnection from their schools, affecting students' motivation to learn. However, posttraumatic development perceived from the outbreak might be a protective factor against stress.¹³

A literature review on health behavior in the COVID-19 situation indicated that few studies were conducted on the leading factors of student health behavior in Thailand. Studies have been performed on the elderly, the general public, and studies of factors related to the COVID-19 pandemic.

However, university students are essential people in the country's future development. Therefore, our results may guide the development of models, management, and promotion of health behavior during COVID-19 crisis in university students, to be in good health and readiness, to learn about the new environment in the age of change.

Patients and Methods

Research Design and Sampling

This cross-sectional predictive study recruited undergraduate students ≥ 18 years of age at a Thai university enrolled in the academic year 2020. Total seven faculty/college were included: (1) Faculty of Education, (2) Faculty Architecture, Urban Design, and Creative Arts, (3) College of Politics and Governance, (4) Faculty of Nursing, (5) Faculty of Informatics, (6) Faculty of Public Health, and (7) Faculty of Tourism and Hotel Management. To calculate the sample group, we used the power analysis using the G * power program.¹⁴ Linear multiple regression analysis revealed the medium effect size was 0.15. Considering alpha of 0.05, power of the test as 0.80, and number of predictive variables was seven, a sample size of 416 participants was considered necessary. The sample was randomly selected from the target population using a convenience sampling technique with proportional allocation.

Research Instrument

Coronavirus Stress for Undergraduate Students (CSUS)

In the present research, we developed the CSUSS based on the literature review, which consisted of 15 items. Participants identify their choices on a 5-point scale from "least serious" (1 point) to "most serious" (5 points); total scores can range from 15 to 75. The interpretation by mean was five levels, which is using the following categories: 1.00–1.49 had the least stress, 1.50–2.49, had a low level of stress, 2.50–3.49 had moderate stress, 3.50–4.49 had a high level of stress, and 4.50–5.00 had the highest level of stress. Cronbach's alpha coefficient was 0.86.

Coronavirus Adversity Quotient

The researchers created the Coronavirus Adversity Quotient (CAQ) based on the literature review, which consisted of 17 items. Participants indicated their choices on a 5-point scale from "least serious" (1 point) to "most serious" (5 points), with total scores ranging from 17 to 85. On the basis of the scores, the adversity quotient was interpreted as follows: 1.00–1.49 indicated extremely low adversity quotient; 1.50–2.49, low

adversity quotient; 2.50–3.49, moderate adversity quotient; 3.50–4.49, high adversity quotient; and 4.50–5.00, extremely high adversity quotient. Cronbach's alpha coefficient was 0.93.

Health Behaviors Level

We used the Health Behaviors Scale (HBS) by the Health Education Division,¹⁵ which comprises 18 items. Participants rated their choices on a 5-point scale from “least serious” (1 point) to “most serious” (5 points), with total scores ranging from 18 to 90. The interpretation by mean was two levels, which is using the following categories: 1.00–2.99 represented a poor level of health behavior, and 3.00–5.00 represented a good level of health behavior. Cronbach's alpha coefficient was 0.82.

Sociodemographic Variables

Socioeconomic characteristics consisted of four items with multiple choices and open-ended questions. The researchers developed this tool based on literature reviews, including sex, academic year, current faculty/college, and family/household income.

Data Collection

We collected information between November 27, 2020, and December 15, 2020, as follows: We sent a letter to the deans of each faculty/college requesting to collect research data. After their and Institutional Review Board (IRB) approval, we met with participants. We explained the study objective and procedures, including sample rights protection. Signed informed consent was obtained from all participants willing to participate. Participants spent approximately 10–15 min completing the self-reported questionnaire. After participants completed the questionnaire, we checked the completeness of the questionnaire information. If we found an incomplete questionnaire, we informed the participants to complete the questionnaire. Participants were free to withdraw anytime, and confidentiality of the participants was ensured. The data sets of the current study are not publicly available due to the information that might be compromised the research participants' privacy.

Ethical Considerations

This research study was approved by the IRB of Mahasarakham University (Research Project Code No. 279/2020), dated 26 November 2020. This study was conducted in accordance with the Declaration of Helsinki.

Statistical Analysis

All data were analyzed using IBM SPSS version 21 (SPSS, Chicago, IL). Characteristics of the students were described using descriptive statistics, including frequency (percentage) or mean \pm standard deviation. The stress, adversity quotient, and health behavior scores were not normally distributed as it has been tested using Kolmogorov–Smirnov test, while the relationship between those factors was tested using non-parametric Chi-square test and Fisher's exact test. The factors predicting those variables were tested using the stepwise multiple linear regression analysis. Statistical significance was set at <0.05 .

Results

Of the 416 participants, most were female ($n = 299$, 71.90%) and freshmen (43.27%). Furthermore, the highest proportion of the participants were studying at the College of Politics and Governance (25.24%). Average monthly income 330–350 USD (33.41%) (Table 1).

Table 1 Sociodemographic Status of the Participants

Demographic Characteristics		Frequency (N = 416)	Percent (%)
Sex	Male	117	28.13
	Female	299	71.87
Academic Year	1st	180	43.27
	2nd	87	20.91
	3rd	64	15.39
	4th	76	18.27
	5th	9	2.16
Faculty/College	College of Politics and Governance	105	25.24
	Faculty of Tourism and Hotel Management	84	20.19
	Faculty of Education	79	18.99
	Faculty of Informatics	75	18.03
	Faculty Architecture, Urban Design, and Creative Arts	38	9.14
	Faculty of Public Health	25	6.01
	Faculty of Nursing	10	2.40
Monthly Income (US dollar)	< 330	68	16.35
	330–350	139	33.41
	350–660	79	18.99
	660–830	29	6.97
	830–1,000	31	7.45
	> 1,000	70	16.83




Figure 1 Distribution of stress and adversity quotient among students.

As shown in Figure 1, the overall stress of the students was at a high level ($M = 3.54, SD = 0.53$) when considered by sex, male students ($M = 3.49, SD = 0.51$) and female students ($M = 3.59, SD = 0.54$) had a high stress level. Moreover, the overall of the students was at a high level ($M = 3.77, SD = 0.63$) when considered by sex, male students ($M = 3.71, SD = 0.56$) had high adversity quotient, compared with female students ($M = 3.79, SD = 0.60$). In addition, the overall health behavior of students had good health behavior ($M = 3.06, SD = 0.53$), when considered by sex male students ($M = 3.38, SD = 0.35$) and female students ($M = 3.37, SD = 0.32$) had good health behavior (Figure 2).

The current study asserted the relationship among the participants' socioeconomic characteristics, such as sex, academic year, current faculty/college, and monthly income. Of them, only current faculty/college was significantly correlated with their health behaviors ($p < 0.05$) (Table 2).

We found that current faculty/college, stress, and adversity quotient by using stepwise multiple regression analysis. Three predictive variables ordered adversity quotient ($\beta = 0.52$), stress ($\beta = -0.32$), and current faculty/college ($\beta = 0.14$). The prediction model was statistically significant, $F(3, 416) = 11.74, p < 0.001$, accounting for approximately 37.2% of the variance of health behaviors ($R^2 = 0.610, \text{adjusted } R^2 = 0.372$) (Table 3).




Figure 2 Distribution of health behavior among students.

Table 2 Association Between Demographic Data and Health Behaviors

Demographic Data	N (%)	Health Behavior		Chi-Square	df	p value
		Poor n (%)	Good n (%)			
Sex						
Male	117 (28.13)	40 (9.62)	77 (18.51)	0.874	1	0.350 ^a
Female	299 (71.87)	117 (28.12)	182 (43.85)			
Academic Year				4.092	4	0.394 ^b
1st	180 (43.27)	72 (17.31)	108 (25.96)			
2nd	87 (20.91)	29 (6.97)	58 (13.94)			
3rd	64 (15.39)	19 (4.57)	45 (10.82)			
4th	76 (18.27)	33 (7.93)	43 (10.34)			
5th	9 (2.16)	4 (0.96)	5 (1.20)			
Faculty/College				19.396	6	0.004 ^b
College of Politics and Governance	105 (25.24)	30 (7.21)	75 (18.03)			
Faculty of Tourism and Hotel Management	84 (20.19)	34 (8.17)	50 (12.02)			
Faculty of Education	79 (18.99)	44 (10.58)	35 (8.41)			
Faculty of Informatics	75 (18.03)	25 (6.01)	50 (12.02)			
Faculty Architecture, Urban Design, and Creative Arts	38 (9.14)	16 (3.85)	22 (5.29)			
Faculty of Public Health	25 (6.01)	5 (1.20)	20 (4.81)			
Faculty of Nursing	10 (2.40)	3 (0.72)	7 (1.68)			
Monthly Income (US dollar)				7.125	5	0.211 ^a
< 330	68 (16.35)	31 (7.45)	37 (8.90)			
330–350	139 (33.41)	50 (12.02)	89 (21.39)			
350–660	79 (18.99)	33 (7.93)	46 (11.06)			
660–830	29 (6.97)	10 (2.40)	19 (4.57)			
830–1,000	31 (7.45)	6 (1.44)	25 (6.01)			
> 1,000	70 (16.83)	27 (6.49)	43 (10.34)			

Notes: ^aChi-squared test; ^bFisher's exact test; significance level $p < 0.05$.

Table 3 Regression of Individual Health Behaviors

Factors	B	SE(b)	β	t	p value	95% CI
Adversity quotient	0.44	0.03	0.52	13.270	<0.001	0.37, 0.50
Stress	-0.30	0.04	-0.30	-7.723	<0.001	-0.38, -0.22
Faculty Sci and health sci vs human ^{*a}	0.43	0.01	0.14	3.426	<0.001	0.02, 0.07
Constant 23.40						

Notes: ^aScience and health science school group vs humanities and social science school group; ^{*}Reference group; $R = 0.610$, $R^2 = 0.372$, $SE = 4.19$, $F = 11.74$, p value < 0.001.

Discussion

In this study, we identified that most of the students had a high stress level, high adversity quotient, and good health behavior. Our results indicated that only current faculty/college was significantly associated with their health behaviors ($p < 0.05$). We also determined factors

predicting health behavior in undergraduate students during the COVID-19 pandemic and found them to be adversity quotient, stress, and current faculty/college that could predict health behavior in undergraduate students.

Our findings demonstrated the first predicted health behavior in undergraduate students was adversity quotient.

Hulaikah et al¹⁶ reported that the ability to face problems and overcome obstacles is a person's ability to respond to events during times of adversity. Endure problems or obstacles when obstacles arise in life and respond to the environment.¹⁷ Moreover, the ability to face and overcome human obstacles is directly related to their physical and mental health because mental strength and self-control affect the body's immune resistance and the individual's self-care behaviors.^{18,19} Kurniawan et al²⁰ explained that facing challenges and overcoming obstacles is indicated by a person's ability to get up and act. Whether it is positive or negative for various situations, it depends on the individual whether there is more or less.²⁰ Phoolka et al²¹ also stated that healthy behaviors and adversity quotient, which are individuals, groups, organizations' actions, and the relational factors and consequences of these actions. For instance, social change policy development and skill improvement actions coping and improving quality of life.²¹

The second factor that predicted health behavior in students was stress. The COVID-19 pandemic has caused tremendous stress and anxiety in people, including students^{22,23} because of various reasons, including lockdown and fear of infection.²⁴ Most people, including students, experience stress, fear, and anxiety when facing emerging diseases and feel overwhelmed.²⁵ Government policies such as maintaining social distance and lockdown can make people feel alone, stressed, and anxious, as well as affect their physical and mental health. It is essential for individuals to prepare themselves to protect against COVID-19.¹³ Our results reveal that both male and female undergraduate students had severe stress (15.87% and 44.47%, respectively). Implementing health behaviors to protect oneself against COVID-19 can reduce students' stress related to the disease.²⁶ Li et al²⁷ examined the relationship between health behaviors and perceived stress in Chinese residents; their results revealed that health behaviors were negatively associated with perceived stress, with 39.89% of residents at risk of stress.²⁷

Additionally, a factor that can predict health behavior in undergraduate students of this study was current faculty/college. Students from science and health science schools had better health behaviors than those from humanities and social science schools. Students from health sciences schools, such as nursing student, may have higher knowledge and awareness of the COVID-19 outbreak and the

relevant self-care behaviors, partly because they may be taught that in their curriculums, such as good self-management of physical and mental health.²⁸ In line with Raingruber²⁹ who found that highly educated individuals practice more health-promoting behaviors than those with low literacy levels, this finding is consistent with Gallè et al,³⁰ who found understanding knowledge and behaviors associated with the COVID-19 outbreak in Italian undergraduate students. A previous study also indicated that students in non-health-related fields had a lower level of knowledge compared with students in health sciences.³⁰ However, the study of Alzoubi et al,³¹ who indicated the inexistence of differences in COVID-19 knowledge between medical and nonmedical colleges. They found that the knowledge of medical and nonmedical students about social media, the Internet, and television had no significant difference ($p > 0.05$). Therefore, undergraduate students still required more information and directed measures, good welfare, and awareness campaigns from government and related agencies to improve the knowledge, attitude, and skills in some critical components to contain the COVID-19 crisis.

For this study's implications, the results indicated that the adversity quotient, stress (negative direction) and current faculty/college are predictors of health behaviors in university students during the COVID-19 crisis. Therefore, when planning to promote and support students' healthy behaviors and impact these patients' psychological status. Health care providers should not overlook the students' mental health; otherwise, they may not achieve their study plan's goals. Additionally, future studies that address interventions to improve adversity quotient and decrease stress in university students during the COVID-19 pandemic are recommended, particularly for students in humanities and social science schools.

This study has some limitations. First, we included only undergraduate students from seven faculties/schools of one university; thus, our results may not be generalizable to the entire student population. Moreover, our cohort had a female predominance (71.87%). Future studies should therefore include a more diverse population, including different sexes, cultures, faculties, and universities to better understand better how COVID-19 affects them. Second, we predicted socioeconomic factors, stress, and adversity quotient influencing university students' health behavior during the COVID-19 pandemic with only a self-reported questionnaire. Longitudinal or intervention studies are warranted to clarify students' long-term outcomes. Third, our study

assessed socio-demographic characteristics such as sex, academic year, current faculty/college, and family/household income; it would be more informative if the residence condition, and the level of social support could be included, which a significant effect on the students' level of stress experience. Finally, our study used convenience sampling; the lack of random sampling may contribute to sample selection bias and also limit the findings' generalization. Future studies should examine students with other schools and also characteristics of people who are not students across time to identify how the behaviors and perspectives in those populations change as the COVID-19 outbreak shifts in scope.

Conclusion

Our study including a cohort of students from a Thai university revealed that the COVID-19 outbreak has a significant negative impact on students' psychological health, particularly on adversity quotient, stress, and health behaviors. The most critical thing for instructors, professors, and students is preparation—both psychically and psychologically—to cope with the new normal of education that changes the way we all learn, including educational management that is no longer the same.

Abbreviations

COVID-19, nCoV, novel coronavirus; coronavirus disease 2019; MERS-CoV, Middle East Respiratory Syndrome; SARS-CoV, Severe acute respiratory syndrome; CSUS, Coronavirus Stress for Undergraduate Students; CAQ, Coronavirus Adversity Quotient; HBS, Health Behaviors Scale.

Data Sharing Statement

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

We would like to thank the participants in our online survey for their valuable data and the reviewers for their helpful comments and suggestions to improve our study.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to

submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

No funding was received.

Disclosure

The authors declare that they have no conflicts of interest for this work.

References

- Hou W, Zhang W, Jin R, Liang L, Xu B, Hu Z. Risk factors for disease progression in hospitalized patients with COVID-19: a retrospective cohort study. *Infect Dis.* 2020;52(7):498–505. doi:10.1080/23744235.2020.1759817
- Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Eng J Med.* 2020;382(8):727–733. doi:10.1056/NEJMoa2001017
- World Health Organization [WHO]. Naming the coronavirus disease (COVID-19) and the virus that causes it. 2021. Available from: [https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-\(covid-2019\)-and-the-virus-that-causes-it](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it). Accessed February 8, 2021.
- Mei X, Zhang Y, Zhu H, et al. Observations about symptomatic and asymptomatic infections of 494 patients with COVID-19 in Shanghai, China. *Am J Infect Control.* 2020;48(9):1045–1050. doi:10.1016/j.ajic.2020.06.221
- Hui DS, Azhar EI, Memish ZA, Zumla A. Human coronavirus infections—severe acute respiratory syndrome (SARS), Middle east respiratory syndrome (MERS), and SARS-CoV-2. *Ref Module Biomed Sci.* 2020. doi:10.1016/B978-0-12-801238-3.11634-4
- Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis.* 2020;20(6):669–677. doi:10.1016/S1473-3099(20)30243-7
- WHO. Pneumonia of unknown cause – China. 2020. Available from: <https://www.who.int/csr/don/05-january-2020-pneumonia-of-unknown-cause-china/en/>. Accessed February 8, 2021.
- Worldometers. COVID-19 Coronavirus pandemic. 2021. Available from: <https://www.worldometers.info/coronavirus/>. Accessed February 8, 2021.
- Choompunuch B, Suksatan W, Sonsroem J, Kutawan S, In-udom A. Stress, adversity quotient, and health behaviors of undergraduate students in a Thai university during COVID-19 outbreak. *BNJ.* 2021;7(1):1–7. doi:10.33546/bnj.1276
- Dechsupa S, Assawakosri S, Phakham S, Honsawek S. Positive impact of lockdown on COVID-19 outbreak in Thailand. *Travel Med Infect Dis.* 2020;36:101802. doi:10.1016/j.tmaid.2020.101802
- Vlachopoulos D. COVID-19: threat or opportunity for online education? *High Learn Res Commun.* 2020;10(1):2. doi:10.18870/hlrc.v10i1.1179
- Savitsky B, Findling Y, Erel A, Hendel T. Anxiety and coping strategies among nursing students during the COVID-19 pandemic. *Nurse Educ Pract.* 2020;46:102809. doi:10.1016/j.nepr.2020.102809
- Wang C, Zhao H. The impact of COVID-19 on anxiety in Chinese university students. *Front Psychol.* 2020;11: 1168. doi:10.3389/fpsyg.2020.01168
- Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods.* 2007;39(2):175–191. doi:10.3758/BF03193146

15. Health Education Division; Health Service Support Department. Operational guidelines for changing health behavior at provincial level. Thailand: Health Education Division, Health Service Support Department; 2013.
16. Hulaikah M, Degeng I. The effect of experiential learning and adversity quotient on problem solving ability. *Int J Instr.* 2020;13(1):869–884. doi:10.29333/iji.2020.13156a
17. Tian Y, Fan X. Adversity quotients, environmental variables and career adaptability in student nurses. *J Vocat Behav.* 2014;85(3):251–257. doi:10.1016/j.jvb.2014.07.006
18. Siphai S. Influences of moral, emotional and adversity quotient on good citizenship of Rajabhat University's Students in the Northeast of Thailand. *Educ Res Rev.* 2015;10(17):2413–2421. doi:10.5897/ERR2015.2212
19. Tumka K, Kongdang K, Suksatan W. Mental health and family relationship of nursing students after the practice in nursing. *J Nurs Health.* 2019;1(2):111–124.
20. Kurniawan W, Daharnis D, Karneli Y. Contribution of adversity quotient, self awareness and demographic factors to student career maturity. *Int J Res Couns Educ.* 2020;4(1):70–75. doi:10.24036/00261za0002
21. Phoolka ES, Kaur N. Adversity quotient: a new paradigm to explore. *Contemporary Business Studies.* 2012;3(4):67–78.
22. Adnan M, Anwar K. Online learning amid the COVID-19 pandemic: students' perspectives. *Online Submission.* 2020;2(1):45–51.
23. Li HY, Cao H, Leung DY, Mak YW. The psychological impacts of a COVID-19 outbreak on college students in China: a longitudinal study. *Int J Environ Res Public Health.* 2020;17(11):3933. doi:10.3390/ijerph17113933
24. Mahato S, Pal S, Ghosh KG. Effect of lockdown amid COVID-19 pandemic on air quality of the megacity Delhi, India. *Sci Total Environ.* 2020;730:139086. doi:10.1016/j.scitotenv.2020.139086
25. Meeter M, Bele T, den Hartogh C, Bakker T, de Vries RE, Plak S. College students' motivation and study results after COVID-19 stay-at-home orders. 2020. doi:10.31234/osf.io/kn6v9
26. Al-Rabiaah A, Temsah M-H, Al-Eyadhy AA, et al. Middle east respiratory syndrome-corona virus (MERS-CoV) associated stress among medical students at a university teaching hospital in Saudi Arabia. *J Infect Public Health.* 2020;13(5):687–691. doi:10.1016/j.jiph.2020.01.005
27. Li Y, Yao L, Luo Y, Yuan F, Yan L. Perceived stress and its impact on health behavior of Chinese residents during the epidemic of COVID-19: an internet survey. 2020. doi:10.21203/rs.3.rs-27180/v1
28. Suksatan W, Ruamsook T, Prabsangob K. Factors influencing health literacy of students in health science curriculum: a cross-sectional study. *Indian J Public Health Res Dev.* 2020;11(7):1469–1474. doi:10.37506/ijphrd.v11i7.10303
29. Raingruber B. *Contemporary Health Promotion in Nursing Practice.* Jones & Bartlett Publishers; 2014.
30. Gallè F, Sabella EA, Da Molin G, et al. Understanding knowledge and behaviors related to COVID–19 epidemic in Italian undergraduate students: the EPICO study. *Int J Environ Res Public Health.* 2020;17(10):3481. doi:10.3390/ijerph17103481
31. Alzoubi H, Alnawaiseh N, Al-Mnayyis A, Lubad M, Aqel A, Al-Shagahin H. COVID-19-knowledge, attitude and practice among medical and non-medical University Students in Jordan. *J Pure Appl Microbiol.* 2020;14(1):17–24. doi:10.22207/JPAM.14.1.04

Journal of Multidisciplinary Healthcare

Dovepress

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal

covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>