

Time trends in incidence, treatment, and outcome in acute myocardial infarction in Norway 2013–19

Jarle Jortveit ^{1,*}, Are Hugo Pripp ², Jørund Langørgen ³,
and Sigrun Halvorsen ⁴

¹Sørlandet Hospital, Arendal, Box 416, Lundsiden, 4604 Kristiansand, Norway; ²Oslo Centre of Biostatistics and Epidemiology, Oslo University Hospital, Oslo, Norway; ³Haukeland University Hospital, Bergen, Norway; and ⁴Oslo University Hospital Ullevaal and University of Oslo, Oslo, Norway

Received 1 May 2022; revised 26 June 2022; online publish-ahead-of-print 10 August 2022

Handling Editor: Maciej Banach

Aims

Acute myocardial infarction (AMI) is a common cause of morbidity and mortality. The aim of the present study was to assess time trends in the incidence, treatment, and outcome of AMI in a nationwide registry-based cohort of patients.

Methods and results

All patients with a first AMI registered in the Norwegian Myocardial Infarction Registry between 2013 and 2019 were included in this cohort study. The number of patients admitted to Norwegian hospitals with a first AMI decreased from 8933 in 2013 to 8383 in 2019. The proportion of patients with ST-elevation myocardial infarction (STEMI) was stable at 30% throughout the period, and the percentage of STEMI undergoing coronary angiography was stable at 87%. The proportion of patients with non-STEMI undergoing coronary angiography increased by 2.4% per year (95% confidence interval 1.6–3.3) from 58% in 2013 to 68% in 2019. More patients were discharged with secondary preventive medication at the end of study period. Age-adjusted 1-year mortality was reduced from 16.4% in 2013 to 15.1% in 2018. The changes over time were primarily seen in the oldest patient groups.

Conclusion

In the period 2013–19 in Norway, we found a reduction in hospitalizations due to a first AMI. Both the percentage of patients undergoing coronary angiography as well as the percentage discharged with recommended secondary preventive therapy increased during the period, and the age-adjusted 1-year mortality after AMI decreased. A national AMI register provides important information about trends in incidence, treatment, and outcome, and may improve adherence to guideline recommendations.

* Corresponding author. Tel: +47 37 07 57 43, Email: jarle.jortveit@sshf.no

© The Author(s) 2022. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Graphical Abstract





Time trends in incidence, treatment, and outcome of acute myocardial infarction, Norway 2013–2019

AIM

To assess time trends in the incidence, treatment, and outcome of acute myocardial infarction (AMI) in a nationwide registry-based cohort of patients

Median age:
69 years (2013) –
70 years (2019)

Daily smokers:
31% (2013) –
26% (2019)



CONCLUSION

In the time period 2013 to 2019, we found a reduction in hospitalizations for a first AMI in Norway. The percentage of patients being treated according to guidelines increased, and the age-adjusted one-year mortality after AMI decreased. A national AMI register provides important information about trends in incidence, treatment, and outcome, and may improve adherence to guideline recommendations.

Keywords

Myocardial infarction • Time trends • Incidence • Outcome

Introduction

Cardiovascular disease is the most common cause of death in most European countries.¹ Acute myocardial infarction (AMI) is a common and serious presentation of cardiovascular disease and is associated with a high rate of mortality.^{2,3}

The Norwegian Myocardial Infarction Registry (NORMI) was established as a national medical quality register in 2013 and is among the few national registries in Europe with ongoing data registration and high case coverage.^{4,5} The register enables monitoring of incidence, treatment, and outcome after AMI. Through annual reports and scientific publications, the register has highlighted adherence to guideline recommendations for treatment of AMI.^{6–11}

The European recommendations for the treatment of AMI have been updated several times during the study period, but the main recommendations for invasive assessment and secondary preventive treatment after AMI have remained unchanged.^{12–15}

The aim of the present nationwide cohort study was to investigate time trends in hospital admissions, treatment, and outcome in patients with a first AMI in Norway in the period 2013–19.

Methods

The Norwegian Myocardial Infarction Registry

The NORMI, a part of the Norwegian Cardiovascular Disease Registry, is a national quality register. Registration into NORMI is mandatory

without requiring patient's consent. For definition of AMI, the NORMI adhered to the Third and Fourth Universal Definition of Myocardial Infarction during the study period.^{16,17} NORMI contains information on gender, age, cardiovascular risk factors, previous diseases and medication, symptoms and clinical findings, electrocardiogram (rhythm and ischaemic changes), in-hospital therapy and complications including death, as well as drugs prescribed at hospital discharge. The registration and quality of the information in the register have been described previously.^{18,19} Causes of death were obtained from the Norwegian Cause of Death Register.

Study population

All patients admitted to hospitals in Norway with a first AMI between 1 January 2013 and 31 December 2019 and registered in the NORMI were included in this cohort study.

Outcomes and follow up

The outcomes of the study were the percentage of patients receiving coronary angiography during hospitalization, the prescription rate of secondary preventive therapy at hospital discharge, and 1-year mortality in patients with a first AMI. Follow-up data were available through NORMI until 31 December 2019.

Patient and public involvement

This study used existing data from Norwegian national health registries. Registration into these registries is mandatory (the Norwegian Cardiovascular Disease Registry Regulation and the Norwegian Health Register Act), and consent by the patient was not required.

Statistics

Continuous variables are presented as the mean \pm standard deviation or median (25th percentile, 75th percentile). Categorical variables are presented as numbers and percentages. Age-adjusted mortality rates were calculated using direct standardization. Time trends were analysed (log-linear model) using the Joinpoint Regression Program (version 4.0; SEER software, National Cancer Institute, USA) and are presented as the expected annual per cent changes with a 95% confidence interval (CI). Other data were analysed using STATA version 17 (StataCorp LLC, College Station, TX, USA). A P-value of <0.05 was regarded as statistically significant.

Ethics

The Regional Committee for Medical and Health Research Ethics North approved the study (REK 2016/170).

Results

Hospital admissions

A total of 61 625 patients with a first AMI were registered in the NORMI in the period 2013–19. A total of 18 499 (30%) patients were classified as ST-elevation myocardial infarction (STEMI), and 40 709 (66%) patients as non-STEMI. The rest was unclassifiable. The number of registered patients per year with a first AMI was reduced from 8933 in 2013 to 8383 in 2019. The decline was minimal in the period 2013–15, but from 2016, we found a significant reduction of 4.4% (95% CI 4.1, 4.6) per year.

The number of admissions per year by gender and age group are presented in [Figure 1](#). For both genders <90 years of age, there were statistically significant reduction during the period.

Clinical characteristics and risk factors

Clinical characteristics are presented in [Table 1](#). A total of 22 581 patients (37%) with a first AMI were women. The proportion of women decreased from 37% in 2013 to 35% in 2019 (change per year: -0.9% , 95% CI $-1.7, -0.2$). The median age for men increased from 65 years (interquartile range 56–76 years) in 2013 to 67 years (interquartile range 57–76 years) in 2019. The median age in women was 76 years (interquartile range 66–85 years) throughout the study period. The proportion of smokers was reduced by 2.7% (95% CI 2.2, 3.2) per year, while the proportions of patients with diabetes and lipid-lowering therapy increased by 2.4% (95% CI 1.0, 3.8) and 2.0% (95% CI 0.4, 3.6) per year, respectively.

Coronary angiography and secondary preventive therapy

Invasive coronary angiography and percutaneous coronary intervention (PCI) were performed in 16 160 (87%) and 15 006 (81%) patients admitted with a first STEMI, and were performed in 25 561 (63%) and 17 101 (42%) patients admitted with a first non-STEMI, respectively. The proportion of patients with STEMI who underwent coronary angiography was stable during the period (change per year: 0.5%, 95% CI $-0.4, 1.4$), while the proportion of non-STEMI





Figure 1 Patients admitted to hospitals with first acute myocardial infarction in Norway 2013–19 and registered in the Norwegian myocardial infarction register.

Table 1 Clinical characteristics in patients admitted to hospitals with first acute myocardial infarction in Norway 2013–19

	2013		2014		2015		2016		2017		2018		2019	
	<i>n</i> = 8933		<i>n</i> = 9012		<i>n</i> = 9020		<i>n</i> = 8926		<i>n</i> = 8773		<i>n</i> = 8578		<i>n</i> = 8383	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Men	5609	63	5656	63	5640	63	5652	63	5562	63	5454	64	5471	65
Median age (year, interquartile range)	69	(59–81)	69	(59–81)	70	(59–81)	70	(59–81)	70	(60–80)	70	(60–80)	70	(59–79)
Smoking	2777	31	2677	30	2642	29	2501	28	2393	27	2315	27	2204	26
Antihypertensive therapy	3984	45	3881	43	4005	44	3882	43	3863	44	3880	45	3779	45
Diabetes	1434	16	1403	16	1485	16	1491	17	1579	18	1535	18	1482	18
Lipid-lowering therapy	2287	26	2164	24	2203	24	2265	25	2343	27	2318	27	2319	28
Previous percutaneous coronary intervention	466	5	463	5	444	5	465	5	510	6	488	6	450	5
Previous coronary artery bypass grafting	397	4	391	4	370	4	394	4	393	4	349	4	308	4
Previous stroke (all types)	713	8	643	7	670	7	603	7	640	7	600	7	518	6

**Figure 2** Proportion of patients admitted with acute non-ST-elevation myocardial infarction (first myocardial infarction) examined with coronary angiography in Norway 2013–19.

patients who underwent coronary angiography increased by 2.4% per year (95% CI 1.6, 3.3), from 58% in 2013 to 68% in 2019.

The proportion of women with non-STEMI who underwent coronary angiography increased by 3.2% per year (95% CI 1.6, 4.7) from 47% in 2013 to 56% in 2019, while the proportion of men with non-STEMI who underwent coronary angiography increased by 1.9% per year (95% CI 1.4, 2.4) from 68% in 2013 to 76% in 2019.

The proportion of patients with non-STEMI undergoing coronary angiography in relation to age is presented in [Figure 2](#) and [Table 2](#).

The prescription of dual antiplatelet therapy (acetylsalicylic acid and P₂Y₁₂ inhibitor) and statins in patients discharged alive increased from 76 to 77% for dual antiplatelet therapy (change per year: 0.4%, 95% CI 0.0, 0.7), and from 82 to 85% (change per year: 0.7%, 95% CI 0.4, 1.1) for statins during the study period.

Table 2 Coronary angiography and percutaneous coronary intervention in patients with acute non-ST-elevation myocardial infarction (first myocardial infarction) in Norway 2013–19

Age (years)	2013			2014			2015			2016										
	Patients		PCI	Patients		PCI	Patients		PCI	Patients		PCI								
	n	%	n	n	%	n	%	n	%	n	%	n								
18–49	425	363	85	243	57	414	355	86	244	59	395	342	87	241	61	451	380	84	254	56
50–66	1771	1502	85	1021	58	1771	1508	85	1056	60	1687	1450	86	1025	61	1712	1467	86	1065	62
67–79	1750	1192	68	731	42	1898	1343	71	821	43	1843	1262	68	792	43	1940	1373	71	864	45
80–89	1472	419	28	268	18	1447	414	29	259	18	1428	445	31	289	20	1393	497	36	305	22
≥90	445	17	4	13	3	517	17	3	8	2	579	22	4	17	3	497	37	7	26	5

Age (years)	2017			2018			2019								
	Patients		Coronary angiography	Patients		PCI	Patients		Coronary angiography	Patients		PCI			
	n	%	n	n	%	n	%	n	%	n	%	n			
18–49	348	303	87	196	56	368	318	86	214	58	352	293	83	200	57
50–66	1677	1465	87	1058	63	1586	1394	88	1026	65	1644	1446	88	979	60
67–79	2002	1459	73	926	46	1887	1410	75	918	49	1902	1422	75	932	49
80–89	1336	519	39	339	25	1281	493	38	335	26	1126	507	45	343	30
≥90	484	42	9	35	7	480	53	11	34	7	398	32	8	24	6




Figure 3 One-year mortality after admission for first acute myocardial infarction in Norway 2013–19.

Outcome

All-cause mortality after a first AMI in different age groups is presented in [Table 3](#) and [Figure 3](#). We found no changes in age-adjusted 30-day mortality during the period, but age-adjusted 1-year mortality was reduced by 1.6% (95% CI 0.4, 2.7) per year from 16.4% in 2013 to 15.1% in 2018. The reduction in mortality was found in patients with non-STEMI: age-adjusted 1-year mortality after non-STEMI was reduced from 17.8% in 2013 to 15.2% in 2018, while it was unchanged at 14% after STEMI. Information regarding 1-year mortality after STEMI and non-STEMI in different age groups is presented in [Supplementary material online, Table S1](#). With respect to gender differences, the reduction was seen only in men; age-adjusted 1-year mortality for men was reduced by 2.6% (95% CI 0.4, 4.8) per year, but we found no corresponding change for women.

Ischaemic heart disease (ICD-10 code I20-I25) was reported as the cause of death in 47% of patients who died within 1 year in 2013. The proportion was reduced to 43% in 2018.

Discussion

This nationwide study of patients with a first AMI admitted to hospitals in Norway from 2013 to 2019 revealed a reduction in hospitalizations due to a first AMI and a gradual increase in the proportion of patients who were examined with coronary angiography and prescribed guideline-recommended secondary preventive therapy. The age-adjusted 1-year mortality decreased during the period. The changes over time were primarily seen in the oldest groups.

A reduced number of hospital admissions and a reduced incidence of AMI in Norway in the period 1991–2014 has been described previously.²⁰ Our study from a national medical quality register shows a further reduction in the number of admissions for incident AMIs until 2019. The relatively high proportion of STEMI vs. non-STEMI patients can be explained by inclusion of patients with first-time AMI only.

In another Norwegian study, the Tromsø study, the declining incidence of acute coronary syndrome was largely attributed to changes in coronary risk factors such as lower cholesterol and blood pressure levels, fewer smokers and more physical activity in the population.²¹ The present study includes only patients admitted to hospitals with an AMI and cannot be used to assess changes in risk factors in the general population. However, in our study, the proportion of patients with a first AMI who smoked was higher than in the general population [2013: 15%; 2019: 9% (age 16–74 years)], suggesting that smoking is still an important modifiable risk factor for AMI, especially in younger patients.^{11,22}

The guidelines from the European Society of Cardiology recommended early coronary angiography in most patients with non-STEMI.^{12–15} Although the proportion of patients with non-STEMI who underwent coronary angiography increased from 58 to 68% during the period, the proportion is still lower in Norway than in other countries in Europe with national AMI registries.⁴ Although the proportion undergoing coronary angiography increased most in the older age groups, it was still significantly lower in these groups compared with younger age groups. Increasing comorbidity with increasing age may have had an impact on the choice of treatment strategy. The gender difference in the proportion of women and men who were

examined with coronary angiography is also noteworthy and persisted during the study period.⁶ Possible differences in symptoms and clinical findings in suspected AMI in younger and older patients and in women and men cannot explain the differences, since only patients with the diagnosis of AMI were registered in the NORMI and consequently were included in this study. We have not investigated geographical differences in treatment strategy, but distance to hospitals offering coronary angiography has probably also been of importance.

Secondary preventive drugs such as antiplatelet therapy and statins are important in preventing new cardiovascular events and are recommended as secondary prevention after AMI.^{12–15} The proportions of patients prescribed these drugs after AMI in Norway were comparable with other countries in Europe.^{4,5} However, several studies have demonstrated a remaining gap between the guidelines and the achievement of recommended targets for cardiovascular risk factors and medication use after myocardial infarction.^{8,23–29} Early combination of statin, ezetimibe, and in some cases also inhibitors of proprotein convertase subtilisin/kexin type 9 is recommended in patients with high risk of new events.³⁰

Mortality after AMI has shown a declining trend in Norway for many years.³¹ The present study shows a further reduction in all-cause mortality after AMI for the period 2013–19. The NORMI does not have follow-up data for secondary preventive therapy, but more favourable risk profiles, improved acute treatment with an increasing proportion of patients examined with coronary angiography and treated with PCI, and increased prescription of secondary preventive drugs may have contributed to the improved survival. Changes in the general mortality in the population as well as changes in the use of medical diagnostic codes and procedure codes may also have affected the results, which must therefore be interpreted with caution.

The main strengths of this study are the large and unselected population comprising nearly all patients hospitalized with a first AMI in Norway from 2013 to 2019, and a nearly complete follow up. However, there are some important limitations associated with the study design and the NORMI. This study was an observational study, making it impossible to demonstrate causal associations between treatment and outcomes. Only AMIs that led to hospitalization were registered in the NORMI. A few hospitals did not deliver complete data for the whole period, but the coverage compared with the Norwegian Patient Register was >90%.¹⁸ We only obtained deidentified data from the NORMI and the Norwegian Cardiovascular Disease Registry and could not verify the information through medical records at the individual patient level. Nevertheless, the degree of completeness and correctness of most variables in the NORMI have been shown to be high.³² The findings of the study must be interpreted with caution and generalization of the results should be avoided.

Conclusion

In conclusion, in the time period 2013–19 in Norway, we found a reduction in the number of patients admitted with a first AMI. Furthermore, more patients with a first AMI underwent coronary angiography, more patients were prescribed secondary preventive therapy at discharge, and the age-adjusted 1-year mortality after

AMI was reduced in this period. The national myocardial infarction registry provides important information about AMI and may have contributed to better adherence to guideline recommendations.

Lead author biography



Jarle Jortveit is a senior researcher and cardiologist at Sørlandet Hospital Arendal, Norway. He is also co-founder and chief medical officer in ECG247 Smart Heart Sensor.

Authors' contributions

J.J. and S.H. were responsible for the conception of the study, and the analysis and interpretation of data. J.J. drafted the manuscript. A.H.P. contributed to the analysis and interpretation of the data and critically revised the manuscript. J.L. and S.H. critically revised the manuscript. All gave final approval and agreed to be accountable for all aspects of work ensuring its integrity and accuracy.

Data availability

The data underlying this article were provided by the Norwegian Institute of Public Health under license/by permission. Data will be shared on request to the corresponding author with permission of the Norwegian Institute of Public Health.

Supplementary material

Supplementary material is available at *European Heart Journal Open* online.

Funding

This work was supported by South-Eastern Norway Regional Health Authority, Hamar, Norway. The funder has no role in the design and conduct of the study, in the collection, analysis, and interpretation of the data, and in the preparation, review, or approval of the manuscript.

Conflict of interest: J.J. has received speaking fees from Amgen, AstraZeneca, BMS, Boehringer Ingelheim, Novartis, Pfizer, and Sanofi. He is a chief medical officer in ECG247 Smart Heart Sensor. S.H. has received speaking fees from Boehringer Ingelheim, BMS, Pfizer, and Sanofi. All other authors no conflict of interest.

References

1. Timmis A, Vardas P, Townsend N, Torbica A, Katus H, De Smedt D, Gale CP, Maggioni AP, Petersen SE, Huculeci R, Kazakiewicz D, de Benito Rubio V, Ignatiuk B, Raisi-Estabragh Z, Pawlak A, Karagiannis E, Treskes R, Gaita D, Beltrame JF, McCannachie A, Bardinet I, Graham I, Flather M, Elliott P, Mossialos EA, Weidinger F, Achenbach S, Atlas Writing Group, European Society of Cardiology.

- European Society of Cardiology: cardiovascular disease statistics 2021. *Eur Heart J* 2022;**43**:716–799.
2. McManus DD, Gore J, Yarzebski J, Lessard D, Goldberg RJ. Recent trends in the incidence, treatment, and outcomes of patients with STEMI and NSTEMI. *Am J Med* 2011;**124**:40–47.
 3. Timmis A, Townsend N, Gale C, Grobbee R, Maniadas N, Flather M, Wilkins E, Wright L, Vos R, Bax J, Blum M, Pinto F, Vardas P, Goda A, Demiraj AF, Weidinger F, Metzler B, Ibrahimov F, Pasquet AA, Claeys M, Thornton Y, Kusljagic Z, Smajic E, Velchev V, Ivanov N, Antoniadou L, Agathangelou P, Táborský M, Gerdes C, Viigima M, Juhani PM, Juilliere Y, Cattani S, Aladashvili A, Hamm C, Kuck K-H, Papoutsis K, Bestehorn K, Fousas S, Giannoulidou G, Varounis C, Kallikazaros I, Kiss RG, Czétényi T, Becker D, Gudnason T, Kearney P, McDonald K, Rozenman Y, Ziv B, Bolognese L, Luciolli P, Boriani G, Berkinbayev S, Rakisheva A, Mirrahimov E, Erglis A, Jegere S, Marinakis G, Beissel J, Marchal N, Kedev S, Xuereb RG, Tilney T, Felice T, Popovici M, Bax J, Mulder B, Simoons ML, Elsendoorn M, Steigen TK, Atar D, Kalarus Z, Tendera M, Cardoso JS, Ribeiro J, Mateus C, Tatu-Chitoiu G, Seferovic P, Beleslin B, Simkova I, Durcikova P, Belicova V, Fras Z, Radelj S, Gonzalez Juanatey JR, Legendre S, Braunschweig F, Kaufmann UP, Rudiger-Sturchler M, Tokgozoglu L, Unver A, Kovalenko V, Nesukay E, Naum A, de Courtelary PT, Martin S, Sebastiao D, Ghislain D, Bardinet I, Logstrup S. European Society of Cardiology: cardiovascular disease statistics 2017. *Eur Heart J* 2018;**39**:508–579.
 4. Edfors R, Jernberg T, Lewinter C, Blöndal M, Eha J, Löveke P, Marandi T, Ainla T, Saar A, Veldre G, Ferenci T, Andr eka P, J nosi A, Jortveit J, Halvorsen S. Differences in characteristics, treatments and outcomes in patients with non-ST-elevation myocardial infarction – novel insights from four national European continuous real-world registries. *Eur Heart J Qual Care Clin Outcomes* 2022;**8**:429–436.
 5. Blondal M, Ainla T, Eha J, L veke P, Marandi T, Saar A, Veldre G, Edfors R, Lewinter C, Jernberg T, Jortveit J, Halvorsen S, Becker D, Csan di Z, Ferenci T, Andr eka P, J nosi A. Comparison of management and outcomes of ST-segment elevation myocardial infarction patients in Estonia, Hungary, Norway and Sweden according to national ongoing registries. *Eur Heart J Qual Care Clin Outcomes* 2021;**7**:307–314.
 6. Jortveit J, Govatsmark RE, Langorgen J, Hole T, Mannsverk J, Olsen S, Ris e C, Halvorsen S. Gender differences in the assessment and treatment of myocardial infarction. *Tidsskr Nor Laegeforen* 2016;**136**:1215–1222.
 7. Jortveit J, Halvorsen S. Geographical differences in prescription of secondary preventive drug therapy after acute myocardial infarction in Norway. *Eur Heart J Cardiovasc Pharmacother* 2017;**3**:132–133.
 8. Jortveit J, Halvorsen S, Kaldal A, Pripp AH, Govatsmark RES, Langorgen J. Unsatisfactory risk factor control and high rate of new cardiovascular events in patients with myocardial infarction and prior coronary artery disease. *BMC Cardiovasc Disord* 2019;**19**:71.
 9. Jortveit J, Pripp AH, Halvorsen S. Outcomes after delayed primary percutaneous coronary intervention versus pharmaco-invasive strategy in ST-segment elevation myocardial infarction in Norway. *Eur Heart J Cardiovasc Pharmacother* 2021:pvb041.
 10. Jortveit J, Pripp AH, Langorgen J, Halvorsen S. Poor adherence to guideline recommendations among patients with atrial fibrillation and acute myocardial infarction. *Eur J Prev Cardiol* 2019;**26**:1373–1382.
 11. Jortveit J, Pripp AH, Langorgen J, Halvorsen S. Incidence, risk factors and outcome of young patients with myocardial infarction. *Heart* 2020;**106**:1420–1426.
 12. Steg PG, James SK, Atar D, Badano LP, Lindqvist CB, Borger MA, Di Mario C, Dickstein K, Ducrocq G, Fernandez-Aviles F, Gershlick AH, Giannuzzi P, Halvorsen S, Huber K, Juni P, Kastrati A, Knuuti J, Lenzen MJ, Mahaffey KW, Valgimigli M, van't Hof A, Widimsky P, Zahger D, Bax JJ, Baumgartner H, Ceconi C, Dean V, Deaton C, Fagard R, Funck-Brentano C, Hasdai D, Hoes A, Kirchhof P, Knuuti J, Kolh P, McDonagh T, Moulin C, Popescu BA, Reiner Z, Sechtem U, Sirnes PA, Tendera M, Torbicki A, Vahanian A, Windecker S, Hasdai D, Astin F,  str m-Olsson K, Budaj A, Clemmensen P, Collet J-P, Fox KA, Fuat A, Gustiene O, Hamm CW, Kala P, Lancellotti P, Maggioni AP, Merkely B, Neumann F-J, Piepoli MF, Van de Werf F, Verheugt F, Wallentin L. ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2012;**33**:2569–2619.
 13. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, Caforio ALP, Crea F, Goudevenos JA, Halvorsen S, Hindricks G, Kastrati A, Lenzen MJ, Prescott E, Roffi M, Valgimigli M, Varenhorst C, Vranckx P, Widimsky P, ESC Scientific Document Group. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: the task force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2018;**39**:119–177.
 14. Hamm CW, Bassand JP, Agewall S, Bax J, Boersma E, Bueno H, Caso P, Dudek D, Gielen S, Huber K, Ohman M, Petrie MC, Sonntag F, Uva MS, Storey RF, Wijns W, Zahger D, Bax JJ, Auricchio A, Baumgartner H, Ceconi C, Dean V, Deaton C, Fagard R, Funck-Brentano C, Hasdai D, Hoes A, Knuuti J, Kolh P, McDonagh T, Moulin C, Poldermans D, Popescu BA, Reiner Z, Sechtem U, Sirnes PA, Torbicki A, Vahanian A, Windecker S, Windecker S, Achenbach S, Badimon L, Bertrand M, Botker HE, Collet J-P, Crea F, Danchin N, Falk E, Goudevenos J, Gulba D, Hambrecht R, Herrmann J, Kastrati A, Kjeldsen K, Kristensen SD, Lancellotti P, Mehilli J, Merkely B, Montalescot G, Neumann F-J, Neyses L, Perk J, Roffi M, Romeo F, Ruda M, Swahn E, Valgimigli M, Vrints CJ, Widimsky P. ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: the task force for the management of acute coronary syndromes (ACS) in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2011;**32**:2999–3054.
 15. Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, Bax JJ, Borger MA, Brotons C, Chew DP, Gencer B, Hasenfuss G, Kjeldsen K, Lancellotti P, Landmesser U, Mehilli J, Mukherjee D, Storey RF, Windecker S, ESC Scientific Document Group. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: task force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2016;**37**:267–315.
 16. Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD, Writing Group on the Joint ESC/ACC/AHA/WHF Task Force for the Universal Definition of Myocardial Infarction, Thygesen K, Alpert JS, White HD, Jaffe AS, Katus HA, Apple FS, Lindahl B, Morrow DA, Chaitman BA, Clemmensen PM, Johanson P, Hod H, Underwood R, Bax JJ, Bonow RO, Pinto F, Gibbons RJ, Fox KA, Atar D, Newby LK, Galvani M, Hamm CW, Uretsky BF, Gabriel Steg P, Wijns W, Bassand J-P, Menasch  P, Ravkilde J, Ohman EM, Antman EM, Wallentin LC, Armstrong PW, Simoons ML, Januzzi JL, Nieminen MS, Gheorghiade M, Filippatos G, Luepker RV, Fortmann SP, Rosamond WD, Levy D, Wood D, Smith SC, Hu D, Lopez-Sendon J-L, Robertson RM, Weaver D, Tendera M, Bove AA, Parkhomenko AN, Vasilieva EJ, Mendis S, ESC Committee for Practice Guidelines (CPG). Third universal definition of myocardial infarction. *Eur Heart J* 2012;**33**:2551–2567.
 17. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, White HD, ESC Scientific Document Group. Fourth universal definition of myocardial infarction (2018). *Eur Heart J* 2019;**40**:237–269.
 18. Jortveit J, Govatsmark RE, Digre TA, Ris e C, Hole T, Mannsverk J, Sl rdahl SA, Halvorsen S. Myocardial infarction in Norway in 2013. *Tidsskr Nor Laegeforen* 2014;**134**:1841–1846.
 19. Govatsmark RE, Sneeggen S, Karls ne H, Sl rdahl SA, B naa KH. Interrater reliability of a national acute myocardial infarction register. *Clin Epidemiol* 2016;**8**:305–312.
 20. Sulo G, Iglund J, Vollet SE, Ebbing M, Egeland GM, Ariansen I, Tell GS. Trends in incident acute myocardial infarction in Norway: an updated analysis to 2014 using national data from the CVDNOR project. *Eur J Prev Cardiol* 2018;**25**:1031–1039.
 21. Mannsverk J, Wilsgaard T, Mathiesen EB, L chen M-L, Rasmussen K, Thelle DS, Nj lstad I, Hopstock LA, B naa KH. Trends in modifiable risk factors are associated with declining incidence of hospitalized and nonhospitalized acute coronary heart disease in a population. *Circulation* 2016;**133**:74–81.
 22. R yk, alkohol og andre rusmidler. <https://www.ssb.no/helse/helseforhold-og-levevaner/statistikk/royk-alkohol-og-andre-rusmidler> (19 December 2021).
 23. Kotseva K, Wood D, De Bacquer G, Ryd n L, Jennings C, Gyberg V, Amouyel P, Bruthans J, Castro Conde A, Cifkov  R, Deckers JW, De Sutter J, Dilic M, Dolzhenko M, Erglis A, Fras Z, Gaita D, Gotcheva N, Goudevenos J, Heuschmann P, Laucevicius A, Lehto S, Lovic D, Milicic D, Moore D, Nicolaides E, Oganov R, Pajak A, Pogossova N, Reiner Z, Stagno M, St rk S, Tokgoz glu L, Vucic D. EUROASPIRE IV: a European Society of Cardiology survey on the lifestyle, risk factor and therapeutic management of coronary patients from 24 European countries. *Eur J Prev Cardiol* 2016;**23**:636–648.
 24. Ferrari R, Ford I, Greenlaw N, Tardif JC, Tendera M, Abergel H, Fox K, Hu D, Shalnova S, Steg PG, CLARIFY Registry Investigators. Geographical variations in the prevalence and management of cardiovascular risk factors in outpatients with CAD: data from the contemporary CLARIFY registry. *Eur J Prev Cardiol* 2014;**22**:1056–1065.
 25. Cacoub PP, Zeymer U, Limbourg T, Baumgartner I, Poldermans D, R ther J, Bhatt DL, Steg PG, REACH Registry Investigators. Effects of adherence to guidelines for the control of major cardiovascular risk factors on outcomes in the REDuction of Atherothrombosis for Continued Health (REACH) registry Europe. *Heart* 2011;**97**:660–667.
 26. Ray KK, Molemans B, Schoonen WM, Giovanna P, Bray S, Kiru G, Murphy J, Banach M, De Servi S, Gaita D, Gouni-Berthold I, Hovingh GK, Jozwiak JJ, Jukema JW, Kiss RG, Kownator S, Iversen HK, Maher V, Masana L, Parkhomenko A, Peeters A, Clifford P, Raslova K, Siostrzonek P, Romeo S, Tousoulis D, Vlachopoulos C, Vrablik M, Catapano AL, Poulter NR. EU-wide cross-sectional observational study of lipid-modifying therapy use in secondary and primary care: the DA VINCI study. *Eur J Prev Cardiol* 2021;**28**:1279–1289.
 27. Dyrbus K, Gasior M, Desperak P, Nowak J, Osadnik T, Banach M. Characteristics of lipid profile and effectiveness of management of dyslipidaemia in patients with acute coronary syndromes - data from the TERCET registry with 19,287 patients. *Pharmacol Res* 2019;**139**:460–466.

28. Vallejo-Vaz AJ, Bray S, Villa G, Brandts J, Kiru G, Murphy J, Banach M, De Servi S, Gaita D, Gouni-Berthold I, Kees Hovingh G, Jozwiak JJ, Jukema JW, Gabor Kiss R, Kownator S, Iversen HK, Maher V, Masana L, Parkhomenko A, Peeters A, Clifford P, Raslova K, Siostrzonek P, Romeo S, Tousoulis D, Vlachopoulos C, Vrablik M, Catapano AL, Poulter NR, Ray KK, DA VINCI Study Investigators. Implications of ACC/AHA versus ESC/EAS LDL-C recommendations for residual risk reduction in ASCVD: a simulation study from DA VINCI. *Cardiovasc Drugs Ther* 2022; May 14. doi:10.1007/s10557-022-07343-x. Epub ahead of print. PMID:35567726.
29. Kaldal A, Tonstad S, Jortveit J. Long-term hospital-based secondary prevention of coronary artery disease: a randomized controlled trial. *BMC Cardiovasc Disord* 2021;**21**:600.
30. Banach M, Penson PE, Vrablik M, Bunc M, Dyrbus K, Fedacko J, Gaita D, Gierlotka M, Jarai Z, Magda SL, Margetic E, Margoczy R, Durak-Nalbantic A, Ostadal P, Pella D, Trbusic M, Udroi CA, Vlachopoulos C, Vulic D, Fras Z, Dudek D, Reiner Ž. Optimal use of lipid-lowering therapy after acute coronary syndromes: a position paper endorsed by the international lipid expert panel (ILEP). *Pharmacol Res* 2021;**166**: 105499.
31. Sulo E, Vollset SE, Nygard O, Sulo G, Iglund J, Egeland GM, Ebbing M. Trends in 28-day and 1-year mortality rates in patients hospitalized for a first acute myocardial infarction in Norway during 2001–2009: a “cardiovascular disease in Norway” (CVDNOR) project. *J Intern Med* 2015;**277**:353–361.
32. Govatsmark RES, Janszky I, Slordahl SA, Ebbing M, Wiseth R, Grenne B, Vesterbeekmo E, Bønaa KH. Completeness and correctness of acute myocardial infarction diagnoses in a medical quality register and an administrative health register. *Scand J Public Health* 2020;**48**:5–13.