


REVIEW



## Dominance and fitness costs of insect resistance to genetically modified *Bacillus thuringiensis* crops

Fangneng Huang 

Department of Entomology, Louisiana State University Agricultural Center, Baton Rouge, LA, USA

### ABSTRACT

Evolution of resistance to genetically modified *Bacillus thuringiensis* (Bt) crops in pest populations is a major threat to the sustainability of the technology. Incidents of field resistance that have led to control problems of Bt crops or significantly reduced susceptibility of individual Bt proteins in pyramided plants have increased dramatically across the world, especially in recent years. Analysis of globally published data showed that 61.5% and 60.0% of the cases of resistance with major alleles that allowed homozygous resistant genotypes to survival on Bt crops were functionally non-recessive and did not involve fitness costs, respectively. Dominance levels ( $D_{FLS}$ ) measured on Bt plants ranged from  $-0.02$  to  $1.56$  with a mean ( $\pm$  sem) of  $0.35 \pm 0.13$  for the 13 cases of single-gene resistance to Bt plants that have been evaluated. Among these, all six cases with field control problems were functionally non-recessive with a mean  $D_{FL}$  of  $0.63 \pm 0.24$ , which was significantly greater than the  $D_{FL}$  ( $0.11 \pm 0.07$ ) of the seven cases without field resistance. In addition, index of fitness costs (IFC) of major resistance was calculated for each case based on the fitness of resistant (R'R') and heterozygous (R'S') genotypes on non-Bt plants divided by the fitness of their susceptible (S'S') counterparts. The estimated IFCs for 15 cases of single-gene resistance were similar for R'R' and R'S', and for the cases with and without field resistance; and the values averaged  $1.10 \pm 0.12$  for R'R' and  $1.20 \pm 0.18$  for R'S'. Limited published data suggest that resistance of insects to dual/multiple-gene Bt crops is likely to be more recessive than the related single-gene resistance, but their IFCs are similar. The quantitative analysis of the global data documents that the prevalence of non-recessive resistance has played an essential role in the widespread evolution of resistance to Bt crops, while the lack of fitness costs is apparently not as critical as the non-recessive resistance. The results suggest that planting of 'high dose' traits is an effective method for Bt crop IRM and more comprehensive management strategies that are also effective for functionally non-recessive resistance should be deployed.

### ARTICLE HISTORY

Received 2 September 2020  
Revised 12 November 2020  
Accepted 12 November 2020



### KEYWORDS


GM Bt crops; insect resistance management; dominance level; fitness costs of resistance

### Introduction

The year of 1996 marked the first year that genetically modified (GM) crops expressing *Bacillus thuringiensis* (Bt) genes were officially commercialized in the US and several other countries.<sup>1</sup> Since then, Bt crops have gained widespread acceptance throughout the world. By 2018, a total of more than one billion hectares of Bt crops had been planted in more than 20 countries.<sup>1</sup> The predominant Bt crops planted are maize, cotton, and soybean. Bt crop traits are usually highly effective in controlling some major insect pests; thus, considerable economic, environmental, and social benefits have been gained from planting Bt crops in both industrial and developing countries.<sup>1–5</sup>

However, evolution of resistance in target insect populations is a great threat to the sustainability of the Bt crop technology.<sup>6–8</sup> To delay resistance evolution, a 'high dose/refuge' (HDR) insect resistance management (IRM) strategy has been recommended in the U.S. and several other countries.<sup>9,10</sup> This strategy requires crop growers to plant a portion of the crop with 'high dose' Bt plants that can kill almost all homozygous-susceptible (SS) individuals and heterozygous-resistant (RS) individuals of the target pest species, while the remaining portion of the crop is planted with non-Bt plants to serve as a 'refuge' for SS populations. In this scenario, abundant SS individuals from the refuge plants will be available to mate with the rare homo-

**CONTACT** Fangneng Huang  [fhuang@agcenter.lsu.edu](mailto:fhuang@agcenter.lsu.edu)  Department of Entomology, Louisiana State University AgCenter, Baton Rouge, LA 70803, USA. This paper reports research results only. Mention of a proprietary product name does not constitute an endorsement for its use by Louisiana State University Agricultural Center.

 Supplemental data for this article can be accessed on the [publisher's website](#).

© 2020 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

zygous-resistant (RR) survivors from the Bt plants. In this way, the majority of the offspring, if they possess resistance alleles, should be heterozygous, and these RS individuals will be killed by the 'high dose' Bt plants. As a result, resistance evolution should be greatly delayed.<sup>9,11</sup> The current HDR for Bt crop IRM was largely based on knowledge generated from earlier laboratory studies of resistance to purified Bt proteins and Bt insecticides. Several earlier studies showed that high-level resistance to purified Bt proteins or Bt insecticides was typically recessive.<sup>11–14</sup> In addition, it was also thought that Bt plants might be able to make 'genetically' non-recessive resistance functionally recessive because GM plants could be engineered to consistently express high levels of Bt proteins that would be capable of killing a large portion of 'genetically' non-recessive RS individuals.<sup>11,15</sup> Thus, functionally recessive resistance is a key foundation for the success of HDR strategy.<sup>14,16,17</sup> Another IRM strategy for Bt crops is 'gene pyramiding' of two or more Bt genes with dissimilar modes of action.<sup>18–20</sup> In the gene pyramiding, if individuals in insect populations carrying resistance alleles to one Bt protein are rare, the individuals simultaneously possess resistance alleles to two or more Bt proteins must be very rare. No significant cross-resistance that allows survival of resistant insects to one Bt protein on pyramided plants is a key assumption for the success of this strategy.

Fitness costs of resistance refers to a reduced fitness (e.g. delayed development, higher mortality, lower reproduction) of RR or RS individuals relative to SS individuals in the absence of selection. If both RR and RS individuals show fitness costs, the phenomenon is called non-recessive fitness costs. Otherwise, if only RR individuals show fitness costs, but RS performs similarly to SS, the fitness cost is considered recessive.<sup>21–23</sup> Both the dominance level and fitness costs of resistance are important factors in resistance evolution. Fitness costs of resistance could result in declines in resistance and even reversion to susceptibility after selection pressure is removed.<sup>21–23</sup> Earlier studies with purified Bt proteins or Bt insecticides showed that Bt resistance, especially high-level resistance, was often associated with fitness costs.<sup>12,23,24</sup> Thus, fitness costs are also considered a positive factor

that may elevate the effectiveness of refuge planting for Bt crop IRM.<sup>23</sup>

After 20+ years of global Bt crop use, field resistance that has resulted in reduced efficacy of Bt crops or significantly reduced susceptibility of individual Bt proteins in pyramided Bt plants has been documented in at least 20 cases involving seven major pest species of maize and cotton in six countries across four continents.<sup>6–8</sup> In addition, major resistance alleles that allow RR individuals to survive and complete their life cycle on Bt plants,<sup>25</sup> while susceptible individuals of the species are controlled by the Bt plants, have also been isolated in several cases.<sup>6,7</sup> To avoid any confusions with the survival due to natural tolerance, in this review, the term 'major resistance allele' for Bt plants is defined as RR individuals (populations, colonies, strains) possessing homozygous resistance alleles should exhibit a significantly greater rate to survive and complete their life cycles on the Bt plants, relative to their susceptible counterparts. It should be pointed out that there is variability in Bt susceptibility within populations of a same pest species even before Bt crops are commercialized. Information that was used to judge if a case of resistance qualified as 'a major resistance allele' in this review was based on only the resistant and susceptible insect populations used in the peer-reviewed studies. Based on this definition, 'field resistance' certainly qualifies as 'major resistance'. However, finding major resistance alleles in a pest population does not necessarily indicate an immediate threat of field resistance to the Bt plants.<sup>26</sup> Field resistance can occur when the frequency of major resistance alleles becomes sufficiently common to reduce efficacy of the Bt crop in field.<sup>6,17</sup> During the last two decades, many studies have been conducted in the world to characterize the resistance (e.g. dominance, fitness costs, etc.) for some of these cases involving major resistance alleles including field resistance. Several previous reviews have analyzed the general conditions that are associated with the evolution of resistance to Bt proteins and/or Bt plants.<sup>6,7,11,12,14,17,23,27–30</sup> In this mini review, I focus on only two important aspects that are closely related to resistance management: dominance level and fitness costs of resistance to Bt plants. More specifically, in the current review I will first extend the methods that are used

to measure dominance levels/fitness costs of resistance in insect on single-gene Bt plants to pyramided Bt plants and then use the related globally published data to quantitatively analyze the relationship between dominance levels/fitness costs and the evolution of resistance to Bt crops. In addition, variations in the dominance levels/fitness costs among pest populations, Bt proteins, test methods, and crop-pest systems are also discussed. Knowledge generated from this analysis should be useful in understanding the reasons that have led to the recent wide occurrence of field resistance to Bt crops in the world and in refining current IRM strategies for the sustainable use of Bt crop technology.

### Criteria for literature selection and cases of resistance to Bt crops

Google Scholar (<https://scholar.google.com/>) was used to search the related literature published before January 31, 2020. Because Bt proteins expressed in GM plants can be different from the proteins produced by *B. thuringiensis* bacteria, information generated from the studies that used Bt protoxin, activated proteins, or microbial insecticides may not directly represent the status of insect resistance to Bt crops.<sup>31</sup> To ensure data used in this review more accurately reflect the real situation of resistance to Bt crops, the following three criteria were used in selection of literature. First, only articles published in peer-reviewed journals were used. Second, resistant insects used in

studies must have possessed major resistant genes to the Bt plants as defined above.<sup>17,25</sup> Third, biological parameters measured in the studies must have been obtained from assays using whole plants/plant tissues of maize or cotton, because, by far, field resistance to Bt crops has been found only in Bt maize or Bt cotton.<sup>7</sup> In other words, those studies using the ‘resistant’ populations that had not been documented to carry major resistance alleles to Bt plants, as well as data generated from assays on meridic diet or Bt protein-treated diet, or plants other than maize or cotton, were not included in this review. Use of published data was approved by the senior or corresponding author of each selected publication. To facilitate the analysis, the definition of a ‘case’ of resistance described in reference<sup>7</sup> was adopted, which means that each case of single-gene resistance represents a resistance of one pest species in one country to one Bt protein in the crop plant. In addition, in this review, the definition of ‘case’ of single-gene resistance was also extended to include dual-/multiple-gene resistance. A case of resistance to dual/multiple-gene Bt gene plants means a resistance of one pest species in one country to the dual-/multiple-Bt proteins expressed in a pyramided crop trait. A pyramided crop trait is a GM plant product that contains two or more Bt genes with dissimilar modes of action for a target pest species.<sup>32</sup>

Based on the literature selection criteria described above, dominance level of resistance to Bt plants was evaluated in a total of 26 studies, which involved 17 cases of major resistance in

**Table 1.** Dominance levels ( $D_{FL}$ ) of 17 cases of major resistance to Bt crops in seven target pest species.

Case of resistance	Field resistance	No. populations	$D_{FL}$	Reference
<b>Resistance to single-gene Bt crops</b>				
<i>B. fusca</i> to Cry1Ab maize in S. Africa	Yes	1	1.56	33
<i>S. frugiperda</i> to Cry1F maize in Brazil	Yes	5	0.23	34–36
<i>S. frugiperda</i> to Cry1F maize in U.S.	Yes	2	0.10	37,38
<i>D. virgifera virgifera</i> to Cry3Bb1 maize in U.S.	Yes	5	0.41	39–41
<i>D. virgifera virgifera</i> to eCry3.1Ab maize in U.S.	Yes	1	1.16	42
<i>S. frugiperda</i> to Cry1A.105 maize in U.S.	Yes	2	0.34	43
<i>O. nubilalis</i> to Cry1F maize in U.S.	No	1	0.04	44
<i>S. frugiperda</i> to Cry2Ab2 maize in U.S.	No	1	–0.02	45
<i>S. frugiperda</i> to Vip3A maize in Brazil	No	1	0.00	46,47
<i>S. frugiperda</i> to Vip3A maize in U.S.	No	1	0.00	48
<i>H. armigera</i> to Cry1Ac cotton in Australia	No	2	0.33	49,50
<i>P. gossypiella</i> to Cry1Ac cotton in U.S.	No	1	0.00	51
<i>D. saccharalis</i> to Cry1Ab maize in U.S.	No	1	0.41	32,52,53
<b>Resistance to dual/multiple-gene Bt crops</b>				
<i>S. frugiperda</i> to Cry1A.105/Cry2Ab maize in Brazil	No	2	0.00	54,55
<i>S. frugiperda</i> to Cry1A.105/Cry2Ab maize in U.S.	No	1	0.20	56,57
<i>S. frugiperda</i> to Cry1Ab/Vip3A maize in Brazil	No	1	0.00	54
<i>S. frugiperda</i> to Cry1A.105/Cry2Ab2/Cry1F in Brazil	No	1	0.00	54,58

seven target insect species to eight Bt proteins in maize or cotton (Table 1 and Supporting information: Table A1). The seven insect species comprised almost all global major target pests of Bt maize and Bt cotton, and the eight Bt proteins included almost all the Bt proteins expressed in Bt crops currently available in the global market. Among the 17 cases, 13 cases were resistant to single-gene Bt crops and the rest four were associated with resistance to dual/multiple-gene Bt crops. Resistance accompanied by field control problems, defined as field resistance mentioned above, has been reported in six of the 13 cases of single-gene resistance. The term ‘field resistance’ in this review means that the resistance has resulted in field control problems of a Bt crop, or significantly reduced susceptibility of individual Bt proteins in pyramided Bt plants, which is similar to the ‘practical resistance’ defined in the reference.<sup>6</sup> Besides the condition of field control problem, the criteria for ‘practical resistance’ also include that >50% of individuals in a population are resistant.<sup>6,60</sup> Because resistance allele frequencies for some cases that qualify as ‘field resistance’ as described above have not been reported, or the rate of the resistant individuals for some cases was <50%, the term ‘field resistance’ is used in this review. The six cases with field resistance are the resistance of *Busseola fusca* to Cry1Ab maize in South Africa<sup>61</sup>; *Spodoptera frugiperda* to Cry1F maize in Brazil<sup>59</sup> and in the U.S.<sup>62,63</sup>; *Diabrotica virgifera virgifera* to Cry3Bb1 maize in the U.S.<sup>64</sup>; *D. virgifera virgifera* to eCry3.1Ab maize in the U.S.<sup>65</sup>; and *S. frugiperda* to Cry1A.105 maize in the U.S.<sup>63,66</sup> The documented high resistance allele frequency in *S. frugiperda* to Cry1A.105 maize and the observed high cross-resistance of the insect between Cry1F and Cry1A.105 maize<sup>63,66</sup> were similar to the results reported in the resistance of *Diatraea saccharalis* to Cry1A.105 maize in Argentina<sup>67</sup> which was listed as a case of ‘practical resistance’ in the reference.<sup>7</sup> Thus, the resistance of *S. frugiperda* to Cry1A.105 maize in the U.S. was also considered a case of field resistance in the current review. Major resistance of the seven cases without field control problems was usually isolated through laboratory selections. Three of the seven laboratory cases were established using massive-selections, while the rest four were isolated with F<sub>2</sub> screen. As mentioned above, these laboratory

selections may not exactly reflect the real situation of the field selections, especially for those cases established from long-term and massive-selections on Bt protein-treated diet.<sup>31</sup> Thus, it is possible that the selection methods might confound the analysis in this review.

On the other hand, fitness costs of resistance to Bt plants have been investigated in a total of 28 studies in the world, which involved 20 cases of major resistance in eight insect species to eight Bt proteins in maize or cotton (Table 2 and Supporting information: Table A2). The eight insect species included all the seven species described above in which the dominance level of resistance has been investigated, plus *Trichoplusia ni*, a secondary target species of Bt cotton in the U. S. In addition, the eight Bt proteins are the same as those evaluated in the studies of dominance levels. Among the 20 cases, 15 cases involved single-gene resistance and five cases were associated with resistance to dual/multiple-gene Bt plants. The 15 cases of single-gene resistance also included all the six cases with field resistance mentioned above, while field resistance has not been documented for all other cases.

### Measurement and calculation of dominance levels of resistance to Bt plants

As described in the reference<sup>14</sup>, dominance of a single gene resistance can be measured in three ways: dominance of insecticide resistance (e.g. D<sub>LC</sub>), which is based on the dose-mortality response curves of RR, RS and SS genotypes; effective dominance (D<sub>ML</sub>), which is based on the mortality levels of the three genotypes at a given toxin concentration; and dominance of relative fitness in the treated area (D<sub>WT</sub>), which is based on the fitness of the three genotypes at a given toxin concentration. These three measurements are related, but they are not the same. Among the three, D<sub>WT</sub> provides the most useful information for resistance management.<sup>14</sup> However, measurement of D<sub>WT</sub> is usually more difficult than measurements of D<sub>LC</sub> and D<sub>ML</sub>. For this reason, most of the early studies of Bt resistance measured only D<sub>LC</sub> or D<sub>ML</sub>. In this study, I extend the methods for calculating the dominance levels of single-gene resistance described in the reference<sup>14</sup> to also include the

**Table 2.** Index of fitness costs (IFCs) of 20 cases of major resistance to Bt crops in eight target pest species.

Case of resistance	Field resistance	No. populations investigated	IFC <sub>RR'</sub> <sup>a</sup>	IFC <sub>RS'</sub>	Reference
<b>Resistance to single-gene Bt crops</b>					
<i>B. fusca</i> to Cry1Ab maize in S. Africa	Yes	1	1.91	n/a	68
<i>S. frugiperda</i> to Cry1F maize in Brazil	Yes	4	0.94	0.98	35,36,54
<i>S. frugiperda</i> to Cry1F maize in U.S.	Yes	3	0.68	0.94	69,70
<i>D. virgifera virgifera</i> to Cry3Bb1 maize in U.S.	Yes	8	1.08	n/a	40,41,71–73
<i>D. virgifera virgifera</i> to eCry3.1Ab maize in U.S.	Yes	1	1.64	n/a	74
<i>S. frugiperda</i> to Cry1A.105 maize in U.S.	Yes	2	1.65	1.99	43
<i>O. nubilalis</i> to Cry1F maize in U.S.	No	1	0.77	0.96	75
<i>S. frugiperda</i> to Cry2Ab2 maize in U.S.	No	1	1.87	2.39	45
<i>S. frugiperda</i> to Vip3A maize in Brazil	No	1	0.80	0.95	46
<i>S. frugiperda</i> to Vip3A maize in U.S.	No	1	1.03	1.06	76
<i>H. armigera</i> to Cry1Ac cotton in Australia	No	2	0.71	0.95	49,77
<i>H. armigera</i> to Cry1Ac cotton in China	No	1	0.77	n/a	78
<i>P. gossypiella</i> to Cry1Ac cotton in U.S.	No	1	0.48	0.52	51
<i>D. saccharalis</i> to Cry1Ab maize in U.S.	No	2	1.28	1.23	32,52,53
<i>T. ni</i> to Cry1Ac cotton in U.S.	No	1	0.94	n/a	79
<b>Resistance to dual/multiple-gene Bt crops</b>					
<i>S. frugiperda</i> to Cry1A.105/Cry2Ab maize in Brazil	No	2	1.00	1.05	54,55
<i>S. frugiperda</i> to Cry1A.105/Cry2Ab maize in U.S.	No	1	0.73	1.15	56,57
<i>T. ni</i> to Cry1Ac/Cry2A cotton in U.S.	No	1	0.81	n/a	79
<i>S. frugiperda</i> to Cry1Ab/Vip3A maize in Brazil	No	1	0.86	1.08	54
<i>S. frugiperda</i> to Cry1A.105/Cry2Ab2/Cry1F in Brazil	No	2	1.00	0.98	54,58

<sup>a</sup>Developmental delay could be a more relevant fitness factor for pests that have multiple generations per cropping cycle than univoltine insects. Six of the 20 cases listed in the table involved the use of the parameter 'insect developmental time' in estimation of fitness costs. Except for the case related to the resistance of *D. virgifera virgifera* to Cry3Bb1 maize, all other five cases were associated with insects having multiple generations per year. For the resistance of the univoltine *D. virgifera virgifera* to Cry3Bb1 maize, two of the five related studies considered 'days of development to adults' in the fitness calculations. Because the developmental time on non-Bt plants reported in the references<sup>40,41</sup> was almost same between RR and SS, IFC<sub>RR'</sub> estimated with/without considering the parameter 'developmental time' was virtually identical (data not shown). To be consistent, the method described in the text was used to calculate IFCs for all the cases regardless of the number of generations per cropping season for the insect evaluated.

cases of resistance to dual/multiple-gene Bt plants. More specifically, the dominance for single- or dual/multiple-gene resistance to Bt plants can be calculated as:

$$D'_{ML} = (M_{R'S'} - M_{S'S'}) / (M_{R'R'} - M_{S'S'}) \text{ or } D'_{WT} \\ = (W_{TR'S'} - W_{TS'S'}) / (W_{TR'R'} - W_{TS'S'})$$

Here,  $D'_{ML}$  is the effective dominance of single- or dual/multiple-gene resistance to Bt plants based on the mortality levels of the three genotypes (R'R', R'S', and S'S') on Bt plants; and  $D'_{WT}$  is the dominance of relative fitness in the treated area based on the fitness of the three genotypes on Bt plants.  $M_{R'R'}$ ,  $M_{R'S'}$ , and  $M_{S'S'}$  are the mortality levels of the single- or dual/multiple-gene homozygous-resistant (R'R'), heterozygous (R'S'), and homozygous-susceptible (S'S') genotypes on the corresponding single- or dual/multiple-gene Bt plants, respectively. For examples, If A, B, and C represent three different resistant alleles and a, b, and c refers to the three corresponding susceptible alleles of the three genes, R'R', R'S', and S'S' represent AA, Aa, and aa for a single-gene resistance; AABB, AaBb, and aabb for a dual-gene resistance; or AABBCC, AaBbCc, and aabbcc for a triple-gene resistance.

The measurement of  $D'_{ML}$  or  $D'_{WT}$  described here can also be used to calculate the dominance levels for other genotypes in dual/multiple-gene resistance (e.g. AABB, AaBB, AABBCC, etc.) as described in reference.<sup>56</sup> Among the 22 studies that evaluated the dominance level of resistance to single-gene Bt crops, five studies measured  $D'_{WT}$ , while the other 17 measured  $D'_{ML}$ s that were based on survivorship of S'S', R'S', and R'R' individuals on whole Bt plants or plant tissues using exposure times from 7 d to a period encompassing neonate-to adult development (Table A1). Among the five studies that evaluated dominance level of dual/multiple-gene resistance to Bt plants, one study evaluated  $D'_{WT}$  and the other four measured  $D'_{ML}$  (Table A1).

In this review, dominance level (functionally) (hereafter referred to as  $D_{FL}$ ) was calculated for each case of single- or dual/multiple-gene resistance to Bt plants based on the values of  $D'_{ML}$  or  $D'_{WT}$  reported in each study. Similarly, as described in the reference<sup>14</sup>,  $D_{FL}$  values normally vary from 0 to 1 ( $D_{FL} = 0$ , functionally completely recessive;  $D_{FL} = 1$ , functionally completely dominant). In the situations in which >1 study was conducted, >1 insect population was evaluated, or >1 trial was

performed for a case, the  $D_{FL}$  for the case was calculated as the average of  $D'_{MLS}$  or  $D'_{WTS}$ , or the mixed  $D'_{MLS}$  and  $D'_{WTS}$  across studies, populations, or trials. Data sources and calculations of  $D_{FL}$ s of the 17 cases are listed in the Supporting Information (Appendix Table A1) linked to this publication.

### Measurement and calculation of index of fitness costs of resistance to Bt plants

To facilitate quantitative analysis of the fitness costs of resistance to Bt plants, a term, index of fitness cost (IFC), is used in this review. IFC for both single- and dual/multiple-gene resistance is calculated using the formula<sup>81</sup>:

$$IFC_{R'R'} = F_{R'R'}/F_{S'S'} \text{ and } IFC_{R'S'} = F_{R'S'}/F_{S'S'}$$

Here  $IFC_{R'R'}$  and  $IFC_{R'S'}$  refer to the index of fitness costs of resistant-homozygous ( $R'R'$ ) and - heterozygous ( $R'S'$ ) genotypes, respectively.  $R'R'$ ,  $R'S'$ , and  $S'S'$  represent the three genotypes as described in the measurement of  $D'_{ML}$  or  $D'_{WT}$ .  $F_{S'S'}$ ,  $F_{R'S'}$ , and  $F_{R'R'}$  refer to the fitness of  $S'S'$ ,  $R'S'$ , and  $R'R'$  genotypes on non-Bt plants or non-Bt plant tissues, respectively.  $IFC < 1$  means that fitness costs are associated with the resistance;  $IFC = 1$  suggests lack of fitness costs; and  $IFC > 1$  indicates that there are fitness advantages. If  $IFC_{R'R'} < 1$  but  $IFC_{R'S'} = 1$  for a resistance, fitness costs are recessive, while if both  $IFC_{R'R'}$  and  $IFC_{R'S'}$  are  $< 1$ , fitness costs are non-recessive. Non-recessive fitness costs are considered more important in resistance management than recessive fitness costs, because  $R'S'$  individuals are usually much more abundant than  $R'R'$  individuals in the absence of Bt selection.<sup>23</sup> Similarly as mentioned for  $D'_{ML}$  or  $D'_{WT}$ , the measurement of IFC described here could also be used to calculate the fitness costs of other genotypes in dual/multiple-gene resistance to Bt crops (e.g.  $AaBB$ ,  $AABbCc$ , etc.).

In the review of the 28 studies, only one<sup>71</sup> reported the IFC value directly, while all others showed various fitness parameters. These biological parameters included insect survivorship with a wide range of exposure period, insect development, growth (e.g. larval and/or pupal body mass), sex ratio, egg production, and egg hatching rate (Supporting Information Appendix Table A2).

One study evaluated fitness for both field-collected parental (F0) and F1 generations.<sup>68</sup> In this review, a 'combined fitness index' was used to measure the fitness ( $F_{S'S'}$ ,  $F_{R'S'}$ , or  $F_{R'R'}$ ) of each insect genotype on plants or plant tissue. Combined  $F_{S'S'}$ ,  $F_{R'S'}$ , or  $F_{R'R'}$  values were calculated based on the most comprehensive measurements reported in each study with the methods described below:

- (a) If the intrinsic rate of population increase,  $r_m$ , was available in a study,  $r_m$  was used as the combined fitness index and no other parameters were considered in IFC calculation for the study.
- (b) If  $r_m$  was not available, but insect survivorship, developmental time, egg production, and egg hatching rate were reported, the combined fitness index was calculated as: (insect survivorship x egg production x egg hatching rate)/insect developmental time. No other parameters were considered in IFC calculation for the study.
- (c) If any of the parameters in the formula described in b was not available, the item for that parameter was excluded in calculation of the combined fitness index.

IFC was calculated for both  $R'R'$  and  $R'S'$  (if data available) for each population in each study. Similarly, as described for the  $D_{FL}$  calculation, in situations in which  $>1$  study was conducted,  $>1$  insect population was evaluated, or  $>1$  trial was performed for a case, the IFC of the case was calculated as the average of IFCs across studies, populations, or trials. Data sources and detailed IFC calculations for the 20 cases are listed in the Supporting Information (Appendix Table A2).

### Dominance, $D_{FL}$ , of Resistance to Bt crops

Analysis of global studies showed that  $D_{FL}$ s of the 13 cases of major resistance to single-gene Bt crops ranged from  $-0.02$  to  $1.56$  with a mean of  $0.35 \pm 0.13$  (Table 1). Among the 13 cases, functionally recessive resistance was reported in only five cases (or 38.5% of the total) with a  $D_{FL}$  of zero or close to zero. These five cases were the resistance of *Ostrinia nubilalis* to Cry1F maize in the U.S. ( $D_{FL} = 0.04$ ), *S. frugiperda* to Cry2Ab2 maize in

the U.S. ( $D_{FL} = -0.02$ ), *S. frugiperda* to Vip3A maize in Brazil and the U.S. ( $D_{FL} = 0$  for both cases), and *Pectinophora gossypiella* to Cry1Ac cotton in the U.S. ( $D_{FL} = 0$ ). To date, field resistance has not been reported for any of these five cases. Resistance in the other eight cases (61.5%) was functionally non-recessive with a  $D_{FL}$  of 0.10 or greater (Table 1).

All of the six cases with field resistance were functionally non-recessive with a  $D_{FL}$  ranging from 0.10 to 1.56 (Table 1). The resistance in two of the six cases was completely or even over-completely dominant: resistance in *B. fusca* to Cry1Ab maize in South Africa, with a  $D_{FL}$  of 1.56, and *D. virgifera virgifera* to eCry3.1Ab maize in the U.S. with a  $D_{FL}$  of 1.16. Resistance in the other four cases of field resistance ranged from incompletely recessive to co-dominant. The mean  $D_{FL}$  for the six cases with field resistance was  $0.63 \pm 0.24$  (mean  $\pm$  sem), while it was  $0.11 \pm 0.07$  for the seven cases without practical field resistance (Table 3). The difference in  $D_{FL}$ s between the cases with and without field resistance was significant (SAS PROC NPAR1WAY Wilcoxon,  $P = .0309$ ) (Table 3). The four cases of dual/multiple-gene resistance in which  $D_{FL}$  has been evaluated involved only the resistance of *S. frugiperda* to Bt maize in Brazil and the U.S. These limited data suggest that dual/multiple-gene resistance is more likely to be recessive than the related single-gene resistance. Three of the four dual/multiple-gene resistance cases were functionally recessive with a  $D_{FL}$  of zero and the remainder was incompletely recessive with a  $D_{FL}$  of 0.20 (Table 1).

As mentioned above, because it is usually difficult to measure  $D'_{WT}$ , 10 of the 13 single-gene resistance cases actually measured only  $D'_{MLS}$ , which is calculated based on mortality only. It is believed that the dominance level can be over-estimated using only  $D'_{ML}$ , because R'S' survivors can

be less fit than R'R' survivors.<sup>14</sup> GM Bt plants are usually very effective against S'S' and thus S'S' individuals rarely survive on Bt crops. In this situation, if R'S' survivors are less fit than R'R' survivors, the actual  $D'_{WT}$  will be lower than  $D'_{ML}$ . However, data from the 13 cases of single-gene resistance analyzed in this review do not provide any evidence to indicate that  $D'_{ML}$  is greater than  $D'_{WT}$ . For example, both  $D'_{ML}$  and  $D'_{WT}$  were estimated for the Brazilian case of Cry1F resistance in *S. frugiperda*. Based on neonate-to-adult survivorships on Cry1F leaf tissue, Farias *et al.*<sup>34</sup> reported a  $D'_{ML}$  of 0.15 for the population BR25R. In another study, Leite *et al.*<sup>35</sup> assessed the dominance levels of two populations (IrmaF and IrmaD) based on a 'fitness index' on Cry1F maize leaf tissue. The 'fitness index' was calculated using the formula, fitness index = (neonate-to-pupal survival  $\times$  pupal weight)/neonate-to-pupal development time. Using this method, the dominance levels for IrmaF and IrmaD were estimated to be 0.36. In addition, Santos-Amaya *et al.*<sup>36</sup> also examined the dominance levels of two other populations (MTH and MRH) on Cry1F plants using the same 'fitness index' as described in reference.<sup>35</sup> The estimated dominance levels for MTH and MRH were 0.12 and 0.17, respectively. Studies have shown that pupal body weight is usually highly correlated to reproduction in many lepidopteran species.<sup>82</sup> Thus, the estimated dominance levels in the four populations evaluated in references<sup>35,36</sup> could be considered a close estimate to the true  $D_{WT}$ . The average dominance level (or  $D'_{WT}$ ) of the four populations was 0.25, which was somewhat greater than the  $D'_{ML}$  (0.15) estimated in reference.<sup>34</sup> In addition, three of the five cases of single-gene resistance that were identified to be completely recessive or nearly completely recessive were based on the measurement of  $D'_{MLS}$ . More importantly, the significantly greater overall  $D_{FL}$  values for the six cases with field resistance, relative

**Table 3.** Comparison of dominance levels ( $D_{FL}$ s) and index of fitness costs (IFC) of single-gene resistance between cases with and without field resistance to Bt crops.

Resistance status	Dominance level		Index of fitness costs			
	No. case	$D_{FL}$	R'R'		R'S'	
			No. case	IFC <sub>R'R'</sub>	No case	IFC <sub>R'S'</sub>
Cases with field resistance occurred	6	$0.63 \pm 0.24$	6	$1.32 \pm 0.20$	3	$1.30 \pm 0.34$
Cases with field resistance not occurred yet	7	$0.11 \pm 0.07$	9	$0.96 \pm 0.14$	7	$1.15 \pm 0.22$
Wilcoxon non-parametric test	$P = .0309$		$P = .1941$		$P = 1.000$	

to the seven cases without field resistance, are particularly telling and document that  $D_{FLS}$  estimated in these studies were closely correlated to the resistance evolution in the field.

There were a few cases in which  $D_{FL}$  of a case have been evaluated for multiple populations. In some cases,  $D_{FL}$  values among populations within a case were consistent. For example, the resistance of *S. frugiperda* to Cry1F maize was incompletely recessive in all five Brazilian populations examined (Table A1). Similarly, the resistance of *D. virgifera virgifera* to Cry3Bb1 maize in five U.S. populations was all incompletely recessive or codominant with  $D_{FL}$  values ranging from 0.27 to 0.59 (Table A1). However, in some cases, notable variations were observed. For example, the resistance of *Helicoverpa armigera* to Cry1Ac cotton in Australia was completely recessive on 4-week old cotton,<sup>49</sup> while it was incompletely dominant on 14-week cotton.<sup>50</sup> Variations in  $D_{FLS}$  among populations in a case were also observed for the resistance of *S. frugiperda* to Cry1F maize and Cry1A.105 maize in the U.S. (Table A1).<sup>37,38,43</sup> In addition, differences in  $D_{FLS}$  were noted among cases of a same pest-Bt crop system, but the differences were relatively small. For example, both cases of *S. frugiperda* resistance to Cry1F maize in Brazil and the U.S. were incompletely recessive. Similarly, both cases of resistance to Vip3A maize were completely recessive in the two countries. In contrast, variation in  $D_{FLS}$  for a target pest species appeared to be greater among different Bt protein-crop systems. For example,  $D_{FLS}$  of *S. frugiperda* resistance varied from -0.02 on Cry2Ab2 maize to 0.34 on Cry1A.105 maize, and  $D_{FLS}$  of *D. virgifera virgifera* resistance differed from 0.41 on Cry3Bb1 maize to 1.16 on eCry3.1Ab maize. Nevertheless, the published data showed that all three cases of resistance associated with maize plants expressing the Vip3A protein were completely recessive with a  $D_{FL}$  of zero which provides evidence that GM plants containing Vip3A gene most likely produce the necessary 'high dose' as required for the HDR strategy. The observed variation in  $D_{FLS}$  among populations or among cases within the same species could be due to genetic differences in resistant genes

and/or differences in test conditions, such as differences in plant growth stages or tissues used in bioassays. Thus, experiments testing with multiple insect populations under different environmental conditions are necessary in order to generate robust  $D_{FLS}$ .

It should be pointed out that the estimated dominance of resistance for dual/multiple-gene resistance to Bt plants may not only reflect the inheritance of survival/fitness to each Bt protein in a pyramid, but also can be associated with the interactions of different Bt proteins in plants, such as effects of cross-resistance and the extent of redundant killing. Exploring such relations in detail is beyond the scope of this review. Nevertheless, information on the dominance of dual/multiple-gene resistance to Bt plants should also be useful in resistance management as for the single-gene resistance. Additional studies are necessary to analyze the dominance levels of dual-/multiple-gene resistance to Bt plants. However, the lower dominance levels observed from the limited cases of dual-/multiple-gene resistance relative to single-gene resistance are an encouraging sign for the use of pyramided Bt crop traits for IRM.<sup>10</sup> The results suggest that pyramiding with dissimilar Bt proteins could make a non-recessive resistance to single-gene Bt plants functionally more recessive. In the U.S., single-gene Bt cotton has already been completely phased out of the market and replaced by pyramided varieties. Pyramided Bt maize was first commercialized in 2010 and since then pyramided Bt maize traits have been widely planted in the U.S. and several other countries. However, individual Bt proteins in all current pyramided crop traits have been used sequentially. In the sequential use of Bt proteins, there is possibility as only one active Bt gene being introduced in each 'new pyramided trait' if the target insects already become resistant to all other Bt proteins after being used for many years. In such cases, a new 'pyramided trait' essentially functionally just like a single-gene trait, which would dramatically reduce the effectiveness of pyramiding for IRM.<sup>6,81,83-86</sup> It is believed that sequential use of Cry1 than Cry1 + Cry2 proteins in Bt maize and cotton could be a key factor that has contributed to the recent widespread occurrence of the field




resistance of *H. zea* to pyramided Cry1A/Cry2A maize and cotton in the U.S. and the field resistance of *P. gossypiella* to Cry1A/Cry2A cotton in India.<sup>80</sup>

### Fitness costs, IFCs, of resistance to Bt crops

Global data analysis of the 28 studies showed that only six (or 40.0% of the total) of the 15 cases of major resistance to single-gene Bt crops were likely associated with fitness costs (Table 2). These six cases were the resistance of *S. frugiperda* to Cry1F maize in the U.S. ( $IFC_{R'R'} = 0.68$ ), *O. nubilalis* to Cry1F maize in U.S. ( $IFC_{R'R'} = 0.77$ ), *S. frugiperda* to Vip3A maize in Brazil ( $IFC_{R'R'} = 0.80$ ), *H. armigera* to Cry1Ac cotton in Australia ( $IFC_{R'R'} = 0.71$ ) and China ( $IFC_{R'R'} = 0.77$ ), and *P. gossypiella* to Cry1Ac cotton in U.S. ( $IFC_{R'R'} = 0.48$ ) (Table 2). Five cases (33.3%) of single-gene resistance showed some level of fitness advantage; these were the resistance of *B. fusca* to Cry1Ab maize in South Africa ( $IFC_{R'R'} = 1.91$ ), *D. virgifera virgifera* to eCry3.1Ab maize in the U.S. ( $IFC_{R'R'} = 1.64$ ), *S. frugiperda* to Cry1A.105 ( $IFC_{R'R'} = 1.65$ ) and Cry2Ab2 ( $IFC_{R'R'} = 1.87$ ) maize in the U.S., and *D. saccharalis* to Cry1Ab maize in the U.S. ( $IFC_{R'R'} = 1.28$ ). The rest of the four cases (33.3%) exhibited  $IFC_{R'R'}$  values from 0.94 to 1.08, indicating lack of fitness costs or advantages. The 15 cases of

single-gene resistance had a mean  $IFC_{R'R'}$  of  $1.10 \pm 0.12$  (mean  $\pm$  sem) and the  $IFC_{R'R'}$  ( $1.32 \pm 0.20$ ) of the six cases with field resistance was statistically similar to that ( $0.96 \pm 0.14$ ) of the nine cases without field resistance (SAS PROC NPAR1WAY Wilcoxon,  $P = .1941$ ) (Table 3).

In addition, fitness costs of R'S' were also evaluated for 10 of the 15 single-gene resistance cases, which included three cases with field resistance and seven cases without field resistance (Table 2). Fitness costs of R'S' were clearly observed in only one of the ten cases, which was the resistance of *P. gossypiella* to Cry1Ac cotton in the U.S. with an  $IFC_{R'S'}$  of 0.51. In contrast, R'S' individuals in three cases had a greater fitness than SS individuals; these cases were the resistance of *S. frugiperda* to Cry1A.105 ( $IFC_{R'S'} = 1.99$ ) and Cry2Ab2 ( $IFC_{R'S'} = 2.39$ ) maize and *D. saccharalis* to Cry1Ab maize ( $IFC_{R'S'} = 1.23$ ) in the U.S.  $IFC_{R'S'}$  values of the remaining six cases ranged from 0.94 to 1.06, suggesting lack of fitness costs for R'S'. The mean  $IFC_{R'S'}$  ( $1.30 \pm 0.34$ ) of the three cases with field resistance was not different compared to that ( $1.15 \pm 0.22$ ) of the seven cases without field resistance (SAS PROC NPAR1WAY Wilcoxon,  $P = 1.000$ ) (Table 3).  $IFC_{R'S'}$  is not independent of  $IFC_{R'R'}$ ; analysis of the ten cases in which both R'R' and R'S' were available showed a strong linear relationship between the two indices ( $IFC_{R'S'} = -0.035 + 1.207 IFC_{R'R'}$ ;  $R^2 = 0.9329$ ,



**Figure 1.** Correlation analysis on index of fitness costs (IFC) of ten single-gene major resistance cases to single-gene Bt plants between resistant-homozygous (R'R') and -heterozygous (R'S') genotypes. Analysis was performed by treating the index of fitness costs for R'R' ( $IFC_{R'R'}$ ) of a case as the independent variable and the index of fitness costs for R'S' ( $IFC_{R'S'}$ ) of the case as the dependent variable.

$P < .0001$ ) (Figure 1). In addition, a paired  $t$ -test with ‘case’ as the subject factor also showed that the mean  $IFC_{R'S'}$  ( $1.20 \pm 0.18$ ) was significantly greater than the  $IFC_{R'R'}$  ( $1.02 \pm 0.14$ ) for the ten cases ( $t = -3.23$ ,  $df = 9$ ,  $P = .0103$ ).

Fitness costs of dual/multiple-gene resistance were evaluated for only five cases and involved only two pest species, *S. frugiperda* and *T. ni* (Table 2). Fitness costs were likely associated with R'R' in three of the five cases, which were the resistance of *S. frugiperda* to Cry1A.105/Cry2Ab2 maize in the U.S. ( $IFC_{R'R'} = 0.73$ ) and to Cry1Ab/Vip3A maize in Brazil ( $IFC_{R'R'} = 0.86$ ), and *T. ni* to Cry1Ac/Cry2A cotton in the U.S. ( $IFC_{R'R'} = 0.81$ ). Fitness costs were not associated with the remaining two cases ( $IFC_{R'R'} = 1.0$  for both cases). The mean  $IFC_{R'R'}$  for the five cases of dual-/multiple-gene resistance was  $0.88 \pm 0.05$ .  $IFC_{R'S'}$  has been evaluated for four of the five cases of dual-/multiple-gene resistance and none of the four cases showed any fitness costs with an  $IFC_{R'S'}$  ranging from 0.98 to 1.15 and an average of  $1.07 \pm 0.04$  (Table 2).

Fitness costs of resistance can vary depending on Bt protein, crop, insect species and population, test conditions, etc.<sup>23</sup> As observed for the  $D_{FLS}$ , variations in IFCs were also reported among populations and cases within the same pest-crop system. For example, in the Brazilian Cry1F-resistant populations, fitness costs were likely associated with the MRH population on non-Bt maize in the study<sup>36</sup> and a population on non-Bt cotton evaluated in the reference<sup>54</sup>, while lack of fitness costs was observed for other populations and test conditions (Table A1). Similarly, notable fitness costs were observed in two U.S. Cry3Bb1-resistant populations of *D. virgifera virgifera*, but lack of fitness costs was detected for other populations (Table A1). In addition, fitness costs were observed in the resistance of *S. frugiperda* to Vip3A maize for both the Brazilian and U.S. populations on non-Bt maize plants, but not for the U.S. population on non-Bt cotton.<sup>46,76</sup> In contrast, in some cases, fitness costs were consistent among populations within a pest-crop system. For examples, all of the three U.S. Cry1F-resistant populations of *S. frugiperda* exhibited some level of fitness costs (Table A1). Considerable fitness costs were also observed in the two Australian Cry1Ac-resistant populations of *H. armigera*. In addition, both U.S. Cry1A.105-resistant populations of *S. frugiperda* studied in the reference<sup>43</sup> performed similarly on non-Bt maize and both did not show any fitness costs.

The significant number (33.3%) of cases of single-gene resistance with fitness advantages observed in this review is a surprise. Fitness advantage of Bt resistance has been supposed to be very rare.<sup>87</sup> However, it is also believed that selection for Bt resistance could be linked to some genes that are favorable for insect growth and development.<sup>72,87</sup> In addition, some of the comparisons might involve the use of unrelated resistant and susceptible strains that may have differed for reasons unrelated to resistance. Unfortunately the available data listed in Tables 1 and Tables 2 could not clarify the situation. Further studies are warranted to understand the biological mechanisms or other factors behind the observed ‘fitness advantages’ of Bt resistance. The greater IFC for RS than RR in the ten cases of single-gene resistance is not surprised, because the fitness costs in four of five cases were recessive. In addition, in the two cases of *S. frugiperda* resistance to Cry1A.105 and Cry2Ab2 maize in which the resistance showed a fitness advantage ( $IFC_{RR} > 1$ ), the  $IFC_{RSS}$  were still somewhat greater than the corresponding  $IFC_{RRS}$  (Table 2). A possible reason for the greater performance of the RS relative to both RR and SS may be hybrid vigor, which could occur when crossing two populations that were inbred and had different genetic backgrounds.<sup>23</sup> Differences between resistant and susceptible strains of an insect species that are unrelated to the Bt resistance could be caused by many factors such as sources of insect strains, lab adaptation, diet adaptation, or isolation in resistance selections. To ensure a similar genetic background between SS and RR, the RR populations used in the two studies had been backcrossed with SS for at least two times and then reselected for resistance before they were used for the crosses to generate RS genotypes. If the better performance of the RS genotypes reported in the two cases was truly caused by hybrid vigor, it suggests that additional backcrossing would be necessary to ensure a more similar genetic background between RR and SS to avoid any possible confounding effect of ‘hybrid vigor’. The use of a susceptible comparator with different genetic bases could undermine the importance of fitness costs in the field.<sup>23</sup> In addition, choice of susceptible insect strain could also affect estimation of dominance because hybrid vigor would artificially inflate the apparent fitness

of heterozygotes on Bt plants, as well as on non-Bt plants, and therefore could affect the calculated functional dominance values. However, a linear regression analysis (SAS PROC REG) with the ten cases that both  $IF_{RS}$  ( $x$ ) and  $D_{FL}$  ( $y$ ) data are available failed to find any linkage between the fitness advantage of RS and the functional dominance levels of the resistance ( $y = 0.106 + 0.031 x$ , correlation coefficient  $R = 0.10$  ( $P = .7762$ )). Nevertheless, as described above, it is critical in study of fitness costs and dominance levels of resistance to ensure a similar genetic basis among insect populations. A common method used to achieve similar genetic bases is to backcross the resistant populations to their susceptible comparator and reselect the resistance in the backcrossed populations. Theoretically, the similarity in genetic background among insect populations increases as the number of backcrosses increases.

## Conclusion

It should be noted that major resistance genes for some insect-Bt crop systems, such as resistance of *O. nubilalis* to Cry1Ab maize, have not been identified yet. However, the 13 (for  $D_{FL}$ ) and 15 (for IFC) cases of major resistance to single-gene Bt crops analyzed in this review have included almost all the global major target pest species and all Bt proteins expressed in the world market of GM Bt crops. The surprisingly high rates of functionally non-recessive resistance (61.5%) and lack of fitness costs (60.0%) of resistance reported in this review clearly documented that high levels of resistance to Bt crops are usually non-recessive with no fitness costs. Use of insect populations with similar genetic background is critical in study of fitness costs and dominance levels of resistance. Otherwise, use of a susceptible comparator with different genetic background could undermine the importance of fitness costs in the field. Limited available data suggest that dual/multiple-gene Bt resistance is likely to be more recessive than the related single-gene resistance, but their IFCs are similar. Many factors can influence the speed of resistance development, but the documentations that all six cases of field resistance are functionally non-recessive, as well as the significantly greater  $D_{FL}$ s of the cases with field resistance than those without field resistance provide

clear evidence that the prevalence of non-recessive resistance has certainly played an essential role in the widespread occurrence of field resistance to Bt crops. In addition, the documented high rate of non-recessive resistance also provides solid counterevidence against a general application of the assumption of functionally recessive resistance for the recommended HDR strategy, at least for single-gene Bt crops. The lack of fitness costs might be associated with the widespread of the field resistance. However, the similar IFCs observed between the cases with and without field resistance suggest that the role of the lack of fitness costs is apparently not as critical as the non-recessive resistance. Information generated from this review suggests that planting of 'high dose' traits is an effective method for Bt crop IRM and more comprehensive management strategies that are also effective for functionally non-recessive resistance should be deployed.

## Acknowledgments

The author thanks Drs. Mike Stout, Jian-Zhou Zhao, and Ying Niu for reviewing an earlier draft of the manuscript. This publication has been approved by the Associate Vice President & Program Leader of the Louisiana State University Agricultural Center as manuscript No. 2020-234-34269. This study was sponsored by Hatch funds from the USDA National Institute of Food and Agriculture, and the USDA Multistate Research Project NC-246.

## Disclosure of potential conflict of interest

I have read the journal's policy and the author declares no competing financial interests.

## ORCID

Fangneng Huang  <http://orcid.org/0000-0003-0956-1958>

## References

1. ISAAA. Global status of commercialized biotech/GM crops in 2018: biotech crops continue to help meet the challenges of increased population and climate change. Ithaca (NY); ISAAA Brief 2018; No. 54.
2. Wu KM, Lu YH, Feng HQ, Jiang YY, Zhao JZ. Suppression of cotton bollworm in multiple crops in China in areas with Bt toxin-containing cotton. *Science*. 2008;321:1676–78. doi:10.1126/science.1160550.

3. Hutchison WD, Burkness EC, Mitchell PD, Moon RD, Leslie TW, Fleischer SJ, Abrahamson M, Hamilton KL, Steffey KL, Gray ME, et al. Areawide suppression of European corn borer with Bt maize reaps savings to non-Bt maize growers. *Science*. 2010;330:222–25. doi:10.1126/science.1190242.
4. Kathage J, Qaim M. Economic impacts and impact dynamics of Bt (*Bacillus thuringiensis*) cotton in India. *Proc Natl Acad Sci USA*. 2012;109:11652–56. doi:10.1073/pnas.1203647109.
5. Dively GP, Venugopal PD, Bean D, Whalen J, Holmstrom K, Kuhar TP, Doughty HB, Patton T, Cissel W, Hutchison WD. Regional pest suppression associated with widespread Bt maize adoption benefits vegetable growers. *Proc Natl Acad Sci USA*. 2018;115:3320–25. doi:10.1073/pnas.1720692115.
6. Tabashnik BE, Carrière Y. Surge in insect resistance to transgenic crops and prospects for sustainability. *Nat Biotech*. 2017;35:926–35. doi:10.1038/nbt.3974.
7. Tabashnik BE, Carrière Y. Global patterns of resistance to Bt crops highlighting pink bollworm in the United States, China, and India. *J Econ Entomol*. 2019;112:2513–23. doi:10.1093/jee/toz173.
8. Smith JL, Farhan Y, Schaafsma AW. Practical resistance of *Ostrinia nubilalis* (Lepidoptera: crambidae) to Cry1F *Bacillus thuringiensis* maize discovered in Nova Scotia. *Can Sci Rep*. 2019;9:18247. doi:10.1038/s41598-019-54263-2.
9. Ostlie KR, Hutchison WD, Hellmich RL, editors. Bt-Corn & European corn borer, Long term success through resistance management. St. Paul: North Central Region Extension Publication 602. University of Minnesota; 1997.
10. Matten SR, Frederick RJ, Reynolds AH. United states environmental protection agency insect resistance management programs for plant-incorporated protectants and use of simulation modeling. In: Wozniak CA, McHughen A editors. Regulation of agricultural biotechnology: the United States and Canada. Springer; New York, USA: 2012. p. 175–267.
11. Gould F. Sustainability of transgenic insecticidal cultivars: integrating pest genetics and ecology. *Ann Rev Entomol*. 1998;43:701–26. doi:10.1146/annurev.ento.43.1.701.
12. Tabashnik BE. Evolution of resistance to *Bacillus thuringiensis*. *Ann Rev Entomol*. 1994;39:47–79. doi:10.1146/annurev.en.39.010194.000403.
13. Alstad D, Andow DA. Managing the evolution of insect resistance to transgenic plants. *Science*. 1995;268:1894–96. doi:10.1126/science.268.5219.1894.
14. Bourguet D, Genissel A, Raymond M. Insecticide resistance and dominance levels. *J Econ Entomol*. 2000;93:1588–95. doi:10.1603/0022-0493-93.6.1588.
15. Metz TD, Roush RT, Tang JD, Shelton AM, Earle ED. Transgenic broccoli expressing a *Bacillus thuringiensis* insecticidal crystal protein: implications for pest resistance management strategies. *Mol Breed*. 1995;1:309–17. doi:10.1007/BF01248408.
16. Huang F, Buschman LL, Higgins RA, McGaughy WH. Inheritance of resistance to *Bacillus thuringiensis* toxin (Dipel ES™) in European corn borer. *Science*. 1999;284:965–67. doi:10.1126/science.284.5416.965.
17. Huang F, Andow DA, Buschman LL. Success of the high-dose/refuge resistance management strategy after 15 years of Bt crop use in North America. *Entom Exp App*. 2011;140:1–16. 10.1111/j.1570-7458.2011.01138.x
18. Roush RT. Two-toxin strategies for management of insecticidal transgenic crops: can pyramiding succeed where pesticide mixtures have not? *Phil Trans R Soc Lond B*. 1998;353:1777–86. doi:10.1098/rstb.1998.0330.
19. Zhao JZ, Cao J, Li Y, Collins HL, Roush RT, Earle ED, Shelton AM. Transgenic plants expressing two *Bacillus thuringiensis* toxins delay insect resistance evolution. *Nat Biotech*. 2003;21:1493–97. doi:10.1038/nbt907.
20. Carriere Y, Crickmore N, Tabashnik BE. Optimizing pyramided transgenic Bt crops for sustainable pest management. *Nat Biotech*. 2015;33:161–68. doi:10.1038/nbt.3099.
21. Carriere Y, Tabashnik BE. Reversing insect adaptation to transgenic insecticidal plants. *Proc R Soc London B*. 2001;268:1475–80. doi:10.1098/rspb.2001.1689.
22. Gould F, Cohen MB, Bentur JS, Kennedy GG, VanDuyn J. Impact of small fitness costs on pest adaptation to crop varieties with multiple toxins: a heuristic model. *J Econ Entomol*. 2006;99:2091–99. doi:10.1093/jee/99.6.2091.
23. Gassmann AJ, Carrière Y, Tabashnik BE. Fitness costs of insect resistance to *Bacillus thuringiensis*. *Ann Rev Entomol*. 2009;54:147–63. doi:10.1146/annurev.ento.54.110807.090518.
24. Ferré J, van Rie J. Biochemistry and genetics of insect resistance to *Bacillus thuringiensis*. *Ann Rev Entomol*. 2002;47:501–33. doi:10.1146/annurev.ento.47.091201.145234.
25. Andow DA. The risk of resistance evolution in insects to transgenic insecticidal crops. *ICGEB Collect Biosaf Rev*. 2008;4:142–99.
26. Huang F, Leonard BR, Andow DA. Sugarcane borer resistance to transgenic *Bacillus thuringiensis*-maize. *J Econ Entomol*. 2007;100:164–71. 10.1603/0022-0493(2007)100[164:SBLCRT]2.0.CO;2.
27. Tabashnik BE, Gassmann AJ, Crowder DW, Carrière Y. Insect resistance to Bt crops: evidence versus theory. *Nat Biotech*. 2008;26:199–202. doi:10.1038/nbt1382.
28. Tabashnik BE, Van Rensburg JBJ, Carrière Y. Field-evolved insect resistance to Bt crops: definition, theory, and data. *J Econ Entomol*. 2009;102:2011–25. doi:10.1603/029.102.0601.
29. Tabashnik BE, Brévault T, Carrière Y. Insect resistance to Bt crops: lessons from the first billion acres. *Nat Biotech*. 2013;31:510–21. doi:10.1038/nbt.2597.
30. Tabashnik BE, van Bensburg JBJ, Carrière Y. Field-evolved insect resistance to Bt crops: definition, theory, and data. *J Econ Entomol*. 2019;102:2011–25.

31. National Research Council. Environmental effects of transgenic plants: the scope and adequacy of regulation. Vol. 320. Washington (DC): National Academy; 2002.
32. Ghimire MN, Huang F, Leonard RB, Head GP, Yang Y. Susceptibility of Cry1Ab-susceptible and -resistant sugarcane borer to transgenic corn plants containing single or pyramided *Bacillus thuringiensis* genes. *Crop Prot.* 2011;30:74–81. doi:10.1016/j.cropro.2010.09.008.
33. Campagne P, Kruger M, Pasquet R, le Ru B. and van den Berg J. Dominant inheritance of field-evolved resistance to Bt Corn in *Busseola fusca*. *PLoS ONE.* 2013;8:e69675. doi:10.1371/journal.pone.0069675.
34. Farias JR, Andow DA, Horikoshi RJ, Sorgatto RJ, Dos Santos AC, Omoto C. Dominance of Cry1F resistance in *Spodoptera frugiperda* (Lepidoptera: noctuidae) on TC1507 Bt maize in Brazil. *Pest Manag Sci.* 2016;72:974–79.
35. Leite NA, Mendes SM, Santos-Amaya OF, Santos CA, Teixeira TP, Guedes RN, Pereira EJ. Rapid selection and characterization of Cry1F resistance in a Brazilian strain of fall armyworm. *Entomol Exp Appl.* 2016;158:236–47. doi:10.1111/eea.12399.
36. Santos-Amaya OF, Tavares CS, Monteiro HM, Teixeira TP, Guedes RN, Alves AP, Pereira EJ. Genetic basis of Cry1F resistance in two Brazilian populations of fall armyworm, *Spodoptera frugiperda*. *Crop Prot.* 2016;81:154–62. doi:10.1016/j.cropro.2015.12.014.
37. Niu Y, Meagher JRL, Yang F, Huang F. Susceptibility of field populations of the fall armyworm (Lepidoptera: noctuidae) from Florida and Puerto Rico to purified Cry1F protein and corn leaf tissue containing single and pyramided Bt genes. *Florida Entomol.* 2013;96:701–13.
38. Camargo AM, Castañera P, Farinós GP, Huang F. Comparative analysis of the genetic basis of Cry1F resistance in two strains of *Spodoptera frugiperda* originated from Puerto Rico and Florida. *J Invertebr Pathol.* 2017;146:47–52. doi:10.1016/j.jip.2017.03.013.
39. Petzold-Maxwell JL, Cibils-Stewart X, French BW, Gassmann AJ. Adaptation by western corn rootworm (Coleoptera: chrysomelidae) to Bt maize: inheritance, fitness costs, and feeding preference. *J Econ Entomol.* 2012;105:1407–18. doi:10.1603/EC11425.
40. Ingber DA, Gassmann AJ. Inheritance and fitness costs of resistance to Cry3Bb1 corn by western corn rootworm (Coleoptera: chrysomelidae). *J Econ Entomol.* 2015;108:2421–32. doi:10.1093/jee/tov199.
41. Paolino AR, Gassmann AJ. Assessment of inheritance and fitness costs associated with field-evolved resistance to Cry3Bb1 maize by western corn rootworm. *Toxins.* 2017;9:159.
42. Geisert RW, Ellersieck MR, Hibbard BE. Tolerance of eCry3.1Ab in reciprocal cross offspring of eCry3.1Ab-selected and control colonies of *Diabrotica virgifera virgifera* (Coleoptera: chrysomelidae). *J Econ Entomol.* 2016;109:815–20. doi:10.1093/jee/tov335.
43. Niu Y, Head GP, Price PA, Huang F. Inheritance and fitness costs of Cry1A.105 resistance in two strains of *Spodoptera frugiperda* (J.E. Smith). *Crop Prot.* 2018;110:29–235. doi:10.1016/j.cropro.2017.06.022.
44. Pereira EJG, Storer NP, Siegfried BD. Inheritance of Cry1F resistance in laboratory-selected European corn borer and its survival on transgenic corn expressing the Cry1F toxin. *Bull Entomol Res.* 2008;98:621–29. doi:10.1017/S0007485308005920.
45. Acharya B, Head G, Price P, Huang F. Inheritance and fitness costs of Cry2Ab2 resistance in *Spodoptera frugiperda*. *J Invertebr Pathol.* 2017;149:8–14. doi:10.1016/j.jip.2017.06.014.
46. Bernardi O, Bernardi D, Horikoshi RJ, Okuma DM, Miraldo LL, Fatoletto J, Medeiros FCL, Burd T, Omoto C. Selection and characterization of resistance to the Vip3Aa20 protein from *Bacillus thuringiensis* in *Spodoptera frugiperda*. *Pest Manag Sci.* 2016;72:1794–802. doi:10.1002/ps.4223.
47. Miraldo LL, Bernardi O, Horikoshi RJ, Bernardi FSAAD, Omoto C. Functional dominance of different aged larvae of Bt-resistant *Spodoptera frugiperda* (Lepidoptera: noctuidae) on transgenic maize expressing Vip3Aa20 protein. *Crop Prot.* 2016;88:65–71.
48. Yang F, Morsello S, Head GP, Huang F, Gilreath RT, Kerns DL. F2 screen, inheritance and cross-resistance of field-derived Vip3A resistance in *Spodoptera frugiperda* (Lepidoptera: noctuidae) collected from Louisiana, USA. *Pest Manag Sci.* 2018;74:1769–78. doi:10.1002/ps.4805.
49. Bird LJ, Akhurst RJ. Relative fitness of Cry1A-resistant and -susceptible *Helicoverpa armigera* (Lepidoptera: noctuidae) on conventional and transgenic cotton. *J Econ Entomol.* 2004;97:1699–709. doi:10.1603/0022-0493-97.5.1699.
50. Bird LJ, Akhurst RJ. Fitness of Cry1A-resistant and -susceptible *Helicoverpa armigera* (Lepidoptera: noctuidae) on transgenic cotton with reduced levels of Cry1Ac. *J Econ Entomol.* 2005;98:1311–19. doi:10.1603/0022-0493-98.4.1311.
51. Liu YB, Tabashnik BE, Dennehy TJ, Patin AL, Sims MA, Meyer SK, Carrière Y. Effects of Bt cotton and Cry1Ac toxin on survival and development of pink bollworm (Lepidoptera: gelechiidae). *J Econ Entomol.* 2001;94:1237–42. doi:10.1603/0022-0493-94.5.1237.
52. Wu X, Huang F, Leonard BR, Moore SH. Evaluation of transgenic *Bacillus thuringiensis* corn hybrids against Cry1Ab-susceptible and -resistant sugarcane borer (Lepidoptera: crambidae). *J Econ Entomol.* 2007;100:1880–86. doi:10.1093/jee/100.6.1880.
53. Wangila DS, Leonard BR, Bai Y, Head G, Huang F. Larval survival and plant injury of Cry1Ab-susceptible, -resistant, and -heterozygous genotypes of the sugarcane borer on transgenic corn containing single or pyramided Bt genes. *Crop Prot.* 2012;42:108–15. doi:10.1016/j.cropro.2012.06.005.

54. Horikoshi RJ, Bernardi D, Bernardi O, Malaquias JB, Okuma DM, Miraldo LL, Amaral FS, Omoto C. Effective dominance of resistance of *Spodoptera frugiperda* to Bt maize and cotton varieties: implications for resistance management. *Sci Rep.* 2016;6:34864. doi:10.1038/srep34864.
55. Santos-Amaya OF, Rodrigues JV, Souza TC, Tavares CS, Campos SO, Guedes RN, Pereira EJ. Resistance to dual-gene Bt maize in *Spodoptera frugiperda*: selection, inheritance, and cross-resistance to other transgenic events. *Sci Rep.* 2015;5:18243. doi:10.1038/srep18243.
56. Niu Y, Guo J, Head GP, Price PA, Huang F. Phenotypic performance of nine genotypes of Cry1A.105/Cry2Ab2-dual gene resistant fall armyworm on non-Bt and MON 89034 maize. *Pest Manag Sci.* 2019;75:2124–32. doi:10.1002/ps.5331.
57. Zhu C, Niu Y, Zhou Y, Guo J, Head GP, Price PA, Wen X, Huang F. Survival and effective dominance level of a Cry1A.105/Cry2Ab2-dual gene resistant population of *Spodoptera frugiperda* (J.E. Smith) on common pyramided Bt corn traits. *Crop Prot.* 2019;115:84–91. doi:10.1016/j.cropro.2018.09.008.
58. Bernardi D, Bernardi O, Horikoshi RJ, Salmeron E, Okuma DM, Farias JR, Do Nascimento ARB, Omoto C. Selection and characterization of *Spodoptera frugiperda* (Lepidoptera: noctuidae) resistance to MON89034 × TC1507 × NK603 maize technology. *Crop Prot.* 2017;94:64–68. doi:10.1016/j.cropro.2016.11.026.
59. Farias JR, Andow DA, Horikoshi RJ, Sorgatto RJ, Fresia P, Santos AC, Omoto C. Field-evolved resistance to Cry1F maize by *Spodoptera frugiperda* (Lepidoptera: noctuidae) in Brazil. *Crop Prot.* 2014;64:150–58. doi:10.1016/j.cropro.2014.06.019.
60. Tabashnik BE, Mota-Sanchez D, Whalon ME, Hollingworth RM, Carrière Y. Defining terms for proactive management of resistance to Bt crops and pesticides. *J Econ Entomol.* 2014;107:496–507. doi:10.1603/EC13458.
61. van Rensburg JBJ. First report of field resistance by the stem borer, *Busseola fusca* (Fuller) to Bt-transgenic maize. *S Afr J Plant Soil.* 2007;24:147–51. doi:10.1080/02571862.2007.10634798.
62. Storer NP, Babcock JM, Schlenz M, Meade T, Thompson GD, Bing JW, Huckaba RM. Discovery and characterization of field resistance to Bt maize: *spodoptera frugiperda* in Puerto Rico. *J Econ Entomol.* 2010;103:1031–38. doi:10.1603/EC10040.
63. Huang F, Qureshi JA, Meagher JRL, Reisig DD, Head GP, Andow DA, Ni X, Kerns D, Buntin GD, Niu Y, et al. Cry1F resistance in fall armyworm *Spodoptera frugiperda*: single gene versus pyramided Bt maize. *PLoS ONE.* 2014;9(11):e112958. doi:10.1371/journal.pone.0112958.
64. Gassmann AJ, Petzold-Maxwell JL, Keweshan RS, Dunbar MW. Field-evolved resistance to Bt maize by western corn rootworm. *PLoS ONE.* 2011;6(7):e22629. doi:10.1371/journal.pone.0022629.
65. Jakka SRK, Shrestha RB, Gassmann AJ. Broad-spectrum resistance to *Bacillus thuringiensis* toxins by western corn rootworm (*Diabrotica virgifera virgifera*). *Sci Rep.* 2016;6:27860. doi:10.1038/srep27860.
66. Huang F, Qureshi JA, Head GP, Price PA, Levy R, Yang F, Niu Y. Frequency of *Bacillus thuringiensis* Cry1A.105 resistance alleles in field populations of the fall armyworm, *Spodoptera frugiperda*, in Louisiana and Florida. *Crop Prot.* 2016;83:83–89. doi:10.1016/j.cropro.2016.01.019.
67. Grimi DA, Parody B, Ramos ML, Machado M, Ocampo F, Willse A, Martinelli S, Head G. Field-evolved resistance to Bt maize in sugarcane borer (*Diatraea saccharalis*) in Argentina. *Pest Manag Sci.* 2018;74:905–13. doi:10.1002/ps.4783.
68. Kruger M, Van Rensburg JBJ, Van den Berg J. No fitness costs associated with resistance of *Busseola fusca* (Lepidoptera: noctuidae) to genetically modified Bt maize. *Crop Prot.* 2014;55:1–6. doi:10.1016/j.cropro.2013.09.004.
69. Jakka SRK, Knight VR, Jurat-Fuentes JL. Fitness costs associated with field-evolved resistance to Bt maize in *Spodoptera frugiperda* (Lepidoptera: noctuidae). *J Econ Entomol.* 2014;107:342–51. doi:10.1603/EC13326.
70. Dangal V, Huang F. Fitness costs of Cry1F resistance in two populations of fall armyworm, *Spodoptera frugiperda* (J.E. Smith), collected from Puerto Rico and Florida. *J Invertebr Pathol.* 2015;127:81–86. doi:10.1016/j.jip.2015.03.004.
71. Hoffmann AM, French BW, Hellmich RL, Lauter N, Gassmann AJ. Fitness costs of resistance to Cry3Bb1 maize by western corn rootworm. *J Appl Entomol.* 2015;139:403–15. doi:10.1111/jen.12209.
72. Oswald KJ, French BW, Nielson C, Bagley M. Assessment of fitness costs in Cry3Bb1-resistant and susceptible western corn rootworm (Coleoptera: chrysomelidae) laboratory colonies. *J Appl Entomol.* 2012;136:730–40.
73. Meihls LN, Higdon ML, Ellersieck MR, Tabashnik BE, Hibbard BE. Greenhouse-selected resistance to Cry3Bb1-producing corn in three western corn rootworm populations. *PLoS ONE.* 2012;7(12):e51055. doi:10.1371/journal.pone.0051055.
74. Geisert RW, Hibbard BE. Evaluation of potential fitness costs associated with eCry3.1Ab resistance in *Diabrotica virgifera virgifera* (Coleoptera: chrysomelidae). *J Econ Entomol.* 2016;109:1853–58. doi:10.1093/jee/tow095.
75. Petzold-Maxwell JL, Siegfried BD, Hellmich RL, Abel CA, Coates BS, Spencer TA, Horikoshi RJ, Gassmann AJ. Fitness costs associated with Cry1F resistance in the European corn borer. *J Appl Entomol.* 2017;141:7–79. doi:10.1111/jen.12356.
76. Chen X, Head GP, Kerns DL, Price P, Rice ME, Huang F, Gilreath RT, Yang F. Fitness costs of Vip3A resistance in *Spodoptera frugiperda* on different hosts. *Pest Manage Sci.* 2019;75:1074–80. doi:10.1002/ps.5218.
77. Akhurst RJ, James W, Bird LJ, Beard C. Resistance to the Cry1Ac  $\delta$ -endotoxin of *Bacillus thuringiensis* in the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: noctuidae). *J Econ Entomol.* 2003;96:1290–99. doi:10.1603/0022-0493-96.4.1290.

78. Liu L, Gao M, Yang S, Liu S, Wu Y. Resistance to *Bacillus thuringiensis* toxin Cry2Ab and survival on single-toxin and pyramided cotton in cotton bollworm from China. *Evol Appl.* **2017**;10:170–79. doi:10.1111/eva.12438.
79. Tetreau G, Wang R, Wang P. Fitness of Bt-resistant cabbage loopers on Bt cotton plants. *Plant Biotech J.* **2017**;15:1322–30. doi:10.1111/pbi.12718.
80. Naik VB, Kumbhare S, Kranthi S, Satija U, Kranthi KR. Field evolved-resistance of pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: gelechiidae) to transgenic Bt-cotton expressing Cry1Ac and Cry2Ab in India. *Pest Manag Sci.* **2018**;74:2544–54.
81. Huang F. Resistance of the fall armyworm, *Spodoptera frugiperda* (J. E. Smith), to transgenic *Bacillus thuringiensis* Cry1F corn in the Americas: lessons and implications for Bt corn IRM in China. *Ins Sci.* **2020**. doi:10.1111/1744-7917.12826.
82. Leahy TC, Andow DA. Egg weight, fecundity, and longevity are increased by adult feeding in *Ostrinia nubilalis* (Lepidoptera: pyralidae). *Ann Entomol Soc Am.* **1994**;87:342–49. doi:10.1093/aesa/87.3.342.
83. Zhao JZ, Cao J, Collins HL, Bates SL, Roush RT, Earle ED, Shelton AM. Concurrent use of transgenic plants expressing a single and two *Bacillus thuringiensis* genes speeds insect adaptation to pyramided plants. *Proc Natl Acad Sci USA.* **2005**;102:8426–30. doi:10.1073/pnas.0409324102.
84. Kaur G, Guo J, Brown S, Head GP, Price PA, Paula-Moraes S, Ni X, Dimase M, Huang F. Field-evolved resistance of *Helicoverpa zea* (Boddie) to transgenic maize expressing pyramided Cry1A.105/Cry2Ab2 proteins in northeast Louisiana, the United States. *J Invertebr Pathol.* **2019**;163:11–20. doi:10.1016/j.jip.2019.02.007.
85. Dively GP, Venugopal PD, Finkenbinder C. Field-evolved resistance in corn earworm to Cry proteins expressed by transgenic sweet corn. *PLoS One.* **2016**;11(12):e0169115. doi:10.1371/journal.pone.0169115.
86. Reisig DD, Huseeth AS, Bacheler JS, Aghaee MA, Braswell L, Burrack HJ, Flanders K, Greene JK, Herbert DA, Jacobson A, et al. Long-term empirical and observational evidence of practical *Helicoverpa zea* resistance to cotton with pyramided Bt toxins. *J Econ Entomol.* **2018**;111:1824–33. doi:10.1093/jee/toy106.
87. Zhang L, Leonard BR, Chen M, Clark T, Konasale A, Huang F. Fitness and stability of Cry1Ab resistance in sugarcane borer, *Diatraea saccharalis* (F.). *J Invertebr Pathol.* **2014**;117:26–32. doi:10.1016/j.jip.2014.01.007.
88. Huang F, Chen M, Gowda A, Clark TL, McNulty BC, Yang F, Niu Y. (2015). Identification, inheritance, and fitness costs of Cry2Ab2 resistance in a field-derived population of sugarcane borer, *Diatraea saccharalis* (F.) (Lepidoptera: Crambidae). *J Invertebr Pathol* 130:116 – 123.
89. Brévaulta T, Heuberger S, Zhang M, Ellers-Kirk C, Ni X, Masson L, Li X, Tabashnik BE, Carrière Y. (2013). Potential shortfall of pyramided transgenic cotton for insect resistance management. *Proc Nat Acad Sci USA* 110: 5806 – 5811
90. Carrière Y, DeGain B, Unnithan GC, Harpold VS, Li X, Tabashnik BE. (2019). Seasonal Declines in Cry1Ac and Cry2Ab Concentration in Maturing Cotton Favor Faster Evolution of Resistance to Pyramided Bt Cotton in *Helicoverpa zea* (Lepidoptera: Noctuidae). *J Econ Entomol* 112: 2019, 2907–2914.

## Appendix

**Table A1.** Data sources and/or calculations of the dominance levels ( $D_{FL}$ ) of 17 cases of major resistance to Bt plants\*.

Case	Field resistance	No. populations investigated	Data sources and/or calculations of $D'_{ML}$ or $D'_{WT}$	$D_{FL}$ measured as $D'_{ML}$ or $D'_{WT}$
<b>Resistance to single-gene Bt plants</b>				
<i>B. fusca</i> to Cry1Ab maize in S. Africa	Yes	1	Based on neonate-to-pupa survivorships on whole maize plants, $D'_{ML}$ was estimated to be 1.56. <sup>33</sup>	$D'_{ML} = 1.56$
<i>S. frugiperda</i> to Cry1F maize in Brazil	Yes	5	Five Brazilian populations were evaluated in three studies. Based on neonate-to-adult survivorships of a population (BR25R) on maize leaf tissue, Farias et al. <sup>59</sup> reported a $D'_{ML}$ of 0.15 for BR25R. Leite et al. <sup>35</sup> assessed $D'_{ML}$ of two populations (IrmaF and IrmaD) on maize leaf tissue. Based on a 'fitness index', which was calculated using a formula: fitness index = (neonate-to-pupal survivorship x pupal weight)/neonate-to-pupal development time), $D'_{WT}$ s of the two populations were estimated to be 0.36. In addition, Santos-Amaya et al. <sup>36</sup> examined $D'_{WT}$ s of two other populations (MTH and MRH) using the 'fitness index' as described in Leite et al. <sup>35</sup> The estimated $D'_{WT}$ was 0.12 for MTH and 0.17 for MRH. Thus, the average $D'_{FL}$ of the five populations was 0.23.	Mixed $D_{ML}$ and $D'_{WT} = 0.23$
<i>S. frugiperda</i> to Cry1F maize in the U.S.	Yes	2	$D'_{ML}$ s of two populations, one from Puerto Rico (PR) and another from Florida (FL) were evaluated based on 7-d larval survivorship on maize leaf tissue in two studies <sup>33,37</sup> . $D'_{ML}$ of PR was 0.23 and 0.12 in two Cry1F maize hybrids, respectively with an average of 0.18. <sup>37</sup> $D'_{ML}$ s estimated in the study <sup>33</sup> was 0.07 for both PR and FL. $D_{ML}$ for PR was calculated based on the average $D'_{ML}$ s of the two studies, which was 0.13. Thus, $D'_{ML}$ for this case was calculated as the average $D'_{ML}$ of the two populations, which was 0.10.	$D'_{ML} = 0.10$
<i>D. virgifera virgifera</i> to Cry3Bb1 maize in U.S.	Yes	5	For this case, five populations were evaluated in four studies. Based on neonate-to-3 <sup>rd</sup> instar survivorship of a population on maize seedling mat, Petzold-Maxwell et al. <sup>39</sup> reported a $D'_{ML}$ of 0.51. Ingber and Gassmann <sup>40</sup> evaluated the survival-to-adult of two populations (Hopkinton and Cresco) on seedling-mat and reported a $D'_{ML}$ of 0.37 for Hopkinton and 0.27 for Cresco. Paolino and Gassmann <sup>41</sup> tested the survival-to-adult of another two populations (Elma and Monona) on seedling mat and 14-d survivorship of Monona with single-plant assays. The results showed a $D'_{ML}$ of 0.29 for Elma and 0.45 for Monona on seedling mats, and 0.73 for Monona in single-plant assays. The average $D'_{ML}$ of Monona in the two assay methods was 0.59. Thus, the average $D'_{ML}$ for the five populations was estimated to be 0.41.	$D'_{ML} = 0.41$
<i>D. virgifera virgifera</i> to eCry3.1Ab maize in U.S.	Yes	1	Geisert et al. <sup>42</sup> evaluated the survival of an eCry3.1Ab-resistant population using 10-d seedling bioassays and reported a $D'_{ML}$ of 0.94 and 1.38 for the two reciprocal crosses and thus the average $D'_{ML}$ of this case was 1.16.	$D'_{ML} = 1.16$
<i>S. frugiperda</i> to Cry1A.105 maize in the U.S.	Yes	2	Niu et al. <sup>43</sup> estimated $D'_{ML}$ based on a 7-d survivorship of two populations (RR32 and RR67) on maize leaf tissue and reported a $D'_{ML}$ of 0.58 for RR32 and 0.10 for RR67. Thus, the $D'_{ML}$ for this case was calculated as the average $D'_{ML}$ of the two populations, which was 0.34.	$D'_{ML} = 0.34$
<i>O. nubilalis</i> to Cry1F maize in U.S.	No	1	Based on the combined data of survivorship and weight gain after 15-d release of neonates on maize plants, $D'_{WT}$ was estimated to be 0.07 in vegetative plant stages and 0.00 in reproductive stages. <sup>44</sup> Thus, $D'_{WT}$ for this case was calculated as the average $D'_{WT}$ (0.04) of the two test methods.	$D'_{WT} = 0.04$
<i>S. frugiperda</i> to Cry2Ab2 maize in U.S.	No	1	Acharya et al. <sup>45</sup> estimated $D'_{ML}$ based on a 7-d survivorship of a population on maize leaf tissue and reported a $D'_{ML}$ of -0.02.	$D'_{ML} = -0.02$
<i>S. frugiperda</i> to Vip3A maize in Brazil	No	2	Bernatdi et al. <sup>46</sup> evaluated $D'_{WT}$ of a population on both whole maize plants and maize leaf tissue, and reported $D'_{WT} = 0$ . Miraldo et al. <sup>47</sup> evaluated neonate-to-4 <sup>th</sup> instar survivorship of another population on maize plants and reported $D'_{ML} = 0$ . Thus, $D'_{WT}$ of the case was zero.	$D'_{WT} = 0.00$
<i>S. frugiperda</i> to Vip3A maize in U.S.	No	1	Yang et al. <sup>48</sup> estimated $D'_{ML}$ based on 7-d survivorship of a population on maize leaf tissue and reported a $D'_{ML}$ of 0.00.	$D'_{ML} = 0.00$
<i>H. armigera</i> to Cry1Ac cotton in Australia	No	2	Bird and Akhurst <sup>49</sup> examined $D'_{WT}$ of a population based on intrinsic rate of population increase, $r_m$ , on 4-week old cotton and reported a $D'_{WT}$ of zero. In addition, based on the $r_m$ values presented in Table 3 in the reference, <sup>50</sup> $D'_{WT}$ was recalculated by the author of this review and resulted in a $D'_{WT}$ of 0.68 in Exp 1 and 0.63 in Exp 2, and thus the average $D'_{WT}$ on 14-week cotton was 0.65. Finally, the $D'_{WT}$ for this case was calculated as the average $D'_{WT}$ of the two studies, which was 0.33.	$D'_{WT} = 0.33$
<i>P. gossypiella</i> to Cry1Ac cotton in U.S.	No	1	Liu et al. <sup>51</sup> examined $D'_{ML}$ based on 54-d survivorship of a population on cotton and reported a $D'_{ML}$ of zero.	$D'_{ML} = 0.00$

(Continued)



**Table A1.** (Continued).

Case	Field resistance	No. populations investigated	Data sources and/or calculations of $D'_{ML}$ or $D'_{WT}$	$D_{FL}$ measured as $D'_{ML}$ or $D'_{WT}$
<i>D. saccharalis</i> to Cry1Ab maize in U.S.	No	1	Wu et al. <sup>52</sup> examined the 21-d larval survival of a population on seven Cry1Ab hybrids at vegetative and reproductive plant stages in the greenhouse in 2005 and 2006, respectively. $D'_{ML}$ calculated based on the published data ranged from 0.04 to 0.28 with an average of 0.17. Ghimire et al. <sup>32</sup> conducted two greenhouse trials and evaluated larval survivorship of a population on six Cry1Ab maize hybrids/lines and reported $D'_{MLS}$ ranged from 0.25 to 0.69 with an average of 0.42. Wangila et al. <sup>53</sup> conducted two trials in 2010 and 2011 and evaluated larval survivorship of the same population on Cry1Ab maize plants and reported a $D'_{ML}$ from 0.50 to 0.78 with an average of 0.65. Thus, $D'_{ML}$ for this case was calculated as the average $D'_{ML}$ of the three studies, which was 0.41.	$D'_{ML} = 0.40$
<b>Resistance to dual/multiple-gene Bt plants</b>				
<i>S. frugiperda</i> to Cry1A.105/ Cry2Ab maize in Brazil	No	2	Santos-Amaya et al. <sup>55</sup> evaluated neonate-to-adult survivorship of a population on dual-gene Cry1A.105/Cry2Ab2 maize plants and leaf tissue, and reported a $D_{ML}$ of zero for the resistance. In addition, Horikoshi et al. <sup>54</sup> evaluated 7-d larval survivorship of another population on maize leaf tissue and also reported a $D'_{ML}$ of zero.	$D'_{ML} = 0.00$
<i>S. frugiperda</i> to Cry1A.105/ Cry2Ab maize in U.S.	No	1	Niu et al. <sup>56</sup> conducted two trials and evaluated net reproductive rate ( $R_0$ ) on whole maize plants and leaf tissue, and reported a pooled $D_{WT}$ of 0.12. Zhu et al. <sup>57</sup> evaluated 14-d survival of the same population on whole plants and reported a $D'_{ML}$ of 0.27. Thus, $D'_{FL}$ for this case was calculated as the average $D'_{WT}$ and $D'_{ML}$ of the two studies, which was 0.20.	Mixed $D'_{WT}$ and $D'_{ML} = 0.20$
<i>S. frugiperda</i> to Cry1Ab/Vip3A maize in Brazil	No	1	Horikoshi et al. <sup>54</sup> evaluated 7-d larval survival of a population on maize leaf tissue and reported a $D'_{ML}$ of zero.	$D'_{ML} = 0.00$
<i>S. frugiperda</i> to Cry1A.105/ Cry2Ab2/Cry1F in Brazil	No	2	Horikoshi et al. <sup>54</sup> and Bernardi et al. <sup>58</sup> evaluated 7-d larval survival on maize leaf tissue and both studies reported a $D'_{ML}$ of zero.	$D'_{ML} = 0.00$

\* All references cited in Table A1 are listed in the main article. In addition, dominance levels of a Cry2Ab2-resistant population of *D. saccharalis* were also evaluated based on 7-d survivorship rates on leaf tissue of a Cry2Ab2 maize experimental line.<sup>88</sup> The Cry2Ab2-resistant strain was documented to possess a major resistance allele to the experimental Cry2Ab2 maize line. However, the experimental line used in the study expressed a relatively low level of Cry2Ab2 protein (FH personal communication), and thus, the case of the Cry2Ab2 resistance in *D. saccharalis* was excluded in this review. Larval development and survivorship of a field-collected (GA) and a Cry1Ac-selected (GA-R) populations, and their F1 progeny of *Helicoverpa zea* have been evaluated on plant tissues of non-Bt, Cry1Ac, and pyramided Cry1Ac/Cry2Ab cotton.<sup>89,90</sup> Because GR was collected from Cry1Ab maize plants and it had shown significant resistance ratios to both Cry1Ac and Cry2Ab (e.g. 55-fold to Cry1Ac and 15-fold to Cry2Ab2, relative to a laboratory strain), this case was also excluded in this review.

**Table A2.** Index of fitness costs (IFC) of 20 cases of major resistance to Bt plants in eight target species.\*.

Case	Field resistance	No. population investigated	Sources and calculation of fitness parameters	F <sub>SS</sub> <sup>a</sup>	F <sub>RR</sub> <sup>b</sup>	F <sub>RS</sub> <sup>c</sup>	IFC <sub>RR</sub> <sup>d</sup>	IFC <sub>RS</sub> <sup>e</sup>
<b>Resistance to single-gene Bt plants</b>								
<i>B. fusca</i> to Cry1Ab maize in South Africa								
	Yes	1	Progeny production of field-collected populations (F0) = number of eggs x egg hatching rate in Table 1 in reference <sup>68</sup>	151.1	213.8	n/a	1.41	n/a
		1	Fitness index = 31-d survival rate x larval mass of F1 populations in Table 2 in reference <sup>68</sup>	5.83	11.55	n/a	1.98	n/a
<i>S. frugiperda</i> to Cry1F maize in Brazil								
	Yes	1 (MTH)	61-d survival rate of F1 populations in Table 2 in reference <sup>68</sup>	0.12	0.34	n/a	2.83	n/a
		1 (MRH)	Average IFC of the two measurements for F1	--	--	--	2.41	n/a
		1 (lrmaF)	Average IFC of F0 and F1 for the case	0.40	0.37	0.35	1.91	n/a
		1	Neonate-to-adult survivorship in Fig 4 in reference <sup>36</sup>	0.42	0.37	0.38	0.93	0.88
			Neonate-to-adult survivorship in Fig 4 in reference <sup>36</sup>	0.71	0.72	0.73	0.88	0.90
			14-d larval survivorship in Fig 3 in reference <sup>35</sup>	0.83	0.92	0.98	1.01	1.03
			7-d larval survivorship on maize in Fig 1 in reference <sup>54</sup>	0.86	0.64	0.86	1.11	1.18
			7-d larval survivorship on cotton in Fig 2 in reference <sup>54</sup>	0.098	0.080	0.087	0.74	1.00
			Average IFC of the two test methods in the study <sup>54</sup>				0.93	1.09
	Yes	1 (PR)	Average IFC of the four populations for the case				0.94	0.98
			Fitness index = (larval survivorship x number of egg masses x egg hatching rate)/(neonate-to-adult developmental time) on maize leaf tissue in Figs 1, 4, and Table 1 in reference <sup>69</sup>	0.055	0.048	0.054	0.87	0.98
			Fitness index = (larval survivorship x number of egg masses x egg hatching rate)/(neonate-to-adult developmental time) on cotton leaf tissue in Figs 1, 4, 5 and Table 1 in reference <sup>69</sup>	8.35	5.35	7.62	0.85	0.94
			Average of the PR population based on the study <sup>69</sup>	8.35	4.51	8.0	0.64	0.91
			Fitness index = (neonate-to-adult survivorship x pupal mass)/neonate-to-adult developmental time in Tables 1 and Tables 2 in the reference <sup>70</sup>	84.00	55.80	n/a	0.54	0.96
			Fitness index = (neonate-to-adult survivorship x pupal mass)/neonate-to-adult developmental time in Tables 1 and Tables 2 in reference <sup>70</sup>				0.68	0.94
			Average of the three populations for the case				0.66	n/a
	Yes	1	Population growth rate per generation in Table 1 in reference <sup>73</sup>				1.28	n/a
<i>D. virgifera virgifera</i> to Cry3Bb1 maize in U.S.								
	Yes	1	Fitness index = number of egg x hatching rate x neonate-to-adult survivorship in Table 1 (moderate selected) in reference <sup>72</sup>	n/a	n/a	n/a	1.22	n/a
		1	Fitness index = number of egg x hatching rate x neonate-to-adult survivorship in Table 1 (intense selected) in reference <sup>72</sup>	n/a	n/a	n/a	1.22	n/a
		1	Fitness index = (relative fecundity x relative egg viability x relative survivorship to adult) in Table 4 in reference <sup>71</sup>	1.00	1.22	n/a	1.22	n/a
		1(Hopkinton)	Fitness index = (egg production x egg viability x rate of larval survival to adult)/days of development to adults in Fig. 3 in reference <sup>40</sup>	51.35	55.18	n/a	1.07	n/a
		1(Cresco)	Fitness index = (egg production x egg viability x rate of survival to adult)/days of development to adults in Fig. 4 in inference <sup>40</sup>	11.46	4.32	n/a	0.38	n/a
		1 (Monona)	Fitness index = (egg production x egg viability x rate of survival to adult)/days of development to adults in Fig 3 in reference <sup>41</sup>	29.15(1210)	42.77 (1779)	n/a	1.47	n/a
		1 (Elma in the test with high food availability)	Fitness index = (egg production x egg viability x rate of survival to adult)/days of development to adults in Fig 4 in reference <sup>41</sup>	24.64	33.85	n/a	1.37	n/a
<i>D. virgifera virgifera</i> to eCry3.1Ab maize in U.S.								
	Yes	1	Average of the 8 populations for the case	7180	11753	n/a	1.08	n/a
			Fitness index = (larval recovery rate in 20 d x number of eggs produced per female x egg viability) in Figs 3 and 4 in reference <sup>74</sup>				1.64	n/a
<i>S. frugiperda</i> to Cry1A.105 maize in U.S.								
	Yes	1 (RR32)	Fitness index = (neonate-to-adult survivorship x number of eggs per female)/neonate-to-adult developmental time in Table 5 in reference <sup>43</sup>	15.64	21.31	15.59	1.36	1.00

(Continued)

Table A2. (Continued).

Case	Field resistance	No. population investigated	Sources and calculation of fitness parameters	$F_{SS}$	$F_{RR}$	$F_{RS}$	$IF_{RR}$	$IF_{RS}$
		1(RR67)	Fitness index = (neonate-to-adult survivorship x number of eggs per female)/neonate-to-adult developmental time in Table 5 in reference <sup>43</sup>	15.64	30.22	46.65	1.93	2.98
<i>O. nubilalis</i> to Cry1F maize in U.S.	No	1	Average of the two populations for the case Fitness index = neonate-to-adult survivorship x fertile eggs produced per female x adult mating success rate on three maize lines in Fig 3 and Table 3 in reference <sup>5</sup>	57.6	44.4	55.5	1.65	1.99
<i>S. frugiperda</i> to Cry2Ab2 maize in U.S.	No	1	Fitness index = (egg production x neonate-to-pupal survivorship)/neonate-to-pupal developmental time in Table 3 in reference <sup>45</sup>	1.34	2.51	3.20	1.87	2.39
<i>S. frugiperda</i> to Vip3A maize in Brazil	No	1	Intrinsic rate of increase ( $r_m$ ) in Table 5 in reference <sup>46</sup>	0.20	0.16	0.19	0.80	0.95
<i>S. frugiperda</i> to Vip3A maize in U.S.	No	1	Intrinsic rate of increase ( $r_m$ ) on maize in Fig 8 in reference <sup>76</sup>	0.19	0.16	0.20	0.84	1.05
			Intrinsic rate of increase ( $r_m$ ) on cotton in Fig 8 in reference <sup>76</sup>	0.14	0.17	0.15	1.21	1.07
<i>H. armigera</i> to Cry1Ac cotton in Australia	No	1	Average of the two measurements for the case Pupation rate on cotton plants reported in reference <sup>77</sup>	0.83	0.60	n/a	1.03	1.06
		1	Average intrinsic rate of increase ( $r_m$ ) of experiments 1 and 2 in Table 6 in reference <sup>49</sup>	0.20	0.14	0.19	0.70	0.95
<i>H. armigera</i> to Cry2Ab cotton in China	No	1	Average of the two studies for the case				0.71	0.95
<i>P. gossypiella</i> to Cry1Ac cotton in U.S.	No	1	Neonate-to-pupal survivorship in Table 3 in reference <sup>78</sup>	0.30	0.23	n/a	0.77	n/a
		1	Fitness index = larval survivorship/development period (degree day) in Tables 2 and Tables 3 in reference <sup>51</sup>	0.00071	0.00034	0.00037	0.48	0.52
<i>D. saccharalis</i> to Cry1Ab maize in U.S.	No	1	Average larval survivorship after 21 d on five non-Bt maize hybrids in two trials at two plant stages in Figs 1 and 3 in reference <sup>52</sup>	0.37	0.40	0.40	1.08	1.08
		1	Average larval survivorship after 21 d on two non-Bt maize hybrids in two trials in Figs 2 and 3 in reference <sup>32</sup>	0.44	0.57	0.57	1.30	1.30
		1	Average larval survivorship after 21 d on two non-Bt maize hybrids in two trials in Figs 2 and 4 in reference <sup>53</sup>	0.46	0.58	0.53	1.26	1.15
<i>T. ni</i> to Cry1Ac cotton in U.S.	No	1	Average of the three studies for the case	0.16	0.15	n/a	1.21	1.18
<b>Resistance to dual-/multiple-gene Bt plants</b>			Intrinsic rate of increase ( $r_m$ ) on cotton leaves in Fig 1 in reference <sup>79</sup>				0.94	n/a
<i>S. frugiperda</i> to Cry1A:105/Cry2Ab maize in Brazil	No	1	Neonate-to-adult survivorship in Fig 3 in reference <sup>55</sup>	0.26	0.24	0.27	0.92	1.04
		1	7-d larval survivorship on non-Bt maize in Fig 1 in reference <sup>54</sup>	0.83	0.98	0.94	1.18	1.13
		1	7-d larval survivorship on non-Bt cotton in Fig. 2 in reference <sup>54</sup>	0.86	0.83	0.84	0.97	0.98
		1	Average in the two crops based on the study <sup>54</sup>				1.08	1.06
		1	Average of the two populations for the case				1.00	1.05
<i>S. frugiperda</i> to Cry1A:105/Cry2Ab maize in U.S.	No	1	Fitness index = net reproductive rate/neonate-to-pupa developmental time in Table 2 in reference	29.5	13.9	30.6	0.47	1.04
		1	14-d larval survivorship in Fig 2 in reference	50.5	50.0	63.5	0.99	1.26
		1	Average of the two studies for the case				0.73	1.15
<i>T. ni</i> to Cry1Ac/Cry2A cotton in U.S.	No	1	Intrinsic rate of increase ( $r_m$ ) on cotton leaves in Fig 1 in the reference <sup>79</sup>	0.16	0.13	n/a	0.81	n/a
<i>S. frugiperda</i> to Cry1Ab/Vip3A maize in Brazil	No	1	7-d larval survivorship on maize in Fig 1 in reference <sup>80</sup>	0.83	0.81	0.88	0.98	1.06
		1	7-d larval survivorship on cotton in Fig. 2 in reference <sup>54</sup>	0.86	0.63	0.95	0.73	1.10
		1	Average on the two crops for the case				0.86	1.08

(Continued)

**Table A2.** (Continued).

Case	Field resistance	No. population investigated	Sources and calculation of fitness parameters	$F_{SS'}$	$F_{RR'}$	$F_{RS'}$	$IFC_{RR'}$	$IFC_{RS'}$
<i>S. frugiperda</i> to Cry1A:105/Cry2Ab2/Cry1F in Brazil	No	1	7-d larval survivorship on maize in Fig 1 in reference <sup>54</sup>	0.83	0.96	0.88	1.16	1.06
		1	7-d larval survivorship on cotton in Fig. 2 in reference <sup>54</sup> Average on the two crops based on the study <sup>54</sup> $r_m$ in Table 3 in reference <sup>56</sup> Average of the two populations for the case	0.86	0.89	0.74	1.03	0.86
				0.18	0.16	0.18	0.89	1.00
							1.00	0.98

\*Index of fitness cost (IFC) of a single- or dual/multiple-gene resistance was calculated using the formula:

$$IFC_{RR'} = F_{RR'}/F_{SS'} \text{ and } IFC_{RS'} = F_{RS'}/F_{SS'}$$

Here  $IFC_{RR'}$  and  $IFC_{RS'}$  mean the index of fitness costs for homozygous- (RR') and heterozygous- (RS') resistant genotypes, respectively.  $F_{RR'}$  is the fitness of homozygous-resistant genotype (RR') on non-Bt plants;  $F_{RS'}$  is the fitness of heterozygous genotype (RS') on non-Bt plants; and  $F_{SS'}$  is the fitness of homozygous-susceptible genotype (SS') on non-Bt plants. If A, B, and C represent three different resistant alleles and a, b, and c refers to the three corresponding susceptible alleles of the three genes, RR', RS', and SS' mean AA, Aa, and aa for a single-gene resistance; AABB, AaBb, and aabb for a dual-gene resistance; or AABBCC, AaBbCc, and aabbcc for a triple-gene resistance.  $IFC < 1$  means that there is fitness cost;  $IFC = 1$  suggests that there is no fitness cost;  $IFC > 1$  indicates that there is fitness advantage associated with the resistance.

All references cited in Table A2 are listed in the main publication.