

Review

Systems biology of death receptor networks: live and let die

IN Lavrik^{*,1,2}

The extrinsic apoptotic pathway is initiated by death receptor activation. Death receptor activation leads to the formation of death receptor signaling platforms, resulting in the demolition of the cell. Despite the fact that death receptor-mediated apoptosis has been studied to a high level of detail, its quantitative regulation until recently has been poorly understood. This situation has dramatically changed in the last years. Creation of mathematical models of death receptor signaling led to an enormous progress in the quantitative understanding of the network regulation and provided fascinating insights into the mechanisms of apoptosis control. In the following sections, the models of the death receptor signaling and their biological implications will be addressed. Central attention will be given to the models of CD95/Fas/APO-1, an exemplified member of the death receptor signaling pathways. The CD95 death-inducing signaling complex (DISC) and regulation of CD95 DISC activity by its key inhibitor c-FLIP, have been vigorously investigated by modeling approaches, and therefore will be the major topic here. Furthermore, the non-linear dynamics of the DISC, positive feedback loops and bistability as well as stoichiometric switches in extrinsic apoptosis will be discussed. Collectively, this review gives a comprehensive view how the mathematical modeling supported by quantitative experimental approaches has provided a new understanding of the death receptor signaling network.

Cell Death and Disease (2014) 5, e1259; doi:10.1038/cddis.2014.160; published online 29 May 2014

Subject Category: Cancer

Facts

- Death receptor activation leads to the formation of death receptor signaling platforms, which activate caspase cascade resulting in the demolition of the cell.
- Quantitative regulation of death receptor network until recently has been poorly understood.
- Within the last decade, a number of mathematical models of death receptor signaling were created that changed the understanding of the extrinsic apoptosis.

Open Questions

- Which system properties determine death receptor-mediated apoptotic cell death?
- What are the initial conditions leading a cell into apoptosis?
- What is the role of DISC dynamics in apoptosis initiation?

The extrinsic apoptotic pathway is initiated by signals emanating from the cell-surface death receptors triggered by death ligands.¹ The binding of death ligands results in the

formation of the death receptor signaling platforms and subsequent apoptosis initiation.^{2,3} In the recent years, remarkable progress has been made in decoding the death receptor network. The major players of the death receptor signaling pathways have been extensively characterized: death receptors and their activating platforms, apoptosis initiation mechanisms and their inhibition.⁴ Nevertheless, despite the fact that the death receptor network has been well described, its understanding in quantitative terms has been largely missing.⁵ Indeed, a number of questions have remained unsolved: What are system properties that determine death receptor-mediated apoptotic cell death? What are the initial conditions leading a cell into apoptosis? Activation of which critical nodes, for example, groups of proteins, defines apoptosis initiation? Can the cell be a 'little bit dead'? To answer these questions by addressing the death receptor network on a quantitative level, the emerging field of systems biology was implemented.^{6–8} Systems biology combines mathematical modeling with powerful experimental methodology, providing a quantitative assessment of the signaling pathways.^{5,9} The application of systems biology to the analysis of death receptor networks resulted in new

¹Department of Translational Inflammation Research, Institute of Experimental Internal Medicine, Otto von Guericke University, Magdeburg, Germany and ²Faculty of Fundamental Medicine, MV Lomonosov Moscow State University, Moscow, Russia

*Corresponding author: IN Lavrik, Department of Translational Inflammation Research, Institute of Experimental Internal Medicine, Otto von Guericke University, Magdeburg, Germany. Tel: +49 3916724767; Fax: +49 3916724769; E-mail: inna.lavrik@med.ovgu.de

Keywords: death receptor; systems biology; CD95; DISC; caspase-8; c-FLIP

Abbreviations: DISC, death-inducing signaling complex; CD95L, CD95 ligand; DED, death effector domain; c-FLIP, cellular FLICE-like inhibitory proteins; XIAP, X-linked inhibitor of apoptosis; MOMP, mitochondrial outer membrane polarization; c-FLIP isoforms, Long (L), Short (S) and Raji (R); TNF, tumor necrosis factor; ODE, Ordinary Differential Equation

Received 27.1.14; revised 11.3.14; accepted 13.3.14; Edited by B Zhivotovsky

fascinating insights into the network regulation.^{6–8,10–11} It turned out that as in the famous ‘live and let die’ movie, the cell also controls cell death in a very precise but also dynamic way, and the same ‘agent’ can both kill the cells or let them live, depending on the circumstances, for example, the concentrations of the molecules in the pathway. Below we shall discuss in detail how systems biology studies provide new understanding of the death receptor networks and comment on the identified non-linear responses, stoichiometric switches, feedback loops and bistability behavior in death receptor system defining life or death of the cell.

Signaling of CD95 as a Prototypic Death Receptor

The death receptor family belongs to the tumor necrosis factor (TNF) superfamily.^{1,4} All members of the death receptor family are characterized by the presence of an intracellular ~80 amino-acid long motif, which is termed the death domain playing a key role in the formation of the death receptor platforms.² The major death receptors characterized to date include: CD95 (also named Fas/APO-1), TNF-R1, TRAIL receptor 1 (TRAIL-R1), TRAIL-R2, DR3 and DR6.²

The CD95-mediated apoptotic pathway is one of the best-studied death receptor signaling pathways.³ Stimulation of CD95 with CD95 ligand (CD95L) or with agonistic anti-CD95 antibodies such as anti-APO-1 induces apoptosis in sensitive cells.⁴ The central apoptosis-initiating platform in CD95 signaling is the CD95 death-inducing signaling complex (DISC) comprising CD95, the adaptor protein FADD, procaspase-8, procaspase-10 and cellular FLICE-like inhibitory proteins (c-FLIP;⁴ Figure 1). All interactions at the DISC are based on the so-called homotypic contacts. FADD is recruited to CD95 via death domain interactions, whereas procaspase-8/10 and c-FLIP are recruited to the DISC via death effector

domain (DED) interactions (Figure 1). Several isoforms of the DED proteins were found at the DISC: two isoforms of procaspase-8, procaspase-8a (p55) and procaspase-8b (p53); three isoforms of procaspase-10 and c-FLIP isoforms, including long (L), short (S) and Raji (R)¹² (Figure 1).

A number of studies has demonstrated that procaspase-8 is activated at the DISC via formation of the procaspase-8 homodimers¹³ (Figures 1 and 2a). Furthermore, recently a new paradigm in the activation of caspase-8 at the DISC was unraveled: it was shown that procaspase-8 is activated via the formation of DED chains at the DISC.^{14–16} These chains comprise procaspase-8 molecules interacting via their DEDs, thereby bringing the caspase-like domains into the close proximity to each other (Figure 1). The DED chain enables the formation of the procaspase-8 homodimers, which is, in turn, a prerequisite for procaspase-8 activation. After formation of the active homodimer, procaspase-8a/b (p55/p53) undergoes processing with the generation of the N-terminal cleavage products p43/p41, the prodomains p26/p24, as well as the C-terminal cleavage products p30, p18 and p10^{17,18} (Figure 1). Active caspase-8 heterotetramers p10₂-p18₂ initiate the apoptotic cascade by cleavage and activation of the effector caspases, which in turn leads to the demolition of the cell.

The activation of procaspase-8 at the DISC is inhibited by c-FLIP proteins that are also recruited to the DISC via DED interactions. Short c-FLIP isoforms, c-FLIP_S and c-FLIP_R, comprise only tandem DEDs, whereas the long isoform, c-FLIP_L, also has the C-terminal caspase-like domain that lacks caspase activity¹² (Figure 1). Short c-FLIP isoforms could block caspase-8 activation upon overexpression, whereas the long isoform, c-FLIP_L, can both block and accelerate caspase-8 activation.^{19,20} The inhibition occurs in the presence of high concentrations of c-FLIP_L at the DISC, whereas acceleration of caspase-8 activation at the DISC




Figure 1 The scheme of the CD95 DISC. CD95L stimulation leads to the induction of the CD95 death-inducing signaling complex (DISC) comprising CD95, adaptor protein FADD, procaspase-8/10 and c-FLIP. All interactions at the DISC are based on the so-called homotypic contacts. FADD is recruited to CD95 via death domain (DD) interactions, whereas procaspase-8/10 and c-FLIP are recruited to the DISC via death effector domain (DED) interactions. Procaspase-8 is activated in the DED chains via the formation of procaspase-8 homodimers. Left side: two isoforms of procaspase-8: procaspase-8a (p55) and procaspase-8b (p53) as well as their processing to p43/p41, p26/p24, p18 and p10 are shown. Right side: three isoforms of c-FLIP are depicted: c-FLIP_L, c-FLIP_R and c-FLIP_S




Figure 2 Modeling pro-apoptotic properties of c-FLIP_L. (a) The scheme of formation of procaspase-8/procaspase-8 homodimer (left) and heterodimer procaspase-8/c-FLIP_L (right) at the DISC. Both scenarios could lead to caspase-8 processing to p43/p41 and active caspase-8 heterotetramer. (b) Predictions of the model for cell death upon different stimulation strength and different levels of c-FLIP_L. The lower panel demonstrates concentration-dependent effects of c-FLIP_L on the apoptosis induction upon high c-FLIP_R overexpression. Adapted from Fricker *et al.*¹⁹

takes place upon lower concentrations of c-FLIP_L. The latter has been reported to be mediated by the formation of catalytically active procaspase-8/c-FLIP_L heterodimers in which the procaspase-8-active loop is stabilized by c-FLIP_L, thereby increasing the catalytic activity of procaspase-8^{21–23} (Figure 2a).

Downstream from CD95 DISC, the apoptotic signal transduction is controlled at several levels. Two types of cells have been described.⁴ Type I cells are characterized by high numbers of the CD95 DISCs, and, correspondingly, high amounts of active caspase-8, that leads to the activation of caspase-3 and apoptosis induction. In type II cells, a lower number of CD95 DISCs is formed and the apoptotic signal is mediated via cleavage of Bid by procaspase-8 and translocation of truncated Bid to mitochondria. At the mitochondria, truncated Bid was suggested to interact with the mitochondrial-specific phospholipid cardiolipin, which promotes BAX or BAK oligomerization.²⁴ The latter leads to the disruption of the integrity of the outer mitochondrial membrane, which is followed by mitochondrial outer membrane polarization (MOMP), cytochrome C release from mitochondria and subsequent apoptosome formation followed by caspase-9 and then caspase-3 activation²⁵ (the mitochondrial branch of the apoptotic pathway is reviewed in detail in this issue by Rehm and colleagues). Apoptosis induction in type II cells could be blocked by anti-apoptotic Bcl-2 family members. Activation of initiator caspase-9 and effector caspase-3 and -7 could be inhibited by X-linked inhibitors of apoptosis (XIAPs).^{26,27} Interestingly, the action of XIAPs could be blocked by the SMAC proteins, which are released from mitochondria in the course of apoptosis.²⁷ Thus, a complex network of pro- and anti-apoptotic regulators controls death receptor-induced apoptosis.

Importantly, we and others have shown that CD95 is not only a potent apoptosis inducer but is also capable of activating multiple non-apoptotic pathways controlled by NF-κB, Erk1/2, p38, JNK and AKT.^{2,4,28} Furthermore, recent publications indicate that CD95-induced NF-κB and MAPK activation are also important for phagocyte attraction and generation of the ‘find me’ signal via the secretion of cytokines, making these pathways responsible for the proper execution of apoptosis as well.²⁹

The First Models of CD95 Signaling has Demonstrated that C-FLIP is the Key Molecule Defining the Live or Let Die Decision

Several mathematical formalisms have been used to model death receptor signaling.^{5,9} So far, Ordinary Differential Equations (ODEs) and Boolean modeling have been used the most. ODEs use the change in molecular concentrations over time, which allows to describe kinetics of a particular signaling pathway. Boolean models are more abstract as they use bimodal switches (on/off) to describe the behavior of the systems and do not require the knowledge of the kinetic parameters. Hence, Boolean models could not be applied for studies of the dynamic changes occurring within the signaling network. However, they are extremely valuable in characterizing large signaling networks.

The first theoretical model of apoptosis by Fussenegger *et al.*³⁰ has been build using literature data only. The model was generated using *ad hoc* parameters therefore, its predictive power was limited. It was followed by ODE-based mathematical model of CD95 signaling generated by Bentele *et al.*³¹ The strong advantage of this model that it was trained against experimental data generated with quantitative

western blot data in type I B-lymphoblastoid cells SKW6.4.³¹ The topology of the model included five major subsystems representing key regulatory nodes in CD95 apoptosis: CD95 DISC, effector caspases, mitochondria, XIAPs and degradation, the latter was approximated by PARP cleavage. Altogether, the model comprised 41 components presented in 32 reactions. The model had a strong predictive power, for example, it could well predict the timing of caspase activation.³¹ Importantly, the major finding of the model was the identification of the DISC as a crucial node of CD95-mediated cell death. Furthermore, the model predicted that CD95-mediated apoptosis is characterized by the threshold behavior.^{15,31} Notably, it turned out that this behavior is entirely based on the amounts of the core DISC proteins and their affinities.³¹ The model predicted the threshold mechanism as follows: a low stimulation strength would lead to a low number of activated CD95 that could recruit adaptor protein FADD. FADD associated to CD95 could bind both procaspase-8 and its inhibitor c-FLIP. However, the affinity of c-FLIP to FADD is higher than that of procaspase-8. Consequently, all active sites at the DISC, for example, FADDs, would be occupied by c-FLIP upon low number of active receptors. In this scenario, procaspase-8 could not bind to the DISC and initiate apoptosis. Thus, the amount of c-FLIP bound to FADD would be the rate-limiting step of caspase-8 activation at the DISC. This prediction has been confirmed by vigorous experimental data: a decrease of the amount of c-FLIP led to a decrease of the threshold, for example, the cells were undergoing CD95-induced apoptosis upon stimulation with lower amounts of CD95L.³¹ Furthermore, we have found that upon low CD95 stimulation strength the amount of c-FLIP at the DISC was increased compared with high stimulation strength.³² These experiments ultimately confirmed the threshold mechanism suggested by the model and strongly supported the role of c-FLIP as a guardian of the threshold of extrinsic apoptosis.

The question which arises next is: how can cells undergo apoptosis if the affinity of c-FLIP to the DISC is higher than the affinity of procaspase-8? This would naturally mean that c-FLIP would always be recruited to the DISC first and would not allow procaspase-8 recruitment and subsequent activation. Interestingly, the amount of c-FLIP proteins in a number of the cells is significantly lower than the amount of procaspase-8, which was quantitatively measured, in particular, in SKW6.4 and HeLa cell lines.^{15,19} Thus, upon high stimulation strength, c-FLIP could not block all DISCs any longer because of the increased number of active receptors and correspondingly higher number of procaspase-8 compared with c-FLIP in the cell. Hence, the amount of c-FLIP proteins in the cell defines the threshold behavior in the CD95-mediated apoptosis.³¹

Another prominent feature of c-FLIP proteins is a high turnover rate that allows them to modulate extrinsic apoptosis pathways in the dynamic manner.^{12,33} Indeed, c-FLIP proteins are characterized by a very short half-life time and they quickly undergo proteosomal degradation.³³ This property of c-FLIP has been studied using a mathematical model of the dynamic turnover and stability of c-FLIP isoforms by Toivonen *et al.*³⁴ This model was developed on the basis of the model by Bentele *et al.*³¹ The c-FLIP degradation rates as

a first-order reaction have been added to the model. In this study, changes in the level of c-FLIP at the single-cell level were taken into account and Monte Carlo simulations to investigate apoptotic response within cell populations considering intracellular variations of the c-FLIP level were applied.³⁴ It was shown that amount of c-FLIP in the cell at the moment of death receptor stimulation defines timing of cell death.³⁴

Hence, as a result of these modeling works, it became evident that the amount of c-FLIP proteins is crucial in defining the sensitivity to apoptosis in the death receptor network.

The Expression Level of c-FLIP Might Define its Functions: Lessons from the DISC Models

The model by Bentele *et al.* only considered anti-apoptotic action of c-FLIP proteins, which is not entirely correct, taken that there are three c-FLIP isoforms, c-FLIP_L, c-FLIP_S and c-FLIP_R, described to have different action. Indeed, one of the controversies that existed in CD95 signaling was the role of the long isoform of c-FLIP, c-FLIP_L. c-FLIP_L has been assigned both pro- or anti-apoptotic roles in the pathway.^{20,35} The pro-apoptotic effects of c-FLIP_L as mentioned above are mediated by the formation of catalytically active procaspase-8/c-FLIP_L heterodimers in which the procaspase-8 active loop is stabilized by c-FLIP_L, thereby increasing the catalytic activity of procaspase-8 (Figure 2a).

To define the conditions when c-FLIP_L has a pro-apoptotic action, an ODE-based model of the DISC was developed by Fricker *et al.*¹⁹ The model was validated using quantitative western blot and single-cell analysis based on experiments in HeLa cells overexpressing CD95 (HeLa-CD95 cells). The topology of the model comprised the generation of homo- and hetero-dimers of c-FLIP_{L/S} and procaspase-8 at the DISC. In accordance with the literature, only procaspase-8 homodimers and procaspase-8-c-FLIP_L heterodimers were considered as species leading to apoptosis (Figure 2a). Dimers formed by procaspase-8 and the short isoform of c-FLIP, c-FLIP_S, as well as between different c-FLIP isoforms were considered inactive in terms of apoptosis induction. Consistent with this, the optimal conditions for the proapoptotic effect of c-FLIP_L would be achieved upon the concentration range of c-FLIP_L leading to the optimal amount of c-FLIP_L-procaspase-8 heterodimers formed at the DISC (Figure 2a). Remarkably, the model predicted that this 'ideal case' scenario occurs under conditions when the concentration of c-FLIP_L is 20 times higher than the endogeneous one¹⁹ (Figure 2b).

Another valuable outcome of this model was the prediction that the apoptosis-promoting effect of c-FLIP_L would be stronger upon a very low rate of caspase-8 activation.¹⁹ To generate an experimental setup with a very-low rate of caspase-8 activation, short c-FLIP isoforms were overexpressed in HeLa-CD95 cells.¹⁹ c-FLIP_{S/R} were considered to occupy most of procaspase-8-binding sites at the DISC, thereby creating a limited amount of binding sites for procaspase-8. This naturally leads to a very-low rate of caspase-8 activation up to complete inhibition. It has to be noted that this scenario very much reflects cancer cells that often highly overexpress c-FLIP proteins and, therefore, are resistant toward death receptor-mediated apoptosis.

Consistent with the model predictions, the 20 times increase of c-FLIP_L levels in these HeLa-CD95 cells overexpressing short c-FLIP isoforms, led to a significant sensitization toward CD95-mediated apoptosis¹⁹ (Figure 2b, lower panel). Apparently, the overexpression of short c-FLIP isoforms has created the optimal amount of procaspase-8-c-FLIP_L heterodimers at the DISC resulting in apoptosis sensitization. This also shows that short c-FLIP isoforms can have the pro-apoptotic role upon favorable stoichiometric conditions. Thus, the model by Fricker *et al.* unraveled the specific concentration range of c-FLIP_L and c-FLIP_S at the DISC leading to apoptosis acceleration (Figure 3). This work has also shown that the level of expression of a protein might define its cellular function contributing to the generic understanding of the origin of cell type specificity.

The model by Fricker *et al.* could uncover the concentration range of the c-FLIP_L isoform leading to its proapoptotic effect.¹⁹ This occurs upon intermediate levels of its overexpression, which is about 20 times higher than its endogenous level (Figures 2b and 3). Naturally, high overexpression of c-FLIP_L would lead to inhibition of procaspase-8 activation by outnumbering the procaspase-8 at the DISC and thereby preventing the formation of proapoptotic procaspase-8 homodimers and procaspase-8-c-FLIP_L heterodimers leading to apoptosis (Figure 3). The inhibiting function of c-FLIP_L upon high overexpression was supported by experimental data from a number of groups.^{12,19} Hence, c-FLIP_L has a very complicated action, it could promote or block CD95-induced apoptosis depending upon its concentration at the DISC. The model by Fricker *et al.* allowed to identify these stoichiometric switches in terms of exact numbers of c-FLIP proteins in the cell leading to the different systems behavior (Figure 3).

Interestingly, the concentration-specific action of c-FLIP_L also takes place upon induction of CD95-mediated NF-κB activation.³⁶ Here, we have shown that the generation of the

cleavage product p43-FLIP at the DISC is an essential step for this signal.^{36,37} p43-FLIP is generated at the CD95 DISC by procaspase-8 activity in the non-linear manner arising from the competitive binding of procaspase-8 and c-FLIP to the DISC. The model of the CD95-mediated NF-κB activation generated by Neumann *et al.* could incorporate this non-linear behavior and describe DISC dynamics of CD95-mediated apoptosis and NF-κB activation.³⁶ This ODE model supported by experimental data generated in HeLa-CD95 cells demonstrated three modes of c-FLIP_L action (Figure 4a). These three scenarios included low, intermediate and high levels of c-FLIP_L at the DISC. Low levels of c-FLIP_L disabled NF-κB as no p43-FLIP could be formed. Intermediate levels of c-FLIP_L at the DISC led to p43-FLIP generation and the transduction of the NF-κB signal. High concentrations of c-FLIP_L at the DISC blocked procaspase-8 activity resulting in the inhibition of p43-FLIP cleavage product generation and subsequently NF-κB activity (Figure 4a). Interestingly, similar to predictions of the model by Fricker *et al.* only intermediate levels of c-FLIP_L at the DISC resulted in the proactive function of c-FLIP_L, for example, promotion of CD95-mediated NF-κB activation.

The non-linearity of p43-FLIP generation by procaspase-8 at the DISC (Figure 4a) translates into the non-linearity of CD95-mediated apoptosis/NF-κB activation shown in the phase diagram (Figure 4b). Here, an important discovery has originated from *in silico* analysis. It was shown that the ratio between c-FLIP and procaspase-8 determines the regions of the predominant NF-κB and/or apoptosis induction in a highly non-linear manner (Figure 4b).

Thus, systems biology studies of the DISC have demonstrated the dual role of c-FLIP proteins. Coming back to 'live and let die' c-FLIP proteins have a role similar to 'secret agents': they can both kill or protect the cells from apoptosis depending upon the circumstances, which in this particular case are the concentrations of the molecules in the pathway.




Figure 3 Concentration-dependent effects of c-FLIP_L on apoptosis induction. The diagram presenting concentration-dependent effects of c-FLIP_L on the apoptosis induction. Generation of active caspase-8 heterotetramers p10₂-p18₂ is demonstrated as an indication of apoptosis




Figure 4 Non-linear dynamics of CD95 signaling. (a) The non-linear dynamics of the generation of p43-FLIP and p43/p41 caspase-8, leading to the induction of NF- κ B and apoptosis, respectively. (b) Non-linearity of life/death decisions at CD95 depending on the amounts of caspase-8 and c-FLIP. NF- κ B activity is shown in green, caspase-3 activation as an indication of apoptosis is shown in red. Adapted from Neumann *et al.*³⁶

Dynamics of ‘Chains of Death’: c-FLIP is not the Only Factor Determining Life and Death

Until recently, most modeling studies of the DISC have concentrated on the key role of c-FLIP in mediating life/death decisions at CD95.⁵ However, recent findings have demonstrated that c-FLIP is not the major factor defining apoptosis initiation. It was shown independently by two groups (Professor MacFarlane with coworkers and our group), that the generation of DED chains at the CD95 and TRAIL-R DISCs forms the platform for procaspase-8 activation and is another important constraint for DISC-mediated apoptosis initiation.^{14,15}

In order to understand DED chain dynamics at the DISC, a mathematical model based on agent-based modeling was generated by Schleich *et al.*¹⁵ The cell was modeled as a three-dimensional grid, and the recruitment probabilities of FADD, procaspase-8 and c-FLIP to the DISC were estimated proportional to the distance of these molecules to the receptors. The model could describe well procaspase-8 processing in the chains. Interestingly, in order to re-model experimental data, we had to introduce the parameter of the restricted chain length.¹⁵ Obviously, a non-restricted chain length could lead to cell death upon spontaneous activation of one death receptor via the formation of one rather long DED chain with a high number of procaspase-8 molecules. Therefore, the assumption of the model upon restricted chain length nicely fitted to the biological knowledge. The restriction of the chain length indicates, however, the existence of the unknown mechanism of the chain termination, which might result from some yet undiscovered structural constraints in the receptor vicinity. Another possibility for the chain termination might result from the DED protein numbers. Indeed, upon high stimulation strength, the majority of procaspase-8 molecules might be recruited into the chains. This might create the natural limitation of the chain length based on the procaspase-8 number in the cell.¹⁵ In terms of dynamics, the model predicted that the shorter chains are formed upon high stimulation strength, whereas low stimulation strength would

lead to longer chains. Indeed, in order to efficiently kill the cell upon low stimulation, a higher number of the active caspase-8 molecules is required, which in turn could be achieved by increasing the chain length. These predictions of the model were confirmed by experimental data using quantitative western blot and mass spectrometry analysis.¹⁵ Thus, the study by Schleich *et al.* shows that the CD95 DISC stoichiometry is very dynamic and strongly depends on the stimulation strength. These dynamics also preclude that one universal DISC stoichiometry exists, rather CD95 DISCs comprise DED chains of different length and composition defined by the amount of molecules in the cell and the stimulation strength.

Undoubtedly, the discovery of DED chains provides a new challenge for the DISC studies as many mechanisms have to be revised. The same is true for mathematical models of the DISC. Indeed, previous DISC models did not consider DED chains upon model construction and were based solely on competition binding mechanisms, between procaspase-8 and c-FLIP for binding to FADD at the DISC. Nevertheless, the *in silico* simulations of these models were confirmed quite well by experimental data. This likely reflects the fact that the procaspase-8/c-FLIP ratio is crucial for caspase-8 activation also in the context of the chains. The further development of computational studies of DED chains at the DISC should provide further insights into dynamics of apoptosis initiation. In addition, a number of questions has to be addressed in terms of molecular architecture of the chains. In particular, the role of c-FLIP has to be determined in the context of chains: what is the exact inhibitory mechanism of c-FLIP, whether c-FLIP-binding terminates the DED chain or promotes chain elongation. The other challenging questions are: whether DED chains are stable, how one can define their structure considering different isoforms/cleavage products of procaspase-8, procaspase-10 and c-FLIP; how affinities of different DED proteins in the chain and their composition stoichiometry influence the switch between life and death.

Modeling Extrinsic Pathways Mediated by Other Death Receptors and Downstream from the DISC

Modeling of the extrinsic apoptotic pathway downstream from the DISC for CD95 and other death receptors, in particular, TRAIL-R1/2, has provided further insights into death receptor biology. Here important knowledge came from the combination of single-cell analysis with mathematical modeling^{6,7,38} (reviewed in detail in this issue by Gaudet and co-workers³⁹). The development of the FRET-based caspase activity probes³⁸ and localization-based caspase activity probes⁴⁰ allowed to elegantly monitor caspase activation at the single-cell level. Using FRET-based caspase activity probes, it was demonstrated that in type II cells there is a lag time after death receptor stimulation, which was termed pre-MOMP delay.^{41,42} In this time, no activation of the effector caspases could be detected at the single-cell level; however, directly after the MOMP, a very fast processing of the FRET probe occurs, indicating that effector caspases are only active after MOMP. The same caspase activity probes allowed to shed light on the mechanism of cell-to-cell variability upon TRAIL induction.⁴³ A remarkable outcome of this study was that differences in cell death timing are based primarily on variations of protein expression levels of the cells.

TRAIL signaling provides another level of complexity compared with CD95L signaling because TRAIL binds to two death receptors TRAIL-R1 and TRAIL-R2 as well as to several decoy receptors.⁴⁴ The kinetic modeling of different TRAIL receptor complexes consisting of either hetero- or homotrimers was the subject of another modeling study by Szegezdi *et al.*⁴⁴ that found how to selectively enhance apoptosis by modulating the composition of receptor complexes. The crosstalk of TRAIL apoptotic signaling with non-apoptotic signaling pathways was the subject of a number of modeling studies that revealed the synergistic effects leading to more effective cell death.^{45–47} This is a very important aspect, as enhancement of TRAIL killing and overcoming TRAIL resistance in cancer cells could potentially have a major role in the development of contemporary anti-cancer therapies.

The modeling studies also addressed the mechanisms of bistability and the positive feedback loops in extrinsic apoptosis. The study by Eissing *et al.* developed a minimal model of caspase-8- and caspase-3-positive feedback in type I cells and has demonstrated that caspase activation in extrinsic apoptosis is characterized by a bistability behavior defined by the inhibitors of caspase activation.⁴⁸ Interestingly, bistability has also been found as a feature of ligand/receptor interactions in the theoretical modeling studies of death receptor oligomerization reactions by Ho *et al.*⁴⁹ Furthermore, this study unraveled that depending on local receptor densities the bistability can be characterized by reversible or irreversible behavior.⁴⁹ The hidden positive feedback loop in caspase activation has been discovered by the model of Legewie *et al.*⁵⁰ This model has investigated the interplay of caspases-3, -9 and XIAP and uncovered that activated caspase-3 sequesters XIAP from caspase-9, which, in turn, enhances caspase-9 activity and subsequently caspase-3 activation. An important contribution to the analysis of the positive feedback loop of caspase-8, -3 and -6 has been made in the study of Würstle *et al.*⁵¹ The authors have demonstrated

that caspase-8 dimerization/dissociation has a key role in the control of caspase activation events and avoiding of spontaneous apoptosis.

A central contribution to the field was the analysis of type I and type II signaling by Aldridge *et al.*⁵² The original report on type I and type II signaling has demonstrated that type I cells are characterized by high amount of caspase-8 formed at the DISC resulting in caspase-8-mediated caspase-3 activation and independence of the following apoptosis execution from the MOMP-dependent caspase activation cascade.⁵³ Later, Aldridge *et al.* have also analyzed the downstream modulators in type I and type II cells and have shown that the ratio of caspase-3 to XIAP has a central role in defining type I/type II.⁵² These seemingly contradictory statements in fact fit together quiet well. The caspase-3/XIAP ratio defines whether apoptosis could be executed without mitochondria activation. Indeed, upon high amounts of active caspase-3 activated by caspase-8 apoptosis could occur without MOMP. However, when the XIAP levels in the cell are high, only MOMP would allow to overcome apoptotic threshold, as the latter would lead to SMAC and cytochrome *c* release from mitochondria resulting in the higher amounts of active caspase-3 in the cell and subsequent apoptosis. Aldridge *et al.* have elegantly classified all cell lines using a phase diagram (separatrix) built on the ratio between caspase-3 and XIAP. Importantly, the prototypic type I cell line SKW6.4 appeared at the part of the diagram representing the lowest number of XIAP.⁵² The type II cells lines, in particular, colon carcinoma HCT116 cells, were located at the part of the separatrix corresponding to the higher amounts of XIAP, and, therefore, requiring the mitochondria activation to undergo apoptosis. Furthermore, Aldridge *et al.* have demonstrated that the deletion of XIAP in HCT116 cells turns them into type I cells that are independent from the mitochondria activation. Jost *et al.* have observed a similar switch from a type II to a type I phenotype in the hepatocytes of Bid^{-/-} XIAP^{-/-} mice.⁵⁴ Another interesting conclusion from the Aldridge study was that type I cells do not necessarily need high levels of caspase-8 formed at the DISC but are rather capable of undergoing apoptosis without mitochondrial involvement. This again shows that there is no contradiction between the two studies but the study by Aldridge *et al.* rather represents a more systematic insight into type I/type II problem.

Posttranslational modifications, such as ubiquitylation, phosphorylation, cleavage, glycosylation and others, have a key role in death receptor signaling control.² Cleavage as a central modification of apoptotic signaling has been addressed by most of the models created to date.⁵ Degradation of c-FLIP by ubiquitylation was investigated in a mathematical model of the dynamic turnover and stability of c-FLIP isoforms by Toivonen *et al.*³⁴ Bagci *et al.*⁵⁵ created a model of the bistability switch between apoptosis and survival based on the regulation of nitric oxide species and inhibition of caspase-8 and -3 by nitric oxide species. The crosstalk between phosphorylation and ubiquitylation is a highly relevant topic in contemporary biomedical research. In particular, the switch between apoptosis, necroptosis and survival signaling is highly dependent on this crosstalk.⁵⁶ Not surprisingly this question is currently in the center of systems biology studies. Finally, a remarkable progress has recently

been achieved in decoding the network of ubiquitin-ligases and deubiquitinylation enzymes controlling the ubiquitinylation status of death receptor platform components.^{57,58} Therefore, the next challenge in modeling will be to incorporate ubiquitinylation control into the existing models of death receptor signaling.

An important contribution to the understanding of the extrinsic apoptotic pathway was the development of Boolean models of death receptor signaling, which allowed to understand the mechanism of the crosstalk between TNF-R and CD95 stimulation in hepatocytes on a larger scale.⁵⁹ Indeed, the model by Schlatter *et al.* included 86 nodes and 125 reactions. The model predicted the cellular response upon stimulation with several stimuli: CD95L, TNF- α , UV-B irradiation, interleukin-1 β , insulin, and was validated by experiments in mouse hepatocytes and Jurkat T cells. This large model can reproduce well the apoptotic behavior, which was fascinating. Furthermore, the model has discovered high connectivity of different nodes of the apoptotic network and importance of the feedback loops. Importantly, a number of new ways to regulate the death receptor network were made. Furthermore, the application of Boolean models by Calzone *et al.* allowed to establish the mechanism of crosstalk between apoptosis, necroptosis and NF- κ B signaling pathways, which is yet another challenging question in modern death receptor biology.⁶⁰

Conclusions and Outlook

Systems biology of the extrinsic signaling pathway has developed a number of powerful models, which describe cell death *in silico* with an impressive level of detail. Death receptor-induced apoptosis can now be described in quantitative terms by the amounts of pro- and anti-apoptotic regulators in the apoptotic complexes, kinetic rates of cell death and the global quantitative view on the regulation of the network. Bistability switches, positive feedback loops in caspase activation and non-linear dynamics of apoptosis responses have been identified and thoroughly characterized by computational approaches. The models could perfectly well describe the regulation of the most important steps of death receptor activation and how subtle changes in the concentrations of apoptotic inhibitors can lead to a change in the outcome between life and death. This progress has to be further implemented for studying diseases connected to defects in the death receptor network. Identification of the most sensitive nodes in the death receptor pathways could allow to develop targeted therapies and new drugs that would allow to interfere with these diseases. There are already first impressive efforts in this direction, which so far mostly have addressed the regulation of the intrinsic apoptotic pathway.^{61–63} The future development of similar studies of the extrinsic pathway and connection of these models to personalized medicine could provide new horizons in systems medicine.

Conflict of Interest

The author declares no conflict of interest.

Acknowledgements. We acknowledge the Ministry of Sciences and Economic Affairs of Saxony-Anhalt (Research Centre Dynamic Systems:

Biosystems Engineering, MW -21LMS 5), BMBF (eBIO project “ImmunoQuant” - TPU - 0316170G), the Helmholtz-Russia Joint Research Groups—2008-2 and RFFI 14-04-00699 for supporting our work.

- Wilson NS, Dixit V, Ashkenazi A. Death receptor signal transducers: nodes of coordination in immune signaling networks. *Nat Immunol* 2009; **10**: 348–355.
- Lavrik IN, Krammer PH. Regulation of CD95/Fas signaling at the DISC. *Cell Death Differ* 2012; **19**: 36–41.
- Strasser A, Jost PJ, Nagata S. The many roles of FAS receptor signaling in the immune system. *Immunity* 2009; **30**: 180–192.
- Krammer PH, Arnold R, Lavrik IN. Life and death in peripheral T cells. *Nat Rev Immunol* 2007; **7**: 532–542.
- Lavrik IN. Systems biology of apoptosis signaling networks. *Curr Opin Biotechnol* 2010; **21**: 551–555.
- Rehm M, Prehn JH. Systems modelling methodology for the analysis of apoptosis signal transduction and cell death decisions. *Methods* 2013; **61**: 165–173.
- Spencer SL, Sorger PK. Measuring and modeling apoptosis in single cells. *Cell* 2011; **144**: 926–939.
- Lavrik IN, Eils R, Fricker N, Pforr C, Krammer PH. Understanding apoptosis by systems biology approaches. *Mol Biosyst* 2009; **5**: 1105–1111.
- Schleich K, Lavrik IN. Mathematical modeling of apoptosis. *Cell Commun Signal* 2013; **11**: 44.
- Janes KA, Albeck JG, Gaudet S, Sorger PK, Lauffenburger DA, Yaffe MB. A systems model of signaling identifies a molecular basis set for cytokine-induced apoptosis. *Science* 2005; **310**: 1646–1653.
- Kutumova EO, Kiselev IN, Sharipov RN, Lavrik IN, Kolpakov FA. A modular model of the apoptosis machinery. *Adv Exp Med Biol* 2012; **736**: 235–245.
- Ozturk S, Schleich K, Lavrik IN. Cellular FLICE-like inhibitory proteins (c-FLIPs): fine-tuners of life and death decisions. *Exp Cell Res* 2012; **318**: 1324–1331.
- Lavrik IN, Golks A, Krammer PH. Caspases: pharmacological manipulation of cell death. *J Clin Invest* 2005; **115**: 2665–2672.
- Dickens LS, Boyd RS, Jukes-Jones R, Hughes MA, Robinson GL, Fairall L *et al.* A death effector domain chain DISC model reveals a crucial role for caspase-8 chain assembly in mediating apoptotic cell death. *Mol Cell* 2012; **47**: 291–305.
- Schleich K *et al.* Stoichiometry of the CD95 death-inducing signaling complex: experimental and modeling evidence for a death effector domain chain model. *Mol Cell* 2012; **47**: 306–319.
- Schleich K, Krammer PH, Lavrik IN. The chains of death: a new view on caspase-8 activation at the DISC. *Cell Cycle* 2013; **12**: 193–194.
- Hoffmann JC, Pappa A, Krammer PH, Lavrik IN. A new C-terminal cleavage product of procaspase-8, p30, defines an alternative pathway of procaspase-8 activation. *Mol Cell Biol* 2009; **29**: 4431–4440.
- Golks A, Brenner D, Schmitz I, Watzl C, Krueger A, Krammer PH *et al.* The role of CAP3 in CD95 signaling: new insights into the mechanism of procaspase-8 activation. *Cell Death Differ* 2006; **13**: 489–498.
- Fricker N, Beaudouin J, Richter P, Eils R, Krammer PH, Lavrik IN. Model-based dissection of CD95 signaling dynamics reveals both a pro- and antiapoptotic role of c-FLIPL. *J Cell Biol* 2010; **190**: 377–389.
- Chang DW, Xing Z, Pan Y, Algeciras-Schimmich A, Barnhart BC, Yaish-Ohad S. c-FLIP(L) is a dual function regulator for caspase-8 activation and CD95-mediated apoptosis. *EMBO J* 2002; **21**: 3704–3714.
- Micheau O, Thome M, Schneider P, Holler N, Tschopp J, Nicholson DW *et al.* The long form of FLIP is an activator of caspase-8 at the Fas death-inducing signaling complex. *J Biol Chem* 2002; **277**: 45162–45171.
- Yu JW, Jeffrey PD, Shi Y. Mechanism of procaspase-8 activation by c-FLIPL. *Proc Natl Acad Sci USA* 2009; **106**: 8169–8174.
- Oberst A, Green DR. It cuts both ways: reconciling the dual roles of caspase 8 in cell death and survival. *Nat Rev Mol Cell Biol* 2011; **12**: 757–763.
- Gonzalez F, Pariselli F, Dupaigne P, Budihardjo I, Lutter M, Antonsson B *et al.* tBid interaction with cardiolipin primarily orchestrates mitochondrial dysfunctions and subsequently activates Bax and Bak. *Cell Death Differ* 2005; **12**: 614–626.
- Rehm M, Huber HJ, Hellwig CT, Anguissola S, Dussmann H, Prehn JH. Dynamics of outer mitochondrial membrane permeabilization during apoptosis. *Cell Death. Differ* 2009; **16**: 613–623.
- Kaufmann T, Strasser A, Jost PJ. Fas death receptor signalling: roles of Bid and XIAP. *Cell Death Differ* 2012; **19**: 42–50.
- Rehm M, Huber HJ, Dussmann H, Prehn JH. Systems analysis of effector caspase activation and its control by X-linked inhibitor of apoptosis protein. *EMBO J* 2006; **25**: 4338–4349.
- Kober AM, Legewie S, Pforr C, Fricker N, Eils R, Krammer PH *et al.* Caspase-8 activity has an essential role in CD95/Fas-mediated MAPK activation. *Cell Death Dis* 2011; **2**: e212.
- Cullen SP, Henry CM, Kearney CJ, Logue SE, Feoktistova M, Tynan GA *et al.* Fas/CD95-induced chemokines can serve as “find-me” signals for apoptotic cells. *Mol Cell* 2013; **49**: 1034–1048.
- Fussenegger M, Bailey JE, Varner J. A mathematical model of caspase function in apoptosis. *Nat Biotechnol* 2000; **18**: 768–774.

31. Bentele M, Lavrik I, Ulrich M, Stösser S, Heermann DW, Kalthoff H *et al*. Mathematical modeling reveals threshold mechanism in CD95-induced apoptosis. *J Cell Biol* 2004; **166**: 839–851.
32. Lavrik IN, Golks A, Riess D, Bentele M, Eils R, Krammer PH. Analysis of CD95 threshold signaling: triggering of CD95 (FAS/APO-1) at low concentrations primarily results in survival signaling. *J Biol Chem* 2007; **282**: 13664–13671.
33. Poukkula M, Kaunisto A, Hietakangas V, Denessiouk K, Katajamäki T, Johnson MS *et al*. Rapid turnover of c-FLIPshort is determined by its unique C-terminal tail. *J Biol Chem* 2005; **280**: 27345–27355.
34. Toivonen HT, Meinander A, Asaoka T, Westerlund M, Pettersson F, Mikhailov A *et al*. Modeling reveals that dynamic regulation of c-FLIP levels determines cell-to-cell distribution of CD95-mediated apoptosis. *J Biol Chem* 2011; **286**: 18375–18382.
35. Sharp DA, Lawrence DA, Ashkenazi A. Selective knockdown of the long variant of cellular FLICE inhibitory protein augments death receptor-mediated caspase-8 activation and apoptosis. *J Biol Chem* 2005; **280**: 19401–19409.
36. Neumann L, Pforr C, Beaudouin J, Pappa A, Fricker N, Krammer PH *et al*. Dynamics within the CD95 death-inducing signaling complex decide life and death of cells. *Mol Syst Biol* 2010; **6**: 352.
37. Pforr C, Neumann L, Eils R, Krammer PH, Lavrik IN. Understanding life and death at CD95. *Adv Exp Med Biol* 2011; **691**: 151–161.
38. Rehm M, Dussmann H, Janicke RU, Tavare JM, Kogel D, Prehn JH. Single-cell fluorescence resonance energy transfer analysis demonstrates that caspase activation during apoptosis is a rapid process. Role of caspase-3. *J Biol Chem* 2002; **277**: 24506–24514.
39. Xia X, Owen MS, Lee REC, Gaudet S. Cell-to-cell variability in cell death: can systems biology help us make sense of it all? *Cell Death Dis* 2014; doi:10.1038/cddis.2014.199.
40. Beaudouin J, Liesche C, Aschenbrenner S, Horner M, Eils R. Caspase-8 cleaves its substrates from the plasma membrane upon CD95-induced apoptosis. *Cell Death Differ* 2013; **20**: 599–610.
41. Albeck JG, Burke JM, Aldridge BB, Zhang M, Lauffenburger DA, Sorger PK. Quantitative analysis of pathways controlling extrinsic apoptosis in single cells. *Mol Cell* 2008; **30**: 11–25.
42. Albeck JG, Burke JM, Spencer SL, Lauffenburger DA, Sorger PK. Modeling a snap-action, variable-delay switch controlling extrinsic cell death. *PLoS Biol* 2008; **6**: 2831–2852.
43. Spencer SL, Gaudet S, Albeck JG, Burke JM, Sorger PK. Non-genetic origins of cell-to-cell variability in TRAIL-induced apoptosis. *Nature* 2009; **459**: 428–432.
44. Szegezdi E, van der Sloot AM, Mahalingam D, O'Leary L, Cool RH, Muñoz IG *et al*. Kinetics in signal transduction pathways involving promiscuous oligomerizing receptors can be determined by receptor specificity: apoptosis induction by TRAIL. *Mol Cell Proteomics* 2012; **11**: M111.
45. Laussmann MA, Passante E, Hellwig CT, Tomiczek B, Flanagan L, Prehn JH *et al*. Proteasome inhibition can impair caspase-8 activation upon submaximal stimulation of apoptotic tumor necrosis factor-related apoptosis inducing ligand (TRAIL) signaling. *J Biol Chem* 2012; **287**: 14402–14411.
46. Piras V, Hayashi K, Tomita M, Selvarajoo K. Enhancing apoptosis in TRAIL-resistant cancer cells using fundamental response rules. *Sci Rep* 2011; **1**: 144.
47. Shi Y, Mellier G, Huang S, White J, Pervaiz S, Tucker-Kellogg L. Computational modelling of LY303511 and TRAIL-induced apoptosis suggests dynamic regulation of cFLIP. *Bioinformatics* 2013; **29**: 347–354.
48. Eissing T, Conzelmann H, Gilles ED, Allgöwer F, Bullinger E, Scheurich P. Bistability analyses of a caspase activation model for receptor-induced apoptosis. *J Biol Chem* 2004; **279**: 36892–36897.
49. Ho KL, Harrington HA. Bistability in apoptosis by receptor clustering. *PLoS Comput Biol* 2010; **6**: e1000956.
50. Legewie S, Bluthgen N, Herzl H. Mathematical modeling identifies inhibitors of apoptosis as mediators of positive feedback and bistability. *PLoS Comput Biol* 2006; **2**: e120.
51. Wurstle ML, Laussmann MA, Rehm M. The caspase-8 dimerization/dissociation balance is a highly potent regulator of caspase-8, -3, -6 signaling. *J Biol Chem* 2010; **285**: 33209–33218.
52. Aldridge BB, Gaudet S, Lauffenburger DA, Sorger PK. Lyapunov exponents and phase diagrams reveal multi-factorial control over TRAIL-induced apoptosis. *Mol Syst Biol* 2011; **7**: 553.
53. Scaffidi C, Fulda S, Srinivasan A, Friesen C, Li F, Tomaselli KJ *et al*. Two CD95 (APO-1/Fas) signaling pathways. *EMBO J* 1998; **17**: 1675–1687.
54. Jost PJ, Grabow S, Gray D, McKenzie MD, Nachbur U, Huang DC *et al*. XIAP discriminates between type I and type II FAS-induced apoptosis. *Nature* 2009; **460**: 1035–1039.
55. Bagci EZ, Vodovotz Y, Billiar TR, Ermentrout B, Bahar I. Computational insights on the competing effects of nitric oxide in regulating apoptosis. *PLoS One* 2008; **3**: e2249.
56. Vandenberghe P, Declercq W, Van HF, Vanden Berghe T. The role of the kinases RIP1 and RIP3 in TNF-induced necrosis. *Sci Signal* 2010; **3**: re4.
57. Walczak H, Iwai K, Dikic I. Generation and physiological roles of linear ubiquitin chains. *BMC Biol* 2012; **10**: 23.
58. Emmerich CH, Schmukle AC, Walczak H. The emerging role of linear ubiquitination in cell signaling. *Sci Signal* 2011; **4**: re5.
59. Schlatter R, Schmich K, Avalos Vizcarra I, Scheurich P, Sauter T, Borner C *et al*. ON/OFF and beyond—a boolean model of apoptosis. *PLoS Comput Biol* 2009; **5**: e1000595.
60. Calzone L, Tournier L, Fourquet S, Thieffry D, Zhivotovskiy B, Barillot E *et al*. Mathematical modelling of cell-fate decision in response to death receptor engagement. *PLoS Comput Biol* 2010; **6**: e1000702.
61. Hector S, Rehm M, Schmid J, Kehoe J, McCawley N, Dicker P *et al*. Clinical application of a systems model of apoptosis execution for the prediction of colorectal cancer therapy responses and personalisation of therapy. *Gut* 2012; **61**: 725–733.
62. Passante E, Wurstle ML, Hellwig CT, Leverkus M, Rehm M. Systems analysis of apoptosis protein expression allows the case-specific prediction of cell death responsiveness of melanoma cells. *Cell Death Differ* 2013; **20**: 1521–1531.
63. Murphy AC, Weyhenmeyer B, Schmid J, Kilbride SM, Rehm M, Huber HJ *et al*. Activation of executioner caspases is a predictor of progression-free survival in glioblastoma patients: a systems medicine approach. *Cell Death Dis* 2013; **4**: e629.



Cell Death and Disease is an open-access journal published by Nature Publishing Group. This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>