



Concerns about how to simultaneously determine microvascular invasion and pathological response after transarterial chemoembolization before hepatocellular carcinoma surgery

Kwang Yeol Paik

Department of Surgery, Yeouido St. Mary's Hospital, The Catholic University of Korea College of Medicine, Seoul, Korea

Correspondence to: Kwang Yeol Paik, MD, PhD. Department of Surgery, Yeouido St. Mary's Hospital, The Catholic University of Korea College of Medicine, 62 Yeouido-dong, Yeongdeungpo-gu, Seoul 150-713, Korea. Email: kpaik@outlook.com.

Comment on: Yang Y, Dang Z, Lu P, *et al.* Impact of pathological response after preoperative transcatheter arterial chemoembolization (TACE) on incidences of microvascular invasion and early tumor recurrence in hepatocellular carcinoma: a multicenter propensity score matching analysis. *Hepatobiliary Surg Nutr* 2022;11:386-99.

Keywords: Microvascular invasion (MVI); transarterial chemoembolization (TACE); hepatocellular carcinoma (HCC)

Submitted Aug 28, 2023. Accepted for publication Sep 08, 2023. Published online Sep 18, 2023.

doi: 10.21037/hbsn-23-443

View this article at: <https://dx.doi.org/10.21037/hbsn-23-443>

Microvascular invasion (MVI) is classified as the strongest risk factor for recurrence and poor prognosis after surgical treatment of hepatocellular carcinoma (HCC). The same results after liver transplantation reflect how MVI reflects the poor prognosis of HCC (1,2).

However, MVI can only be evaluated via microscopic examination of the entire surgical specimen, limiting its use as a prognostic factor for treatment allocation in real practice, regardless of its clinical importance (3). Therefore, predicting MVI prior to the treatment stage is very important regarding selection of optimal treatment.

In this study, since the degree of response after transarterial chemoembolization (TACE) can only be confirmed by postoperative pathological results, it may also be difficult to apply it to determine the direction of treatment before surgery. Additionally, TACE as a neoadjuvant therapy for resectable HCC is not recommended, as the therapy is ineffective and the delay in surgery can turn a resectable HCC to unresectable (4,5).

Since both MVI and degree of response after TACE are known predictors of poor outcome after surgery, it does not appear that they are mutually opposed to each other. Of course, I would like to encourage this study to prove the relationship between the two proportionally for the first time.

This is very important finding in this study which is the

incidence of MVI in patients with complete pathological response (PR) was absent and the incidences of MVI gradually increased with decreasing percentages of tumor PR.

As authors suggested, there is still no direct evidence to indicate incomplete TACE induces MVI. Yang *et al.* (6) suggested TACE can be speculated to kill well-differentiated HCC cells while leaving behind those HCC cells which exhibit more aggressive behaviors, such as tumors with MVI and epithelial-to-mesenchymal transition can further promote formation of MVI and enhance HCC invasion and metastasis.

In order for the results of this study to stand out more meaningfully, there are two important points to be addressed in clinical practice. This is to clarify which patients had MVI before surgery and whether it is possible to know in advance which patients would have poor PR if TACE was performed. If the two risk factors coexist, we can assume a scenario in which surgical treatment is excluded or a more aggressive method is selected even if surgery is performed.

On post-TACE computed tomography (CT) imaging, a lack of residual contrast enhancement, a decrease in lesion size, a high lesion density with accumulation of ethiodized oil, and a diffuse distribution of ethiodized oil throughout the lesion correlated with near-complete lesion necrosis

upon histopathological analysis (7-9).

Author (6) suggested that patients with PR of <60% were shown in this study to have poorer long-term survival outcomes after liver resection, these patients should be closely followed-up, and post-operative adjuvant treatment using TACE or molecular targeted agents should be discussed with these patients.

In this respect, there are significant research findings. For patients with MVI, a resection margin of >1 cm resulted in better 5-year recurrence and overall survival than a resection margin under 1 cm (10).

However, it should be recognized that the responsiveness after TACE may have limitations in determining the selection of a patient's surgical treatment, since there are many groups that do not undergo TACE before surgery in the case of resectable HCC.

It is necessary to conduct a study to determine the direction of surgery by selecting a group that responds well by performing TACE in patients in whom resection is difficult or the effect of surgical treatment is ambiguous.

In this case, if MVI can be predicted at the same time, it can be much more useful in selecting patients. Going one step further, we also suggest a study to find out the response by performing TACE on patients with MVI predicted.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Hepatobiliary Surgery and Nutrition*. The article did not undergo external peer review.

Conflicts of Interest: The author has completed the ICMJE uniform disclosure form (available at <https://hbsn.amegroups.com/article/view/10.21037/hbsn-23-443/coif>). The author has no conflicts of interest to declare.

Ethical Statement: The author is accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons

Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Shetty K, Timmins K, Brensinger C, et al. Liver transplantation for hepatocellular carcinoma validation of present selection criteria in predicting outcome. *Liver Transpl* 2004;10:911-8.
- Iguchi T, Shirabe K, Aishima S, et al. New Pathologic Stratification of Microvascular Invasion in Hepatocellular Carcinoma: Predicting Prognosis After Living-donor Liver Transplantation. *Transplantation* 2015;99:1236-42.
- Lee S, Kang TW, Song KD, et al. Effect of Microvascular Invasion Risk on Early Recurrence of Hepatocellular Carcinoma After Surgery and Radiofrequency Ablation. *Ann Surg* 2021;273:564-71.
- Zhou WP, Lai EC, Li AJ, et al. A prospective, randomized, controlled trial of preoperative transarterial chemoembolization for resectable large hepatocellular carcinoma. *Ann Surg* 2009;249:195-202.
- Zhang X, Li J, Shen F, et al. Significance of presence of microvascular invasion in specimens obtained after surgical treatment of hepatocellular carcinoma. *J Gastroenterol Hepatol* 2018;33:347-54.
- Yang Y, Dang Z, Lu P, et al. Impact of pathological response after preoperative transcatheter arterial chemoembolization (TACE) on incidences of microvascular invasion and early tumor recurrence in hepatocellular carcinoma: a multicenter propensity score matching analysis. *Hepatobiliary Surg Nutr* 2022;11:386-99.
- Kwan SW, Fidelman N, Ma E, et al. Imaging predictors of the response to transarterial chemoembolization in patients with hepatocellular carcinoma: a radiological-pathological correlation. *Liver Transpl* 2012;18:727-36.
- Chapiro J, Wood LD, Lin M, et al. Radiologic-pathologic analysis of contrast-enhanced and diffusion-weighted MR imaging in patients with HCC after TACE: diagnostic accuracy of 3D quantitative image analysis. *Radiology* 2014;273:746-58.
- Najmi Varzaneh F, Pandey A, Aliyari Ghasabeh M, et al. Prediction of post-TACE necrosis of hepatocellular

carcinoma using volumetric enhancement on MRI and volumetric oil deposition on CT, with pathological correlation. *Eur Radiol* 2018;28:3032-40.

10. Yang P, Si A, Yang J, et al. A wide-margin liver resection

improves long-term outcomes for patients with HBV-related hepatocellular carcinoma with microvascular invasion. *Surgery* 2019;165:721-30.

Cite this article as: Paik KY. Concerns about how to simultaneously determine microvascular invasion and pathological response after transarterial chemoembolization before hepatocellular carcinoma surgery. *HepatoBiliary Surg Nutr* 2023;12(5):815-817. doi: 10.21037/hbsn-23-443