

Draft Genome Sequence of the Arsenate-Respiring Bacterium *Chrysiogenes arsenatis* Strain DSM 11915

David A. Coil,^a Jonathon R. Lo,^a Roger Chen,^{a*} Naomi Ward,^b Frank T. Robb,^c Jonathan A. Eisen^{a,d}

University of California Davis Genome Center, Davis, California, USA^a; University of Wyoming, Laramie, Wyoming, USA^b; University of Maryland School of Medicine, Baltimore, Maryland, USA^c; University of California Davis, Department of Evolution and Ecology, Department of Medical Microbiology and Immunology, Davis, California, USA^d

* Present address: Roger Chen, Stanford University, Stanford, California, USA.

Here we present the draft genome sequence of *Chrysiogenes arsenatis* strain DSM 11915, only the second genome sequence from the phylum *Chrysiogenetes*. This strictly anaerobic organism was isolated from arsenic-contaminated gold mine wastewater and respire arsenate or nitrate instead of oxygen. The assembly contains 2,824,977 bp in 22 scaffolds.

Received 9 October 2013 Accepted 10 October 2013 Published 14 November 2013

Citation Coil DA, Lo JR, Chen R, Ward N, Robb FT, Eisen JA. 2013. Draft genome sequence of the arsenate-respiring bacterium *Chrysiogenes arsenatis* strain DSM 11915. *Genome Announc.* 1(6):e00953-13. doi:10.1128/genomeA.00953-13.

Copyright © 2013 Coil et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Jonathan A. Eisen, jaisen@ucdavis.edu.

Chrysiogenes arsenatis is a curved, rod-shaped, Gram-negative, strictly anaerobic bacterium that was isolated from arsenic-contaminated mud of a reed bed in the Ballarat Goldfields, Australia (1). Unusually, *C. arsenatis* uses acetate as the electron donor and carbon source, and arsenate as the preferred electron acceptor, producing CO₂ and arsenite (2). Nitrate or nitrite can also be used as electron acceptors, producing NH₄. At the time of isolation, this species was the only member of the *Chrysiogenetes* phylum (3) but has since been joined by members of the genera *Desulfurispira* and *Desulfurispirillum* (4, 5). The only other member of this phylum to be sequenced is *Desulfurispirillum indicum* (6). *C. arsenatis* was selected in 2002 as part of a project at the Institute for Genomic Research (TIGR) to sequence the genomes of representatives of the seven phyla of bacteria that at the time had cultured representatives but no available genome sequence. Sanger clone-based sequencing of the genome failed to produce a high-quality assembly, and the project was delayed pending additional sequencing with Illumina technology.

DNA from *C. arsenatis* strain DSM 11915 was extracted at Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) (German Collection of Microorganisms and Cell Cultures) via lysis with lysozyme and proteinase K, phenol and chloroform extraction, and purification with Prep-A-Gene (7). Sanger libraries (small and medium insert and fosmid) were constructed as previously described (8). Illumina paired-end libraries were made from sonicated DNA using standard Illumina protocols and reagents.

A hybrid genome assembly was generated using MIRA (version 3.9.17) (9), with 33,005 Sanger reads (average length, 1,037 bp) and 4,000,000 Illumina reads (average length, 84 bp) that were randomly subsampled from the 41,433,640 total Illumina reads sequenced. The contigs resulting from this assembly were filtered to remove contigs shorter than 500 bp as well as those with >33% of the average coverage for the assembly. This resulted in 51 contigs which, along with 37,362,734 error-corrected/quality-

trimmed Illumina reads generated by the A5 assembly pipeline (10), were used to build 22 scaffolds in SSPACE (11). These Illumina reads were then aligned to the scaffolds using Bowtie (12) to eliminate artifacts in the assembly. One such region (~700 bp) was found, which by BLAST (13) analysis appeared to be a vector sequence and was removed from the assembly. During the creation of GenBank submission files, some contigs were merged based on short overlaps and read pair information, yielding a final collection of 23 contigs. The final assembly contains 2,824,977 bp and has a GC content of 50% and coverage estimates of 12× (Sanger) and 120× (Illumina).

Completeness of the genome was assessed using the PhyloSift software (A. Darling, G. Jospin, E. Lowe, E. Matsen, H. Bik, and J. Eisen, submitted for publication), which searches for 40 highly conserved, single-copy marker genes (14). All 40 genes were found in this assembly. Automated annotation was performed using the RAST annotation server (15). *C. arsenatis* DSM 11915 contains 2,592 predicted coding sequences and 43 predicted RNAs.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [AWNK00000000](http://www.ncbi.nlm.nih.gov/nuccore/AWNK00000000). The version described in this paper is version [AWNK01000000](http://www.ncbi.nlm.nih.gov/nuccore/AWNK01000000). Complete Illumina and Sanger reads are available at <http://dx.doi.org/10.6084/m9.figshare.799759>.

ACKNOWLEDGMENTS

Illumina sequencing was performed at the DNA Technologies Core facility at the University of California Davis. Sanger sequencing was performed at the Institute for Genomic Research (TIGR) in Rockville, MD.

We thank the many others who contributed to this project over the last decade, including Jenna Lang, Jonathan Badger, Guillaume Jospin, Aaron Darling, Andrew Tritt, Ryan Maples, Russell Neches, Andrew Kassen, Martin Wu, Julie Enticknap, Liz O'Connor, Hoda Khouri, Jan Weidman, Yasmin Mohamoud, Grace Pai, Shannon Smith, Tamara Feldblum, Terry Utterback, Jason Inman, and Mihai Pop.

This work was funded by the National Science Foundation "Assem-

bling the Tree of Life” grant number 0228651, overseen by Jonathan A. Eisen and Naomi Ward.

REFERENCES

- Macy JM, Nunan K, Hagen KD, Dixon DR, Harbour PJ, Cahill M, Sly LI. 1996. *Chrysiogenes arsenatis* gen. nov., sp. nov., a new arsenate-respiring bacterium isolated from gold mine wastewater. *Int. J. Syst. Bacteriol.* 46:1153–1157.
- Krafft T, Macy JM. 1998. Purification and characterization of the respiratory arsenate reductase of *Chrysiogenes arsenatis*. *Eur. J. Biochem.* 255: 647–653.
- Garrity GM, Holt JG. 2001. Phylum BV. *Chrysiogenetes* ph. nov., p 421–425. In Garrity GM, Boone DR, Castenholz RW (ed), *Bergey’s manual of systematic bacteriology*, 2nd ed, vol 1. Springer, New York, NY.
- Sorokin DY, Foti M, Tindall BJ, Muyzer G. 2007. *Desulfurispirillum alkaliphilum* gen. nov. sp. nov., a novel obligately anaerobic sulfur- and dissimilatory nitrate-reducing bacterium from a full-scale sulfide-removing bioreactor. *Extremophiles* 11:363–370.
- Sorokin DY, Muyzer G. 2010. *Desulfurispira natronophila* gen. nov. sp. nov.: an obligately anaerobic dissimilatory sulfur-reducing bacterium from soda lakes. *Extremophiles* 14:349–355.
- Bini E, Rauschenbach I, Narasingarao P, Starovoytov V, Hauser L, Jeffries CD, Land M, Bruce D, Detter C, Goodwin L, Han S, Held B, Tapia R, Copeland A, Ivanova N, Mikhailova N, Nolan M, Pati A, Pennacchio L, Pitluck S, Woyke T, Haggblom M. 2011. Complete genome sequence of *Desulfurispirillum indicum* strain S5(T). *Stand. Genomic Sci.* 5:371–378.
- Willis EH, Mardis ER, Jones WL, Little MC. 1990. Prep-A-Gene: a superior matrix for the purification of DNA and DNA fragments. *Bio-Techniques* 9:92–99.
- Wu D, Daugherty SC, Van Aken SE, Pai GH, Watkins KL, Khouri H, Tallon LJ, Zaborsky JM, Dunbar HE, Tran PL, Moran NA, Eisen JA. 2006. Metabolic complementarity and genomics of the dual bacterial symbiosis of sharpshooters. *PLoS Biol.* 4:e188. doi:10.1371/journal.pbio.0040188.
- Chevreur B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. *In* Computer science and biology: proceedings of the German Conference on Bioinformatics. Research Centre for Biotechnology (GBF), Braunschweig, Germany.
- Tritt A, Eisen JA, Facciotti MT, Darling AE. 2012. An integrated pipeline for de novo assembly of microbial genomes. *PLoS One* 7:e42304.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27: 578–579.
- Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410.
- Wu D, Jospin G, Eisen J. Systematic identification of gene families for use as markers for phylogenetic and phylogeny-driven ecological studies of bacteria and archaea and their major subgroups. *PLoS One*, in press.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.