

3-(4-Chlorophenyl)-1-(2-methyl-4-phenylquinolin-3-yl)prop-2-en-1-one

R. Prasath,^a P. Bhavana,^a Ray J. Butcher^{b*} and Jerry P. Jasinski^c

^aChemistry Group, BITS, Pilani, K. K. Birla Goa Campus, Goa 403 726, India,

^bDepartment of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA, and ^cDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA

Correspondence e-mail: rbutcher99@yahoo.com


Received 30 January 2011; accepted 7 February 2011

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.049; wR factor = 0.160; data-to-parameter ratio = 16.0.

The crystal structure of the title compound, $\text{C}_{25}\text{H}_{18}\text{ClNO}$, shows that the molecules are isolated and not involved in intermolecular $\text{C}-\text{H}\cdots\text{O}$ or $\text{C}-\text{H}\cdots\text{Cl}$ interactions. However, the phenyl and quinoline rings are involved in $\pi-\pi$ interactions [centroid-centroid distance = 3.8829 (9) Å].

Related literature

For background details and the biological activity of quinolines, see: Markees *et al.* (1970); Campbell *et al.* (1998); Bhat *et al.* (2005). For the biological activity of chalcones, see: Wu *et al.* (2006).



Experimental

Crystal data

$\text{C}_{25}\text{H}_{18}\text{ClNO}$	$\gamma = 107.035$ (3)°
$M_r = 383.85$	$V = 976.36$ (6) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 6.5376$ (2) Å	Cu $K\alpha$ radiation
$b = 10.0345$ (4) Å	$\mu = 1.84$ mm ⁻¹
$c = 15.6545$ (6) Å	$T = 295$ K
$\alpha = 90.845$ (3)°	$0.52 \times 0.18 \times 0.12$ mm
$\beta = 95.521$ (3)°	

Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer	7607 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2009)	4065 independent reflections
$T_{\min} = 0.544$, $T_{\max} = 1.000$	3402 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$	254 parameters
$wR(F^2) = 0.160$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\max} = 0.29$ e Å ⁻³
4065 reflections	$\Delta\rho_{\min} = -0.24$ e Å ⁻³

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

RJB wishes to acknowledge the NSF-MRI program (grant CHE-0619278) for funds to purchase the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZZ231).

References

- Bhat, B. A., Dhar, K. L., Puri, S. C., Saxena, A. K., Shanmugavel, M. & Qazi, G. N. (2005). *Bioorg. Med. Chem. Lett.* **15**, 3177–3180.
- Campbell, S. F., Hardstone, J. D. & Palmer, M. J. (1998). *J. Med. Chem.* **31**, 1031–1035.
- Markees, D. G., Dewey, V. C. & Kidder, G. W. (1970). *J. Med. Chem.* **13**, 324–326.
- Oxford Diffraction (2009). *CrysAlis PRO*, *CrysAlis RED* and *CrysAlis CCD*. Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wu, X., Tiekink, E. R. T., Iouri, K., Nikolai, K. & Mei, L. G. (2006). *Eur. J. Pharm. Sci.* **27**, 175–187.

supplementary materials

Acta Cryst. (2011). E67, o621 [doi:10.1107/S1600536811004661]

3-(4-Chlorophenyl)-1-(2-methyl-4-phenylquinolin-3-yl)prop-2-en-1-one

R. Prasath, P. Bhavana, R. J. Butcher and J. P. Jasinski

Comment

Quinoline derivatives are very important compounds because of their wide occurrence in natural products and biologically active compounds (Markees *et al.*, 1970; Campbell *et al.*, 1998). Additionally, chalcone derivatives are attracting the increasing interest of many researchers, because of their bioactivity such as antimicrobial, antimalarial, anticancer, antiviral, antitumor activities (Bhat *et al.*, 2005). Introduction of the quinoline scaffold into chalcone compounds can bring about significant changes in biological effects (Wu *et al.*, 2006).

The crystal structure of the title compound shows that the molecules are isolated and not involved in intermolecular interactions. However, both the phenyl ring and the quinoline rings are involved in π - π interactions (centroid to centroid distances of 3.428 (2) and 3.770 (2) Å, respectively).

Experimental

A mixture of 3-acetyl-2-methyl-4-phenylquinoline (2.61 g, 0.01 M), 4-chlorobenzaldehyde (1.40 g, 0.01 M) and KOH (1.12 g, 0.02 M) in distilled ethanol (20 ml) was stirred for 12 h at room temperature. The resulting mixture was neutralized with dilute acetic acid. The resultant solid was filtered, dried and purified by column chromatography using 1:1 mixture of ethyl acetate and hexane. Re-crystallization was by slow evaporation of acetone solution of (I) which yielded colourless needle type crystals. M.pt. 453-455 K. Yield: 72%.

Refinement

H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with a C—H distances of 0.93 and 0.96 Å $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and 0.98 Å for CH₃ [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$].

Figures




Fig. 1. Diagram of the title compound, C₂₅H₁₈ClNO, showing atom labeling.




Fig. 2. The molecular packing for C₂₅H₁₈ClNO viewed down the *a* axis.

3-(4-Chlorophenyl)-1-(2-methyl-4-phenylquinolin-3-yl)prop-2-en-1-one

Crystal data

$C_{25}H_{18}ClNO$	$Z = 2$
$M_r = 383.85$	$F(000) = 400$
Triclinic, PT	$D_x = 1.306 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$
$a = 6.5376 (2) \text{ \AA}$	Cell parameters from 4929 reflections
$b = 10.0345 (4) \text{ \AA}$	$\theta = 5.3\text{--}77.1^\circ$
$c = 15.6545 (6) \text{ \AA}$	$\mu = 1.84 \text{ mm}^{-1}$
$\alpha = 90.845 (3)^\circ$	$T = 295 \text{ K}$
$\beta = 95.521 (3)^\circ$	Needle, colorless
$\gamma = 107.035 (3)^\circ$	$0.52 \times 0.18 \times 0.12 \text{ mm}$
$V = 976.36 (6) \text{ \AA}^3$	

Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer	4065 independent reflections
Radiation source: Enhance (Cu) X-ray Source graphite	3402 reflections with $I > 2\sigma(I)$
Detector resolution: $10.5081 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.019$
ω scans	$\theta_{\text{max}} = 77.4^\circ$, $\theta_{\text{min}} = 5.3^\circ$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2009)	$h = -3 \rightarrow 8$
$T_{\text{min}} = 0.544$, $T_{\text{max}} = 1.000$	$k = -12 \rightarrow 12$
7607 measured reflections	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.049$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.160$	H-atom parameters constrained
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.1125P)^2 + 0.0628P]$
4065 reflections	where $P = (F_o^2 + 2F_c^2)/3$
254 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.29 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.41045 (10)	-0.24622 (7)	0.99317 (4)	0.1004 (3)
O	0.6524 (2)	0.15628 (12)	0.65464 (9)	0.0678 (3)
N	0.22876 (19)	0.41447 (12)	0.55076 (8)	0.0485 (3)
C1	0.0479 (2)	0.02060 (15)	0.82219 (9)	0.0505 (3)
C2	0.0750 (3)	-0.10510 (17)	0.85056 (12)	0.0603 (4)
H2A	0.1889	-0.1338	0.8340	0.072*
C3	-0.0651 (3)	-0.18756 (18)	0.90290 (12)	0.0671 (4)
H3A	-0.0465	-0.2714	0.9212	0.081*
C4	-0.2326 (3)	-0.14364 (19)	0.92754 (11)	0.0638 (4)
C5	-0.2633 (3)	-0.0196 (2)	0.90095 (13)	0.0696 (4)
H5A	-0.3765	0.0092	0.9181	0.083*
C6	-0.1218 (3)	0.06090 (18)	0.84822 (12)	0.0620 (4)
H6A	-0.1416	0.1444	0.8299	0.074*
C7	0.1907 (3)	0.11066 (14)	0.76588 (10)	0.0515 (3)
H7A	0.1646	0.1949	0.7532	0.062*
C8	0.3530 (3)	0.08465 (14)	0.73113 (10)	0.0543 (4)
H8A	0.3797	0.0001	0.7420	0.065*
C9	0.4934 (2)	0.18093 (14)	0.67652 (10)	0.0488 (3)
C10	0.4416 (2)	0.31192 (13)	0.64783 (8)	0.0434 (3)
C11	0.2682 (2)	0.30333 (14)	0.58322 (9)	0.0463 (3)
C12	0.1198 (3)	0.16483 (16)	0.54764 (11)	0.0588 (4)
H12A	0.0640	0.1744	0.4898	0.088*
H12B	0.0030	0.1345	0.5824	0.088*
H12C	0.1978	0.0973	0.5480	0.088*
C13	0.3573 (2)	0.54308 (14)	0.58070 (8)	0.0447 (3)
C14	0.3076 (3)	0.66120 (16)	0.54642 (10)	0.0547 (4)
H14A	0.1939	0.6494	0.5037	0.066*
C15	0.4265 (3)	0.79222 (17)	0.57609 (12)	0.0622 (4)
H15A	0.3928	0.8693	0.5535	0.075*
C16	0.5992 (3)	0.81211 (15)	0.64035 (12)	0.0596 (4)
H16A	0.6782	0.9020	0.6602	0.072*
C17	0.6522 (3)	0.70013 (15)	0.67399 (10)	0.0511 (3)
H17A	0.7675	0.7145	0.7162	0.061*

supplementary materials

C18	0.5324 (2)	0.56215 (13)	0.64491 (8)	0.0422 (3)
C19	0.5751 (2)	0.44052 (13)	0.67829 (8)	0.0414 (3)
C20	0.7569 (2)	0.45328 (13)	0.74642 (8)	0.0437 (3)
C21	0.9681 (3)	0.50871 (19)	0.72893 (11)	0.0599 (4)
H21A	0.9977	0.5393	0.6744	0.072*
C22	1.1357 (3)	0.5186 (2)	0.79260 (14)	0.0710 (5)
H22A	1.2772	0.5567	0.7807	0.085*
C23	1.0937 (3)	0.4725 (2)	0.87317 (12)	0.0673 (5)
H23A	1.2066	0.4777	0.9153	0.081*
C24	0.8847 (3)	0.41878 (18)	0.89139 (11)	0.0623 (4)
H24A	0.8563	0.3886	0.9461	0.075*
C25	0.7164 (2)	0.40945 (15)	0.82857 (9)	0.0512 (3)
H25A	0.5753	0.3736	0.8414	0.061*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl	0.0863 (4)	0.1060 (5)	0.0916 (4)	-0.0055 (3)	0.0264 (3)	0.0365 (3)
O	0.0731 (7)	0.0532 (6)	0.0893 (8)	0.0308 (6)	0.0285 (6)	0.0164 (6)
N	0.0481 (6)	0.0462 (6)	0.0503 (6)	0.0133 (5)	0.0028 (5)	0.0012 (5)
C1	0.0537 (8)	0.0416 (7)	0.0515 (7)	0.0073 (6)	0.0036 (6)	0.0039 (5)
C2	0.0624 (9)	0.0487 (8)	0.0697 (10)	0.0146 (7)	0.0107 (7)	0.0116 (7)
C3	0.0731 (11)	0.0506 (8)	0.0718 (10)	0.0093 (7)	0.0057 (8)	0.0178 (7)
C4	0.0599 (9)	0.0639 (9)	0.0542 (8)	-0.0025 (7)	0.0051 (7)	0.0109 (7)
C5	0.0627 (10)	0.0746 (11)	0.0719 (11)	0.0178 (8)	0.0168 (8)	0.0097 (8)
C6	0.0667 (10)	0.0536 (8)	0.0677 (9)	0.0189 (7)	0.0126 (7)	0.0111 (7)
C7	0.0600 (8)	0.0358 (6)	0.0576 (8)	0.0121 (6)	0.0061 (6)	0.0067 (5)
C8	0.0636 (9)	0.0368 (6)	0.0634 (8)	0.0146 (6)	0.0107 (7)	0.0104 (6)
C9	0.0551 (8)	0.0377 (6)	0.0540 (7)	0.0135 (6)	0.0082 (6)	0.0043 (5)
C10	0.0478 (7)	0.0374 (6)	0.0460 (6)	0.0120 (5)	0.0112 (5)	0.0043 (5)
C11	0.0475 (7)	0.0404 (6)	0.0497 (7)	0.0102 (5)	0.0079 (5)	-0.0003 (5)
C12	0.0581 (8)	0.0442 (7)	0.0668 (9)	0.0065 (6)	-0.0008 (7)	-0.0035 (6)
C13	0.0488 (7)	0.0419 (6)	0.0454 (6)	0.0152 (5)	0.0093 (5)	0.0047 (5)
C14	0.0580 (8)	0.0511 (8)	0.0578 (8)	0.0214 (6)	0.0024 (6)	0.0100 (6)
C15	0.0742 (10)	0.0447 (7)	0.0734 (10)	0.0255 (7)	0.0080 (8)	0.0132 (7)
C16	0.0712 (10)	0.0368 (7)	0.0684 (9)	0.0124 (6)	0.0065 (7)	0.0008 (6)
C17	0.0579 (8)	0.0413 (7)	0.0522 (7)	0.0121 (6)	0.0040 (6)	0.0016 (5)
C18	0.0492 (7)	0.0374 (6)	0.0411 (6)	0.0127 (5)	0.0099 (5)	0.0036 (5)
C19	0.0466 (6)	0.0390 (6)	0.0403 (6)	0.0134 (5)	0.0099 (5)	0.0039 (5)
C20	0.0491 (7)	0.0386 (6)	0.0454 (6)	0.0153 (5)	0.0068 (5)	0.0021 (5)
C21	0.0529 (8)	0.0722 (10)	0.0563 (8)	0.0185 (7)	0.0134 (6)	0.0028 (7)
C22	0.0461 (8)	0.0879 (13)	0.0803 (12)	0.0219 (8)	0.0073 (8)	-0.0080 (9)
C23	0.0642 (10)	0.0716 (10)	0.0683 (10)	0.0301 (8)	-0.0129 (8)	-0.0077 (8)
C24	0.0742 (10)	0.0602 (9)	0.0508 (8)	0.0198 (8)	-0.0021 (7)	0.0065 (6)
C25	0.0544 (8)	0.0476 (7)	0.0488 (7)	0.0108 (6)	0.0058 (6)	0.0060 (5)

Geometric parameters (\AA , $^\circ$)

Cl—C4	1.7399 (17)	C12—H12C	0.9600
-------	-------------	----------	--------

O—C9	1.2149 (19)	C13—C18	1.4149 (19)
N—C11	1.3147 (18)	C13—C14	1.4173 (19)
N—C13	1.3646 (18)	C14—C15	1.365 (2)
C1—C6	1.380 (2)	C14—H14A	0.9300
C1—C2	1.397 (2)	C15—C16	1.403 (2)
C1—C7	1.467 (2)	C15—H15A	0.9300
C2—C3	1.384 (2)	C16—C17	1.367 (2)
C2—H2A	0.9300	C16—H16A	0.9300
C3—C4	1.379 (3)	C17—C18	1.4196 (19)
C3—H3A	0.9300	C17—H17A	0.9300
C4—C5	1.381 (3)	C18—C19	1.4253 (17)
C5—C6	1.385 (2)	C19—C20	1.4929 (18)
C5—H5A	0.9300	C20—C21	1.385 (2)
C6—H6A	0.9300	C20—C25	1.390 (2)
C7—C8	1.326 (2)	C21—C22	1.388 (2)
C7—H7A	0.9300	C21—H21A	0.9300
C8—C9	1.471 (2)	C22—C23	1.375 (3)
C8—H8A	0.9300	C22—H22A	0.9300
C9—C10	1.5142 (18)	C23—C24	1.374 (3)
C10—C19	1.3770 (18)	C23—H23A	0.9300
C10—C11	1.4267 (19)	C24—C25	1.384 (2)
C11—C12	1.5053 (19)	C24—H24A	0.9300
C12—H12A	0.9600	C25—H25A	0.9300
C12—H12B	0.9600		
C11—N—C13	118.73 (12)	N—C13—C18	122.82 (12)
C6—C1—C2	118.20 (15)	N—C13—C14	117.62 (13)
C6—C1—C7	118.85 (14)	C18—C13—C14	119.55 (13)
C2—C1—C7	122.95 (14)	C15—C14—C13	120.05 (14)
C3—C2—C1	120.99 (16)	C15—C14—H14A	120.0
C3—C2—H2A	119.5	C13—C14—H14A	120.0
C1—C2—H2A	119.5	C14—C15—C16	120.79 (14)
C4—C3—C2	119.06 (16)	C14—C15—H15A	119.6
C4—C3—H3A	120.5	C16—C15—H15A	119.6
C2—C3—H3A	120.5	C17—C16—C15	120.43 (14)
C3—C4—C5	121.40 (16)	C17—C16—H16A	119.8
C3—C4—C1	119.65 (14)	C15—C16—H16A	119.8
C5—C4—C1	118.96 (15)	C16—C17—C18	120.51 (14)
C4—C5—C6	118.56 (17)	C16—C17—H17A	119.7
C4—C5—H5A	120.7	C18—C17—H17A	119.7
C6—C5—H5A	120.7	C13—C18—C17	118.66 (12)
C1—C6—C5	121.79 (16)	C13—C18—C19	117.70 (12)
C1—C6—H6A	119.1	C17—C18—C19	123.63 (13)
C5—C6—H6A	119.1	C10—C19—C18	118.40 (12)
C8—C7—C1	126.86 (13)	C10—C19—C20	121.16 (12)
C8—C7—H7A	116.6	C18—C19—C20	120.42 (11)
C1—C7—H7A	116.6	C21—C20—C25	118.93 (14)
C7—C8—C9	124.21 (13)	C21—C20—C19	120.71 (13)
C7—C8—H8A	117.9	C25—C20—C19	120.37 (13)
C9—C8—H8A	117.9	C20—C21—C22	120.14 (16)

supplementary materials

O—C9—C8	120.17 (13)	C20—C21—H21A	119.9
O—C9—C10	119.62 (13)	C22—C21—H21A	119.9
C8—C9—C10	120.21 (12)	C23—C22—C21	120.37 (16)
C19—C10—C11	119.78 (12)	C23—C22—H22A	119.8
C19—C10—C9	119.63 (12)	C21—C22—H22A	119.8
C11—C10—C9	120.30 (12)	C24—C23—C22	119.90 (16)
N—C11—C10	122.56 (12)	C24—C23—H23A	120.1
N—C11—C12	116.00 (13)	C22—C23—H23A	120.1
C10—C11—C12	121.44 (13)	C23—C24—C25	120.18 (16)
C11—C12—H12A	109.5	C23—C24—H24A	119.9
C11—C12—H12B	109.5	C25—C24—H24A	119.9
H12A—C12—H12B	109.5	C24—C25—C20	120.47 (15)
C11—C12—H12C	109.5	C24—C25—H25A	119.8
H12A—C12—H12C	109.5	C20—C25—H25A	119.8
H12B—C12—H12C	109.5		
C6—C1—C2—C3	0.4 (3)	C13—C14—C15—C16	0.2 (3)
C7—C1—C2—C3	-179.28 (15)	C14—C15—C16—C17	0.4 (3)
C1—C2—C3—C4	-0.5 (3)	C15—C16—C17—C18	-0.5 (3)
C2—C3—C4—C5	0.1 (3)	N—C13—C18—C17	-177.84 (13)
C2—C3—C4—C1	179.88 (13)	C14—C13—C18—C17	0.8 (2)
C3—C4—C5—C6	0.2 (3)	N—C13—C18—C19	0.69 (19)
C1—C4—C5—C6	-179.55 (14)	C14—C13—C18—C19	179.30 (12)
C2—C1—C6—C5	-0.1 (3)	C16—C17—C18—C13	-0.1 (2)
C7—C1—C6—C5	179.63 (16)	C16—C17—C18—C19	-178.57 (13)
C4—C5—C6—C1	-0.2 (3)	C11—C10—C19—C18	1.17 (19)
C6—C1—C7—C8	-176.02 (16)	C9—C10—C19—C18	174.99 (11)
C2—C1—C7—C8	3.7 (3)	C11—C10—C19—C20	179.45 (11)
C1—C7—C8—C9	-178.63 (14)	C9—C10—C19—C20	-6.74 (19)
C7—C8—C9—O	172.12 (16)	C13—C18—C19—C10	-1.45 (18)
C7—C8—C9—C10	-7.4 (2)	C17—C18—C19—C10	177.00 (13)
O—C9—C10—C19	-67.03 (19)	C13—C18—C19—C20	-179.74 (11)
C8—C9—C10—C19	112.49 (15)	C17—C18—C19—C20	-1.3 (2)
O—C9—C10—C11	106.75 (17)	C10—C19—C20—C21	114.65 (16)
C8—C9—C10—C11	-73.73 (18)	C18—C19—C20—C21	-67.11 (18)
C13—N—C11—C10	-0.7 (2)	C10—C19—C20—C25	-65.29 (17)
C13—N—C11—C12	179.73 (13)	C18—C19—C20—C25	112.95 (15)
C19—C10—C11—N	-0.1 (2)	C25—C20—C21—C22	0.6 (2)
C9—C10—C11—N	-173.84 (13)	C19—C20—C21—C22	-179.32 (15)
C19—C10—C11—C12	179.44 (13)	C20—C21—C22—C23	0.6 (3)
C9—C10—C11—C12	5.7 (2)	C21—C22—C23—C24	-1.3 (3)
C11—N—C13—C18	0.4 (2)	C22—C23—C24—C25	0.8 (3)
C11—N—C13—C14	-178.23 (13)	C23—C24—C25—C20	0.5 (2)
N—C13—C14—C15	177.86 (15)	C21—C20—C25—C24	-1.2 (2)
C18—C13—C14—C15	-0.8 (2)	C19—C20—C25—C24	178.80 (13)

Fig. 1




Fig. 2

