

# Amine Functionalization via Oxidative Photoredox Catalysis: Methodology Development and Complex Molecule Synthesis

Published as part of the *Accounts of Chemical Research* special issue "Synthesis, Design, and Molecular Function".


Joel W. Beatty and Corey R. J. Stephenson\*

Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109, United States

**CONSPECTUS:** While the use of visible light to drive chemical reactivity is of high importance to the development of environmentally benign chemical transformations, the concomitant use of a stoichiometric electron donor or acceptor is often required to steer the desired redox behavior of these systems. The low-cost and ubiquity of tertiary amine bases has led to their widespread use as reductive additives in photoredox catalysis. Early use of trialkylamines in this context was focused on their role as reductive excited state quenchers of the photocatalyst, which in turn provides a more highly reducing catalytic intermediate.

In this Account, we discuss some of the observations and thought processes that have led from our use of amines as reductive additives to their use as complex substrates and intermediates for natural product synthesis. Early attempts by our group to construct key carbon–carbon bonds via free-radical intermediates led to the observation that some trialkylamines readily behave as efficient hydrogen atom donors under redox-active photochemical conditions. In the wake of in-depth mechanistic studies published in the 1970s, 1980s and 1990s, this understanding has in turn allowed for a systematic approach to the design of a number of photochemical methodologies through rational tuning of the amine component. Minimization of the C–H bond dissociation energy of the amine additive was found to promote desired C–C bond formation in a number of contexts, and subsequent elucidation of the amine's redox fate has sparked a reevaluation of the amine's role from that of reagent to that of substrate.

The reactivity of tertiary amines in these photochemical systems is complex, and allows for a number of mechanistic possibilities that are not necessarily mutually exclusive. A variety of combinations of single-electron oxidation, C–H abstraction, deprotonation, and  $\beta$ -scission result in the formation of reactive intermediates such as  $\alpha$ -amino radicals and iminium ions. These processes have been explored in depth in the photochemical literature and have resulted in a firm mechanistic grasp of the behavior of amine radical cations in fundamental systems. Harnessing the synthetic potential of these transient species represents an ongoing challenge for the controlled functionalization of amine substrates, because these mechanistic possibilities may result in undesired byproduct formation or substrate decomposition. The presence of tertiary amines in numerous alkaloids, pharmaceuticals, and agrochemicals lends credence to the potential utility of this chemistry in natural product synthesis, and herein we will discuss how these transformations might be controlled for synthetic purposes.



## INTRODUCTION


Historically,  $\text{Ru}(\text{bpy})_3^{2+}$  and similar photoactive complexes have been used for water oxidation<sup>1</sup> and  $\text{CO}_2$  reduction,<sup>2</sup> with examples of the latter often employing tertiary amines as sacrificial electron donors. Recent photochemical methods using  $\text{Ru}(\text{bpy})_3\text{Cl}_2$ <sup>3</sup> reported by MacMillan,<sup>4</sup> Yoon,<sup>5</sup> and our group<sup>6</sup> utilized amines as reductive intermediates or as stoichiometric additives for quenching of the  $\text{Ru}(\text{bpy})_3^{2+*}$  photoexcited state to initiate fundamentally important organic reactions (Scheme 1).<sup>7</sup> The use of amines as reductants for photocatalysis is ideal, because trialkylamines are inexpensive, ubiquitous, and readily oxidized.<sup>8</sup> As the research field has progressed, further practical understanding of amine reactivity in these systems has allowed for a broadening scope of application in photoredox catalysis.<sup>9</sup> Herein, we will discuss the progression of results that have led to our current implementation of amines, first as additives and later as substrates, in photoredox catalysis and share some of the insights gleaned in this process.

The use of tertiary amines as reductive quenchers is not limited to net-reductive transformations, and there are many examples of redox-neutral transformations that utilize amines as stoichiometric additives.<sup>7a</sup> As evidenced by the redox cycle of  $\text{Ru}(\text{bpy})_3^{2+}$  (Scheme 1A), the ground state reducing potential of the complex ( $E_{1/2}^{\text{II/I}} = -1.33$  V vs SCE) is significantly more negative than that of the excited state ( $E_{1/2}^{\text{III/II}*} = -0.81$  V vs SCE).<sup>10</sup> As a consequence,  $\text{Ru}(\text{bpy})_3^{2+}$  mediated photoredox reactions that require a strong reduction potential sometimes incorporate stoichiometric reductive quenchers in order to access the more strongly reducing  $\text{Ru}^{\text{I}}$  species.

While trialkylamines have often been exploited for this purpose, through the years an understanding of further reactivity of the amine radical cation has evolved to the point where the amine can be used as a substrate for controlled photochemical oxidation. Using triethylamine as an example, single-electron

Received: February 6, 2015

Published: May 7, 2015

Scheme 1. Redox Cycle of Ru(bpy)<sub>3</sub><sup>2+</sup> and Further Amine Reactivity


oxidation to the radical cation results in a dramatic estimated acidification of the  $\alpha$ -amino C–H bond (Scheme 1B).<sup>8,11,12</sup> The  $\alpha$ -C–H bond of the aminium ion is also significantly weakened to an estimated  $\sim 42$  kcal/mol.<sup>13</sup> Detailed studies by Lewis,<sup>14</sup> Mariano,<sup>15</sup> and Saveant,<sup>16</sup> to name a few,<sup>17</sup> have elucidated many mechanistic aspects of amine radical cation  $\alpha$ -C–H functionalization in terms of electronics, sterics, and regiochemical outcome. This remarkable activation through the removal of a single electron allows for a number of subsequent mechanistic pathways leading to useful reactive synthetic intermediates (Scheme 1C). We have found a number of instances in which these modes of reactivity can be controlled and will delineate some of the guiding design features of these reactions below.

## NATURAL PRODUCTS AS INSPIRATION FOR REACTION DEVELOPMENT


Our initial interest in the reactivity afforded by photoredox catalysis arose from strategic bond disconnections in the context of complex molecule synthesis. At the start of our research program, the natural product actinophyllic acid **1** served as inspiration for novel bond disconnections in the context of photoredox catalysis (Scheme 2A).<sup>18</sup> Specifically, the ability to

### Scheme 2. Actinophyllic Acid as Inspiration for Radical Indole Malonation

A. Actinophyllic acid as inspiration for intra- or intermolecular indole functionalization chemistry



B. Initial attempt for intermolecular indole malonation using photoredox catalysis



directly functionalize the 2-position of indoles with a malonate equivalent was envisioned to allow for the desired bond disconnections en route to the natural product. Using Ru(bpy)<sub>3</sub>Cl<sub>2</sub> as the photocatalyst and <sup>i</sup>Pr<sub>2</sub>NEt as the reductive quencher, the initial attempt to perform the light-mediated intermolecular functionalization of *N*-methylindole with diethyl bromomalonate **2** resulted in complete hydrodehalogenation of the bromomalonate reactant (Scheme 2B), giving diethyl malonate as the exclusive product.

Further experimentation related to indole functionalization was motivated by another natural product of interest, (+)-gliocladin C, **3** (Scheme 3A). We were curious whether we could access tertiary radical intermediates such as **4** through the photochemical single-electron reduction of related bromopyrroloindoline scaffolds.<sup>19</sup> Again using <sup>i</sup>Pr<sub>2</sub>NEt as the reductive quencher, initial attempts to couple the Boc-protected bromopyrroloindoline substrate **5** with indole resulted in the isolation of the hydrodehalogenated product **6** in 75% yield (Scheme 3B). While the lack of desired intermolecular reactivity represented a setback in terms of our goals of natural product synthesis, we realized that the general efficiency that we observed for the reductive hydrodehalogenation reactivity may be leveraged into a more generalized methodology.<sup>20</sup>

Initial investigations into the reaction conditions were performed using substrate **7**, which could be dehalogenated in high yield using 10 equiv of the formate salt of Hunig's base in DMF in only 4 h (Table 1, entry 1).<sup>6</sup> It was found that substitution of diisopropylethylamine with triethylamine resulted in incomplete conversion (only 25%) after a significantly increased reaction time of 24 h (entry 2). Further experimentation revealed that Hantzsch ester **8** could be used in place of formic acid to significantly decrease the equivalents of additive used (entry 3). The scope of the reaction was found to encompass a number of activated alkyl halides, but unactivated vinyl and aryl halides were unaffected under the reaction conditions.

Mechanistically, it is expected that this reaction proceeds through reductive quenching of the Ru(bpy)<sub>3</sub><sup>2+\*</sup> excited state. As a result of this quenching process, the trialkylammonium formate radical cation is thought to perform the role of the major H atom source in the reaction. The observed difference in reactivity between the triethyl- and diisopropylethylamine additives was valuable information, which we were next able to use as a design principle to minimize hydrodehalogenation byproduct formation; we eagerly applied this knowledge toward our previous goals of indole functionalization. With the hypothesis that the rate of an intramolecular indole functionalization reaction may be sufficient to outcompete intermolecular C–H abstraction from a poor H-atom donor, we began to investigate this chemistry using tethered malonates such as **9** (Scheme 4).<sup>21</sup> As supported by previous observations, the use of Hunig's base as a reductive quencher in an intramolecular radical addition to indole resulted in a significant amount of hydrodehalogenation byproduct **10** (Scheme 4A). Triethylamine was again found to be less promoting of hydrodehalogenation (*vide supra*) and as a result was selected for the more generalized conditions (Scheme 4B).

## Scheme 3. Gliocladin C as Inspiration for Methodology Development




Table 1. Optimization of the Reductive Dehalogenation Reaction

entry	additives	time (h)	yield (%)
1	$t\text{Pr}_2\text{NEt}$ (10 equiv), $\text{HCOOH}$ (10 equiv)	4	90
2	$\text{Et}_3\text{N}$ (10 equiv), $\text{HCOOH}$ (10 equiv)	24	20
3	$t\text{Pr}_2\text{NEt}$ (2 equiv), <b>8</b> (1.1 equiv)	4	95

While hydrodehalogenation was minimized by using triethylamine, a number of additional insights were uncovered during the investigation of this reaction. Most notably, when substrate **11** was subjected to the standard conditions, a mixture of three products was produced, with acetaldehyde incorporated product **12** present in 20% yield (Scheme 5A).<sup>22</sup> The genesis of this material can be rationalized through iminium formation from the triethylamminium radical cation, either through direct C–H abstraction or a sequential deprotonation–oxidation process (Scheme 5B). Tautomerization of the iminium ion provides an enamine equivalent, which is electronically paired with the electron-poor malonyl radical to produce the undesired aldehyde


product **12** after radical addition, oxidation, and hydrolysis of the product iminium ion. The isolation of this material was further evidence of our mechanistic hypotheses involving  $\alpha$ -amino C–H chemistry and provided further confidence in our understanding of the observed reactivity going forward.

## TERTIARY AMINES AS SUBSTRATES



With this insight into the fate of the amine component, the possibility of applying this chemistry to the  $\alpha$ -functionalization of tertiary amine-containing substrates became more appealing to us.<sup>23</sup> The issue of regioselectivity in  $\alpha$ -amino functionalization was vital, because factors determining which C–H bond would react were expected to be governed chiefly by substrate characteristics.<sup>14</sup> With this in mind, efforts to functionalize *N*-aryl tetrahydroisoquinolines were undertaken.<sup>24</sup> In our initial design, we anticipated that we could leverage our prior observations in Scheme 5 to selectively form iminium ions; we expected a dual role for the bromomalonate, where it first would behave as the terminal oxidant before subsequent enlistment as a nucleophile (Scheme 6A).

An early experiment along these lines utilized *N*-phenyl tetrahydroisoquinoline **13** as the substrate (Scheme 6B). Using the photocatalyst  $\text{Ru}(\text{bpy})_3\text{Cl}_2$  and diethyl bromomalonate **2** in DMF, the reaction was run with an aim to produce malonate functionalization at the benzylic position of the substrate.

## Scheme 4. Intramolecular Radical Malonation of Heterocycles



## Scheme 5. Understanding the Fate of the Amine Reductive Quencher

Scheme 6. Oxidative Functionalization of *N*-Aryl Tetrahydroisoquinolines


Curiously, the starting material was consumed in an overnight reaction, but none of the desired product was observed. Instead, after careful analysis of the reaction, **15** was isolated as the sole product, resulting from methanol trapping of iminium **14** during column chromatography.

Interestingly, the reaction was found to proceed with 100% conversion in methanol without the use of diethylbromomalonate; however, later experiments revealed that the reaction slowed significantly in the absence of oxygen, pointing toward oxygen's role as the terminal oxidant.<sup>25</sup> With nitromethane as the solvent, high yields of the aza-Henry product **17** could be obtained (Scheme 6C). We also experimented with the use of the cyclometalated heteroleptic iridium-based catalyst  $\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$  **16**, which we found to accelerate the aerobic aza-Henry reaction significantly and provide the product in higher yield. A slow background reaction was observed, providing 83% conversion of the starting material after 5 days when no catalyst was present. An unexpected challenge associated with this chemistry was encountered upon evaluation of the substrate scope. A wide range of *N*-aryl tetrahydroisoqui-

nolines provided >90% yield in 18 h or less; however, *N*-phenyl pyrrolidine provided 27% yield of the aza-Henry product **18** in only 40% conversion after a 72 h reaction time.

The aerobic oxidation reactions were somewhat slow (10–18 h) compared with reaction rates with terminal organic oxidants such as diethylbromomalonate (2 h). Additionally, byproducts were often isolated from the reactions, including the endocyclic amide **20** and dimer **21**, both presumably arising from an  $\alpha$ -amino radical intermediate (Scheme 7A).<sup>27</sup> We postulated that anaerobic oxidation of the substrate using a suitable oxidant such as bromochloroform may result in direct iminium formation through C–H abstraction from **13** by the resulting trichloromethyl radical **22** (Scheme 7B).


While the anaerobic use of  $\text{BrCCl}_3$  was empirically effective in eliminating the observed byproducts, the profound increase in observed reactivity is likely attributable to an efficient chain propagation mechanism (Scheme 7B). Propagation of the free-radical intermediates through sequential atom-transfer reactions may explain how the reaction is able to proceed with such efficiency. Direct C–H abstraction from the closed shell

Scheme 7. Mechanistic Possibilities for Substrate Oxidation with BrCCl<sub>3</sub>


substrate **13** would form the  $\alpha$ -amino radical **24**, which then can be further oxidized by another equivalent of BrCCl<sub>3</sub> to form the iminium ion **14** while reforming an additional equivalent of trichloromethyl radical. The radical–radical disproportionation between **22** and **23** is a statistically disfavored termination step, since presumably the individual concentrations of the two free-radical intermediates are low. The propagation mechanism is also statistically favored, because the BrCCl<sub>3</sub> is used in stoichiometric excess (3 equiv). Furthermore, the bond dissociation energies (BDE) of this propagation are estimated to align with a thermodynamically favored process, since the experimental C–H BDE of chloroform is 96 kcal/mol<sup>28</sup> while the BDE of a methylene C–H bond of *N,N*-dibenzylaniline is measured at 85 kcal/mol (Scheme 7C).<sup>29</sup> More recent work from our group has provided evidence for a propagation mechanism in a light-mediated atom transfer reaction.<sup>30</sup> One of the benefits of these types of mechanisms is that in the event of a chain termination, reactive intermediates can be continuously produced by the catalyst.

While the use of BrCCl<sub>3</sub> decreased reaction times to 3 h, we have been able to shorten reaction time even further by applying the optimized oxidative conditions in a flow reactor.<sup>31</sup> For

example, the oxidation of *N*-phenyl tetrahydroisoquinoline **13** to the iminium ion **14** proceeds with a residence time<sup>32</sup> of only 0.5 min (Scheme 8), which corresponds to material throughput of 5.75 mmol h<sup>-1</sup>, a roughly 70-fold increase in comparison to batch reaction material throughput. The flow reaction can be eluted into a stirred solution of nucleophile for facile structural diversification, with cyanation, allylation, and alkynylations all proceeding in good yields.


Scheme 8. Flow Functionalization of *N*-Phenyl Tetrahydroisoquinoline

## Scheme 9. Asymmetric Nucleophilic Addition Using Anion Binding Catalysis




## Scheme 10. Selected Reactivity Compatible with Photochemical Amine Oxidation


## A. Select examples of diverse nucleophile trapping (ref. 34)



## B. Xiao's oxidative [3+2] cycloaddition



## C. DiRocco and Rovis' asymmetric acylation of tetrahydroisoquinolines



Further work in our lab to elaborate the chemistry of photochemical amine oxidation has involved the asymmetric alkylation of iminium ions of type **14** through the use of chiral anion-binding catalysis in collaboration with Jacobsen and co-workers (Scheme 9).<sup>33</sup> Because the reductive dehalogenation of  $\text{BrCCl}_3$  or  $\text{CCl}_4$  results in the formation of halide counterions associated with the oxidized substrate, it was postulated that the use of thiourea catalysis would enable stereoselective nucleophilic addition.

Initial reactions focused on the use of silyl ketene acetal **26** for nucleophilic addition to the iminium intermediate. Unfortunately, the photocatalyst  $\text{Ru}(\text{bpy})_3\text{Cl}_2$  was found to be entirely insoluble in methyl *tert*-butyl ether (MTBE) as well as other nonpolar solvents known for providing high enantioselectivities in concert with thiourea catalysis. Unsurprisingly, high yields of racemic products were isolated from reactions performed in DMF,  $\text{CH}_2\text{Cl}_2$ , and MeCN. As a solution to the orthogonal polarity requirements for each mode of catalysis a solvent switch was required; The MeCN was removed upon complete photochemical oxidation of the substrate, and the reaction was reconstituted in MTBE for the nucleophilic addition step.

Since our initial report,<sup>23</sup> oxidative amine photoredox catalysis has become more widely adopted, and many additional examples


of nucleophilic additions to tetrahydroisoquinolines have been published (Scheme 10A).<sup>34</sup> The versatility of these systems is impressive, and there have been many creative additions to the literature in this context. Xiao and co-workers have demonstrated the compatibility of the photochemical tetrahydroisoquinoline oxidation with dipolar [3 + 2] cycloaddition chemistry (Scheme 10B), performing a final oxidation with *N*-bromosuccinimide (NBS) to provide penta-substituted pyrrole products.<sup>35</sup> Additionally, the oxidative conditions are fully compatible with *N*-heterocyclic carbene cocatalysis, which DiRocco and Rovis have demonstrated elegantly (Scheme 10C).<sup>36</sup> These extensions of the amine oxidation highlight the versatility and robust nature of the photochemical tetrahydroisoquinoline oxidation.

### ■ $\alpha$ -AMINO C–H AND C–C FUNCTIONALIZATION OF TERTIARY ALIPHATIC AMINES


Experimentation with *N*-phenyltetrahydroisoquinoline **13** revealed efficient intermolecular Mannich reactivity with indole to provide **29** in 83% yield (Scheme 11A). These developments toward an efficient  $\alpha$ -amino functionalization reaction represented additional opportunity for us in alkaloid synthesis, particularly in the context of an oxidative Mannich reaction en

### Scheme 11. Alkaloids as Further Inspiration for Reaction Development

#### A. Intermolecular oxidative Mannich reaction with a tetrahydroisoquinoline substrate



#### B. Oxidative Mannich reaction as a key bond disconnection



route to actinophyllic acid **1** (Scheme 11B). With a working knowledge of the  $\alpha$ -amino functionalization of tetrahydroisoquinolines and how these processes can be accelerated with flow chemistry, we began developing ways to apply these concepts in a more complex setting. A key challenge was the lack of general substrate scope for the oxidative iminium forming reaction, because early experimentation had shown that even simple dialkyl anilines such as *N*-phenylpyrrolidine were recalcitrant to product formation (*vide supra*).

The commercially available natural product (+)-catharanthine **30** was selected as a starting material for our initial investigations in this area (Scheme 12). A series of reports on the total synthesis of (+)-vinblastine and related natural products by Boger and co-workers detailed the reactivity of **30** upon oxidation by  $\text{FeCl}_3$ .<sup>37</sup> Bolstered by the possibility of promoting carbon–carbon bond fragmentation through reductive quenching of a photocatalyst excited state,<sup>38</sup> we began to investigate the reactivity of catharanthine under photocatalytic conditions.<sup>39</sup>

It was found that light exposure of a solution of catharanthine,  $\text{Ir}(\text{dF}(\text{CF}_3)\text{ppy})_2(\text{dtbbpy})\text{PF}_6$  (2.5 mol %), and trimethylsilyl-cyanide (TMSCN, 2.0 equiv) in methanol provided the cyanated ring-opened product **31** in 93% yield after 3 h.<sup>40</sup> Application of these exact conditions in a flow reactor resulted in the scalable application of this procedure to 2 g of material in 88% yield. Reliable access to significant amounts of this complex material allowed us to investigate further photochemical reactivity in this context.

In an effort to synthesize the natural product (–)-pseudovincadifformine, **33** (Table 2), from the fragmented and cyanated catharanthine, we subjected the material to a short synthetic route involving hydrogenation of the C15–C20 double bond (catharanthine numbering) followed by quenching of the reaction with sodium borohydride to remove the  $\alpha$ -aminonitrile functionality (Scheme 12). The advanced intermediate **32** was

obtained through this reduction procedure and served as an ideal substrate for testing further applications of oxidative photoredox catalysis on a complex tertiary aliphatic amine substrate. Specifically, to synthesize **33** from **32**, we recognized the need for selective C–H functionalization on C3 in preference to the two alternative  $\alpha$ -amino methylenes on C5 and C21. While attempts at aerobic photochemical oxidation of **32** resulted in a complex mixture of decomposition products (Table 2, entry 1), we were excited to find that the use of  $\text{BrCCl}_3$  resulted in the formation of the natural product in 22% yield (entry 2). Further evaluation of oxidants revealed that diethyl bromomalonate and diethyl 2-bromo-2-methylmalonate resulted in successively improved yields of the desired product (entries 3–4). No products of C21 oxidation were observed in the reaction mixtures. Subjection of the reaction to a flow protocol at 50 °C with a 5 min residence time resulted in the highest yields of the product, yielding **32** in 58% yield and an 8:1 diastereomeric ratio in favor of the desired ethyl epimer (entry 5).

There are a number of possibilities that may account for the origin of the observed regioselectivity in this oxidative cyclization process. Iminium formation is thought to be limited to C3 and C21, because geometric constraints prevent favorable overlap of the nitrogen lone pair with the C5–H bond. Additionally, while on first approximation C21 may appear to be less sterically encumbered, a three-dimensional analysis of structure **32** reveals that C3 may be equally if not more accessible to either intermolecular deprotonation or H atom abstraction. The possibilities of iminium or amino-radical isomerization cannot be ruled out, because transannular cyclization could be expected to serve as a thermodynamic trap for such equilibria. A further alternative is that as the yield trends upward with the steric bulk of the oxidant, there may be a matching effect in which the more bulky oxidant provides higher regioselectivity in a possible C–H abstraction step. It is worthy of note that this C–H oxidation exhibits rare efficiency for a photochemical aliphatic amine oxidation. Preliminary experimentation in our group has suggested that the transannular nature of the cyclization is responsible for reaction success; similar cyclization attempts on structures without the ethylene tether between the indole and the nitrogen have resulted in decomposition of the starting material, possibly through enamine intermediates.

In an attempt to accomplish a more generally applicable  $\alpha$ -functionalization of tertiary aliphatic amines, we have further evaluated this chemistry from a pharmaceutical synthesis standpoint.<sup>41</sup> A collaboration with Lilly Research Laboratories brought our attention to the selective JAK2 inhibitor LY2784544, **35** (Scheme 13A). The industrial synthesis of **35**, which was used to produce over one metric ton of the advanced pharmaceutical intermediate **36**, relied upon a vanadium-mediated addition of *N*-methylmorpholine *N*-oxide to the core imidazopyridazine scaffold.<sup>42</sup> While the exact mechanistic course of this reaction has yet to be elucidated, it may proceed through an exocyclic  $\alpha$ -amino radical; consequently, other methods for

### Scheme 12. Photochemical Fragmentation of Catharanthine






Table 2. Photochemical Oxidative Cyclization To Form (–)-Pseudovincadifformine




entry	oxidant	yield (%)
1	air	0
2	BrCCl <sub>3</sub> (3 equiv)	22
3	2 (3 equiv)	34
4	34 (3 equiv)	39
5 <sup>a</sup>	34 (3 equiv)	58

<sup>a</sup>Flow reactor,  $t_R = 5$  min.

## Scheme 13. Aliphatic Amine Functionalization for API Synthesis



## Scheme 14. Photochemical Amidoalkylation of Heterocycles and Arenes



the formation of this radical, including photoredox catalysis, were examined.

Initial experiments revealed that in addition to the desired  $\alpha$ -amino functionalization reaction, several side products were observed in the reaction mixture, including products of double addition, reductive dechlorination, methylation, and a solvent incorporation adduct. Following extensive optimization, by-product formation was minimized and the product **38** was

produced in 56% isolated yield (10:1 *exo/endo*, Scheme 13B). The observed reactivity proved challenging to control, resulting in a reactant scope that was broad for the amine component but limited for the heterocyclic coupling partners.

Of note in the discussion of possible mechanistic pathways for this transformation is the observation of amidoalkylation products arising from solvent reactivity. When the reaction was performed in *N,N'*-dimethylpropylene urea (DMPU) in the



absence of *N*-methyl-morpholine, a mixture of *endo* and *exo* adducts were observed in a 5:1 ratio and combined 53% isolated yield (Scheme 14A). Previous research efforts in our group have revealed that  $\alpha$ -amido C–H functionalization in this manner can be accomplished through an initial C–H abstraction from the amide solvent, followed by oxidation to the *N*-acyliminium ion, which is a potent Friedel–Crafts electrophile (Scheme 14B).<sup>43</sup> While the electronic nature of substrate **37** strongly suggests that radical addition is the operative mechanism of heterocycle addition, the analogous reactivity of these two systems suggests that amidoalkylation of these electron-poor substrates is precipitated by direct C–H abstraction.

## CONCLUSION

This Account has summarized some of our contributions in relation to amine reactivity in light-mediated redox catalysis. Photoredox catalysis has allowed for an environmentally benign approach to the study of amine reactivity; however, significant questions remain to be addressed. The use of aliphatic tertiary amines as substrates is a particularly underexplored area, because previous synthetic work has focused mainly on the use of aniline and tetrahydroisoquinoline substrates. Due to the ubiquity of amine functionality in natural products and commodity chemicals, the ability to controllably oxidize these substrates to access radical and electrophilic functionality is an important goal. There is significant opportunity for this type of C–H oxidation, particularly in complex molecule synthesis, because the efficient formation of  $\alpha$ -amino C–C bonds would provide increased retrosynthetic flexibility. Further study in this regard would be beneficial, because the observed differences in reaction efficiency between aryl and aliphatic amines remain to be elucidated experimentally.

## AUTHOR INFORMATION

### Corresponding Author

\*E-mail: crjsteph@umich.edu

### Notes

The authors declare no competing financial interest.

### Biographies

**Joel Beatty** received a B.S. in Chemistry from Northeastern University in 2011 and an M.A. in Chemistry from Boston University in 2013. He is currently a Ph.D. candidate at the University of Michigan under the direction of Corey Stephenson.

**Corey Stephenson** earned his Ph.D. from the University of Pittsburgh under Professor Peter Wipf before pursuing his postdoctoral studies at ETH Zürich with Professor Erick M. Carreira. He began his independent career at Boston University in 2007 and moved to the University of Michigan in 2013. His research interests are broadly focused on complex molecule synthesis, biomass degradation, and catalysis.

## REFERENCES

- (1) Kärkäs, M. D.; Verho, O.; Johnston, E. V.; Åkermark, B. Artificial Photosynthesis: Molecular Systems for Catalytic Water Oxidation. *Chem. Rev.* **2014**, *114*, 11863–12001.
- (2) Morris, A. J.; Meyer, G. J.; Fujita, E. Molecular Approaches to the Photocatalytic Reduction of Carbon Dioxide for Solar Fuels. *Acc. Chem. Res.* **2009**, *42*, 1983–1994.
- (3) Juris, A.; Balzani, V.; Barigelletti, F.; Campagna, S.; Belser, P.; von Zelewski, A. Ru(II) Polypyridine Complexes: Photophysics, Photo-

chemistry, Electrochemistry, and Chemiluminescence. *Coord. Chem. Rev.* **1988**, *84*, 85–277.

- (4) Nicewicz, D. A.; MacMillan, D. W. C. Merging Photoredox Catalysis with Organocatalysis: The Direct Asymmetric Alkylation of Aldehydes. *Science* **2008**, *322*, 77–80.

- (5) Ischay, M. A.; Anzovino, M. E.; Du, J.; Yoon, T. P. Efficient Visible Light Photocatalysis of [2 + 2] Enone Cycloadditions. *J. Am. Chem. Soc.* **2008**, *130*, 12886–12887.

- (6) Narayanam, J. M. R.; Tucker, J. W.; Stephenson, C. R. J. Electron-Transfer Photoredox Catalysis: Development of a Tin-Free Reductive Dehalogenation Reaction. *J. Am. Chem. Soc.* **2009**, *131*, 8756–8757.

- (7) For reviews, see: (a) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 5322–5363. (b) Wallentin, C. J.; Nguyen, J. D.; Stephenson, C. R. J. Radical Carbon–Carbon Bond Formations Enabled by Visible Light Active Photocatalysts. *Chimia* **2012**, *66*, 394–398. (c) Xuan, J.; Xiao, W. J. Visible-Light Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2012**, *51*, 6828–6838. (d) Shi, L.; Xia, W. Photoredox Functionalization of C–H Bonds Adjacent to a Nitrogen Atom. *Chem. Soc. Rev.* **2012**, *41*, 7687–7697.

- (8) Lindsay Smith, J. R.; Masheder, D. Amine Oxidation. Part IX. The Electrochemical Oxidation of Some Tertiary Amines: The Effect of Structure on Reactivity. *J. Chem. Soc., Perkin Trans. 2* **1976**, 47–51.

- (9) (a) Hu, J.; Wang, J.; Nguyen, T. H.; Zheng, N. The Chemistry of Amine Radical Cations Produced by Visible Light Photoredox Catalysis. *Beilstein J. Org. Chem.* **2013**, *9*, 1977–2001. (b) DeLaive, P. J.; Sullivan, B. P.; Meyer, T. J.; Whitten, D. G. Applications of Light-Induced Electron-Transfer Reactions. Coupling of Hydrogen Generation with Photoreduction of Ruthenium (II) Complexes by Triethylamine. *J. Am. Chem. Soc.* **1979**, *101*, 4007–4008.

- (10) Bock, C. R.; Connor, J. A.; Gutierrez, A. R.; Meyer, T. J.; Whitten, D. G.; Sullivan, B. P.; Nagle, J. K. Estimation of Excited-State Redox Potentials by Electron-Transfer Quenching. Application of Electron-Transfer Theory to Excited-State Redox Processes. *J. Am. Chem. Soc.* **1979**, *101*, 4815–4824.

- (11) For the oxidation potential of Et<sub>3</sub>N vs AgNO<sub>3</sub> in MeCN, converted to SCE in Scheme 1, see: Newman, J. D. S.; Blanchard, G. J. Formation of Gold Nanoparticles Using Amine Reducing Agents. *Langmuir* **2006**, *22*, 5882–5887.

- (12) Estimation of the pK<sub>s</sub> of the  $\alpha$ -C–H bond of the triethylamine radical cation was performed according to eq 19 found in Nicholas, M. de P.; Arnold, D. R. Thermochemical Parameters for Organic Radicals and Radical Ions. Part 1. The Estimation of the pK<sub>s</sub> of Radical Cations Based on Thermochemical Calculations. *Can. J. Chem.* **1982**, *60*, 2165–2179.

- (13) Estimation of the  $\alpha$ -C–H BDE of the triethylamine radical cation was performed with BDE values from ref 29 according to Wayner, D. D. M.; Dannenberg, J. J.; Griller, D. Oxidation Potentials of  $\alpha$ -Amino Radicals: Bond Dissociation Energies for Related Radical Cations. *Chem. Phys. Lett.* **1986**, *131*, 189–191.

- (14) (a) Lewis, F. D.; Ho, T.-I. On the Selectivity of Tertiary Amine Oxidations. *J. Am. Chem. Soc.* **1980**, *102*, 1751–1752. (b) Lewis, F. D.; Ho, T.-I.; Simpson, J. T. Photochemical Addition of Tertiary Amines to Stillbene. Stereoelectronic Control of Tertiary Amine Oxidation. *J. Org. Chem.* **1981**, *46*, 1077–1082.

- (15) Zhang, X.; Yeh, S.-R.; Hong, S.; Freccero, M.; Albini, A.; Falvey, D. E.; Mariano, P. S. Dynamics of  $\alpha$ -CH Deprotonation and  $\alpha$ -Desilylation Reactions of Tertiary Amine Cation Radicals. *J. Am. Chem. Soc.* **1994**, *116*, 4211–4220.

- (16) Anne, A.; Hapoit, P.; Moiroux, J.; Neta, P.; Saveant, J.-M. Dynamics of Proton Transfer from Cation Radicals. Kinetic and Thermodynamic Acidities of Cation Radicals of NADH Analogues. *J. Am. Chem. Soc.* **1992**, *114*, 4694–4701.

- (17) (a) Yoon, U. C.; Su, Z.; Mariano, P. S. The Dynamics and Photochemical Consequences of Aminium Radical Reactions. In *CRC Handbook of Organic Photochemistry and Photobiology*, 2nd ed.; Horspool, W., Lenci, F., Eds.; CRC Press: Boca Raton, FL, 2004; Vol. 2, pp 101-1–101-20. (b) Schmittel, M.; Burghart, A. Understanding

Reactivity Patterns of Radical Cations. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2550–2589.

(18) Carroll, A. R.; Hyde, E.; Smith, J.; Quinn, R. J.; Guymer, G.; Forster, P. I. Actinophyllic Acid, a Potent Indole Alkaloid Inhibitor of the Coupled Enzyme Assay Carboxypeptidase U/Hippurase from the Leaves of *Alstonia actinophylla* (Apocynaceae). *J. Org. Chem.* **2005**, *70*, 1096–1099.

(19) A long-term goal of this chemistry was to perform reductive dimerization of bromopyrroloindoline substrates. This is quite challenging to accomplish with photoredox catalysis, because the reactive radical species are likely short-lived and in low concentration. For select radical reactions of this type, see: (a) Bruncko, M.; Crich, D.; Samy, R. Chemistry of Cyclic Tautomers of Tryptophan: Formation of a Quaternary Center at C3a and Total Synthesis of the Marine Alkaloid (+)-*ent*-Debromoflustramine B. *J. Org. Chem.* **1994**, *59*, 5543–5549. (b) Depew, K. M.; Marsden, S. P.; Zatorska, D.; Zatorski, A.; Bornmann, W. G.; Danishefsky, S. J. Total Synthesis of 5-*N*-Acetylardeemin and Amauromine: Practical Routes to Potential MDR Reversal Agents. *J. Am. Chem. Soc.* **1999**, *121*, 11953–11963. (c) Kim, J.; Ashenhurst, J. A.; Movassaghi, M. Total Synthesis of (+)-11,11'-Dideoxyverticillin A. *Science* **2009**, *324*, 238–241.

(20) (a) Van Bergen, T. J.; Hedstrand, D. M.; Kruizinga, W. H.; Kellogg, R. M. Hydride Transfer from 1,4-Dihydropyridines to sp<sup>3</sup>-Hybridized Carbon in Sulfonium Salts and Activated Halides. Studies with NAD(P)H models. *J. Org. Chem.* **1979**, *44*, 4953–4962. (b) Hironaka, K.; Fukuzumi, S.; Tanaka, T. Tris(bipyridyl)Ruthenium(II)-Photosensitized Reaction of 1-Benzyl-1,4-Dihydronicotinamide with Benzyl bromide. *J. Chem. Soc., Perkin Trans. 2* **1984**, 1705–1709. (c) Fukuzumi, S.; Mochizuki, S.; Tanaka, T. Photocatalytic Reduction of Phenacyl Halides by 9,10-Dihydro-10-Methylacridine. Control Between the Reductive and Oxidative Quenching Pathways of Tris(bipyridine) Ruthenium Complex Utilizing an Acid Catalysis. *J. Phys. Chem.* **1990**, *94*, 722–726.

(21) (a) Tucker, J. W.; Narayanam, J. M. R.; Krabbe, S. W.; Stephenson, C. R. J. Electron Transfer Photoredox Catalysis: Intramolecular Radical Addition to Indoles and Pyrroles. *Org. Lett.* **2010**, *12*, 368–371. (b) Magolan, J.; Kerr, M. A. Expanding the Scope of Mn(OAc)<sub>3</sub>-Mediated Cyclizations: Synthesis of the Tetracyclic Core of Tronocarpine. *Org. Lett.* **2006**, *8*, 4561–4564.

(22) Furst, L.; Matsuura, B. S.; Narayanam, J. M. R.; Tucker, J. W.; Stephenson, C. R. J. Visible Light-Mediated Intermolecular C–H Functionalization of Electron-Rich Heterocycles with Malonates. *Org. Lett.* **2010**, *12*, 3104–3107.

(23) Condie, A. G.; González-Gómez, J. C.; Stephenson, C. R. J. Visible-Light Photoredox Catalysis: Aza-Henry Reactions via C–H Functionalization. *J. Am. Chem. Soc.* **2010**, *132*, 1464–1465.

(24) (a) Murahashi, S. I.; Zhang, D. Ruthenium Catalyzed Biomimetic Oxidation in Organic Synthesis Inspired by Cytochrome P-450. *Chem. Soc. Rev.* **2008**, *37*, 1490–1501. (b) Li, C. J. Cross-Dehydrogenative Coupling (CDC): Exploring C–C Bond Formations Beyond Functional Group Transformations. *Acc. Chem. Res.* **2009**, *42*, 335–344.

(25) Zhu, S.; Das, A.; Bui, L.; Zhou, H.; Curran, D. P.; Rueping, M. Oxygen Switch in Visible-Light Photoredox Catalysis: Radical Additions and Cyclizations and Unexpected C–C-Bond Cleavage Reactions. *J. Am. Chem. Soc.* **2013**, *135*, 1823–1829.

(26) Slinker, J. D.; Gorodetsky, A. A.; Lowry, M. S.; Wang, J.; Parker, S.; Rohl, R.; Bernhard, S.; Malliaras, G. G. Efficient Yellow Electroluminescence from a Single Layer of a Cyclometallated Iridium Complex. *J. Am. Chem. Soc.* **2004**, *126*, 2763–2767.

(27) Byproducts of these type have since been isolated and characterized from similar reactions: (a) Espelt, L. R.; Wiensch, E. M.; Yoon, T. P. Brønsted Acid Cocatalysts in Photocatalytic Radical Addition of  $\alpha$ -Amino C–H Bonds across Michael Acceptors. *J. Org. Chem.* **2013**, *78*, 4107–4114. (b) Mitkina, T.; Stanglmair, C.; Setzer, W.; Gruber, M.; Kisch, H.; König, B. Visible Light Mediated Homo- and Heterocoupling of Benzyl Alcohols and Benzyl Amines on Polycrystalline Cadmium Sulfide. *Org. Biomol. Chem.* **2012**, *10*, 3556–3561.

(28) McMillen, D. F.; Golden, D. M. Hydrocarbon Bond Dissociation Energies. *Annu. Rev. Phys. Chem.* **1982**, *33*, 493–532.

(29) Dombrowski, G. W.; Dinnocenzo, J. P.; Farid, S.; Goodman, J. L.; Gould, I. R.  $\alpha$ -C–H Bond Dissociation Energies of Some Tertiary Amines. *J. Org. Chem.* **1999**, *64*, 427–431.

(30) Wallentin, C. J.; Nguyen, J. D.; Finkbeiner, P.; Stephenson, C. R. J. Visible Light-Mediated Atom Transfer Radical Addition via Oxidative and Reductive Quenching of Photocatalysts. *J. Am. Chem. Soc.* **2012**, *134*, 8875–8884.

(31) Tucker, J. W.; Zhang, Y.; Jamison, T. F.; Stephenson, C. R. J. Visible-Light Photoredox Catalysis in Flow. *Angew. Chem., Int. Ed.* **2012**, *51*, 4144–4147.

(32) “Residence time” refers to the amount of time a single portion of the reaction spends within the flow reactor. In this case, it refers to the amount of time a portion of the reaction is exposed to the light source.

(33) Bergonzini, G.; Schindler, C. S.; Wallentin, C. J.; Jacobsen, E. N.; Stephenson, C. R. J. Photoredox Activation and Anion Binding Catalysis in the Dual Catalytic Enantioselective Synthesis of  $\beta$ -Amino Esters. *Chem. Sci.* **2014**, *5*, 112–116.

(34) Select examples of nucleophilic additions to photochemically activated THIQs: (a) Pan, Y.; Wang, S.; Kee, C. W.; Dubuisson, E.; Yang, Y.; Loh, K. P.; Tan, C. H. Graphene Oxide and Rose Bengal: Oxidative C–H Functionalization of Tertiary Amines Using Visible Light. *Green Chem.* **2011**, *13*, 3341–3344. (b) Rueping, M.; Vila, C.; Koenigs, R. M.; Poschary, K.; Fabry, D. C. Dual Catalysis: Combining Photoredox and Lewis Base Catalysis for Direct Mannich Reactions. *Chem. Commun.* **2011**, *47*, 2360–2362. (c) Rueping, M.; Zhu, S.; Koenigs, R. M. Photoredox Catalyzed C–P Bond Forming Reactions—Visible Light Mediated Oxidative Phosphonylations of Amines. *Chem. Commun.* **2011**, *47*, 8679–8681. (d) Zhao, G.; Yang, C.; Guo, L.; Sun, H.; Chen, C.; Xia, W. Visible Light-Induced Oxidative Coupling Reaction: Easy Access to Mannich-Type Products. *Chem. Commun.* **2012**, *48*, 2337–2339. (e) Fu, W.; Guo, W.; Zou, G.; Xu, C. Selective Trifluoromethylation and Alkynylation of Tetrahydroisoquinolines Using Visible Light Irradiation by Rose Bengal. *J. Fluorine Chem.* **2012**, *140*, 88–94. (f) Xuan, J.; Feng, Z. J.; Duan, S. W.; Xiao, W. J. Room Temperature Synthesis of Isoquino[2,1-*a*][3,1]oxazine and Isoquino[2,1-*a*]pyrimidine Derivatives via Visible Light Photoredox Catalysis. *RSC Adv.* **2012**, *2*, 4065–4068. (g) Mathis, C. L.; Gist, B. M.; Frederickson, C. K.; Midkiff, K. M.; Marvin, C. C. Visible Light Photooxidative Cyclization of Amine Alcohols to 1,3-Oxazines. *Tetrahedron Lett.* **2013**, *54*, 2101–2104. (h) Feng, Z. J.; Xuan, J.; Xia, X. D.; Ding, W.; Guo, W.; Chen, J. R.; Zou, Y. Q.; Lu, L. Q.; Xiao, W. J. Direct sp<sup>3</sup> C–H Acroleination of *N*-aryl-Tetrahydroisoquinolines by Merging Photoredox Catalysis with Nucleophilic Catalysis. *Org. Biomol. Chem.* **2014**, *12*, 2037–2040. (i) Xiao, T.; Li, L.; Lin, G.; Mao, Z.; Zhou, L. Metal-Free Visible-Light Induced Cross-Dehydrogenative Coupling of Tertiary Amines with Diazo Compounds. *Org. Lett.* **2014**, *16*, 4232–4235.

(35) Zou, Y. Q.; Lu, L. Q.; Fu, L.; Chang, N. J.; Rong, J.; Chen, J. R.; Xiao, W. J. Visible-Light-Induced Oxidation/[3 + 2] Cycloaddition/Oxidative Aromatization Sequence: A Photocatalytic Strategy to Construct Pyrrolo[2,1-*a*]isoquinolines. *Angew. Chem., Int. Ed.* **2011**, *50*, 7171–7175.

(36) DiRocco, D. A.; Rovis, T. Catalytic Asymmetric  $\alpha$ -Acylation of Tertiary Amines Mediated by a Dual Catalysis Mode: *N*-Heterocyclic Carbene and Photoredox Catalysis. *J. Am. Chem. Soc.* **2012**, *134*, 8094–8097.

(37) (a) Ishikawa, H.; Colby, D. A.; Boger, D. L. Direct Coupling of Catharanthine and Vindoline to Provide Vinblastine: Total Synthesis of (+)- and *ent*(-)-Vinblastine. *J. Am. Chem. Soc.* **2008**, *130*, 420–421. (b) Ishikawa, H.; Colby, D. A.; Seto, S.; Va, P.; Tam, A.; Kakei, H.; Rayl, T. J.; Hwang, I.; Boger, D. A. Total Synthesis of Vinblastine, Vincristine, Related Natural Products, and Key Structural Analogues. *J. Am. Chem. Soc.* **2009**, *131*, 4904–4916. (c) Gotoh, H.; Sears, J. E.; Eschenmoser, A.; Boger, D. L. New Insights into the Mechanism and an Expanded Scope of the Fe(III)-Mediated Vinblastine Coupling Reaction. *J. Am. Chem. Soc.* **2012**, *134*, 13240–13243.

(38) For select examples of C–C bond scission with photoredox catalysis, see (a) Maity, S.; Zhu, M.; Shinabery, R. S.; Zheng, N. Intermolecular [3 + 2] Cycloaddition of Cyclopropylamines with

Olefins by Visible-Light Photocatalysis. *Angew. Chem., Int. Ed.* **2012**, *51*, 222–226. (b) Cai, S.; Zhao, X.; Wang, X.; Liu, Q.; Li, Z.; Wang, D. Z. Visible-Light-Promoted C–C Bond Cleavage: Photocatalytic Generation of Iminium Ions and Amino Radicals. *Angew. Chem., Int. Ed.* **2012**, *51*, 8050–8053.

(39) (a) Sundberg, R. J.; Desos, P.; Gadamasetti, K. G.; Sabat, M. Photoactive C16-C21 Fragmentation of Catharanthine. *Tetrahedron Lett.* **1991**, *32*, 3035–3038. (b) Cocquet, G.; Rool, P.; Ferroud, C. A Catalytic Versus Stoichiometric Electron Transfer Promoted Selective C16-C21 Bond Cleavage of Catharanthine. *Tetrahedron Lett.* **2001**, *42*, 839–841.

(40) Beatty, J. W.; Stephenson, C. R. J. Synthesis of (–)-Pseudotabersonine, (–)-Pseudovincadifformine, and (+)-Coronaridine Enabled by Photoredox Catalysis in Flow. *J. Am. Chem. Soc.* **2014**, *136*, 10270–10273.

(41) Douglas, J. J.; Cole, K. P.; Stephenson, C. R. J. Photoredox Catalysis in a Complex Pharmaceutical Setting: Toward the Preparation of JAK2 Inhibitor LY2784544. *J. Org. Chem.* **2014**, *79*, 11631–11643.

(42) Mitchell, D.; Cole, K. P.; Pollock, P. M.; Coppert, D. M.; Burkholder, T. P.; Clayton, J. R. Development and a Practical Synthesis of the JAK2 inhibitor LY2784544. *Org. Process Res. Dev.* **2012**, *16*, 70–81.

(43) Dai, C.; Meschini, F.; Narayanam, J. M. R.; Stephenson, C. R. J. Friedel-Crafts Amidoalkylation via Thermolysis and Oxidative Photocatalysis. *J. Org. Chem.* **2012**, *77*, 4425–4431.