

Received:
27 July 2018

Revised:
5 November 2018

Accepted:
21 January 2019

Cite as:
Efraín M. Castro-Alayo,
Guillermo Idrogo-Vásquez,
Raúl Siche,
Fiorella P. Cardenas-Toro.
Formation of aromatic
compounds precursors during
fermentation of Criollo and
Forastero cocoa.
Heliyon 5 (2019) e01157.
doi: [10.1016/j.heliyon.2019.e01157](https://doi.org/10.1016/j.heliyon.2019.e01157)



Review Article

Formation of aromatic compounds precursors during fermentation of Criollo and Forastero cocoa

Efraín M. Castro-Alayo^{a,c,*}, Guillermo Idrogo-Vásquez^a, Raúl Siche^b,
Fiorella P. Cardenas-Toro^c

^a Institute of Research, Innovation and Development for the Agricultural and Agroindustrial Sector of the Amazonas Region (IIDAA - Amazonas), Faculty of Engineering and Agricultural Sciences, Toribio Rodríguez de Mendoza National University of Amazonas, Higos Urco Street 342-350-356, Chachapoyas, Amazonas, Peru

^b Institute of Research and Development, National University of Trujillo, Av. Juan Pablo II s/n, University City, Trujillo, Peru

^c Section of Industrial Engineering, Department of Engineering, Pontifical Catholic University of Peru, Av. Universitaria 1801, San Miguel, Lima 32, Peru

* Corresponding author.

E-mail addresses: efrain.castro@untrm.edu.pe, efrain.castro@pucp.edu.pe (E.M. Castro-Alayo).

Abstract

There are three main genetic varieties of cocoa (*Theobroma cacao* L) used in chocolate making: Forastero, Trinitario and Criollo, which are distinguished by their aroma, an attribute that determines their quality. Criollo cocoa is of the highest quality and is used in the manufacture of fine chocolates because of its fruity aroma. The aroma of Criollo cocoa is defined by volatile compounds such as pyrazines and aldehydes, which are formed during roasting of the bean, from aroma precursors (reducing sugars and free amino acids) that are generated inside the bean via enzymatic reactions during fermentation; for this reason, fermentation is the most important process in the value chain. This review discusses the production of aroma precursors of Criollo and Forastero cocoa by studying the kinetics of spontaneous fermentation and the role of starter cultures

to produce aroma precursors. Fine aroma precursors produced in the pulp during the fermentation phase will migrate into the bean when its permeability is improved and then retained during the drying phase. Diffusion of aroma precursors into the cocoa bean may be possible, this process is mathematically characterized by the coefficient of molecular diffusion D , which describe the process of mass transfer via Fick's Second Law. The current state of knowledge is analyzed based on existing research and reports some gaps in the literature, suggesting future research that will be necessary for a better understanding of cocoa fermentation.

Keywords: Food technology, Food science

1. Introduction

Cocoa (*Theobroma cacao* L) and its most important products, such as chocolate, are known as luxury foods that provide an astringent taste and typical aroma (Pedan et al., 2017). The fruit of this tree is a pod that contains seeds, commonly known as cocoa beans (Kongor et al., 2016); when harvested, they are surrounded by a sweet, acidic, aromatic mucilage that is pleasant to the palate and is known as pulp; this material is difficult to extract by mechanical means. There are three commonly recognized genetic varieties: Forastero, Criollo, and Trinitario; a fourth variety that grows in Ecuador is called Nacional (Afoakwa et al., 2011). Criollo is the finest variety of cocoa. Cocoa varieties classification is largely based on the aspect of the fruit, so the terms associated to Criollo are descriptive of the fruit characters (Ascrizzi et al., 2017), however it is the least studied. The production of cocoa is led by Forastero, and Criollo contributes a smaller amount to world production, approximately 5% (Afoakwa, 2010). In Latin American production, Peruvian cocoa is of interest for its quality; however, its production is insufficient for international demand. In Amazonas, Peru, dry fermented Criollo cocoa bean is produced for the Italian market, and this product has received the designation of the origin "Cacao Amazonas Peru" by the Peruvian state.

The most important attribute for which the cocoa bean is commercially accepted is its aroma, which also determines its quality. The aroma is formed by volatile compounds that are perceived by the smell receptors of the olfactory tissue of the nasal cavity (Belitz et al., 2009). The aroma of the genetic varieties mentioned above is used for commercial classification into divided into two types of cocoa: bulk/ordinary cocoa (Forastero) and cocoa with a fine aroma (Criollo and Trinitario) (Kongor et al., 2016; Chetschik et al., 2017). The volatile compounds of the cocoa aroma and chocolate are formed during the roasting of the bean, transforming the molecules known as aroma precursors, which are generated during fermentation by proteolysis of the proteins stored inside this bean (Janek et al., 2016). Adequate quantities and proportions of the formed precursors are essential for the optimal

production of aromatic volatiles compounds (Afoakwa, 2010). In Criollo cocoa (Frauendorfer and Schieberle, 2008) revealed concentrations above the odor threshold for 22 compounds in the unroasted and 27 compounds in the roasted cocoa beans, respectively. It is also assumed that cocoa pulp has an impact on the development of aroma during fermentation, due to the possible migration of aroma compounds from the pulp to the bean tissue, and is considered a deposit for the fineness of cocoa (Chetschik et al., 2017).

To improve the quality of cocoa, the effects of the use of starter cultures composed of different types of microorganisms that can lead a controlled fermentation is investigated. Although experimental applications in clone cocoa fermentations of these cultures have given good results (Schwan, 1998; Sandhya et al., 2016), the process is still performed spontaneously, at least among small producers (Vázquez-Ovando et al., 2016). There are no studies on the use of starter culture in fermentation of Criollo cocoa, all studies on the subject mentioned in this document refer to cocoa of other varieties. Therefore, this review to discuss about the production of cocoa aroma precursors during spontaneous and controlled fermentation. The role of the microorganisms in the production of aroma precursors, that influence in the final content of aromatics compounds, and its production kinetics in cocoa beans are analyzed. The current state of knowledge is analyzed based on existing research and reports about some gaps in knowledge, suggesting future research that will be necessary for a better understanding of Criollo cocoa fermentation.

2. Main text

2.1. Formation of the aromatic compounds in cocoa beans

2.1.1. Volatile aromatic compounds in Criollo cocoa

The aroma and taste of chocolate depends on the origin and genotype of the cocoa bean (Menezes et al., 2016), which has been used by the International Cocoa Organization in numerous studies to conclude that, in general, cocoas from different origins have different aroma profiles, thus eliminating market competition among them (ICCO, 2017a). Volatile aromatic compounds are present in traces, mainly at levels of a few $\mu\text{g}/\text{kg}$ or no more than a few mg/kg , with approximately 100 different pyrazines present in the predominant aroma fraction (Beckett, 2009).

Criollo cocoa is of high value and is a fine cocoa used to produce high-quality chocolates. The fine aromas include fruit notes (fresh and ripe), floral, herbal, wood, nuts and caramel notes; monoterpenes such as linalool are also part of the compounds responsible for the fine aroma in cocoa; therefore, fine cocoas contain higher amounts of linalool than bulk cocoa (Ziegleder, 1990). Frauendorfer and

Schieberle (2008) analyzed unroasted and roasted Criollo cocoa, they concluded that various compounds contributing to the aroma of roasted cocoa beans, such as 3-methylbutanoic acid, ethyl 2-methylbutanoate and 2-phenylethanol, were already present in unroasted, fermented cocoa beans and were not increased during roasting. Table 1 shows the compounds of the fine aroma identified in Criollo cocoa of different origins and processing stages (fresh, fermented and roasted bean) (Ascrizzi et al., 2017; Tran et al., 2015). In the Peruvian Criollo cocoa, linalool was not found as the main component, the presence of this component will depend on the geographical origin where the cocoa is cultivated; however, other components responsible for the aroma of floral and caramel were founded.

The basic aromas of cocoa beans include pleasant and balanced chocolate notes (ICCO, 2017b), and the aromas of cocoa and chocolate are attributed to 2,3,5,6-tetramethylpyrazine (TMP) and 2,3,5-trimethylpyrazine (TrMP), which form the so-called basic notes (Sukha et al., 2013). In addition to pyrazines, aldehydes also contribute to the chocolate aroma (Diab et al., 2014). Chocolate manufacturers

Table 1. Components of fine aroma identified in samples of Criollo cocoa of different origins.

Aromatic compound	Description	Provided by Amedei's factory (country Perú** not identified)*		
		Raw cocoa beans, fermented and dried, husked, %	Roasted cocoa beans with husk, %	Roasted cocoa beans, %
2-Heptanol	Citrus, fresh, lemon grass-likee	0.7		0.2
Phenylethyl alcohol	Flowery, spicy, honey-like, rose	1.6		0.8
Ethyl octanoate	Fruity, floral, pineapple	0.5		0.3
Ethyl phenylacetate	Fruit, sweet, honey-like	0.5		0.4
Ethyl decanoate	Pear, grape, brandy	0		0.1
Acetophenone	Sweet, almond, flowery, must-like	0.2		0
cis-Linalool oxide (furanoid)	Sweet, nutty	0.2		0
Linalool	Flowery	1.2		0.5
trans-Linalool oxide (pyranoid)	Floral	0.2		0.1
2-Phenylethylacetate	Fruity, sweet, flowery	2.5	1.5	0.73
1,3-Butanediol	Sweet, flowery, caramel			16.21
2,3-Butanediol	Sweet, flowery			5.12

* Ascrizzi et al. (2017).

** Tran et al. (2015).

separate cocoa beans into fines and bulk; fines are characterized by their peculiar notes and are used to produce fine chocolates (Ascrizzi et al., 2017), while bulk beans are used to produce low-quality chocolates and products such as cocoa powder (Vargas Jentzsch et al., 2016). Table 2 shows a comparison between the roasted dry fermented beans of Criollo cocoa obtained from Tumbes (Peru), Grenada and Venezuela, and Forastero cocoa from Ghana (Africa); the named sensory description is accepted by U. S flavor industry (FEMA, 2018). In fact, there is a clear quantitative difference between the volatile aroma compounds of fine cocoa and bulk cocoa (Tran et al., 2015), fines (Peru) have higher levels of pyrazine than bulk beans (Ghana) and a higher percentage of 1,3-butanediol, which gives notes of sweet, floral and caramel (Tran et al., 2015), to mention just one example. In Criollo cocoa beans from Venezuela, 1,3-butanediol was identified (Álvarez, 2016). 3-methylbutanal (malty) is a Strecker aldehyde (Frauendorfer and Schieberle, 2008) and it is crucial aroma compound contributing to the chocolate aroma intensity (Van Durme et al., 2016; Saputro et al., 2018). Excluding the basic aromas of cocoa, the majority of fine aroma features depend on the sensory quality of ripe pod: for Criollo beans, caramel and walnut notes are due to the very sweet nature of their pulp (Ascrizzi et al., 2017). The Criollo cacao from Grenada has a greater amount of 3-methylbutanal than those from the other countries, also this cacao and the one from Venezuela contain linalool as one of its components; this corroborates the fact that the geographical origin influences the aromatic profile of Criollo cocoa; however, research is still scarce.

Table 2. Comparison of aromatic notes of fine and bulk cocoa in dry and roasted fermented bean.

Aroma notes	Identified component	Sensory description	Percentage contribution to aroma			
			Fine (Criollo - Peru) ²	Fine (Criollo - Grenada) ³	Fine (Criollo - Venezuela) ⁴	Bulk (Forastero - Ghana) ²
Fine notes	3-methylbutanal		0.73	8.08	0.5	0.41
	1,3-butanediol	Sweet, floral and caramel ²	13.18	NI	I	11.11
	Linalool	Coriander, Floral, Lavender, Lemon, Rose ¹	NI	0.03	I	NI
	2-phenylethylacetate	Frutal, sweet, floral ¹	0.73	0.22	I	0.27
Basic notes	TrMP	Cocoa, Earth, Must, Potato, Roast ¹	4.4	0.22	I	3.65
	TMP	Cocoa, Coffee, Green, Mocha, Roast ¹	17.28	NI	0.5	5.49

1. FEMA, 2018; 2. Tran et al. (2015); 3. Frauendorfer and Schieberle (2008); 4. Álvarez (2016).
NI: not identified, I: identified, but no value.

2.1.2. Development of volatile aromatic compounds from aroma precursors

Typical aromatic compounds are developed during the roasting stage (Ascrizzi et al., 2017) from aroma precursors by the Maillard reaction (Giacometti et al., 2015). The fermentation of cocoa pulp is crucial for the generation of aroma precursors, and this process is performed by microorganisms such as yeast, lactic acid bacteria (LAB) and acetic acid bacteria (AAB) (John et al., 2016). Fresh beans with low precursor content will have limited commercial use, and fermentation will not be able to rectify this shortfall; therefore, an appropriate amount and ratio of aroma precursors are essential for the optimal production of volatile aromatic compounds in roasting (Afoakwa et al., 2008; Aprotosoai et al., 2016). Tran et al. (2015) related the degree of fermentation of fermented cocoa beans from different origins with the presence of aroma precursors, establishing as well fermented when there is a high concentration of free amino acids and absence of sucrose (Table 3). In Criollo from Peru, a higher concentration of amino acids was found than in Vietnam and Ghana varieties, so it could be said that it has a better potential to generate aromatic compounds during roasting, provided that the fermentation has been carried out efficiently.

The results of previous studies have shown that peptides and free hydrophobic amino acids, such as leucine, alanine, phenylalanine and tyrosine, are precursors that contribute to the aroma formation of cocoa and chocolate (Voigt, 2009; Sukha et al., 2017) and develop during fermentation (Hashim et al., 1998) through the proteolysis of vicilin-class globulin (VCG) (Janek et al., 2016; Kumari et al., 2016), which is induced by acetic and lactic acids (Voigt et al., 2016) and cooperative action of the aspartic endoprotease and carboxypeptidase that are present in

Table 3. Aroma precursors founds in different varieties and origin of cocoa beans.

	Clone/variety		
	Criollo	Tritario × Forastero	Forastero
Hybrid name	Tumbes	TD10	PA7xIFC5
Origin	Peru	Vietnam	Ghana
Sucrose, mg/g	0	0	2.42 ± 0.18
Glucose, mg/g	0	0	0
Fructose, mg/g	4.61 ± 0.98	10.01 ± 0.13	7.18 ± 1.15
Leucine, mg/g	3.14 ± 0.01	2.75 ± 0.02	2.00 ± 0.07
Alanine, mg/g	2.27 ± 0.00	2.05 ± 0.01	1.38 ± 0.06
Phenylalanine, mg/g	2.45 ± 0.00	2.14 ± 0.03	1.58 ± 0.10
Tyrosine, mg/g	1.55 ± 0.01	1.33 ± 0.01	1.09 ± 0.05

Data extracted from Table 2 published by Tran et al. (2015).

mature cocoa beans and those that are not germinated. Hydrophilic peptides and free hydrophobic amino acids contribute to the aroma by their reaction with fructose and glucose (Afoakwa, 2010) during roasting. In the pulp, sucrose is hydrolyzed to glucose and fructose by the invertase activity of the yeast, as well as in the bean by diffusion of acetic acid, lactic acid and ethanol, together with the production of heat (de Melo Pereira et al., 2013) (Fig. 1). Approximately 25% of free amino acids and 70% of glucose and fructose are used (Beckett, 2009). Leucine and glucose produce aromatic notes described as “sweet chocolate” (Afoakwa, 2010). Recent work concludes that there is no clear quantitative correlation between the amounts of precursors and the aromatic compounds formed (Tran et al., 2015; Frauendorfer and Schieberle, 2008), while others claim that Criollo cocoa beans contain high levels of precursors that can produce high levels of pyrazines (Giacometti et al., 2015), however, pyrazines are not responsible for the fine aroma of Criollo cocoa. The compounds 3-methylbutanal and phenylacetaldehyde (honey, green, floral (Crafack et al., 2014)) correspond to the so-called aldehydes of Strecker degradation, which are the products derived from valine, leucine, isoleucine and phenylalanine. The compounds 3-methylbutanal, 2-methylbutanal and 2-methylpropanal provide a fruity aroma and contribute to the sugary taste that is characteristic of Criollo cocoa (Álvarez et al., 2012). In a previous analysis of dry fermented beans of Criollo cocoa




Fig. 1. Aroma precursors produced during cocoa bean fermentation by microbial action and its molecular diffusion (adapted from Beckett, 2009). (*) In addition to the internal production of phenylalanine from the VCG, this amino acid can also be produced by *Bacillus* in the pulp of the bean, then called external precursor, and later enter the interior of the bean by molecular diffusion to be part of the internal precursors.

produced by the Asociación de Productores Cafetaleros y Cacaoteros de Amazonas (APROCAM)-Peru, 28.08% of 3-methylbutanal and 9.17% of 2-methylbutanal were detected (data not yet published). In addition to the endogenous formation of the precursors, there is growing evidence of the exogenous influence of the pulp directly on the development of the aroma (Sukha et al., 2017). It is believed that aromatic compounds directly penetrate from the pulp into the bean tissue during fermentation and can be retained during the drying process (Eskes et al., 2012).

There are few studies on volatiles compounds of Criollo cocoa, (Rodriguez-Campos et al., 2011, 2012; Ramos et al., 2014) characterized the dynamics of volatile compounds during fermentation of cacao varieties, including the bulk Forastero and various cacao hybrids, but the fine-flavor cultivars Criollo were not assessed. The volatile aromatic compounds present in Criollo, Forastero, and Trinitario cocoa beans from the same origin have been characterized (Qin et al., 2016; Tran et al., 2015; Counet et al., 2004) but changes in aroma precursors occurring during fermentation Criollo cocoa were not reported.

2.2. Action of the microorganisms in the production of aroma precursors and aromatics compounds

In spontaneous fermentation of cocoa beans Forastero hybrid from Ivory Coast (Visintin et al., 2016), isolated 106 yeasts, 105 LAB and 82 AAB which were identified by means of rep-PCR grouping and sequencing of the rRNA genes. Oxygen is one of the factors that determines the microbial succession. Yeasts and LAB are the predominating microbial species at the initial phase of the fermentation process (Romanens et al., 2018). Jamili (2016); Ardhana (2003) states that the yeast at most the number and the type found in 24–36 hours after fermentation of cocoa because of their role in the process of degradation the various types of sugar that contained in the cocoa bean pulp. Visintin et al. (2017) found that in fermentation of two different cocoa hybrids (PS1319 and SJ02), the yeast (*Saccharomyces cerevisiae* and *Torulopsis delbrueckii*) metabolism results in ethanol production. In region called Agneby-Tiassa, species of *Pichia kudriazevii* and *Candida nitrativorans* was isolated; they are resistant to high temperatures (40 °C) and high ethanol concentrations (20%). Moreover, they exhibited pectin-hydrolysing enzymatic activities, suggesting its key role in the degradation of cocoa bean pulp during fermentation (Samagaci et al., 2016).

Spontaneous Malaysian cocoa bean box fermentations were carried out with high-quality raw material, resulting in successfully fermented dry cocoa beans and good chocolates produced thereof, it is likely that the prevailing species *Hanseniaspora opuntiae*, *S. cerevisiae*, *Lactobacillus fermentum* and *Acetobacter pasteurianus* were responsible for it (Papalexandratou et al., 2013). In cocoa beans fermentation performed in Abidjan (South East of Ivory Coast) (Koné et al., 2016), identified

yeasts that produced a total of 33 aroma compounds, among all yeasts involved *P. kudriavzevii*, *S. cerevisiae*, *Galactomyces geotrichum* and *Wickerhamomyces anomalus* could be considered as the most important contributors to the formation of cocoa specific aromatic compounds. Yeasts produced ethanol from sugars, and LAB produced lactic acid, acetic acid, ethanol, and mannitol from sugars and/or citrate (Camu et al., 2007; Papalexandratou et al., 2013). The experiment conducted by (Magalhães da Veiga Moreira et al., 2016) in Porto Híbrido 9, Porto Híbrido 15 and Porto Híbrido 16 cocoa beans at the Vale do Juliana farm in Igrapiúna, Bahia, Brazil, *Hanseniaspora uvarum*, *P. kluyveri*, *P. caribbica* and *S. cerevisiae* were the predominant yeasts in the three fermentations. *L. plantarum*, *A. pasteurianus*, *Bacillus cereus* and *Lysinibacillus fusiformis* were the common bacterial species isolated from the fermentations. Twenty-seven volatile compounds were identified during the cocoa fermentation of all hybrids (Koffi et al., 2018). Investigated 743 yeast strains, two strains *S. cerevisiae* YB14 and *P. kudriavzevii* YP13 were able on one hand to resist to most parameters such as temperature, pH, ethanol, organics acid, and on the other hand they were able to produce specific enzymes like pectinase, β -glucosidase, protease necessary to have a good cocoa fermentation process, and finally to produce acetoin which is desirable for flavor development in the fermentation process. These two strains could therefore be used as starter cultures which may contribute to the control of cocoa fermentation in Ivory Coast. Other results shown that *P. kudriavzevii* and *Candida nitrativorans* are key players in cocoa bean fermentation in the Agneby-Tiassa region and they are promising candidates for developing starter cocktails that could be used to improve the overall efficiency of cocoa fermentation in Ivory Coast (Samagaci et al., 2016). Furthermore some specific yeast isolates could be used as biological markers to predict the determining of chocolate sensorial characteristics and to indicate the geographical origin or processing story of cocoa bean batches (Koné et al., 2016). Actually, it was concluded that the growth of LAB and AAB may not be essential for the fermentation of cocoa beans (Ho et al., 2018).

The experiment conducted by (de Melo Pereira et al., 2013) at a cocoa farm located in the city of Itajuípe, Bahia State, Brazil, the dominant species of major physiological roles were *S. cerevisiae* and *Hanseniaspora sp.* in the yeast group; *L. fermentum* and *L. plantarum* in the LAB group; *A. tropicalis* belonging to the AAB group; and *Bacillus subtilis* in the *Bacillaceae* family. It demonstrated that at least three species of *Bacillus* are involved in cocoa fermentation, namely *B. subtilis*, *B. flexus* and *B. megaterium* (de Melo Pereira et al., 2013). Aerobic spore-forming bacteria, such as *Bacillus* strains, produce a variety of chemical compounds, including 2,3-butanediol, pyrazines, acetic acid and lactic acid, under fermentative conditions, which may contribute to the acidity and perhaps, at times, to off-flavors of fermented cocoa beans (Schwan, 1998). Increased aeration, increased pH value (3.5–5.0) of cocoa pulp, and a rise in temperature to about 45 °C in the cocoa mass in the later stages

of fermentation are associated with the development of bacteria of the genus *Bacillus* (Schwan and Wheals, 2004).

In Forastero and Trinitario cultivars at commercial fermentations in East Java, Indonesia was investigated. The later stages of fermentation were dominated by the presence of *Bacillus* species, mostly, *B. pumilus* and *B. licheniformis* (Ardhana, 2003; Nielsen et al., 2007). With regard to the fate of *Bacillus spp.* towards the end of the fermentation, this constitutes a microbial group that persists until the end (Lima et al., 2011). *Bacillus spp.* has also been classified as responsible for the production of TMP (de Melo Pereira et al., 2013) and 2,3-butanediol (sweet, floral (Crafack et al., 2014; Ardhana, 2003)) during cocoa fermentation; a temperature of 50–55 °C assists in this formation (Selamat et al., 1994).

The role of bacilli in cocoa fermentation is not well known (Nielsen et al., 2007). However (Ouattara et al., 2008) suggest that *Bacillus sp.* is liable to produce at least one enzyme during cocoa fermentation, so the degradation of the pulp during cocoa fermentation might not be only due to pectinolytic enzymes produced by yeasts but rather by a combined action of enzymes produced by both yeast and *Bacillus* strains. Ouattara et al. (2017) assumed that, during the latter stage of cocoa fermentation when simple sugars are depleted, production of pectate lyases (Pel) in *Bacillus* is stimulated by DegU gene to supply microbial cells with carbon source from polymeric pectic compounds. It is possible that *Bacillus spp.* play a bigger role in box fermentations than in heap fermentations, but more fermentations need to be investigated before firm conclusions can be drawn (Nielsen et al., 2007).

In other raw materials, the result showed that *Bacillus sp.* KR-8104 growth and the production of α -amylase on a wheat bran substrate could successfully (Hashemi et al., 2011). The production of L-phenylalanine (phenylalanine) is related to strains of *B. subtilis* (De Boer and Dijkhuizen, 1990). In brewery wastes, *B. subtilis* UO-01 was studied. Protease production delay seemed to be related to the consumption of non-protein and protein nitrogen, the maximum protease level (9.77 EU/mL) was obtained at pH 7.1 and 37.8 °C (Sánchez Blanco et al., 2016). In soya roasted cotyledons fermented for 18 h and 36 h at 35 °C, growth of *B. subtilis* led to the formation of volatile compounds: acetoin, 2,5-dimethylpyrazine (2,5-DMP) and TrMP (Owens et al., 1997). 2,5-DMP and TMP were produced using *B. subtilis* IFO 3013 grown on soybeans (Besson et al., 1997).

In a previous analysis of dry fermented beans of Criollo cocoa from APROCAM, considerable amounts of TMP, TrMP, 2,3-butanediol and acetoin (butter, creamy, green pepper (FEMA, 2018)) were founds (data not yet published). The presence of these compounds would lead us to think that in addition to the yeasts, LAB and AAB there may be certain *Bacillus* strains during the spontaneous fermentation of Criollo cocoa, however there is no information to corroborate this hypothesis, since the data so far found only they refer to other varieties or clones of cocoa.

Likewise, in previous studies (not in cocoa) the production of phenylalanine by *B. subtilis* has been reported, which gives the possibility of finding this amino acid in spontaneous cocoa fermentation and we would be facing a process of production of “external aroma precursors” by microbial action on the outside of the cocoa bean (Fig. 1).

2.3. Solid-state fermentation (SSF) of cocoa beans

2.3.1. Spontaneous fermentation

Fresh cocoa beans are very astringent and do not develop the delicious and typical chocolate aromas alone (Papalexandratou and Nielsen, 2016) because they do not contain specific aroma precursors (Kadow et al., 2015); thus, the reason for the fermentation of cocoa is to reach biochemical reactions inside the bean that lead to the formation of aroma precursors, flavor and color, reduction of bitter and astringent tastes, and improvement of the physical appearance of cocoa (Apriyanto et al., 2017). Well-conducted fermentation is a prerequisite for the production of high-quality chocolates; if performed improperly, the production of specific aroma compounds will fail during the post-processing stages (Crafack et al., 2013). A 5-day fermentation period is common for the Forastero cultivar while shorter periods of about 2–3 days are required for Criollo and Nacional cultivars (Cevallos-Cevallos et al., 2018). In Peru, fermentation systems are used in wooden fermentation boxes, forming a batch process. These boxes act as bioreactors; the process begins at room temperature (27 °C) and ends at approximately 50 °C in seven days, with the transfer of the beans from one box to another from the second day to incorporate air. Cocoa beans are fermented by small producers in a traditional and uncontrolled process known as spontaneous fermentation (Magalhães da Veiga Moreira et al., 2017). During fermentation, small amounts of free water are present, and this process is characterized as SSF (Nielsen et al., 2014).

Until the pods open, the beans are sterile microbiologically. Once the pod is opened, the beans are exposed to numerous sources of microorganisms, including the hands of farmers and utensils. The immediate effect is the initiation of microbiological attack on the pulp sugars by the yeasts, and this action is followed by the LAB and AAB (Afoakwa, 2010). This succession is directed by yeasts, which dominate the microbial population during the first hours, followed by the LAB, which decrease after 48 hours of fermentation, and finally the AAB increase in number (Ardhana, 2003). The fermentation of cocoa beans occurs in two phases. The first involves reactions that occur in the pulp, and the second involves several hydrolytic reactions that occur within the bean (Pereira et al., 2012). While the first phase is performed at 25–45 °C, the second phase is at 42–52 °C (Kadow et al., 2015). Changes in pH, temperature, sugar content and fermentation products or metabolites exert a selection pressure on existing natural biotypes, favoring the strains that are better adapted to

this environment (Pereira et al., 2012). These metabolites kill the bean and trigger a series of biochemical reactions in their interior (Pereira et al., 2016), generating aroma precursors (Fernández Maura et al., 2016); the pulp is solubilized, giving rise to a liquid material (exudate) that drains through the holes of the fermenter boxes (Ardhana, 2003), and this material serves as a way of transport for the metabolites of fermentation.

The pulp is the base substrate for microorganisms during fermentation. It is important to define the best state of the pulp for a successful fermentation according to the best results of quality determining components (Voigt and Lieberei, 2014). In fine cocoas, the level of the basic aromatic components (basic notes) of cocoa (and chocolate) is usually low due to the short duration of the fermentation (2–4 days), compared to bulk cocoa (5–7 days) (Torres-Moreno et al., 2012). The findings of previous studies show that linalool is a constituent of the pulp that is transferred to the beans during fermentation (Voigt and Lieberei, 2014). Linalool can be produced by *S. cerevisiae* from the catabolism of leucine (Carrau et al., 2005). It is possible that glycosidase enzymes (α -arabinosidase, β -galactosidase, α -mannosidase) play an important role in the release of this compounds (Voigt and Lieberei, 2014). For example, 1,020 ppb of linalool was found in dry fermented Criollo cocoa beans (Chetschik et al., 2017), or, in another study, 130 $\mu\text{g}/\text{kg}$ (Frauendorfer and Schieberle, 2008).

The quality of beans is strongly dependent on the degree and time of acidification of the cotyledon during the fermentation process (Voigt and Lieberei, 2014). Cocoa fermentation at a pH of approximately 5 shows a cocoa with the highest potential for specific aromas (Janek et al., 2016). It is known that strong acidification of the cotyledon ($\text{pH} < 5$) during the fermentation produces a poor potential of aroma in the cocoa beans, because the activity of carboxypeptidase is optimal at approximately pH 5.5 and is reduced to $\text{pH} < 5$. At pH values >5.5 , the aspartic endoprotease is inactivated (Voigt et al., 1994).

Spontaneous fermentation is an empirical procedure, which may not yield beans of consistent quality, which requires the chocolate industry to continuously modify its formulations (de Melo Pereira et al., 2013). Common problems involve acidity levels and incomplete fermentation, resulting in deficient or unpleasant aromas (Schwan, 1998). The studies carried out in a simulated medium of cocoa pulp and indicate that there could be competition among specific strains called *L. plantarum*, *L. fermentum* and *A. pasteurianus* during the spontaneous fermentation process, improving the aroma of the bean (Lefeber et al., 2010); however, there is no further information on how this has improved the fermentation process of the cocoa bean under field conditions (Ghosh, 2016), and thus producers still use spontaneous fermentation for their commercial cocoa. As a alternative (de Melo Pereira et al., 2013), described the first experimental validation of the stainless steel tank method

for cocoa bean fermentation. They concluded that the use of stainless steel tanks may be of great interest for those who seek improved control over the cocoa fermentation process and/or to optimize cocoa fermentation through the use of starter cultures. In accord (Mota-Gutierrez et al., 2018), found that the microbial dynamics and associations between the bacteria, yeast and metabolites were found to depend on the type of fermentation. With the growth of niche markets for chocolate manufacturers, a better understanding of the factors that contribute to flavor variations will have significant commercial implications (Afoakwa et al., 2008). As a consequence of the increasing implementation of traceability systems in the food industry, detailed information about how products are handled could be transferred to the producers and could be used to optimize the processes (Saltini et al., 2013).

Although it is true that cocoa fermentation has been widely studied, there are few works on spontaneous fermentation of Criollo cocoa, in the work made by (Cevallos-Cevallos et al., 2018), spontaneous fermentation was carried out in a greenhouse at ambient temperature (about 35 °C) for Forastero, Criollo and Nacional cultivars; fine-flavor cultivars were characterized by the presence of significant amounts of hydrocarbons such as valencene, aromadendrene, germacrene, 1,3 cyclohexadiene and octacosane while fermented beans of the Forastero bulk cocoa showed negligible levels of these volatiles. In Amazonas, the fermentation process is the same for both Criollo cocoa and CCN 51 clone (bulk cocoa); the same boxes are used and a product is obtained in 7 days, which leads to obtaining cocoa beans of different qualities. This is a problem that has not yet been solved due to the lack of research.

2.3.2. Fermentation controlled by starter culture

As far as we know, the use of starter cultures in Criollo cocoa fermentation has not been studied; therefore, the studies analyzed here generally correspond to cocoa clones. Studies of fermentation and chocolate using PS1319 hybrid cocoa and starter cultures can provide important results for improving the fermentation of cocoa and the quality of chocolate (Batista et al., 2016). The use of LAB starter cultures could lead to a controlled fermentation process and thus the possibility of controlling the aroma (Saltini et al., 2013). SSF was used to ferment Forastero cocoa with a starter culture; it was found that a 10% culture composed of *S. cerevisiae*, *L. plantarum* and *A. aceti* was adequate to produce chocolate with 7.5 out of 10 points on a hedonic scale for acceptability of the aroma of chocolates produced from cocoa beans fermented with starter culture (Sandhya et al., 2016). Other researchers have inoculated starter culture composed of *S. cerevisiae*, *L. plantarum* and *A. pasteurianus* in the fermentation of cocoa hybrids; the fermentation process was accelerated, and the chocolates made with inoculated beans were bitter, sweet and cocoa-flavored, with considerable amounts of 3-methylbutanal, 2-phenylethylacetate (Magalhães

da Veiga Moreira et al., 2017). In the roasted liquor of cocoa produced from fermentations of hybrids inoculated with *L. fermentum* L18, *A. pasteurianus* A149, *P. kluyveri* and *Kluyveromyces marxianus*, 1,3-butanediol was detected (Crafack et al., 2014). Thus, inoculation affects the profiles of volatile compounds and their relative concentrations, which influence the sensory characteristics of chocolate (Batista et al., 2016).

Results reported by (Pereira et al., 2017) indicated that inoculated fermentations with *P. kudriavzevii* LPB06 and *P. kudriavzevii* LPB07 generated cocoa beans with better color development and richer aroma composition, suggesting that cocoa-associated yeast diversity at strain level can be exploited for flavor modulation of cocoa beans. In fermentation inoculated with *S. cerevisiae* FNCC 3056, *Lactococcus lactis* FNC 0086 and *A. aceti* FNCC 0016, an increase of hydrophobic amino acids was obtained, demonstrating that the addition of inoculum can gradually degrade protein to produce more hydrophobic amino acids as aroma precursors (Apriyanto et al., 2017). LAB can also metabolize sugars and organic acids to produce various aldehydes, ketones and other volatile components that can impact the sensory quality of the bean (Ardhana, 2003). It is known that some compounds come from the metabolism of yeasts, such as phenylethylacetate (fruit, sweet, honey (Magalhães da Veiga Moreira et al., 2016)), 2-phenylethanol (honey, floral (Magalhães da Veiga Moreira et al., 2016)), 2-phenylacetic acid and acetoin (butter, cream (Batista et al., 2016; Visintin et al., 2017)). Table 4 shows that in the fermentation of cocoa hybrids controlled by a starter culture, it is possible to obtain aromatic compounds identified in Criollo cocoa (Crafack et al., 2014; Batista et al., 2016; Visintin et al., 2017; Magalhães da Veiga Moreira et al., 2017). It may not be necessary to use starter cultures to improve the aromatic quality of Criollo cacao, which is why this type of research has not been carried out with this variety, since its own genetics and origin would determine its fine aroma. However, we consider it important to study the process of fermentation of Criollo cocoa to have a standard process based on the amount of aroma precursors produced. Then, so that the Forastero cocoa fermentation reaches this standard, a type and concentration of the starter culture would be necessary. From the isolation of the best strains found in the Criollo cocoa fermentation, this culture would be developed and used in Forastero cocoa fermentation.

2.4. SSF kinetics through mathematical modeling and its application in cocoa

A mathematical model serve as tools for engineers and scientists to develop an understanding of important systems and processes using mathematical equations (Ribas-García et al., 2011; Rasmuson, 2014), is an important instrument which allows understanding the behavior of food, predict results (Pinheiro et al., 2017),

Table 4. Fine aroma compounds produced by starter culture.

Starter culture	Sample	Raw material	Fine aroma components identified	Source
<i>S. cerevisiae</i> UFLA CA11 <i>P. kluyveri</i> CCMA0237 <i>Hanseniaspora uvarum</i> CCMA0236	Chocolate and dry fermented beans	Hybrid PS1319, Brazil	2-Heptanol, 2-pentanol, Phenylethyl alcohol, 2-Methylbutanal, 2-Phenyl-2-butenal, Ethyl octanoate, Ethyl phenylacetate, 2-Heptanone, 2-Phenylethyl acetate, 2-pentanone, 2-nonanone, 2-undecanone, Acetophenone	Batista et al. (2016)
<i>S. cerevisiae</i> UFLA CCMA 0200 <i>L. plantarum</i> CCMA 0238 <i>Acetobacter pasteurianus</i> CCMA 0241	Chocolate	PH15 cocoa hybrid, Brazil	2-Heptanol, 2-pentanol, 2,3-Butanediol, Phenylethyl alcohol, 2-Phenyl-2-butenal, 2-Heptanone, Acetophenone, Linalool, trans-Linalool oxide (pyranoid), cis-Linalool oxide (furanoid), 2-nonanone	Magalhães da Veiga Moreira et al. (2017)
<i>S. cerevisiae</i> <i>Torulaspora delbrueckii</i>	Chocolate	Hybrids of cocoa: PS1319 and SJ02 de Brazil	2,3-Butanediol	Visintin et al. (2017)
<i>P. kluyveri</i> <i>Kluyveromyces marxianus</i>	Chocolate	Forastero from Ghana	2-pentanol, 1,3-Butanediol, 2,3-Butanediol, 2-Methylbutanal, 2-Phenyl-2-butenal, Ethyl phenylacetate, Ethyl decanoate, Ethyl phenylacetate, 2-Heptanone, 2-nonanone, Linalool, trans-Linalool oxide (pyranoid), β -myrcene	Crafack et al. (2014)

generalize the processes evaluating the optimal conditions that lead them (Kulov and Gordeev, 2014) and generate insurance control mechanisms for process quality (Balbinoti et al., 2018). A realistic kinetic model is required for the design, optimization and control of processes (Ccopa Rivera et al., 2017). The SSF is driven by a complex interaction between various microorganisms and their metabolites. To control this process, and consequently the quality of the final product, a deep

understanding of this interaction is necessary (Nielsen et al., 2014). Then, the optimal parameters to control the process can be determined through the development of kinetic models for the process (Liu et al., 2016). For example, progress has already been made in describing the process of SSF conducted by *S. cerevisiae* in the production of rice wine, for which a first-order kinetic model based on the analysis of the main metabolites of fermentation has been developed (Liu et al., 2015). In the case of cocoa, there have been some attempts to study the transition of the microbial population in heap fermentation, but there are no detailed studies of its kinetics (Ghosh, 2016). Recently (Moreno-Zambrano et al., 2018), has constructed a mathematical model using ordinary differential equations with two distinct types of state variables: (i) metabolite concentrations of glucose, fructose, ethanol, lactic acid and acetic acid and (ii) population sizes of yeast, LAB and AAB. However, it is worth mentioning that the spontaneous fermentation process will be specific for each batch of cocoa, therefore it will be necessary to develop a controlled fermentation that is a standard reference for the batches and thus be able to produce cocoa of the most uniform quality possible.

The study of SSF involves the determination of fermentation kinetics (e.g., biomass production, substrate consumption and production of metabolites). The data collection comprises the measurement of microbial growth quantified by standard methods of isolation and metabolic activity with high-performance liquid chromatography (HPLC) (Nielsen et al., 2014). Kinetic models use parameters that describe how quickly a microorganism can grow, such as the duration of the lag phase and generation time, and those models are used as the response variable for various conditions of pH, temperature, and water activity (Adams and Moss, 2008). Studies have demonstrated the utility of kinetic models that can predict cell growth and ethanol production from sugarcane fermentations (Ccopa Rivera et al., 2017). The influence of fermentation practices on the dynamics of the microbial population over 3 days in National cocoa (a type of fine cocoa) has been studied; analyzing the pulp and bean metabolites and sensory analysis of the chocolates produced from both fermentations (Papalexandratou et al., 2011); however, they did not develop a kinetic model that allows predicting the behavior of the process. In SSF the water content of the solid substrate is usually maintained between 12 and 80% (Chen, 2013). The most important consideration for kinetics is the formation of biomass, either linear (equation 1), exponential (equation 2), logarithmic (equation 3) or two-phase formation (equation 4) (Ghosh, 2016). Where K is the linear growth rate, μ is the specific growth rate, x is the biomass at a time t , x_m is the maximum biomass, t_a is the time when the deceleration phase begins, k is the first order constant and L is a ratio between the growth rate in the deceleration phase and the specific growth rate during the exponential phase. It is still unknown to extent these calculations are valid in food processing SSF (Ghosh, 2016).

$$dx/dt = K \quad (1)$$

$$dx/dt = \mu x \quad (2)$$

$$dx/dt = \mu x [1 - (x/x_m)] \quad (3)$$

$$dx/dt = [\mu L e^{-k(t-t_a)}] x, t > t_a \quad (4)$$

Previous studies suggest that the cotyledon of cocoa absorbs the aromatic components of the pulp during the fermentation process (Eskes et al., 2012). This cotyledon is considered a reservoir for fine notes (Chetschik et al., 2017); however, the exact mechanism by which this situation is reached and the role of the fermentation microflora in this process has not been investigated (Ali et al., 2013). The fine aroma precursors produced in the pulp during the fermentation phase will migrate into the bean when the permeability of the bean is improved and are then retained during the drying phase (Ascrizzi et al., 2017); Then, the diffusion of aroma precursors into the Criollo and Forastero cocoa bean is possible; for example, during the process of fermentation of cocoa, the amino acid phenylalanine (external precursor) produced by bacillus in the pulp of cocoa can enter the bean by molecular diffusion and accumulate inside to join the amino acids (internal precursors) already produced during the internal reaction of the VCG, this way increase the amount of aroma precursors (Fig. 1). The bean shell would be the barrier for this diffusion and its structure would determine the speed of this phenomenon and therefore the speed of the process to obtain an optimal value. Theoretically, this process is characterized by the coefficient of molecular diffusion D . The experimental determination of this coefficient is indispensable to describe the process of mass transfer via Fick's Second Law (Koukouch et al., 2017) in a non-stationary state.

In other studies, it has been demonstrated that fermentations with added passion fruit pulp showed more pronounced aromas of interest, namely, fruit, acid and floral notes in cocoa beans (Ali et al., 2013). In the cucumber fermentation process, the exchange of malic acid, lactic acid, NaCl and sugar between the cucumber and its brine was monitored; an exponential equation that described the movement of the solutes during the fermentation was described, and the diffusion coefficient of the sugar was estimated between 1.80×10^{-9} and 9.18×10^{-9} m²/s, which was used to determine the optimal concentration of sugar within the cucumber at any time; to determine processing parameters in manufacturing, this process was modeled mathematically (Fasina et al., 2002). If this phenomenon occurs in the aforementioned food systems, it will also happen in the process of fermentation of Criollo and Forastero cocoa, where it will be necessary to characterize the phenomenon of transfer of aroma precursors to the interior of the bean to establish an optimal process time. In the case of Criollo de Amazonas cocoa, it is possible that the current fermentation time of 7 days is excessive and it would be risking the aromatic quality of the cocoa to obtain an acceptable percentage of fermentation.

During the last decade, the analysis and the modeling of the kinetics of the convective drying of cocoa beans were the scope of several papers (Hii et al., 2008; Hii et al., 2009b; Hii et al., 2009a; Clement et al., 2009; MacManus Chinenye et al., 2010; Teh et al., 2016; Nwakuba et al., 2017; Olatinjo et al., 2017; Herman et al., 2018) leaving a side the study of the kinetic of formation and diffusion of aroma precursors through mathematical models that allow to improve processes, even more so in Criollo cacao there have been no advances.

3. Conclusions

The cocoa market has divided this product into fine aroma and bulk. Because bulk beans are more productive and resistant to diseases, studies are being carried out with the aim of improving their quality. The technology developed for this purpose is the use of starter cultures that help the fermentation process making it more efficient, this culture will be mainly composed of yeasts, since they would be the most important microorganisms. However, this technology has not been widely adopted by cocoa farmers, at least in Peru the traditional process of spontaneous fermentation has been carried out for all varieties of cocoa, including Criollo and Forastero. To achieve the expected quality results, it is necessary to study the production of external aroma precursors by the microorganisms and the diffusion of these precursors towards the interior of the bean and how they come to be added to the internally generated precursors until reaching optimum concentrations. The study of this phenomenon of molecular diffusion must be carried out through mathematical modeling in order to predict these concentrations in a determined time. We consider it necessary to know in depth the process of fermentation of Criollo cocoa, the process of diffusion of external precursors and isolate the microorganisms founded in the process, from this, develop starter cultures that can be applied to the process of fermentation of cocoa in Forastero (bulk). Undoubtedly, the both origin and genetic factors will play an important role that should not be left aside.

Declarations

Author contribution statement

All authors listed have significantly contributed to the development and the writing of this article.

Funding statement

This work was supported by Project N°16808-2016/PROYECTO-CACAO, financed by the National Agrarian Innovation Program (PNIA) of the Peruvian

government, the Toribio Rodríguez de Mendoza National University of Amazonas and the Pontifical Catholic University of Peru.

Competing interest statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

References

Adams, M.R., Moss, M.O., 2008. Food Microbiology, third ed. RSC Publishing, Cambridge, UK, pp. 20–52.

Afoakwa, E.O., 2010. Chocolate Science and Technology. Wiley-Blackwell, Chichester, U.K.; Ames, Iowa, pp. 58–71.

Afoakwa, E.O., Paterson, A., Fowler, M., Ryan, A., 2008. Flavor formation and character in cocoa and chocolate: a critical review. Crit. Rev. Food Sci. Nutr. 48, 840–857.

Afoakwa, E.O., Quao, J., Budu, A.S., Takrama, J., Saalia, F.K., 2011. Effect of pulp preconditioning on acidification, proteolysis, sugars and free fatty acids concentration during fermentation of cocoa (*Theobroma cacao*) beans. Int. J. Food Sci. Nutr. 62, 755–764.

Ali, N.A., Baccus-Taylor, G.S.H., Sukha, D.A., Umaharan, P., 2013. The effect of cacao (*Theobroma cacao* L.) pulp on final flavour. In: III International Conference on Postharvest and Quality Management of Horticultural Products of Interest for Tropical Regions 1047, pp. 245–254.

Álvarez, C., Pérez, E., Boulanger, R., Lares, M., Ssemat, As, Davrieux, F., Emile, C., 2012. Identificación de los compuestos aromáticos en el cacao criollo de Venezuela usando microextracción en fase sólida y cromatografía de gases. Vitae 19, S370–S372. Disponible en: <http://www.redalyc.org/articulo.oa?id=169823914116>.

Álvarez, C., 2016. Identification of the volatile compounds in the roasting Venezuela criollo cocoa beans by gas chromatography-spectrometry mass. J. Nutr. Health Food Eng. 5 (4), 00178.

Apriyanto, M., Sutardi, S., Supriyanto, S., Harmayani, E., 2017. Amino acid analysis of cocoa fermented by high performance liquid chromatography (HPLC). Asian J. Dairy Food Res. 36, 156–160.

Aprotosoae, A.C., Luca, S.V., Miron, A., 2016. Flavor chemistry of cocoa and cocoa products-An overview. *Compr. Rev. Food Sci. Food Saf.* 15, 73–91.

Ardhana, M., 2003. The microbial ecology of cocoa bean fermentations in Indonesia. *Int. J. Food Microbiol.* 86, 87–99.

Ascrizzi, R., Flamini, G., Tessieri, C., Pistelli, L., 2017. From the raw seed to chocolate: volatile profile of blanco de Criollo in different phases of the processing chain. *Microchem. J.* 133, 474–479.

Balbinoti, T.C.V., Jorge, L.M. de M., Jorge, R.M.M., 2018. Mathematical modeling of paddy (*Oryza sativa*) hydration in different thermal conditions assisted by Raman spectroscopy. *J. Cereal. Sci.* 79, 390–398.

Batista, N.N., Ramos, C.L., Dias, D.R., Pinheiro, A.C.M., Schwan, R.F., 2016. The impact of yeast starter cultures on the microbial communities and volatile compounds in cocoa fermentation and the resulting sensory attributes of chocolate. *J. Food Sci. Technol.* 53, 1101–1110.

Beckett, S.T. (Ed.), 2009. *Industrial Chocolate Manufacture and Use*, fourth ed. Wiley-Blackwell, Chichester, U.K.; Ames, Iowa, pp. 169–188.

Belitz, H.-D., Grosch, W., Schieberle, P., 2009. *Food Chemistry*, fourth ed. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 340–400.

Besson, I., Creuly, C., Gros, J.B., Larroche, C., 1997. Pyrazine production by *Bacillus subtilis* in solid-state fermentation on soybeans. *Appl. Microbiol. Biotechnol.* 47, 489–495.

Camu, N., De Winter, T., Verbrugge, K., Cleenwerck, I., Vandamme, P., Takrama, J.S., Vancanneyt, M., De Vuyst, L., 2007. Dynamics and biodiversity of populations of lactic acid bacteria and acetic acid bacteria involved in spontaneous heap fermentation of cocoa beans in Ghana. *Appl. Environ. Microbiol.* 73, 1809–1824.

Carrau, F.M., Medina, K., Boido, E., Farina, L., Gaggero, C., Dellacassa, E., Versini, G., Henschke, P.A., 2005. De novo synthesis of monoterpenes by *Saccharomyces cerevisiae* wine yeasts. *FEMS Microbiol. Lett.* 243, 107–115.

Cevallos-Cevallos, J.M., Gysel, L., Maridueña-Zavala, M.G., Molina-Miranda, M.J., 2018. Time-related changes in volatile compounds during fermentation of bulk and fine-flavor cocoa (*Theobroma cacao*) beans. *J. Food Qual.* 2018, 1–14.

Ccopa Rivera, E., Yamakawa, C.K., Saad, M.B.W., Atala, D.I.P., Ambrosio, W.B., Bonomi, A., Nolasco Junior, J., Rossell, C.E.V., 2017. Effect of temperature on

sugarcane ethanol fermentation: kinetic modeling and validation under very-high-gravity fermentation conditions. *Biochem. Eng. J.* 119, 42–51.

Chen, H., 2013. *Modern Solid State Fermentation: Theory and Practice*. Springer Science & Business Media, pp. 23–74.

Chetschik, I., Kneubühl, M., Chatelain, K., Schlüter, A., Bernath, K., Hühn, T., 2017. Investigations on the aroma of cocoa pulp (*Theobroma cacao* L.) and its influence on the odor of fermented cocoa beans. *J. Agric. Food Chem.* 66 (10), 2467–2472.

Clement, A.D., Emmanuel, A.N., Patrice, K., Benjamin, Y.K., 2009. Mathematical modelling of sun drying kinetics of thin layer cocoa (*Theobroma cacao*) beans. *J. Appl. Sci. Res.* 5, 1110–1116. https://www.researchgate.net/publication/228508629_Mathematical_Modelling_of_Sun_Drying_Kinetics_of_Thin_Layer_Cocoa_Theobroma_Cacao_Beans.

Counet, C., Ouwerx, C., Rosoux, D., Collin, S., 2004. Relationship between pro-cyanidin and flavor contents of cocoa liquors from different origins. *J. Agric. Food Chem.* 52, 6243–6249.

Crafack, M., Keul, H., Eskildsen, C.E., Petersen, M.A., Saerens, S., Blennow, A., Skovmand-Larsen, M., Swiegers, J.H., Petersen, G.B., Heimdal, H., Nielsen, D.S., 2014. Impact of starter cultures and fermentation techniques on the volatile aroma and sensory profile of chocolate. *Food Res. Int.* 63, 306–316.

Crafack, M., Mikkelsen, M.B., Saerens, S., Knudsen, M., Blennow, A., Lowor, S., Takrama, J., Swiegers, J.H., Petersen, G.B., Heimdal, H., Nielsen, D.S., 2013. Influencing cocoa flavour using *Pichia kluyveri* and *Kluyveromyces marxianus* in a defined mixed starter culture for cocoa fermentation. *Int. J. Food Microbiol.* 167, 103–116.

De Boer, L., Dijkhuizen, L., 1990. Microbial and enzymatic processes for L-phenylalanine production. *Microb. Bioprod.* 1–27.

de Melo Pereira, G.V., Magalhães, K.T., de Almeida, E.G., da Silva Coelho, I., Schwan, R.F., 2013. Spontaneous cocoa bean fermentation carried out in a novel-design stainless steel tank: influence on the dynamics of microbial populations and physical–chemical properties. *Int. J. Food Microbiol.* 161, 121–133.

Diab, J., Hertz-Schünemann, R., Streibel, T., Zimmermann, R., 2014. Online measurement of volatile organic compounds released during roasting of cocoa beans. *Food Res. Int.* 63, 344–352.

Eskes, A., Ahnert, D., Assemat, S., Seguine, E., 2012. Evidence for the Effect of the cocoa Bean Flavor Environment during Fermentation on the Final Flavor Profile of

cocoa Liquor and Chocolate. INGENIC Newsletter. Available on: https://agritrop.cirad.fr/568108/1/document_568108.pdf (Accessed 30 October 2017).

Fasina, O., Fleming, H., Thompson, R., 2002. Mass transfer and solute diffusion in brined cucumbers. *J. Food Sci.* 67, 181–187.

FEMA, 2018. About FEMA | FEMA [WWW Document]. <https://www.femaflavor.org/about> (Accessed 27 September 2018).

Fernández Maura, Y., Balzarini, T., Clapé Borges, P., Evrard, P., De Vuyst, L., Daniel, H.-M., 2016. The environmental and intrinsic yeast diversity of Cuban cocoa bean heap fermentations. *Int. J. Food Microbiol.* 233, 34–43.

Fraudendorfer, F., Schieberle, P., 2008. Changes in key aroma compounds of Criollo cocoa beans during roasting. *J. Agric. Food Chem.* 56, 10244–10251.

Ghosh, J.S., 2016. Solid state fermentation and food processing: a short review. *J. Nutr. Food Sci.* 06, 1–7.

Giacometti, J., Jolić, S.M., Josić, D., 2015. Cocoa processing and impact on composition. In: Preedy, V. (Ed.), *Processing and Impact on Active Components in Food*. Elsevier, pp. 605–612.

Hashemi, M., Mousavi, S.M., Razavi, S.H., Shojaosadati, S.A., 2011. Mathematical modeling of biomass and α -amylase production kinetics by *Bacillus* sp. in solid-state fermentation based on solid dry weight variation. *Biochem. Eng. J.* 53, 159–164.

Hashim, P., Selamat, J., Muhammad, S., Kharidah, S., Ali, A., 1998. Changes in free amino acid, peptide-N, sugar and pyrazine concentration during cocoa fermentation. *J. Sci. Food Agric.* 78, 535–542.

Herman, C., Spreutels, L., Turomzsa, N., Konagano, E.M., Haut, B., 2018. Convective drying of fermented Amazonian cocoa beans (*Theobroma cacao* var. Forasteiro). Experiments and mathematical modeling. *Food Bioprod. Process.* 108, 81–94.

Hii, C.L., Law, C.L., Cloke, M., 2009a. Modeling using a new thin layer drying model and product quality of cocoa. *J. Food Eng.* 90, 191–198.

Hii, C.L., Law, C.L., Cloke, M., 2008. Modelling of thin layer drying kinetics of cocoa beans during artificial and natural drying. *J. Eng. Sci. Technol.* 3, 1–10. <https://www.revistavirtualpro.com/biblioteca/modelado-de-cinetica-de-secado-de-capa-fina-de-granos-de-cacao-durante-secado-artificial-y-natural>.

Hii, C.L., Law, C.L., Cloke, M., Suzannah, S., 2009b. Thin layer drying kinetics of cocoa and dried product quality. *Biosyst. Eng.* 102, 153–161.

Ho, V.T.T., Fleet, G.H., Zhao, J., 2018. Unravelling the contribution of lactic acid bacteria and acetic acid bacteria to cocoa fermentation using inoculated organisms. *Int. J. Food Microbiol.* 279, 43–56.

ICCO, 2017a. Study of the Chemical, Physical and Organoleptic Parameters to Establish the Difference between Fine and Bulk Cocoa [WWW Document]. URL. <https://www.icco.org/about-cocoa/fine-or-flavour-cocoa/10-projects/144-study-the-chemical-physical-and-organoleptic-parameters-to-establish-the-difference-between-fine-and-bulk-cocoa.html> (Accessed 17 May 2017).

ICCO, 2017b. Fine or Flavour Cocoa [WWW Document]. <https://www.icco.org/about-cocoa/fine-or-flavour-cocoa.html> (Accessed 17 May 2017).

Jamili, J., 2016. Diversity and the role of yeast in spontaneous cocoa bean fermentation from Southeast Sulawesi, Indonesia. *Biodiversitas J. Biol. Divers.* 17, 90–95.

Janek, K., Niewianda, A., Wöstemeyer, J., Voigt, J., 2016. The cleavage specificity of the aspartic protease of cocoa beans involved in the generation of the cocoa-specific aroma precursors. *Food Chem.* 211, 320–328.

John, W.A., Kumari, N., Böttcher, N.L., Koffi, K.J., Grimbs, S., Vrancken, G., D'Souza, R.N., Kuhnert, N., Ullrich, M.S., 2016. Aseptic artificial fermentation of cocoa beans can be fashioned to replicate the peptide profile of commercial cocoa bean fermentations. *Food Res. Int.* 89, 764–772.

Kadow, D., Niemenak, N., Rohn, S., Lieberei, R., 2015. Fermentation-like incubation of cocoa seeds (*Theobroma cacao* L.) – reconstruction and guidance of the fermentation process. *LWT - Food Sci. Technol.* 62, 357–361.

Koffi, O., Samagaci, L., Goualie, B., Niamke, S., 2018. Screening of potential yeast starters with high ethanol production for a small-scale cocoa fermentation in Ivory Coast. *Food Environ. Saf.* 17, 113–130. <http://www.fia.usv.ro/fiajournal/index.php/FENS/article/view/572>.

Koné, M.K., Guéhi, S.T., Durand, N., Ban-Koffi, L., Berthiot, L., Tachon, A.F., Brou, K., Boulanger, R., Montet, D., 2016. Contribution of predominant yeasts to the occurrence of aroma compounds during cocoa bean fermentation. *Food Res. Int.* 89, 910–917.

Kongor, J.E., Hinnéh, M., de Walle, D.V., Afoakwa, E.O., Boeckx, P., Dewettinck, K., 2016. Factors influencing quality variation in cocoa (*Theobroma cacao*) bean flavour profile — a review. *Food Res. Int.* 82, 44–52.

Koukouch, A., Idlimam, A., Asbik, M., Sarh, B., Izrar, B., Bostyn, S., Bah, A., Ansari, O., Zegaoui, O., Amine, A., 2017. Experimental determination of the

effective moisture diffusivity and activation energy during convective solar drying of olive pomace waste. *Renew. Energy* 101, 565–574.

Kulov, N.N., Gordeev, L.S., 2014. Mathematical modeling in chemical engineering and biotechnology. *Theor. Found. Chem. Eng.* 48, 225–229.

Kumari, N., Kofi, K.J., Grimbs, S., D'Souza, R.N., Kuhnert, N., Vrancken, G., Ullrich, M.S., 2016. Biochemical fate of vicilin storage protein during fermentation and drying of cocoa beans. *Food Res. Int.* 90, 53–65.

Lefeber, T., Janssens, M., Camu, N., De Vuyst, L., 2010. Kinetic analysis of strains of lactic acid bacteria and acetic acid bacteria in cocoa pulp simulation media toward development of a starter culture for cocoa bean fermentation. *Appl. Environ. Microbiol.* 76, 7708–7716.

Lima, L.J.R., Almeida, M.H., Nout, M.J.R., Zwietering, M.H., 2011. *Theobroma cacao* L., “The Food of the Gods”: quality determinants of commercial cocoa beans, with particular reference to the impact of fermentation. *Crit. Rev. Food Sci. Nutr.* 51, 731–761.

Liu, D., Zhang, H., Lin, C.-C., Xu, B., 2016. Optimization of rice wine fermentation process based on the simultaneous saccharification and fermentation kinetic model. *Chin. J. Chem. Eng.* 24, 1406–1412.

Liu, D., Zhang, H., Xu, B., Tan, J., 2015. Development of a kinetic model structure for simultaneous saccharification and fermentation in rice wine production: kinetic model structure for simultaneous saccharification and fermentation. *J. Inst. Brew.* 121, 589–596.

MacManus Chinenye, N., Ogunlowo, A., Olukunle, O., 2010. Cocoa bean (*Theobroma cacao* L.) drying kinetics. *Chil. J. Agric. Res.* 70, 633–639.

Magalhães da Veiga Moreira, I., de Figueiredo Vilela, L., da Cruz Pedroso Miguel, M., Santos, C., Lima, N., Freitas Schwan, R., 2017. Impact of a microbial cocktail used as a starter culture on cocoa fermentation and chocolate flavor. *Molecules* 22, 766.

Magalhães da Veiga Moreira, I., Gabriela da Cruz Pedroso Miguel, M., Lacerda Ramos, C., Ferreira Duarte, W., Efraim, P., Freitas Schwan, R., 2016. Influence of cocoa hybrids on volatile compounds of fermented beans, microbial diversity during fermentation and sensory characteristics and acceptance of chocolates: cocoa hybrids influenced chocolate quality. *J. Food Qual.* 39, 839–849.

Menezes, A.G.T., Batista, N.N., Ramos, C.L., Silva, A.R. de A. e, Efraim, P., Pinheiro, A.C.M., Schwan, R.F., 2016. Investigation of chocolate produced from

four different Brazilian varieties of cocoa (*Theobroma cacao* L.) inoculated with *Saccharomyces cerevisiae*. *Food Res. Int.* 81, 83–90.

Moreno-Zambrano, M., Grimbs, S., Ullrich, M.S., Hütt, M.T., 2018. A mathematical model of cocoa bean fermentation. *R. Soc. Open Sci.* 5, 180964.

Mota-Gutierrez, J., Botta, C., Ferrocino, I., Giordano, M., Bertolino, M., Dolci, P., Cannoni, M., Cocolin, L., 2018. Dynamics and biodiversity of bacterial and yeast communities during fermentation of cocoa beans. *Appl. Environ. Microbiol.* 84.

Nielsen, D.S., Arneborg, N., Jespersen, L., 2014. Mixed microbial fermentations and methodologies for their investigation. In: Schwan, R.F., Fleet, G. (Eds.), *Cocoa and Coffee Fermentations*. CRC Press/Taylor & Francis, pp. 1–42.

Nielsen, D.S., Teniola, O.D., Ban-Koffi, L., Owusu, M., Andersson, T.S., Holzapfel, W.H., 2007. The microbiology of Ghanaian cocoa fermentations analysed using culture-dependent and culture-independent methods. *Int. J. Food Microbiol.* 114, 168–186.

Nwakuba, N.R., Ejeku, P.K., Okafor, V.C., 2017. A mathematical model for predicting the drying rate of cocoa bean (*Theobroma cacao* L.) in a hot air dryer. *Agric. Eng. Int. CIGR J* 19 (3), 195–202. https://www.researchgate.net/publication/320336317_A_mathematical_model_for_predicting_the_drying_rate_of_cocoa_bean_Theobroma_cacao_L_in_a_hot_air_dryer.

Olabinjo, O.O., Olajide, J.O., Olalusi, A.P., 2017. Mathematical modeling of sun and solar drying kinetics of fermented cocoa beans. *Int. J. Environ. Agric. Biotechnol.* 2, 2419–2426.

Ouattara, H.G., Koffi, B.L., Karou, G.T., Sangaré, A., Niamke, S.L., Diopoh, J.K., 2008. Implication of *Bacillus* sp. in the production of pectinolytic enzymes during cocoa fermentation. *World J. Microbiol. Biotechnol.* 24, 1753–1760.

Ouattara, H.G., Reverchon, S., Niamke, S.L., Nasser, W., 2017. Regulation of the synthesis of pulp degrading enzymes in *Bacillus* isolated from cocoa fermentation. *Food Microbiol.* 63, 255–262.

Owens, J.D., Allagheny, N., Kipping, G., Ames, J.M., 1997. Formation of volatile compounds during *Bacillus subtilis* fermentation of soya beans. *J. Sci. Food Agric.* 74, 132–140.

Papalexandratou, Z., Falony, G., Romanens, E., Jimenez, J.C., Amores, F., Daniel, H.-M., De Vuyst, L., 2011. Species diversity, community dynamics, and metabolite kinetics of the microbiota associated with traditional Ecuadorian spontaneous cocoa bean fermentations. *Appl. Environ. Microbiol.* 77, 7698–7714.

Papalexandratou, Z., Lefeber, T., Bahrim, B., Lee, O.S., Daniel, H.-M., De Vuyst, L., 2013. *Hanseniaspora opuntiae*, *Saccharomyces cerevisiae*, *Lactobacillus fermentum*, and *Acetobacter pasteurianus* predominate during well-performed Malaysian cocoa bean box fermentations, underlining the importance of these microbial species for a successful cocoa bean fermentation process. *Food Microbiol.* 35, 73–85.

Papalexandratou, Z., Nielsen, D.S., 2016. It's gettin' hot in here: breeding robust yeast starter cultures for cocoa fermentation. *Trends Microbiol.* 24, 166–168.

Pedan, V., Fischer, N., Bernath, K., Hühn, T., Rohn, S., 2017. Determination of oligomeric proanthocyanidins and their antioxidant capacity from different chocolate manufacturing stages using the NP-HPLC-online-DPPH methodology. *Food Chem.* 214, 523–532.

Pereira, G.V.M., Alvarez, J.P., Neto, D.P. de C., Soccol, V.T., Tanobe, V.O.A., Rogez, H., Góes-Neto, A., Soccol, C.R., 2017. Great intraspecies diversity of *Pichia kudriavzevii* in cocoa fermentation highlights the importance of yeast strain selection for flavor modulation of cocoa beans. *LWT - Food Sci. Technol.* 84, 290–297.

Pereira, G.V. d. M., Miguel, M.G. d. C.P., Ramos, C.L., Schwan, R.F., 2012. Microbiological and physicochemical characterization of small-scale cocoa fermentations and screening of yeast and bacterial strains to develop a defined starter culture. *Appl. Environ. Microbiol.* 78, 5395–5405.

Pereira, G.V. de M., Soccol, V.T., Soccol, C.R., 2016. Current state of research on cocoa and coffee fermentations. *Curr. Opin. Food Sci.* 7, 50–57.

Pinheiro, Á.D.T., da Silva Pereira, A., Barros, E.M., Antonini, S.R.C., Cartaxo, S.J.M., Rocha, M.V.P., Goncalves, L.R.B., 2017. Mathematical modeling of the ethanol fermentation of cashew apple juice by a flocculent yeast: the effect of initial substrate concentration and temperature. *Bioproc. Biosyst. Eng.* 40, 1221–1235.

Qin, X.-W., Lai, J.-X., Tan, L.-H., Hao, C.-Y., Li, F.-P., He, S.-Z., Song, Y.-H., 2016. Characterization of volatile compounds in Criollo, Forastero, and Trinitario cocoa seeds (*Theobroma cacao* L.) in China. *Int. J. Food Prop.* 1–15.

Ramos, C.L., Dias, D.R., Miguel, M.G. da C.P., Schwan, R.F., 2014. Impact of different cocoa hybrids (*Theobroma cacao* L.) and *S. cerevisiae* UFLA CA11 inoculation on microbial communities and volatile compounds of cocoa fermentation. *Food Res. Int.* 64, 908–918.

Rasmuson, A., 2014. *Mathematical Modeling in Chemical Engineering*. Cambridge University Press, United Kingdom ; New York.

- Ribas-García, M., Hurtado-Vargas, R., Garrido-Carralero, N., Domenech-López, F., Sabadí-Díaz, R., 2011. Metodología para la modelación matemática de procesos. Caso de estudio, fermentación alcohólica. ICIDCA Sobre Los Deriv. Caña Azúcar 45.
- Rodriguez-Campos, J., Escalona-Buendía, H.B., Contreras-Ramos, S.M., Orozco-Avila, I., Jaramillo-Flores, E., Lugo-Cervantes, E., 2012. Effect of fermentation time and drying temperature on volatile compounds in cocoa. Food Chem. 132, 277–288.
- Rodriguez-Campos, J., Escalona-Buendía, H.B., Orozco-Avila, I., Lugo-Cervantes, E., Jaramillo-Flores, M.E., 2011. Dynamics of volatile and non-volatile compounds in cocoa (*Theobroma cacao* L.) during fermentation and drying processes using principal components analysis. Food Res. Int. 44, 250–258.
- Romanens, E., Näf, R., Lobmaier, T., Pedan, V., Leischtfeld, S.F., Meile, L., Schwenninger, S.M., 2018. A lab-scale model system for cocoa bean fermentation. Appl. Microbiol. Biotechnol. 102, 3349–3362.
- Saltini, R., Akkerman, R., Frosch, S., 2013. Optimizing chocolate production through traceability: a review of the influence of farming practices on cocoa bean quality. Food Control 29, 167–187.
- Samagaci, L., Ouattara, H., Niamké, S., Lemaire, M., 2016. *Pichia kudrazevii* and *Candida nitrativorans* are the most well-adapted and relevant yeast species fermenting cocoa in Agneby-Tiassa, a local Ivorian cocoa producing region. Food Res. Int. 89, 773–780.
- Sandhya, M.V.S., Yallappa, B.S., Varadaraj, M.C., Puranaik, J., Rao, L.J., Janardhan, P., Murthy, P.S., 2016. Inoculum of the starter consortia and interactive metabolic process in enhancing quality of cocoa bean (*Theobroma cacao*) fermentation. LWT - Food Sci. Technol. 65, 731–738.
- Sánchez Blanco, A., Palacios Durive, O., Batista Pérez, S., Díaz Montes, Z., Pérez Guerra, N., 2016. Simultaneous production of amylases and proteases by *Bacillus subtilis* in brewery wastes. Braz. J. Microbiol. 47, 665–674.
- Saputro, A.D., Van de Walle, D., Hinneh, M., Van Durme, J., Dewettinck, K., 2018. Aroma profile and appearance of dark chocolate formulated with palm sugar–sucrose blends. Eur. Food Res. Technol. 244, 1281–1292.
- Schwan, R.F., 1998. Cocoa fermentations conducted with a defined microbial cocktail inoculum. Appl. Environ. Microbiol. 64, 1477–1483. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC106173/>.

- Schwan, R.F., Wheals, A.E., 2004. The microbiology of cocoa fermentation and its role in chocolate quality. *Crit. Rev. Food Sci. Nutr.* 44, 205–221.
- Selamat, J., Harun, S.M., others, 1994. Formation of methyl pyrazine during cocoa bean fermentation. *Pertanika J. Trop. Agric. Sci.* 17, 27–32. <http://www.pertanika.upm.edu.my/Pertanika%20PAPERS>.
- Sukha, D.A., Bharath, S.M., Ali, N.A., Umaharan, P., 2013. An assessment of the quality attributes of the Imperial College Selections (ICS) cacao (*Theobroma cacao* L.) clones. In: III International Conference on Postharvest and Quality Management of Horticultural Products of Interest for Tropical Regions, pp. 237–243.
- Sukha, D.A., Umaharan, P., Butler, D.R., 2017. The impact of pollen donor on flavor in cocoa. *J. Am. Soc. Hortic. Sci.* 142, 13–19.
- Teh, Q.T.M., Tan, G.L.Y., Loo, S.M., Azhar, F.Z., Menon, A.S., Hii, C.L., 2016. The drying kinetics and polyphenol degradation of cocoa beans: cocoa drying and polyphenol degradation. *J. Food Process. Eng.* 39, 484–491.
- Torres-Moreno, M., Tarrega, A., Costell, E., Blanch, C., 2012. Dark chocolate acceptability: influence of cocoa origin and processing conditions. *J. Sci. Food Agric.* 92, 404–411.
- Tran, P.D., Van de Walle, D., De Clercq, N., De Winne, A., Kadow, D., Lieberei, R., Messens, K., Tran, D.N., Dewettinck, K., Van Durme, J., 2015. Assessing cocoa aroma quality by multiple analytical approaches. *Food Res. Int.* 77, 657–669.
- Van Durme, J., Ingels, I., De Winne, A., 2016. Inline roasting hyphenated with gas chromatography–mass spectrometry as an innovative approach for assessment of cocoa fermentation quality and aroma formation potential. *Food Chem.* 205, 66–72.
- Vargas Jentzsch, P., Ciobotă, V., Salinas, W., Kampe, B., Aponte, P.M., Rösch, P., Popp, J., Ramos, L.A., 2016. Distinction of Ecuadorian varieties of fermented cocoa beans using Raman spectroscopy. *Food Chem.* 211, 274–280.
- Vázquez-Ovando, A., Ovando-Medina, I., Adriano-Anaya, L., Betancur-Ancona, D., Salvador-Figueroa, M., 2016. Alcaloides y polifenoles del cacao, mecanismos que regulan su biosíntesis y sus implicaciones en el sabor y aroma. *Arch. Latinoam. Nutr.* 66, 239–254. <https://medes.com/publication/117717>.
- Visintin, S., Ramos, C.L., Batista, N.N., Dolci, P., Schwan, R.F., Cocolin, L., 2017. Impact of *Saccharomyces cerevisiae* and *Torulaspota delbrueckii* starter cultures on cocoa beans fermentation. *Int. J. Food Microbiol.* 257, 31–40.

Visintin, S., Alessandria, V., Valente, A., Dolci, P., Cocolin, L., 2016. Molecular identification and physiological characterization of yeasts, lactic acid bacteria and acetic acid bacteria isolated from heap and box cocoa bean fermentations in West Africa. *Int. J. Food Microbiol.* 216, 69–78.

Voigt, J., 2009. Origin of the chocolate-specific flavour notes: essential precursors are generated by proteolysis of a cocoa storage protein. *Agro Food Ind. Hi-Tech* 20 (6), 26–28. https://www.researchgate.net/publication/286122942_Origin_of_the_chocolatespecific_flavour_notes_Essential_precursors_are_generated_by_proteolysis_of_a_cocoa_storage_protein.

Voigt, J., Biehl, B., Heinrichs, H., Kamaruddin, S., Marsoner, G.G., Hugi, A., 1994. In-vitro formation of cocoa-specific aroma precursors: aroma-related peptides generated from cocoa-seed protein by co-operation of an aspartic endoprotease and a carboxypeptidase. *Food Chem.* 49, 173–180.

Voigt, J., Janek, K., Textoris-Taube, K., Niewianda, A., Wöstemeyer, J., 2016. Partial purification and characterisation of the peptide precursors of the cocoa-specific aroma components. *Food Chem.* 192, 706–713.

Voigt, J., Lieberei, R., 2014. Biochemistry of cocoa fermentation. In: *Cocoa and Coffee Fermentations*. CRC Press/Taylor & Francis, pp. 193–206.

Ziegleder, G., 1990. Linalool contents as characteristic of some flavor grade cocoas. *Z. Für Leb. -Forsch. A* 191, 306–309. <https://link.springer.com/article/10.1007/BF01202432>.