



Changes in the expression of interleukin-10 in myocardial infarction and its relationship with macrophage activation and cell apoptosis

Wenqi Yang¹, Shuming Li², Yang Zhao³, Ying Sun², Yuling Huang⁴, Xinying Cao⁴, Cainai Xing⁴, Fang Yang², Wenbo Liu⁴, Xuan Zhao⁴, Xuejing Dong⁴, Hua Bai⁴, Xiaoming Shang¹

¹Department of Internal Medicine, Hebei Medical University, Shijiazhuang 050017, China; ²School of Basic Medical Sciences, North China University of Science and Technology, Tangshan 065299, China; ³Department of Medical imageology, Hebei North University, Zhangjiakou 075132, China; ⁴Department of Cardiology, North China University of Science and Technology Affiliated Hospital, Tangshan 065299, China

Contributions: (I) Conception and design: W Yang, X Shang, F Yang; (II) Administrative support: X Shang; F Yang; (III) Provision of study materials: S Li, Y Zhao, Y Sun; (IV) Collection and assembly of data: W Yang, S Li, Y Zhao, Y Huang, X Cao, C Xing, W Liu, X Zhao, X Dong, H Bai; (V) Data analysis and interpretation: W Yang, S Li, F Yang; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Xiaoming Shang. Department of Internal Medicine, Hebei Medical University, 361 Zhongshan East Road, Shijiazhuang 050017, China. Email: yangxiaots@163.com.

Background: Currently, the role of interleukin-10 (IL-10) as an anti-inflammatory factor in the occurrence and development of heart disease is still unclear. This study aimed to observe the dynamic changes in the expression of IL-10 in serum and myocardial tissues and to investigate the relationship of IL-10 expression with macrophage activation and cardiomyocyte apoptosis during the occurrence of myocardial infarction (MI).

Methods: Mice models with MI were prepared by ligating the anterior descending branch of the coronary artery. The animals were classified into the sham operation group (the control group), and the day 1, 7, 14, and 28 MI groups based.

Results: On days 7 and 14, the cells with positive IL-10 expression were largely distributed in the infarct areas, while cells with positive IL-10 expression were decreased on day 28. Serum IL-10 was significantly positively correlated with IL-10 protein expression in myocardial tissues. Moreover, Bcl-2 and Bax protein expression in myocardial tissues, along with the ratio of Bcl-2/Bax proteins, were gradually elevated with prolonged time of infarction. The expression of arginase protein increased gradually too. There were positive correlations between IL-10 and arginase expressions, and between the expressions of Bcl-2 and Bax proteins.

Conclusions: After the occurrence of MI, the expression of IL-10 first increased and then decreased in serum and myocardial tissues, with this likely affecting macrophage activation, phenotypic transformation, and the occurrence of cardiomyocyte apoptosis.

Keywords: Myocardial infarction (MI); interleukin-10 (IL-10); macrophage; cell apoptosis

Submitted Apr 13, 2020. Accepted for publication Apr 26, 2020.

doi: 10.21037/atm-20-3349

View this article at: <http://dx.doi.org/10.21037/atm-20-3349>

Introduction

Coronary atherosclerosis, also known as coronary heart disease, is caused by myocardial ischemia, hypoxia and necrosis due to stenosis or complete occlusion of the vascular lumen secondary to the atherosclerosis of the coronary artery. Myocardial infarction (MI) is the

most common type of heart disease (1,2). A widespread hypothesis among scholars is that immune and inflammatory responses are important causes for the onset of coronary heart diseases. Several types of inflammatory cells secrete and release various proinflammatory factors at the site of inflammation, and these are involved in the pathogenesis

of coronary heart disease (2,3). Myocardial cells undergo apoptosis in addition to ischemia and necrosis in patients with coronary heart diseases (4,5). Currently, numerous apoptotic proteins are being studied, of which the B cell lymphoma-2 (Bcl-2) protein family, being the most classical apoptotic protein, is the most commonly investigated (5,6).

Recent studies have revealed interleukin-10 (IL-10) to be an anti-inflammatory factor that contributes prominently to the stability maintenance of coronary atherosclerotic plaque, tissue repair, etc., and also plays a protective role in the occurrence and development of coronary heart disease (7-9). Most inflammatory cells (including macrophages) can synthesize and secrete IL-10 (9). Nevertheless, it is currently unclear whether IL-10 expression is internally correlated with macrophage activation and cardiomyocyte apoptosis in coronary heart disease (including acute MI).

We present the following article in accordance with the ARRIVE reporting checklist (available at <http://dx.doi.org/10.21037/atm-20-3349>).

Methods

Animals and experiment design

Studies were performed with 16-to-20-week-old, male C57BL/6 mice (purchased from the Medical Experimental Center of North China University of Science and Technology) fed in a temperature (23–25 °C) and humidity (40–60%)-controlled room, and kept in a 12-h light–dark cycle with access to normal chow diet and water. All animal protocols were approved by the Institutional Review and Ethics Board of North China University of Science and Technology (LX2019035). The total of 25 mice were divided into 5 groups in the experiment at random, which included the control group (C) (n=5), the group of mice with MI for 1 day (1 d) (n=5), the group of mice with MI for 7 days (7 d) (n=5), the group of mice with MI for 14 days (14 d) (n=5) and the group of mice with MI for 28 days (28 d) (n=5).

The mice were anesthetized by intraperitoneal injection of 1% pentobarbital sodium, followed by tracheal intubation, and thus a ventilator was connected. A 2 cm incision was made at the left sternal margin to expose the heart between the third and fourth rib. The anterior descending branch of the left coronary artery was ligated to prepare the MI model while the sham group underwent the same surgical procedure except with the exclusion of coronary artery ligation. When the color of left ventricular changed from red to white, we could judge that the animal model is successful.

A total of 4 mice died (2 mice due to pneumothorax and 2 mice due to hemorrhage in the thoracic cavity) during the animal experiment, and 4 mice respectively died at 3, 11, 24, 70 hours (due to hemothorax) after the operation. The survival rate of mice models was 81%.

Material collection

After the operation, all mice in model group presented distress, ruffled fur, contracture, reduced mobility and eating. Based on these behaviors, at the specified time points, the mice were anesthetized by 1% pentobarbital sodium (50 mg/kg, i.p.) and euthanized by blood collection from the eyeball. Blood was collected and stored in a refrigerator at 4 °C for 2 h. The blood was centrifuged to collect the supernatant and then stored in a refrigerator at –80 °C for later use. Exposed chest cavity, then, the heart was extracted and the left ventricle was cut into 2 parts longitudinally. One part was fixed using 4% paraformaldehyde, while the other part was stored in a refrigerator at –80 °C.

Immunohistochemistry

The heart tissues of the mice were fixed with 4% paraformaldehyde for 24 h. The tissues were then soaked and embedded in paraffin after gradient dehydration using ethanol and de-ethanolization using xylene. The paraffin tissues were cut into 4 μm slices. Pathological changes were observed by routine hematoxylin and eosin (HE) staining, and collagen deposition was observed by Masson staining. Distributions of IL-10 and arginase in myocardial tissues were observed by immunohistochemical staining. Cell apoptosis was observed under a fluorescence microscope after terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) staining (Millipore, USA).

Serum IL-10 determination

The expression of IL-10 in venous blood of mice was detected by enzyme-linked immunosorbent assay (ELISA) kit. Serum samples were allowed to clot 2 hours at 4 °C before centrifugation at 3,000 ×g, 4 °C for 15 min. The supernatant was harvested immediately and stored at –80 °C for later analysis. Then, the IL-10 (E-10155); (Andy Gene Biotechnology Co., Ltd., 301 University Village Drive, Richardson, TX75081, USA) concentration in the serum were detected with the ELISA kits according to the manufacturer's instructions. Peptide concentration was

expressed in pg/mL.

Western blot

A total of 50 mg of myocardial tissues were weighed, and 1 mL of RIPA lysate containing protease inhibitor was added. After that, centrifugation was performed to extract the supernatant protein. The protein standard curve was plotted by Bradford method. The required 15 µg of protein that needed to be tested was loaded, and analysis of electrophoresis and electrolytic membrane was done then performed. Primary antibodies (IL-10, tubulin, MCP-1, arginase, iNOS, Bcl-2 and Bax; dilution performed at 1:1,000) and secondary antibodies (HRP-labeled goat anti-rabbit or goat anti-mouse IgG; dilution performed at 1:5,000) were added. Color development was conducted by electrochemiluminescence (ECL) reagent. The bands were analyzed using Image J image processing software. The ratio of optical density of the target protein to the optical density of internal (tubulin) protein was used as the relative expression of the protein.

Statistical analysis

SPSS 20.0 software was used for statistical analysis. Normality was tested for quantitative data, and the data that met normal distribution were represented by mean ± standard deviation. The mean values of multiple groups were compared using one-way analysis of variance (ANOVA), while pair-wise comparison was performed using the least significance difference (LSD) test. A P value <0.05 was considered to be statistically significant.

Results

Pathomorphological changes in mice with MI

On day 1 of MI, severe degeneration and necrosis in the in the infarct areas of the left ventricle, accompanied by inflammatory cell infiltration were observed in the mice. On day 7 of MI, there were more inflammatory cell infiltrates in addition to degenerated and necrotic cardiomyocytes, the hyperplastic collagen fibers (blue in Masson staining) were extended until the infarcted myocardial tissues, and the infarcted myocardial tissues were separated. On day 14 of MI, there were still degenerated and necrotic cardiomyocytes, accompanied by more inflammatory cell infiltrates. Moreover, collagen hyperplasia was more significant, separating the infarcted areas and the myocardial

tissues in the adjacent areas into “islands”. On day 28 of MI, the inflammatory cell infiltrates had reduced when compared to their extent on day 14, and the tissues in the infarct areas were replaced by hyperplastic collagen fibers, while a small amount of residual cardiomyocytes were observed during that period (the results of Masson staining and HE staining were omitted from *Figure 1*).

Distribution and expression of IL-10 in the heart tissues of mice

Immunohistochemical staining demonstrated the existence of a few IL-10-positive expression cells in the left ventricle in the control group. There was a small amount of cells with positive IL-10 expression in the infarcted areas on day 1 of MI, a large number of cells with positive IL-10 expression on day 7 of MI, a large number of cells with positive IL-10 expression remaining on day 14, but a reduced number of cells with positive IL-10 expression on day 28. IL-10 was mainly expressed in the inflammatory cells in the lesion areas (*Figure 2*).

Serum IL-10 expression

The results of ELISA demonstrated that the serum IL-10 expression was significantly increased when compared to the control group on days 1, 7, and 14 of MI, which were 1.19 times, 1.22 times and 1.18 times that of the control group, respectively, with these differences being statistically significant (P<0.05). However, on day 28 of infarction, the expression was decreased to a level approaching that of the control group (*Figure 3*).

IL-10 expression in myocardial tissues

The results of western blotting revealed that the expression of IL-10 protein in myocardial tissues was increased to a certain degree when compared to the control group on day 1 of MI, and was significantly increased on day 7 of MI, and was 1.43 times that of the control group (P<0.05). Moreover, IL-10 expression peaked on day 14 of MI, and was 1.7 times that of the control group. However, on day 28 of MI, the expression of IL-10 protein was decreased significantly when compared to that on day 14, and was 74.48% of that found on day 14 (P<0.05, *Figure 4*).

Correlation analysis of serum IL-10 and IL-10 expression in myocardial tissues showed that r was 0.543 and P was




Figure 1 Pathological changes in mice with MI (Masson staining). Note C: the control group; 1 d, the group of myocardial infarction for day 1; 7 d, the group of myocardial infarction for day 7; 14 d, the group of myocardial infarction for day 14; 28 d, the group of myocardial infarction for day 28. MI, myocardial infarction.




Figure 2 Changes of myocardial protein expression of MCP-1, iNOS and Arginase in mice with MI (western blot). Note C, the control group; 1 d, the group of myocardial infarction for day 1; 7 d, the group of myocardial infarction for day 7; 14 d, the group of myocardial infarction for day 14; 28 d, the group of myocardial infarction for day 28. IL-10, interleukin-10; MI, myocardial infarction.




Figure 3 Changes of expression of IL-10 in serum of mice with MI across different times ($\bar{x} \pm s$, pg/mL, ELISA assay), *, $P < 0.05$ vs. C; #, $P < 0.05$ vs. 14 d. IL-10, interleukin-10; ELISA, enzyme-linked immunosorbent assay. MI, myocardial infarction.




Figure 4 Changes of myocardial IL-10 protein expression in mice with MI (western blot). *, $P < 0.05$ vs. C; #, $P < 0.05$ vs. 14 d. IL-10, interleukin-10; MI, myocardial infarction.

0.005, demonstrating that the 2 were positively correlated.

Changes in the expressions of IL-10 messenger ribonucleic acid (mRNA) in mice with MI

Moreover, IL-10 mRNA expressions were detected via real time polymerase chain reaction (PCR) in mice with MI. The results of real time PCR revealed that the expression of IL-10 mRNA in myocardial tissues was increased to a certain degree when compared to the control group on day 1 of MI, and was significantly increased on day 7




Figure 5 Changes of expression of IL-10 mRNA in serum of mice with MI across different times ($\bar{x} \pm s$, pg/mL, real time PCR). *, $P < 0.05$ vs. C; #, $P < 0.05$ vs. 14 d. IL-10, interleukin-10; MI, myocardial infarction.

of MI. Moreover, IL-10 expression peaked on day 14 of MI. However, on day 28 of MI, the expression of IL-10 protein was decreased significantly when compared to that on day 14, ($P < 0.05$, Figure 5). The trend of IL-10 mRNA expressions were the same as the protein of the IL-10.

Distribution and changes of M2 macrophages in the infarcted areas of mice with MI

Immunohistochemical staining results showed that there were no cells with a significantly positive expression of arginase (i.e., M2 macrophages) distributed in the myocardial tissues in the control group. There were some cells with positive expression of arginase occasionally visible in the infarcted areas on day 1 of MI, there were some cells with positive expression of arginase on day 7, there were more cells with positive expression of arginase on day 14 when compared to day 7, and there were still many cells with positive expression of arginase on day 28 of MI (Figure 2).

Changes in the expressions of MCP-1, iNOS, and arginase in mice with MI

The results of western blotting revealed that there was an increasing trend first and then a decreasing expression of MCP-1 and iNOS proteins in the myocardial tissues. Compared to the control group, the expressions of MCP-1 protein were 1.08 times, 2.73 times, 2.29 times, and 1.39 times that of the control group on days 1, 7, 14, and 28 of MI, respectively, while the expressions of iNOS




Figure 6 Changes of protein expression of MCP-1, iNOS, and arginase in myocardial tissue of mice with MI (western blot). *, $P < 0.05$ vs. C; #, $P < 0.05$ vs. 14 d. MI, myocardial infarction.

protein were 1.5 times, 4.48 times, 3.28 times, and 1.43 times higher, respectively. However, there was a gradual increasing trend in the arginase protein expression. Compared to the control group, the expression of arginase protein was 1.54 times, 2.42 times, 3.78 times, and 4.12 times that of the control group on days 1, 7, 14, and 28 of MI, respectively (Figure 6), with these differences being statistically significant ($P < 0.05$).

Cardiomyocyte apoptosis of mice with MI

The results of TUNEL staining showed that individual apoptotic cells were occasionally visible in the myocardial tissues in the control group. On day 1 of MI, the number of apoptotic cells in the infarcted areas was increased to some extent, and the number of apoptotic cells was increased gradually with the prolonged time of infarction (Figure 7).

Changes in the expressions of Bax and Bcl-2 proteins in the myocardial tissues of mice with MI

Western blotting revealed the expressions of Bcl-2 and Bax proteins, along with the ratio of Bcl-2/Bax proteins, were gradually increased with prolonged time of infarction. Compared with the control group, the expressions of Bax protein on days 1, 7, 14, and 28 of MI were 1.19 times, 1.50 times, 1.70 times, and 2.23 times those of the control group, while the expressions of Bcl-2 protein were 1.44 times, 3.08 times, 4.54 times, and 6.04 times those of the control

group (Figure 8). The ratios of Bcl-2/Bax were 1.22 times, 2.03 times, 2.67 times, and 2.74 times those of the control group (Figure 9) with these differences being statistically significant ($P < 0.05$). Tubulin is an internal control protein. As fold as Tubulin means the ratio of the one protein to Tubulin

Correlation analysis of various proteins in myocardial tissues

Correlation analysis suggested a positive correlation among IL-10, Bcl-2, Bax, and arginase protein expressions (Table 1).

Discussion

The inflammatory responses after MI act as an important pathological process in MI. A large number of inflammatory factors with different functions are significantly involved in the inflammatory responses (2,10). IL-10 is an important anti-inflammatory factor, and plays a strong anti-inflammatory role. Studies have demonstrated that IL-10 is closely related to the occurrence and prognosis of cardiovascular diseases, and has an important regulatory effect on the remodeling of tissue functions and structures after ischemic injury (8,9,11). Recently, it has been reported that the target cells in which IL-10 plays anti-inflammatory effects are mainly macrophages. Mononuclear macrophages secrete IL-10 after activation by various endogenous and exogenous mediators, causing IL-10 genetic transcription




Figure 7 Morphology changes of apoptosis in mice with MI (Tunel staining). Note C: the control group; 1 d, the group of myocardial infarction for 1 day; 7 d, the group of myocardial infarction for 7 days; 14 d, the group of myocardial infarction for 14 days; 28 d, the group of myocardial infarction for 28 days. MI, myocardial infarction.

via lipopolysaccharide activation pathway and protein kinase A-dependent activation pathway, among others (12,13). Additionally, mononuclear macrophages also secrete IL-10 during the process of clearing apoptotic cells, and this process depends on the activation of p38 mitogen-activated protein kinase (MAPK) (14). It has been reported that the expression of IL-10 is increased during ischemia, and acts as a protective response to ischemia (15,16). Some researchers have found that serum IL-10 is significantly increased on the day of acute MI. Moreover, IL-10 remains at a higher level on day 1 after percutaneous coronary intervention

(PCI) therapy, but it decreases to some extent on day 7 after therapy, suggesting that IL-10 is associated with the time and severity of myocardial ischemia to some extent (17). Our study results showed that serum IL-10 expression was increased significantly on day 1 of MI in mice, and remained at a higher level on days 7 and 14 of MI, but was decreased after that, continuing to decline to a level comparable to that of the control group on day 28. There was a small number of cells with positive IL-10 expression in the infarcted areas on day 1 of MI. Moreover, the majority of the cells with positive IL-10 expression were visible on days 7 and 14




Figure 8 Changes of myocardial protein expression of Bcl-2 and Bax in mice with MI (western blot). MI, myocardial infarction.




Figure 9 The ratio changes of Bcl-2 to Bax in myocardial protein expression of mice with MI (western blot). MI, myocardial infarction.

of MI, while the cells with positive IL-10 expression were decreased on day 28. However, the expression of myocardial IL-10 protein was significantly elevated on day 7 of MI, and peaked on day 14, while it decreased significantly on day 28. Correlation analysis showed that the expression of serum IL-10 was positively correlated with the expression of IL-10 protein in myocardial tissues. These results suggested an increasing trend initially and then a decrease in IL-10 expression (regardless of in serum or myocardial

tissues) with the prolonged time of infarction in the mice models with MI. On day 1, MI was considered as the initial stage of acute inflammation, thus a greater necrosis of cardiomyocytes, accompanied by exudation and infiltrates of inflammatory cells, was observed. However, day 7 of MI was the critical stage of acute inflammation, in which a large amount of exudation and infiltrates of inflammatory cells, accompanied by the release of a large number of proinflammatory factors were observed. In response to these, the body releases anti-inflammatory factors (such as IL-10) to maintain a balance in the biological functions (11,18). Therefore, at around days 1 and 7 of MI, IL-10 was highly expressed in serum and/or myocardial tissues, and this continued until week 2. After that, tissue injury entered the repair stage, with inflammation also degrading gradually, and there was more extracellular matrix (including collagen) to replace the infarcted myocardial tissues gradually (3,19). At that time, synthesis and secretion of IL-10 gradually decreased, suggesting that the dynamic changes of IL-10 in serum and myocardial tissues during the process of MI were closely associated with the occurrence and development of inflammation, as well as the outcomes of MI. These might be caused by the defense and anti-inflammatory effects of the body to inflammatory responses during MI.

In inflammatory responses, macrophages are activated and phenotypic transformation occurs with the progression

Table 1 Correlation analysis between several protein expressions in myocardial tissue

Factor	IL-10		Bcl-2		Bax		Bcl-2/Bax		Arginase	
	r	P	r	P	r	P	r	P	r	P
IL-10	–		0.690	0.004	0.546	0.035	0.588	0.021	0.669	0.006
Bcl-2	0.690	0.004	–		0.901	0.000	0.951	0.000	0.965	0.000
Bax	0.546	0.035	0.901	0.000	–		0.912	0.000	0.973	0.000
Bcl-2/Bax	0.588	0.021	0.951	0.000	0.912	0.000	–		0.951	0.000
Arginase	0.669	0.006	0.956	0.000	0.973	0.000	0.951	0.000	–	

IL-10, interleukin-10.

of inflammation. In the initial and early stages of inflammation, macrophages act as “proinflammatory” macrophages (M1 phenotype, secreting specific marker proteins such as TNF- α , iNOS, and IL-1 β , etc.), which in turn promote the progression of inflammation. However, in the middle and advanced stages, macrophages act as “repairing” macrophages (M2 phenotype, secreting specific marker proteins such as arginase and CD206), effecting anti-inflammation and tissue repair (19,20). In addition, *in vitro* experiment showed that fibroblasts can be activated and strengthened proliferation and migratory abilities after IL-10 intervention, and there were more collagens that were closely related to M2 phenotypic transformation of macrophages (20). In the present study, degeneration and necrosis of cardiomyocytes were observed on day 1 of MI in mice, the expression of inflammatory factor monocyte chemoattractant protein-1 (MCP-1) began to increase, and there was inflammatory cell infiltrates in the infarcted areas. Results of immunohistochemical staining showed the existence of only a small number of M2 macrophages in these inflammatory cells. Meanwhile, iNOS, which represents the marker protein of MI macrophages, was increased to some extent. On day 7 of MI, inflammatory cell infiltration in the infarcted areas was more significant. Among them, the expression of MCP-1 protein peaked, and iNOS expression achieved the highest value. However, the number of cells (M2 macrophages) with positive arginase expression and the expression of arginase protein were both increased significantly. On day 14 of MI, the iNOS expression showed declination with decreasing expression of MCP-1 protein. Until day 28, the expressions of MCP-1 and iNOS proteins declined back to levels comparable to those in the control group, while the number of cells with positive arginase expression still remained high, and the expression of arginase protein was still elevated. This

suggested that the macrophages were activated and there was a phenotypic transformation from M1 macrophages in the initial stage of inflammation to M2 macrophages in the middle and advanced stages of inflammation during the process of MI. Correlation analysis showed that IL-10 expression in myocardial tissues was positively correlated with the expression of Arginase protein during the occurrence and development of MI. This indicated that the dynamic changes of IL-10 after MI might affect or regulate macrophage activation and promote the transformation of proinflammatory macrophages to repair-type macrophages.

Cardiomyocyte apoptosis also occurs with the degeneration and necrosis of cells during the process of MI. Bax and Bcl-2 are important cell apoptosis-related genes and cell apoptosis regulatory proteins. Bax plays a positive regulatory effect during cell apoptosis, promoting the occurrence of cell apoptosis, promoting the occurrence of cell apoptosis. Bcl-2 plays a negative regulatory role, blocking the apoptotic pathway and inhibiting the occurrence of apoptosis (5,21). However, the ratio of Bcl-2/Bax is closely related to cell apoptosis and survival (6,22). In this study, the results showed that the number of apoptotic cells in myocardial tissues was increased gradually with prolonged time of infarction, and the expression of Bax protein was also increased gradually, indicating that cardiomyocyte apoptosis was gradually aggravated with prolonged time of infarction. With prolonged time of infarction, the expression of Bcl-2 protein was elevated gradually, and the ratio of Bcl-2/Bax was gradually increased in particular. Correlation analysis showed that the expression of IL-10 protein in myocardial tissues was positively correlated in a significant way with the expressions of Bax and Bcl-2 proteins, especially the ratio of Bcl-2/Bax proteins. These dynamic changes suggest that IL-10 expression can affect the occurrence of cardiomyocyte apoptosis during the emergence and development of MI.

Due to the gradual increase in the ratio of Bcl-2 to Bax, the effects of IL-10 might regulate cardiomyocyte apoptosis by controlling the balance between the anti-apoptotic gene (Bcl-2) and the apoptotic gene (Bax), thereby affecting remodeling of functions and structures of myocardial tissues.

In addition, further correlation analysis showed a positive correlation between IL-10 and arginase in myocardial tissues, and between the expressions of Bcl-2 and Bax proteins. This suggests that IL-10, arginase, and apoptosis-regulatory proteins show close associations and play an important role in the remodeling of tissue structures and damage repair processes after MI.

Acknowledgments

The authors would like to thank the collaborators at the School of Public Health, North China University of Science and Technology. A special thanks goes to Miss Sun Ying, head of the laboratory of Yang fang and his team who conducts IL-10 analyses.

Funding: None.

Footnote

Reporting Checklist: The authors have completed the ARRIVE reporting checklist. Available at <http://dx.doi.org/10.21037/apm-20-3349>

Data Sharing Statement: Available at <http://dx.doi.org/10.21037/atm-20-3349>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/atm-20-3349>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was approved by North China University of Science and Technology (2019052).

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with

the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Yuepeng J, Zhao X, Zhao Y, et al. Gene polymorphism associated with TNF- α (G308A) IL-6 (C174G) and susceptibility to coronary atherosclerotic heart disease: A meta-analysis. *Medicine (Baltimore)* 2019;98:e13813.
2. Kirolos I, Yakoub D, Pendola F, et al. Cardiac physiology in post myocardial infarction patients: the effect of cardiac rehabilitation programs—a systematic review and update meta-analysis. *Ann Transl Med* 2019;7:416.
3. Prabhu SD, Frangiannis NG. The Biological Basis for Cardiac Repair After Myocardial Infarction: From Inflammation to Fibrosis. *Circ Res* 2016;119:91-112.
4. Zhao L, Yang XR, Han X. MicroRNA-146b induces the PI3K/Akt/NF- κ B signaling pathway to reduce vascular inflammation and apoptosis in myocardial infarction by targeting PTEN. *Exp Ther Med* 2019;17:1171-81.
5. Lv FH, Yin HL, He YQ, et al. Effects of curcumin on the apoptosis of cardiomyocytes and the expression of NF- κ B, PPAR- γ and Bcl-2 in rats with myocardial infarction injury. *Exp Ther Med* 2016;12:3877-84.
6. Naseri MH, Mahdavi M, Davoodi J, et al. Up regulation of Bax and down regulation of Bcl2 during 3-NC mediated apoptosis in human cancer cells. *Cancer Cell Int* 2015;15:55.
7. Kamaly N, Fredman G, Fojas JJ, et al. Targeted interleukin-10 nanotherapeutics developed with a microfluidic chip enhance resolution of inflammation in advanced atherosclerosis. *ACS Nano* 2016;10:5280-92.
8. Shi J, Li J, Guan H, et al. Anti-fibrotic actions of interleukin-10 against hypertrophic scarring by activation of PI3K/AKT and STAT3 signaling pathways in scar-forming fibroblasts. *PLoS One* 2014;9:e98228.
9. Jung M, Ma YG, Iyer RP, R et al. IL-10 improves cardiac remodeling after myocardial infarction by stimulating M2 macrophage polarization and fibroblast activation. *Basic Res Cardiol* 2017;112:33.
10. Frangiannis NG. The inflammatory response in myocardial injury, repair, and remodelling. *Nat Rev Cardiol* 2014;11:255-65.
11. de Waal Malefyt R, Abrams J, Bennett B, et al. Interleukin 10(IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J*

- Exp Med 1991;174:1209-20.
12. Ernst O, Glucksam-Galnoy Y, Athamna M, et al. The cAMP pathway amplifies early myD88-dependent and type I interferon-independent LPS-induced interleukin-10 expression in mouse macrophages. *Mediators Inflamm* 2019;2019:3451461.
 13. MacKenzie KF, Clark K, Naqvi S, et al. PGE (2) induces macrophage IL-10 production and a regulatory-like phenotype via a protein kinase A-SIK-CRTC3 pathway. *J Immunol* 2013;190:565-77.
 14. Raza A, Crothers JW, McGill MM, et al. Anti-inflammatory roles of p38 α MAPK in macrophages are context dependent and require IL-10. *J Leukoc Biol* 2017;102:1219-27.
 15. Shibata M, Endo S, Inada K, et al. Elevated plasma levels of interleukin-1 receptor antagonist and interleukin-10 in patients with acute myocardial infarction. *J Interferon Cytokine Res* 1997;17:145-50.
 16. Bakhta O, Blanchard S, Guihot AL, et al. Cardioprotective role of colchicine against inflammatory injury in a rat model of acute myocardial infarction. *J Cardiovasc Pharmacol Ther* 2018;23:446-55.
 17. Wang HJ, Zhao XH. Changes and its significance of serum interleukin-6 and interleukin-10 in AMI patients after percutaneous coronary intervention. *Chin J Cardiovasc Rehabil Med* 2012;21:372-4.
 18. Peng Y, Cai X, Zhang G, et al. Molecular characterization and expression of interleukin-10 and interleukin-22 in golden pompano (*Trachinotus ovatus*) in response to *Streptococcus agalactiae* stimulus. *Fish Shellfish Immunol* 2017;65:244-55.
 19. Mills CD. M1 and M2 macrophages: oracles of health and disease. *Crit Rev Immunol* 2012;32:463-88.
 20. Gombozhapova A, Rogovskaya Y, Shurupov V, et al. Macrophage activation and polarization in post-infarction cardiac remodeling. *J Biomed Sci* 2017;24:13.
 21. Uzu M, Sato H, Shimizu A, et al. Connexin 43 enhances Bax activation via JNK activation in sunitinib-induced apoptosis in mesothelioma cells. *J Pharmacol Sci* 2017;134:101-7.
 22. Vucicevic K, Jakovljevic V, Colovic N, et al. Association of Bax expression and Bcl2/Bax ratio with clinical and molecular prognostic markers in chronic lymphocytic leukemia. *J Med Biochem* 2016;35:150-7.

Cite this article as: Yang W, Li S, Zhao Y, Sun Y, Huang Y, Cao X, Xing C, Yang F, Liu W, Zhao X, Dong X, Bai H, Shang X. Changes in the expression of interleukin-10 in myocardial infarction and its relationship with macrophage activation and cell apoptosis. *Ann Transl Med* 2020;8(10):643. doi: 10.21037/atm-20-3349