



Brief Report

Impulsive Personality Traits Predicted Weight Loss in Individuals with Type 2 Diabetes after 3 Years of Lifestyle Interventions

Giulia Testa^{1,2,3,†}, Lucía Camacho-Barcia^{1,2,3,†}, Carlos Gómez-Martínez^{3,4,5}, Bernat Mora-Maltas^{1,2}, Rafael de la Torre^{3,6,7}, Xavier Pintó^{8,9}, Dolores Corella^{3,10}, Roser Granero^{1,3,11}, Aida Cuenca-Royo^{3,6}, Susana Jiménez-Murcia^{1,2,3,9}, Nancy Babio^{3,4,5}, Rebeca Fernández-Carrión^{3,10}, Virginia Esteve-Luque⁸, Laura Forcano^{3,6}, Jiaqi Ni^{4,5}, Mireia Malcampo¹², Sara De las Heras-Delgado^{4,5}, Montse Fitó^{3,12}, Jordi Salas-Salvadó^{3,4,5,*} and Fernando Fernández-Aranda^{1,2,3,9,*}



Citation: Testa, G.; Camacho-Barcia, L.; Gómez-Martínez, C.; Mora-Maltas, B.; de la Torre, R.; Pintó, X.; Corella, D.; Granero, R.; Cuenca-Royo, A.; Jiménez-Murcia, S.; et al. Impulsive Personality Traits Predicted Weight Loss in Individuals with Type 2 Diabetes after 3 Years of Lifestyle Interventions. *J. Clin. Med.* **2022**, *11*, 3476. <https://doi.org/10.3390/jcm11123476>

Academic Editor: Katrien Benhalima

Received: 29 April 2022

Accepted: 14 June 2022

Published: 16 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

- ¹ Psychoneurobiology of Eating and Addictive Behaviors Group, Institut d'Investigació Biomèdica de Bellvitge (IDIBELL), 08907 Barcelona, Spain; gtesta@idibell.cat (G.T.); lcamacho@idibell.cat (L.C.-B.); bernatmoramaltas@gmail.com (B.M.-M.); roser.granero@uab.cat (R.G.); sjimenez@bellvitgehospital.cat (S.J.-M.)
 - ² Department of Psychiatry, University Hospital of Bellvitge, 08907 Barcelona, Spain
 - ³ CIBER Physiology of Obesity and Nutrition (CIBEROBN), Carlos III Health Institute, 28029 Madrid, Spain; carlos.gomez@urv.cat (C.G.-M.); rtorre@imim.es (R.d.l.T.); dolores.corella@uv.es (D.C.); acuenca@imim.es (A.C.-R.); nancy.babio@urv.cat (N.B.); rebeca.fernandez@uv.es (R.F.-C.); lforcano@imim.es (L.F.); mfito@imim.es (M.F.)
 - ⁴ Universitat Rovira i Virgili, Departament de Bioquímica i Biotecnologia, Unitat de Nutrició Humana, 43201 Reus, Spain; jiaqi.ni@alumni.urv.cat (J.N.); sara.delasheras@urv.cat (S.D.I.H.-D.)
 - ⁵ Institut d'Investigació Sanitària Pere Virgili (IISPV), Unitat de Nutrició Humana, 43201 Reus, Spain
 - ⁶ Neurosciences Program, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM), 08003 Barcelona, Spain
 - ⁷ Medicine and Life Sciences Department, Universitat Pompeu Fabra (UPF), 08002 Barcelona, Spain
 - ⁸ Lipids and Vascular Risk Unit, Internal Medicine, University Hospital of Bellvitge-IDIBELL, Hospitalet de Llobregat, 08907 Barcelona, Spain; xpinto@bellvitgehospital.cat (X.P.); virginia.esteveluque@gmail.com (V.E.-L.)
 - ⁹ Department of Clinical Sciences, School of Medicine and Health Sciences, University of Barcelona, Hospitalet de Llobregat, 08907 Barcelona, Spain
 - ¹⁰ Department of Preventive Medicine, University of Valencia, 46010 Valencia, Spain
 - ¹¹ Department of Psychobiology and Methodology, Autonomous University of Barcelona, 08193 Barcelona, Spain
 - ¹² Unit of Cardiovascular Risk and Nutrition, Hospital del Mar Institute for Medical Research (IMIM), 08003 Barcelona, Spain; mireiamalcampo@gmail.com
- * Correspondence: jordi.salas@urv.cat (J.S.-S.); ffernandez@bellvitgehospital.cat (F.F.-A.)
† These authors contributed equally to this work.

Abstract: Impulsivity has been associated with type 2 diabetes (T2D) and may negatively impact its management. This study aimed to investigate impulsive personality traits in an older adult population with T2D and their predicting role in long-term weight control and glycemic management, through glycosylated hemoglobin (HbA_{1c}), following 3 years of intervention with a Mediterranean diet. The Impulsive Behavior Scale (UPPS-P) was administered as a measure of impulsive traits at baseline. Results showed higher total baseline scores of UPPS-P, and higher positive urgency in individuals with T2D, compared with those without T2D. The regression analysis in patients with T2D showed that sensation seeking and lack of perseverance predicted weight loss at follow-up. By contrast, impulsive traits did not predict follow-up levels of HbA_{1c}. In conclusion, the present findings suggest that higher impulsive traits in individuals with T2D seem to affect long-term weight control, but not glycemic control.

Keywords: impulsivity; type 2 diabetes; weight loss; glycosylated hemoglobin

1. Introduction

Type 2 Diabetes (T2D) is a metabolic disorder that represents a major concern in the global healthcare system, as its prevalence has rapidly increased worldwide. This increment is especially observable in older adults, aged 65 or older, who constitute about half of the total population with this condition [1]. Several factors associated with the ageing process increase the susceptibility to T2D, including the reduced β -cell sensitivity to glucose, diminution of skeletal muscle tissue and increment of central adiposity [2]. The presence of T2D increases the risk of several metabolic comorbidities, as well as the development of cognitive decline [3], factors that constitute a challenge in the management of T2D [4].

Impulsive behaviors, defined as the tendencies to act rashly without giving adequate forethought to the consequences of the behaviors, could contribute to diabetes mismanagement in terms of diet, physical activity, medication adherence and blood sugar control. So far, impulsivity has been conceived as a multidimensional construct, which includes personality traits and behavioral and cognitive factors [5]. Obesity has been strongly linked to impulsivity at several levels, and the underlying mechanisms have begun to be elucidated [6–12]. In terms of impulsivity-related cognitive functions, studies in individuals with T2D have shown poor decision-making and reduced inhibitory control [13,14], possibly due to the impact of insulin resistance and impaired glycemic control on the brain dopaminergic pathways [15,16].

Impulsivity has also been conceptualized in terms of more stable traits by multiple models of personality [5]. Previous studies that explored personality traits in individuals with T2D [17–19] mostly adopted the Big Five factors model of personality [20], which is a global measure of personality that might be suboptimal for exploring impulsive traits. A widely accepted instrument used to conceptualize the multivariable structure of impulsivity is the impulsive behavior scale (UPPS) [21], which includes five dimensions related to various impulsive behaviors (i.e., sensation seeking, negative urgency, positive urgency, lack of premeditation and lack of perseverance). To date, no studies have been conducted in patients with T2D using the UPPS model, which could help to better characterize impulsive traits in this population. Considering the evidence for impulsivity-related cognitive dysfunctions shown in the individuals with T2D [13,14], it could also be expected to find further impulsive personality traits in these individuals.

Given that impulsivity leads to a lack of planning and insufficient regulation of behavior, impulsive traits may be expected to predispose to poor weight control and reduced self-management of T2D in the long term. In line with this view, a recent study in patients with T2D showed a negative correlation between impulsive traits and diabetes self-management, and a positive correlation with self-reported glycated hemoglobin (HbA_{1c}) levels, an indicator of glycemic control [22]. In addition, impulsive personality has been shown to indirectly and negatively predict exercise and diet adherence via diabetes management self-efficacy [23]. Nevertheless, the cross-sectional designs of these studies did not provide a conclusion as to whether the impulsive personality could have a long-term impact on diabetes self-management.

Regarding the relationship between BMI and impulsive traits in T2D, a study in a large population of individuals with T2D showed that higher scores in the Barratt Impulsiveness Scale (BIS-11) were related to higher BMI [24]. Still, no studies have explored the relationship between impulsive traits and difficulties in long-term weight control in individuals with T2D. However, in a recent study performed in an older adult population with obesity, individuals with lower impulsive traits achieved higher BMI decreases at one-year follow-ups [25]. Longitudinal studies are needed in patients with T2D to elucidate the impact of impulsive personality traits on long-term weight management.

The present study first aimed to compare impulsive personality traits between older Mediterranean adult individuals with obesity in the presence or absence of T2D, and to explore the relationship between impulsivity and diabetic metabolic indicators among the population with T2D.

A second aim was to identify in patients with T2D whether impulsive personality traits were predictors of weight loss and diabetes control after 3 years of lifestyle interventions. We hypothesized that patients with T2D would be characterized by more impulsive personality traits, according to the UPPS model, and impulsivity was expected to negatively affect weight loss and glycemic control after an intervention with a Mediterranean diet (MedDiet).

2. Materials and Methods

2.1. Study Design and Participants

This longitudinal analysis was performed in the framework of the PREDIMED-Plus-Cognition sub-study ($n = 487$) (see Figure S1. Flowchart for the sampling, Supplementary Materials), a subset sample of the clinical trial PREDIMED-Plus, a large, multicentric, randomized study ($n = 6874$) that aimed to assess the effect of an energy-restricted Mediterranean diet, physical activity promotion and behavioral intervention on the primary prevention of cardiovascular disease. The design, procedure and population was reported previously [26], and its full protocol is available in the RCT website (<https://predimedplus.com>, accessed on 1 April 2022). The trial was registered at the International Standard Randomized Controlled Trial in the year 2014, with the number ISRCTN89898870. Criteria for selecting the subjects were as follows: community-dwelling adults with Metabolic Syndrome (with any 3 of 5 risk factors [27]), aged between 55 and 75 years, with a body mass index (BMI) between 27 and 40 kg/m². Participants with diabetes constituted approximately 25% of the final total PREDIMED-Plus sample. Individuals were recruited from 23 Spanish health centers that participated in the study, between October 2013 and December 2016, and randomly assigned to either the intervention group or the control group. The intervention is based on an energy restricted MedDiet, with behavioral support and physical activity. The individuals randomized into the control group were advised to follow an unrestricted MedDiet without any further indication.

2.2. Procedure

The Cognition sub-study had a specific objective, to perform an in-depth assessment of psychological and cognitive performance in this sub-sample. For this purpose, four of the recruiters' sites: Bellvitge University Hospital (Barcelona); Universitat Rovira i Virgili (Reus); University of Valencia (Valencia); and Hospital del Mar Medical Research Institute (Barcelona); assessed cognitive function and trait impulsivity through a battery of questionnaires and tests at baseline (T0) and at three years of follow-up (T3). During these visits, investigators also collected additional information, such as clinical and anthropometric variables.

2.3. Measures

2.3.1. Anthropometric and Metabolic Measurements

At baseline, and in each yearly follow-up visit thereafter, trained personnel measured different anthropometric outcomes, including weight and height, that were then used to establish the body mass index [BMI = weight(kg)/height(m)²]. The BMI was classified using the World Health Organization (WHO)'s nutritional status ranges [28]. Fasting blood samples were taken at baseline, 1-year and 3-year of follow-up to determine the levels of fasting blood glucose, glycated hemoglobin (HbA_{1c}), insulin and lipid profile. Insulin levels measured at baseline were used to estimate the Homeostasis Model Assessment of Insulin Resistance index (HOMA-IR= fasting glucose levels [mg/dL] × fasting insulin levels [μU/mL]/405.13) [29].

2.3.2. Trait Impulsivity Assessment

This study used the UPPS-P framework [21], assessing five impulsive behavior pathways to measure impulsivity traits: (1) sensation seeking, described as the tendency to seek sensory pleasure and excitement; (2) lack of premeditation, described as the tendency to

act without forethought and planning; (3) lack of perseverance, defined as the tendency to not finish tasks; (4) negative urgency, defined as the tendency to act rashly in negative emotional states; and (5) positive urgency, defined as the tendency to act rashly in positive emotional states. Each item of the UPPS-P is rated on a 4-point Likert scale ranging from 1—“strongly agrees” to 4—“strongly disagree”. Scores for each dimension and a total score of the UPPS-P were calculated, with higher scores indicating higher levels of impulsivity. Previous studies have shown that these traits share between 6% and 27% of their variance, with negative and positive urgency sharing the largest proportion of variance [30]. The measurement of separate aspects of impulsivity can clarify discrete relationships that might be masked or watered-down when such constructs are combined [31]. The Spanish adaptation of this instrument, which was used in the present study, showed adequate psychometric properties [32]. In our sample, internal consistency ranged from adequate ($\alpha = 0.75$ for lack of perseverance) to excellent ($\alpha = 0.92$ for the negative urgency subscale).

2.4. Statistical Analysis

Stata17.0 was used for the statistical analysis [33]. The comparison between the groups (presence/absence of T2D) at baseline (T0) was based on chi-square tests (χ^2) for categorical variables and T-tests for independent samples for continuous variables. Partial correlation coefficients (adjusted by sex, age and education levels) assessed the association at baseline (T0) between the impulsive measures (UPPS-P) and the insulin-related metabolic conditions. Multiple regression was used to assess the predictive capacity of the impulsivity levels (UPPS-P scores) at baseline (T0) on the BMI, weight and HbA_{1c} measures at the 3-year follow-up (T3). These regressions were performed in two blocks: (a) in the first block, the covariates sex, age, education and intervention group were entered and fixed (ENTER method); (b) in the second block, the significant predictors of the criteria were automatically selected by the stepwise method (this selection was used due the large set of potential predictors and the exploratory nature of the analysis) [34]. The final models generated by the regression procedures were next considered adequate if they allowed for reliable clinical interpretation. The ANOVA and regression models were adjusted by sex, age, education and the intervention group. The level of significance was $p < 0.05$.

3. Results

3.1. Description of the Sample

Table 1 contains the descriptive for the total sample at baseline, as well as the comparison of the participants with (T2D+) and without (T2D−) diabetes at baseline. No differences between the groups were found for the sociodemographic characteristics, but the participants in the T2D+ group were older and also had higher levels of HOMA-IR, glucose and HbA_{1c}. Regarding the UPPS-P scores, positive urgency and total UPPS-P score were higher within T2D+ participants.

Table 1. Descriptive of the sample at baseline.

	Total (n = 447)		T2D (–)(n = 308)		T2D(+) (n = 139)		p
	n	%	n	%	n	%	
Sex							
Men	216	48.3%	146	47.4%	70	50.4%	0.563
Women	231	51.7%	162	52.6%	69	49.6%	
Civil status							
Single	17	3.8%	12	3.9%	5	3.6%	0.886
Married	351	78.5%	244	79.2%	107	77.0%	
Divorced-separated	29	6.5%	20	6.5%	9	6.5%	
Widowed	50	11.2%	32	10.4%	18	12.9%	
School							
University (high)	42	9.4%	30	9.7%	12	8.6%	0.985
University (grade)	36	8.1%	25	8.1%	11	7.9%	
Secondary	130	29.1%	89	28.9%	41	29.5%	
Primary	239	53.5%	164	53.2%	75	54.0%	
Employment							
Unemployed	81	18.1%	61	19.8%	20	14.4%	0.307
Work at home	47	10.5%	35	11.4%	12	8.6%	
Retired	287	64.2%	189	61.4%	98	70.5%	
Unemployed	32	7.2%	23	7.5%	9	6.5%	
Group weight							
Overweight	124	27.7%	91	29.5%	33	23.7%	0.145
Obesity I (BMI 30–35)	217	48.5%	150	48.7%	67	48.2%	
Obesity II (BMI 35–40)	102	22.8%	63	20.5%	39	28.1%	
Obesity III (BMI > 40)	4	0.9%	4	1.3%	0	0.0%	
Intervention group							
Control	228	51.0%	161	70.6%	67	29.4%	0.425
Experimental	219	49.0%	147	67.1%	72	32.9%	
	Mean	SD	Mean	SD	Mean	SD	p
Age (years-old)	65.19	4.60	64.83	4.59	65.97	4.53	0.015 *
	Mean	SD	Mean	SD	Mean	SD	p
HOMA-IR	5.43	3.05	4.86	2.51	6.69	3.71	<0.001 *
Glucose	116.60	29.68	103.61	12.95	145.37	35.55	<0.001 *
Glycated hemoglobin (HbA _{1c})	6.15	0.78	5.80	0.39	6.91	0.87	<0.001 *
	Mean	SD	Mean	SD	Mean	SD	p
UPPS-P Lack of premeditation	19.85	5.60	19.69	5.59	20.18	5.62	0.397
UPPS-P Lack of perseverance	18.92	5.01	18.83	5.06	19.14	4.89	0.547
UPPS-P Sensation seeking	20.51	7.14	20.29	7.22	20.99	6.97	0.338
UPPS-P Positive urgency	23.10	8.28	22.52	8.08	24.38	8.60	0.028 *
UPPS-P Negative urgency	24.57	7.35	24.21	7.11	25.38	7.84	0.118
UPPS-P Total score	106.95	22.62	105.55	22.40	110.07	22.86	0.049 *

Note. T2D(–): diabetes absent. T2D(+): diabetes present. SD: standard deviation. BMI: body mass index. HOMA-IR: Homeostasis Model Assessment of Insulin Resistance index. UPPS-P: impulsive behavior scale. * Bold: significant comparison.

3.2. Association of Impulsivity with Markers of Glucose Metabolism

No relevant associations were found between the UPPS-P scores with the HOMA-IR, plasma-fasting glucose or HbA_{1c} levels at baseline. All of the partial correlations obtained non-significant results ($p < 0.05$), and coefficients were within the low-effect size range (in the study, the |R| were between 0.003 and 0.098) (see Table S1, Supplementary Materials).

3.3. Predictive Capacity of the Impulsivity at Baseline on BMI, Weight and HbA_{1c}

Table 2 shows the results for the final regressions obtained in the multiple regressions, using the subsample of participants with T2D. Higher scores in the lack of perseverance subscale at baseline were associated with higher BMI levels at the end of the study, while a higher total score of UPPS-P predicted a higher weight at year 3. Higher scores of sensation

seeking predicted lower decreases in the BMI and the weight. The impulsivity levels at baseline were not associated with the HbA_{1c} levels at 3 years of follow-up (mean = 6.74, SD = 1.00), nor with the decrease in the HbA_{1c} comparing the values at the end of the study with the baseline.

Table 2. Significant predictors at 3 year follow up, considering impulsivity measures at baseline; final models adjusted by sex, age, education and intervention group (results for the subsample of patients with T2D).

Criteria (3 Years)	UPPS-P (at Baseline)	B	SE	Beta	p	95%CI B	
BMI	Lack of perseverance	0.122	0.064	0.160	0.048	0.005	0.250
Changes in BMI	Sensation seeking	−0.067	0.022	−0.259	0.002	−0.110	−0.025
	Total score	0.125	0.048	0.196	0.010	0.030	0.219
Changes in weight	Sensation seeking	−0.218	0.058	−0.305	<0.001	−0.333	−0.103
	No significant predictors						
Changes in Hb1Ac	No significant predictors						

Note. UPPS-P: impulsive behavior scale. B: non-standardized coefficient. SE: error standard. Beta: standardized coefficient. 95%CI: 95% confidence interval. BMI: body mass index. HbA_{1c}: Glycated hemoglobin.

4. Discussion

The present study investigated impulsive personality traits in individuals with T2D as predictors of BMI changes and diabetes control, measured by the levels of glycated hemoglobin after 3 years of lifestyle interventions.

Baseline evaluations of metabolic parameters and personality traits showed differences between individuals with T2D and those without it. As expected, higher levels of glucose, HOMA-IR and HbA_{1c} were characteristic of individuals with T2D. Concerning impulsive personality traits: although the UPPS scores were sub-clinical in both groups, patients with T2D showed significantly higher total scores and positive urgency scores at baseline. Positive urgency is defined as the tendency to act rashly in response to positive affective states [35], which has been strongly associated with emotional factors [32]. Neuroimaging studies have suggested that emotional-based dispositions to act impulsively are linked to orbitofrontal cortex–amygdala connections in the brain, regulated by dopamine and serotonin neurotransmitters [30]. Individuals with T2D showed impaired central insulin signaling, which has a direct impact on brain dopamine systems [15,36–38], indirectly contributing to impulsivity. This is exemplified in the impulsive decision making shown in individuals with T2D [13,14]. According to this view, the current results suggest that impulsive tendencies in individuals with T2D may be particularly affected during emotional states, mainly those involving positive emotions.

When focusing on the T2D group, no correlations at baseline were shown between the impulsivity measures and metabolic parameters. However, a higher UPPS-P total score at baseline predicted higher weight at follow-up. Likewise, baseline scores in sensation seeking predicted a lower decrease in weight/BMI after 3 years of lifestyle intervention. This result could be explained by the link between sensation seeking and behaviors associated with the prospect of reward or novel stimuli [39], including high rewarding foods. Furthermore, a higher BMI at 3 years was predicted by baseline scores in lack of perseverance. This reflects a reduced ability to remain focused on difficult and boring tasks. Associations between lack of perseverance and BMI have been described in transversal studies in individuals with obesity [40]. The present results suggest that low perseverance affects weight control over time in individuals with T2D. These findings could have significant implications for T2D management and prevention interventions, since overweight and obesity represent major risk factors for a poor T2D prognosis and the development of diabetic complications [2]. Lifestyle interventions in older adults have shown significant reductions in body weight and HbA_{1c} [41], so knowing a differential prognosis in the presence of impulsivity traits could help in the planning of a more effective therapeutic intervention or treatment.

Concerning diabetes management, in our population with T2D, impulsive traits were not predictors of glycemic control after 3-years of follow-up. By contrast, another study found that impulsivity negatively correlated with diabetes self-management and positively correlated with self-reported HbA_{1c} levels during a three-months period [22]. However, discrepancies in the results could be related by different timings for the assessment of the HbA_{1c}, with a three-year follow-up considered in the present study. This would suggest that even though impulsive personality traits may correlate with glycemic control, it may not have a long-term effect on the control of diabetes. These results, however, must be interpreted with caution, as our sample presented sub-clinical levels of impulsivity. Therefore, it might be expected that, in individuals with T2D characterized by high impulsivity, this personality trait could have a negative effect on diabetes control. The results of this study should be interpreted in light of some limitations. Firstly, even though the statistical model was adjusted for the intervention group and other potential confounders to diminish the residual effect, we cannot completely discard that these factors are affecting the observed associations. However, it is important to acknowledge that the individuals randomized into the “control” group are also recommended to follow a healthy Mediterranean diet that has previously shown beneficial effects [42,43]. A second limitation is related to the generalizability of these results. For instance, given the samples’ sub-clinical scores in the UPPS-P and the moderate statistical effect, results should be interpreted with caution. To overcome this limitation, future studies in individuals with T2D presenting highly impulsive traits should be conducted. Thirdly, it should be mentioned that self-report measures are not free from response bias, and a more comprehensive evaluation of impulsivity using more objective measures (e.g., cognitive tests) could be useful to better characterize impulsive profiles in individuals with T2D. Additionally, given that our population represent a sample of older Mediterranean adults with metabolic syndrome, further research is needed to determine the presence of these associations in a younger sample without the coexistence of other metabolic disorders. Despite these limitations, several strengths should be highlighted, including the length of the follow-up (i.e., 3 years) that was conducted in a large sample of individuals with T2D. In clinical terms, these results suggest the importance of considering impulsivity as an additional factor that may negatively affect patients’ adherence, opening future research to the development of ad hoc intervention for individuals with T2D characterized by more impulsive personality traits.

5. Conclusions

Overall, our study results showed that, in this older adult population, there are differences in personality traits in individuals with or without T2D, where UPPS-P scores, positive urgency and total factor registered a higher mean within individuals with T2D, indicating that higher scores on the impulsivity scale are associated with T2D, and these are predictors of weight loss but not of diabetes control after 3 years of lifestyle intervention. Further research is required to determine whether a clinic impulsive trait could have an effect on diabetes control.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/jcm11123476/s1>, Table S1: Cross-sectional association between impulsivity levels (UPPS-P scores) with insulin-related metabolic disorders in TD2+ individuals at baseline: partial correlations adjusted by education (analysis within the TD2+ group, $n = 139$). Figure S1: Flowchart for the sampling.

Author Contributions: Conceptualization, F.F.-A., J.S.-S., G.T. and L.C.-B.; methodology, R.d.I.T., X.P., D.C., M.F., J.S.-S. and F.F.-A.; formal analysis, R.G.; investigation, G.T., L.C.-B., C.G.-M., B.M.-M., R.d.I.T., X.P., D.C., R.G., A.C.-R., S.J.-M., N.B., R.F.-C., V.E.-L., L.F., J.N., M.M., S.D.I.H.-D., M.F., J.S.-S. and F.F.-A.; resources, R.d.I.T., X.P., D.C., M.F., J.S.-S. and F.F.-A.; data curation, R.G.; writing—original draft preparation, G.T., L.C.-B., C.G.-M., B.M.-M., R.G., J.S.-S. and F.F.-A.; writing—review and editing, G.T., L.C.-B., C.G.-M., B.M.-M., R.d.I.T., X.P., D.C., R.G., A.C.-R., S.J.-M., N.B., R.F.-C., V.E.-L., L.F., J.N., M.M., S.D.I.H.-D., M.F., J.S.-S. and F.F.-A.; supervision, R.d.I.T., X.P., D.C., M.F., J.S.-S. and F.F.-A.; project administration, R.d.I.T., X.P., D.C., M.F., J.S.-S. and F.F.-A.; funding acquisition,

R.d.I.T., X.P., D.C., M.F., J.S.-S. and F.F.-A. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the official Spanish Institutions for funding scientific biomedical research, CIBER Fisiopatología de la Obesidad y Nutrición (CIBEROBN) and Instituto de Salud Carlos III (ISCIII), through the Fondo de Investigación para la Salud (FIS), which is co-funded by the European Regional Development Fund (six coordinated FIS projects led by JS-S and JVi, including the following projects: PI13/00673, PI13/00492, PI13/00272, PI13/01123, PI13/00462, PI13/00233, PI13/02184, PI13/00728, PI13/01090, PI13/01056, PI14/01722, PI14/00636, PI14/00618, PI14/00696, PI14/01206, PI14/01919, PI14/00853, PI14/01374, PI14/00972, PI14/00728, PI14/01471, PI16/00473, PI16/00662, PI16/01873, PI16/01094, PI16/00501, PI16/00533, PI16/00381, PI16/00366, PI16/01522, PI16/01120, PI17/00764, PI17/01183, PI17/00855, PI17/01347, PI17/00525, PI17/01827, PI17/00532, PI17/00215, PI17/01441, PI17/00508, PI17/01732, PI17/00926, PI19/00957, PI19/00386, PI19/00309, PI19/01032, PI19/00576, PI19/00017, PI19/01226, PI19/00781, PI19/01560, PI19/01332, PI20/01802, PI20/00138, PI20/01532, PI20/00456, PI20/00339, PI20/00557, PI20/00886, PI20/01158); the Especial Action Project entitled: Implementación y evaluación de una intervención intensiva sobre la actividad física Cohorte PREDIMED-Plus grant to JS-S; the Recercaixa (number 2013ACUP00194) grant to JS-S; grants from the Consejería de Salud de la Junta de Andalucía (PI0458/2013, PS0358/2016, PI0137/2018); the PROMETEO/2017/017 grant from the Generalitat Valenciana; the SEMERGEN grant. We thank CERCA Programme/Generalitat de Catalunya for institutional support. This research was also partially funded by EU-H2020 Grants (Eat2beNICE/H2020-SFS-2016-2; Ref 728018; and PRIME/H2020-SC1-BHC-2018-2020; Ref: 847879). J.S.-S. is partially supported by ICREA under the ICREA Academia program. J.N. is supported by a pre-doctoral grant from Ministerio de Ciencia, Innovación y Universidades (FPU 20/00385). RG is supported by the Catalan Institution for Research and Advanced Studies (ICREA-Academia, 2021-Programme). None of the funding sources took part in the design, collection, analysis, or interpretation of the data or in the decision to submit the manuscript for publication.

Institutional Review Board Statement: According to the ethical standards of the Declaration of Helsinki by the Research Ethics Committees, all the participating institutions approved the study protocol and procedures (PR240/13; H1509263926814; 2013/5276/I; 13-07-25/7proj2).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Due to signed consent agreements regarding data sharing, there are restrictions on data availability for the PREDIMED-Plus trial. These only allow access to external researchers for studies following the project purposes. Requestors wishing to access the PREDIMED-Plus trial data used in this study can make a request to the PREDIMED-Plus trial Steering Committee chair: jordi.salas@urv.cat. The request will then be passed to members of the PREDIMED-Plus Steering Committee for deliberation.

Acknowledgments: We thank all the volunteers for the participation in and personnel for the contributions to the PREDIMED-Plus trial. CIBEROBN, CIBERESP, and CIBERDEM are initiatives of Instituto de Salud Carlos III (ISCIII), Madrid, Spain. The authors also thank the PREDIMED-Plus Biobank Network as a part of the National Biobank Platform of the ISCIII for storing and managing the PREDIMED-Plus biological samples.

Conflicts of Interest: F.F.-A. reports consultation fees and travel expenses from Novo Nordisk, but also editorial honoraria as EIC from Wiley. J.S.-S. serves on the board of the International Nut and Dried Fruit Council and received grant support through this institution. He also served on the Executive Committee of the Instituto Danone, Spain, and on the Scientific Committee of the Danone International Institute. He has received research support from the Patrimonio Comunal Olivarero, Spain, and Borges S.A., Spain. He receives consulting fees or travel expenses from Eroski Foundation, the Instituto Danone, Spain, Mundipharma and Abbot Laboratories. The rest of authors declare no conflict of interest.

References

1. Khan, M.A.B.; Hashim, M.J.; King, J.K.; Govender, R.D.; Mustafa, H.; Kaabi, J. Al Epidemiology of Type 2 diabetes—Global burden of disease and forecasted trends. *J. Epidemiol. Glob. Health* **2020**, *10*, 107–111. [[CrossRef](#)] [[PubMed](#)]
2. Kirkman, M.S.; Briscoe, V.J.; Clark, N.; Florez, H.; Haas, L.B.; Halter, J.B.; Huang, E.S.; Korytkowski, M.T.; Munshi, M.N.; Odegaard, P.S.; et al. Diabetes in older adults. *Diabetes Care* **2012**, *35*, 2650–2664. [[CrossRef](#)] [[PubMed](#)]

3. Mallorquí-Bagué, N.; Lozano-Madrid, M.; Toledo, E.; Corella, D.; Salas-Salvadó, J.; Cuenca-Royo, A.; Vioque, J.; Romaguera, D.; Martínez, J.A.; Wärnberg, J.; et al. Type 2 diabetes and cognitive impairment in an older population with overweight or obesity and metabolic syndrome: Baseline cross-sectional analysis of the PREDIMED-plus study. *Sci. Rep.* **2018**, *8*, 16128. [[CrossRef](#)] [[PubMed](#)]
4. Bellary, S.; Kyrou, I.; Brown, J.E.; Bailey, C.J. Type 2 diabetes mellitus in older adults: Clinical considerations and management. *Nat. Rev. Endocrinol.* **2021**, *17*, 534–548. [[CrossRef](#)]
5. Cyders, M.A.; Coskunpinar, A. Measurement of constructs using self-report and behavioral lab tasks: Is there overlap in nomothetic span and construct representation for impulsivity? *Clin. Psychol. Rev.* **2011**, *31*, 965–982. [[CrossRef](#)]
6. Bartholdy, S.; Dalton, B.; O'Daly, O.G.; Campbell, I.C.; Schmidt, U. A systematic review of the relationship between eating, weight and inhibitory control using the stop signal task. *Neurosci. Biobehav. Rev.* **2016**, *64*, 35–62. [[CrossRef](#)]
7. VanderBroek-Stice, L.; Stojek, M.K.; Beach, S.R.H.; VanDellen, M.R.; MacKillop, J. Multidimensional assessment of impulsivity in relation to obesity and food addiction. *Appetite* **2017**, *112*, 59–68. [[CrossRef](#)]
8. Mobbs, O.; Crépin, C.; Thiéry, C.; Golay, A.; Van der Linden, M. Obesity and the four facets of impulsivity. *Patient Educ. Couns.* **2010**, *79*, 372–377. [[CrossRef](#)]
9. Aiello, M.; Ambron, E.; Situlin, R.; Foroni, F.; Biolo, G.; Rumiati, R.I. Body weight and its association with impulsivity in middle and old age individuals. *Brain Cogn.* **2018**, *123*, 103–109. [[CrossRef](#)]
10. Meule, A.; Blechert, J. Interactive and indirect effects of trait impulsivity facets on body mass index. *Appetite* **2017**, *118*, 60–65. [[CrossRef](#)]
11. Meule, A.; Blechert, J. Trait impulsivity and body mass index: A cross-sectional investigation in 3073 individuals reveals positive, but very small relationships. *Health Psychol. Open* **2016**, *3*, 1–6. [[CrossRef](#)] [[PubMed](#)]
12. Lavagnino, L.; Arnone, D.; Cao, B.; Soares, J.C.; Selvaraj, S. Inhibitory control in obesity and binge eating disorder: A systematic review and meta-analysis of neurocognitive and neuroimaging studies. *Neurosci. Biobehav. Rev.* **2016**, *68*, 714–726. [[CrossRef](#)] [[PubMed](#)]
13. Testa, G.; Mora-Maltas, B.; Camacho-Barcia, L.; Granero, R.; Lucas, I.; Agüera, Z.; Jiménez-Murcia, S.; Baños, R.; Bertaina-Anglade, V.; Botella, C.; et al. Transdiagnostic Perspective of Impulsivity and Compulsivity in Obesity: From Cognitive Profile to Self-Reported Dimensions in Clinical Samples with and without Diabetes. *Nutrients* **2021**, *13*, 4426. [[CrossRef](#)] [[PubMed](#)]
14. Sun, D.M.; Ma, Y.; Sun, Z.B.; Xie, L.; Huang, J.Z.; Chen, W.S.; Duan, S.X.; Lin, Z.R.; Guo, R.W.; Le, H.B.; et al. Decision-making in primary onset middle-age type 2 diabetes mellitus: A BOLD-fMRI study. *Sci. Rep.* **2017**, *7*, 10246. [[CrossRef](#)] [[PubMed](#)]
15. Kleinridders, A.; Cai, W.; Cappellucci, L.; Ghazarian, A.; Collins, W.R.; Vienberg, S.G.; Pothos, E.N.; Kahn, C.R. Insulin resistance in brain alters dopamine turnover and causes behavioral disorders. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 3463–3468. [[CrossRef](#)] [[PubMed](#)]
16. Fiory, F.; Perruolo, G.; Cimmino, I.; Cabaro, S.; Pignalosa, F.C.; Miele, C.; Beguinot, F.; Formisano, P.; Oriente, F. The Relevance of Insulin Action in the Dopaminergic System. *Front. Neurosci.* **2019**, *13*, 868. [[CrossRef](#)] [[PubMed](#)]
17. Čukić, I.; Weiss, A. Personality and diabetes mellitus incidence in a national sample. *J. Psychosom. Res.* **2014**, *77*, 163–168. [[CrossRef](#)]
18. Skinner, T.C.; Bruce, D.G.; Davis, T.M.E.; Davis, W.A. Personality traits, self-care behaviours and glycaemic control in Type 2 diabetes: The Fremantle Diabetes Study Phase II. *Diabet. Med.* **2014**, *31*, 487–492. [[CrossRef](#)]
19. Čukić, I.; Möttus, R.; Realo, A.; Allik, J. Elucidating the links between personality traits and diabetes mellitus: Examining the role of facets, assessment methods, and selected mediators. *Personal. Individ. Differ.* **2016**, *94*, 377–382. [[CrossRef](#)]
20. Costa, P.; McCrae, R. *Neo Personality Inventory-Revised (NEO PI-R)*; Psychological Assessment Resources: Odessa, FL, USA, 1992.
21. Whiteside, S.P.; Lynam, D.R. The five factor model and impulsivity: Using a structural model of personality to understand impulsivity. *Personal. Individ. Differ.* **2001**, *30*, 669–689. [[CrossRef](#)]
22. Hadj-Abo, A.; Enge, S.; Rose, J.; Kunte, H.; Fleischhauer, M. Individual differences in impulsivity and need for cognition as potential risk or resilience factors of diabetes self-management and glycemic control. *PLoS ONE* **2020**, *15*, e0227995. [[CrossRef](#)]
23. Wainwright, K.; Romanowich, P.; Crabtree, M.A. Associations between impulsivity and self-care adherence in individuals diagnosed with Type 2 or prediabetes. *PLoS ONE* **2022**, *17*, e0263961. [[CrossRef](#)]
24. Raymond, K.L.; Lovell, G.P. Food addiction symptomology, impulsivity, mood, and body mass index in people with type two diabetes. *Appetite* **2015**, *95*, 383–389. [[CrossRef](#)] [[PubMed](#)]
25. Mallorquí-Bagué, N.; Lozano-Madrid, M.; Vitró-Alcaraz, C.; Forcano, L.; Díaz-López, A.; Galera, A.; Fernández-Carrión, R.; Granero, R.; Jiménez-Murcia, S.; Corella, D.; et al. Effects of a psychosocial intervention at one-year follow-up in a PREDIMED-plus sample with obesity and metabolic syndrome. *Sci. Rep.* **2021**, *11*, 9144. [[CrossRef](#)]
26. Martínez-González, M.A.; Buil-Cosiales, P.; Corella, D.; Bulló, M.; Fitó, M.; Vioque, J.; Romaguera, D.; Alfredo Martínez, J.; Wärnberg, J.; López-Miranda, J.; et al. Cohort profile: Design and methods of the PREDIMED-Plus randomized trial. *Int. J. Epidemiol.* **2019**, *48*, 387–388. [[CrossRef](#)]
27. Alberti, K.G.M.M.; Eckel, R.H.; Grundy, S.M.; Zimmet, P.Z.; Cleeman, J.I.; Donato, K.A.; Fruchart, J.C.; James, W.P.T.; Loria, C.M.; Smith, S.C. Harmonizing the metabolic syndrome: A joint interim statement of the international diabetes federation task force on epidemiology and prevention; National heart, lung, and blood institute; American heart association; World heart federation; International. *Circulation* **2009**, *120*, 1640–1645. [[CrossRef](#)]

28. WHO/Europe Nutrition—Body Mass Index—BMI. Available online: <https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi> (accessed on 28 February 2022).
29. Matthews, D.R.; Hosker, J.P.; Rudenski, A.S.; Naylor, B.A.; Treacher, D.F.; Turner, R.C. Homeostasis model assessment: Insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* **1985**, *28*, 412–419. [[CrossRef](#)]
30. Cyders, M.A.; Smith, G.T. Emotion-Based Dispositions to Rash Action: Positive and Negative Urgency. *Psychol. Bull.* **2008**, *134*, 807–828. [[CrossRef](#)]
31. Smith, G.T.; Fischer, S.; Fister, S.M. Incremental Validity Principles in Test Construction. *Psychol. Assess.* **2003**, *15*, 467–477. [[CrossRef](#)]
32. Verdejo-García, A.; Lozano, Ó.; Moya, M.; Alcázar, M.Á.; Pérez-García, M. Psychometric properties of a spanish version of the UPPS-P impulsive behavior scale: Reliability, validity and association with trait and cognitive impulsivity. *J. Personal. Assess.* **2010**, *92*, 70–77. [[CrossRef](#)]
33. *StataCorp Stata Statistical Software*; StataCorp: College Station, TX, USA, 2021.
34. Kleinbaum, D.G.; Kupper, L.L.; Nizam, A.; Rosenberg, E.S. *Applied Regression Analysis and Other Multivariable Methods*; Cengage Learning: Boston, MA, USA, 2013; ISBN 1-28-505108-4.
35. Cyders, M.A.; Smith, G.T. Mood-based rash action and its components: Positive and negative urgency. *Personal. Individ. Differ.* **2007**, *43*, 839–850. [[CrossRef](#)]
36. Schoffeleer, A.N.M.; Drukarch, B.; De Vries, T.J.; Hogenboom, F.; Schetters, D.; Pattij, T. Insulin modulates cocaine-sensitive monoamine transporter function and impulsive behavior. *J. Neurosci.* **2011**, *31*, 1284–1291. [[CrossRef](#)]
37. Könnner, A.C.; Hess, S.; Tovar, S.; Mesáros, A.; Sánchez-Lasheras, C.; Evers, N.; Verhagen, L.A.W.; Brönneke, H.S.; Kleinridders, A.; Hampel, B.; et al. Role for insulin signaling in catecholaminergic neurons in control of energy homeostasis. *Cell Metab.* **2011**, *13*, 720–728. [[CrossRef](#)]
38. Sevak, R.J.; Koek, W.; Galli, A.; France, C.P. Insulin replacement restores the behavioral effects of quinpirole and raclopride in streptozotocin-treated rats. *J. Pharmacol. Exp. Ther.* **2007**, *320*, 1216–1223. [[CrossRef](#)]
39. Joseph, J.E.; Liu, X.; Jiang, Y.; Lynam, D.; Kelly, T.H. Neural correlates of emotional reactivity in sensation seeking. *Psychol. Sci.* **2009**, *20*, 215–223. [[CrossRef](#)]
40. Murphy, C.M.; Stojek, M.K.; MacKillop, J. Interrelationships among impulsive personality traits, food addiction, and Body Mass Index. *Appetite* **2014**, *73*, 45–50. [[CrossRef](#)]
41. Valabhji, J.; Barron, E.; Bradley, D.; Bakhai, C.; Fagg, J.; O'Neill, S.; Young, B.; Wareham, N.; Khunti, K.; Jebb, S.; et al. Early outcomes from the English National Health Service Diabetes prevention programme. *Diabetes Care* **2020**, *43*, 152–160. [[CrossRef](#)]
42. Martínez-González, M.A.; Salas-Salvadó, J.; Estruch, R.; Corella, D.; Fitó, M.; Ros, E. Benefits of the Mediterranean Diet: Insights From the PREDIMED Study. *Prog. Cardiovasc. Dis.* **2015**, *58*, 50–60. [[CrossRef](#)]
43. Salas-Salvadó, J.; Bulló, M.; Babio, N.; Martínez-González, M.Á.; Ibarrola-Jurado, N.; Basora, J.; Estruch, R.; Covas, M.I.; Corella, D.; Arós, F.; et al. Reduction in the incidence of type 2 diabetes with the Mediterranean diet: Results of the PREDIMED-Reus nutrition intervention randomized trial. *Diabetes Care* **2011**, *34*, 14–19. [[CrossRef](#)]