

# Benign Mesenteric Lesion Presenting with Features of a Malignant Mass

## Abstract

Encapsulated fat necrosis (EFN), most commonly, is an asymptomatic entity and is often found incidentally in images. However, in the abdomen, it may present as an acute abdomen. Mesenteric fat necrosis is part of a larger disease spectrum called collectively mesenteric sclerosis. It results in forming of a mass that can be confused with other pathologies such as liposarcoma, carcinoma of the cecum, and other more benign conditions such as appendagitis of the epiploon. We present the case of an 82-year-old male who presented with an asymptomatic right lower quadrant mass with concerning computed tomography findings with no previous abdominal surgery or trauma history. Diagnosing EFN is crucial as it can mimic bowel cancer and immune-related mesenteric pathology such as sclerosing mesenteritis, the management of which is far more extreme and aggressive than EFN.

**Keywords:** EFN, encapsulated fat necrosis, mesenteric pathology

## Introduction

Fat necrosis is a benign non-suppurative inflammatory process of adipose tissue.<sup>[1]</sup>

Encapsulated fat necrosis was first described in the breast in 1975, but it can happen anywhere in the body. Most commonly, it is an asymptomatic entity and appears as an incidental finding in images. However, In the abdomen, it may present with symptoms such as a vague and dull pain or as an acute abdomen mimicking appendicitis or diverticulitis.

Mesenteric fat involvement often times results in the formation of a mass that can be confused with other pathologies such as liposarcoma, carcinoma of the cecum and other more benign conditions such as appendagitis of the epiploon. This creates the need for a differential diagnosis from a potential malignant and invasive condition, making the exploratory laparoscopic surgery one of the most common approaches.

We report the case of an incidental finding during a laparoscopic resection in a patient with an acute presentation mimicking appendicitis.

## Case Report

We present the case of an 82-year-old male who presented to our surgery center with

a right lower quadrant mass. The patient states that he noticed the mass a year ago during a routine physical check-up and that it is nonprogressive and denies abdominal pain, fever, weight loss, nausea, or vomiting. On physical examination, his abdomen was soft, nontender, and nondistended, and bowel sounds were heard and did not reveal any mass. He has daily bowel movements and denies dysuria. He has a history of diverticulosis of the descending and sigmoid colon, and his medication consists of daily atorvastatin. The patient denies any history of trauma or abdominal surgery. A previous colonoscopy was clear of intraluminal anomalies. Computed tomography (CT) scan performed 10 months prior showed a 4.5 cm mass right lower quadrant with densely calcified rim and mixed-density internal components, including calcifications. The appearance was nonspecific. It was interpreted as a possible cecal tumor/liposarcoma and referred for surgery. Blood investigations were unremarkable.

The decision was made to carry out a laparoscopic exploration with a possible cecectomy. During surgery, an encapsulated mass was found to be at the mesentery of the terminal ileum near the ileocecal valve, very close to the base of the appendix. The mass appeared to be exophytic, and there was a small pedicle coming from the

**Josefina Principe,  
Anna Mary Jose<sup>1</sup>,  
Santiago Niño  
Basto<sup>2</sup>,  
Isabela Castañeda<sup>3</sup>,  
Abdul Rafay Pasha<sup>4</sup>**

*Department of Surgery,  
Pontifical Catholic University  
of Argentina, Buenos Aires,  
Argentina, <sup>1</sup>Department of  
Surgery, Acharya Vinoba  
Bhave Rural Hospital,  
Wardha, Maharashtra, India,  
<sup>2</sup>Department of Surgery, Del  
Rosario University, <sup>3</sup>Department  
of Surgery, Javeriana University,  
Bogota, Colombia, <sup>4</sup>Allama  
Iqbal Medical College, Lahore,  
Pakistan*

**Submitted:** 19-Feb-2023  
**Revised:** 15-Jul-2023  
**Accepted:** 11-Aug-2023  
**Published:** 25-Sep-2023

### Address for correspondence:

*Dr. Josefina Principe,  
Pontifical Catholic  
University of Argentina,  
Buenos Aires, Argentina.  
E-mail: joselaprincede@  
gmail.com*

### Access this article online

**Website:**  
<https://journals.ww.com/IJABMR>

**DOI:**  
10.4103/ijabmr.ijabmr\_77\_23

### Quick Response Code:



**How to cite this article:** Principe J, Jose AM, Basto SN, Castañeda I, Pasha AR. Benign mesenteric lesion presenting with features of a malignant mass. Int J App Basic Med Res 2023;13:186-8.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com



**Figure 1: Macroscopic view of the mass. Gross view of the encapsulated fat necrosis**

mesentery. It was a 4–5 cm calcified mass that was hard and fibrous [Figure 1]. The appendix was slightly dilated, so both the mass and the appendix were removed without bleeding or abdominal injury. The patient's postoperative course was uneventful, and he was discharged 2 days later. Histological examination of the mass showed a collection of atypical cells with clear cytoplasm consistent with lipoblast mixed with mitotic figures and extensive necrosis, revealing the culprit to be mesenteric fat necrosis [Figure 2].

## Discussion

Fat necrosis is a benign nonsuppurative inflammatory process of adipose tissue. It is commonly due to accidental or surgical trauma to the adipose tissue with extracellular liberation of fat or enzymatic lysis of fat due to the release of lipases.<sup>[1]</sup> Encapsulated fat necrosis (EFN) is an entity that is rarely diagnosed at first glance; there are few case reports about the same in the literature.<sup>[2–6]</sup> Mesenteric fat necrosis is part of a larger disease spectrum called collectively as mesenteric sclerosis. It was first described in 1924 as retractile mesenteritis.<sup>[7]</sup>

Mesenteric fat necrosis belongs to a group of idiopathic disorders of the mesentery and peritoneum referred to as “sclerosing mesenteritis.” It is pathologically characterized by the succession of different events that can be classified into three stages: Inflammation (mesenteric panniculitis), fat necrosis (described as mesenteric lipodystrophy), and fibrosis (known as retractile or sclerosing mesenteritis). This spectrum of disorders is notorious for being indistinguishable from each other from a clinical, radiographic, and microscopic angle. The majority of the cases are incidental findings in abdominal CT or ultrasound. Due to the lack of clearly defining terms to differentiate one from the other, these pathological terms are often used interchangeably, creating much confusion. Having a sense of clarity on what separates these pathologies is critical for their management. Emory *et al.* reviewed cases 84 cases



**Figure 2: Histological report of the mass showing collection of atypical cells with clear cytoplasm mixed with mitotic figures and extensive necrosis (red circle)**

of mesenteric lipodystrophy, mesenteric panniculitis, and sclerosing mesenteritis. They found that in most patients, the extent of fibrosis, inflammation, and fat necrosis were too mixed to differentiate the three mesenteric pathologies.<sup>[2]</sup>

Our patient presented with an asymptomatic right lower quadrant swelling with a calcified rim on the CT scan, increasing the suspicion of malignancy with no history of trauma or previous abdominal surgeries. The proximity of the mass to the cecum and the patient's age brought to mind the possibility of a cecal tumor. Most of the tumors involving the cecum are asymptomatic and diagnosed incidentally at the time of appendectomy. A malignancy is found in approximately 1% of appendectomy specimens. The most malignant neoplasm of the cecum is the adenocarcinoma which can present with abdominal pain, ascites, abdominal mass, or an acute presentation similar to appendicitis. Most tumors are diagnosed postoperatively, and thus, staging procedures mostly occur in postoperative management. Therefore, a surgical approach is preferred in diagnosing and treating unexplained abdominal masses. Multiple studies found the average age in the 4–5<sup>th</sup> decade.<sup>[3,6,8]</sup> Abdominal pain was the most commonly described symptom in most studies.<sup>[2,3,8]</sup> History of trauma or surgery was infrequently found but still an attributing factor in many studies.<sup>[3,6]</sup>

EFN can demonstrate a mild mass effect on adjacent structures, and its fibrous capsule may slightly enhance after administration of intravenous contrast material: findings that raise suspicion of a malignant lesion. Its encapsulated asymptomatic presentation and long history aroused the suspicion of liposarcoma. Liposarcoma appears to arise from precursors of adipocytes (fat cells) and is most commonly found in the extremities and retroperitoneum. The reported characteristics of liposarcoma on CT images are (1) homogeneity, (2) infiltration or poor margination, (3) CT numbers more significant than normal fat, and

(4) Contrast enhancement. However, unlike liposarcoma, fat necrosis does not show organ invasion and may be focally tender at palpation.

Another differential that overlaps with EFN is sclerosing mesenteritis, a rare, nonneoplastic inflammatory and fibrotic disease that affects the mesentery.

A presumptive diagnosis of sclerosing mesenteritis can be based on abdominal CT findings of a fat ring or halo sign and pseudo capsule in contrast to EFN that appears as a round, encapsulated with predominantly fat attenuation. However, a definitive diagnosis of sclerosing mesenteritis requires histologic evaluation to rule out other etiologies.

In our case, perioperatively, the mass was found to be attached by a pedicle to the cecum and did not directly arise from it. In the face of an asymptomatic patient with regular bowel habits, the possibility of a cecal tumor at this point was questionable. In addition, the mass was encapsulated and noninvasive, making liposarcoma a less likely diagnosis. In the face of such a dilemma, the next best step was to surgically excise the mass and send the specimen for pathological analysis for a definitive diagnosis. Pathology confirmed that it was a case of EFN of the mesentery. Although the diagnosis was benign, it is clinically and radiographically misleading and alarming. Since the diagnosis is almost always made postoperatively, it warrants the need for EFN to be a differential in cases concerning a suspicious abdominal mass. Although our patient was asymptomatic, De Kock and Delrue in their case report, presented a 33-year-old female with a 36-h history of periumbilical pain associated with anorexia and nausea, and a tender right lower quadrant and CT showed no calcifications within the mass.<sup>[5]</sup>

EFN becomes crucial in patients with abdominal cancer who can present with acute abdomen or signs of bowel obstruction, creating an impression of a recurrent tumor.<sup>[3,4]</sup>

The rarity with which EFN presents can blindside the surgeon and the patient, mainly because of its ability to

present in a varied manner clinically and radiographically. EFN is known to decrease in size, and the most appropriate way to manage it is to excise and have a histopathologically confirmed diagnosis.

### Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

### References

1. Goepel J. Responses to cellular injury. *Gen Syst Pathol*;2:117.
2. Emory TS, Monihan JM, Carr NJ, Sobin LH. Sclerosing mesenteritis, mesenteric panniculitis and mesenteric lipodystrophy: A single entity? *Am J Surg Pathol* 1997;21:392-8.
3. Watanabe J, Osaki T, Tatebe S, Goto K, Endo K, Nakamura S, *et al.* Encapsulated fat necrosis mimicking abdominal liposarcoma: A case report and literature review. *Clin Case Rep* 2020;8:2255-8.
4. Yeh HJ, Chang CC, Kuo LJ. Right pelvis wall encapsulated fat necrosis mimics malignancy change: A case report. *Clin Surg* 2019;4:2480.
5. De Kock I, Delrue L. Encapsulated mesenteric fat necrosis. *J Belg Soc Radiol* 2016;100:53.
6. Kiryu H, Rikihisa W, Furue M. Encapsulated fat necrosis – A clinicopathological study of 8 cases and a literature review. *J Cutan Pathol* 2000;27:19-23.
7. Sulla JV. Mesenterite retractile sclerosante. *Policlinico (sezprat)* 1924;31:575-81.
8. Available from: <http://www.chinadot.cn/portal/mr.action?doi=10.3760/cma.j.issn.0578-1426.2017.02.006>. [Last accessed on 2023 Feb 12].