

REVIEW

Physiologic roles of P2 receptors in leukocytes

Anael Viana Pinto Alberto¹ | Natiele Carla da Silva Ferreira¹ |
 André Gustavo Calvano Bonavita² | Oscar Kenji Nihei³ | Fernando Pires de Farias⁴ |
 Rodrigo da Cunha Bisaggio^{1,5} | Cristóvão de Albuquerque⁶ | Wilson Savino^{7,8} |
 Robson Coutinho-Silva⁹ | Pedro Muanis Persechini¹⁰ | Luiz Anastacio Alves¹

¹Laboratory of Cellular Communication, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro, RJ, Brazil

²Laboratory of Bioactive Products, Multidisciplinary Center, Federal University of Rio de Janeiro, Macaé, RJ, Brazil

³Center of Education and Letter, State University of the West of Paraná, Foz do Iguaçu, PR, Brazil

⁴Souza Marques School of Medicine, Souza Marques College, Rio de Janeiro, RJ, Brazil

⁵Federal Institute of Education, Science, and Technology of Rio de Janeiro, Rio de Janeiro, RJ, Brazil

⁶PositivaMente Consulting, São Paulo, SP, Brazil

⁷Laboratory on Thymus Research, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro, RJ, Brazil

⁸Brazilian National Institute of Science and Technology on Neuroimmunomodulation, Rio de Janeiro Research Network on Neuroinflammation, Rio de Janeiro, RJ, Brazil

⁹Laboratory of Immunophysiology, Carlos Chagas Filho Biophysics Institute, Federal University of Rio de Janeiro, Rio de Janeiro, RJ, Brazil

¹⁰Laboratory of Immunobiophysics, Carlos Chagas Filho Biophysics Institute, Federal University of Rio de Janeiro, Rio de Janeiro, RJ, Brazil

Correspondence

Luiz Anastacio Alves, Av. Brasil 4365,
 Manginhos, Rio de Janeiro, RJ, Brazil.
 Email: alveslaa30@gmail.com

(in memory)

Abstract

Since their discovery in the 1970s, purinergic receptors have been shown to play key roles in a wide variety of biologic systems and cell types. In the immune system, purinergic receptors participate in innate immunity and in the modulation of the adaptive immune response. In particular, P2 receptors, which respond to extracellular nucleotides, are widely expressed on leukocytes, causing the release of cytokines and chemokines and the formation of inflammatory mediators, and inducing phagocytosis, degranulation, and cell death. The activity of these receptors is regulated by ectonucleotidases—expressed in these same cell types—which regulate the

Abbreviations: 2-AG, 2-arachidonoylglycerol; 2meSATP, 2-methylthio-adenosine triphosphate; AA, arachidonic acid; AC, adenylate cyclase; AD, Alzheimer disease; ADA, adenosine deaminase; ADO, adenosine; ADP, adenosine diphosphate; AMP, adenosine monophosphate; ATP, adenosine triphosphate; ATP_γS, adenosine-5'-o-(3-thio-triphosphate); Aβ, amyloid β; BAL, bronchoalveolar lavage; BALF, bronchoalveolar lavage fluid; BCG, bacilli Calmette-Guerin; BD, Behçet's disease; BMDMs, bone marrow derived-macrophages; BzATP, 2'-(3')-o-(4-benzoylbenzoyl) adenosine triphosphate; CALHM1, calcium homeostasis modulating channel 1; CFA, complete Freund's adjuvant; CFTR, cystic fibrosis transmembrane conductance regulator; CLL, chronic lymphocytic leukemia; COX-2, cyclooxygenase-2; cP2X7, chicken P2X7; Cx43, connexin 43; DAMPs, damage-associated molecular patterns; DCs, dendritic cells; ECP, eosinophil cationic protein; E-NPP, ectonucleotide pyrophosphate/phosphodiesterase; E-NTPDase, ectonucleoside triphosphate diphosphohydrolase; GALT, gut-associated lymphoid tissue; GM-CSF, granulocyte-macrophage colony stimulating factor; GPCR, G protein-coupled receptor; gpP2X7, giant panda P2X7; GTP, guanosine triphosphate; HDM, house dust mite; HIV, human immunodeficiency virus; HLA-G, human leukocyte antigen G; HMGB1, high mobility group box 1; hP2X3, human P2X3; IBD, inflammatory bowel disease; IP₃, inositol triphosphate; KC, keratinocyte chemoattractant; KO, knockout; LPS, lipopolysaccharide; LTB₄, leukotriene B₄; LTC₄, leukotriene C₄; MACs, maxi-anion channels; MCP-1, monocyte chemoattractant protein-1; MGCs, multinucleated giant cells; MIP-1α, macrophage inflammatory protein-1α; MIP-1MIP-2, macrophage inflammatory protein 2, ted giant cells; MIP-3α, macrophage inflammatory protein-3α; MPS, mononuclear phagocytic system; NAADP, nicotinic acid adenine dinucleotide phosphate; NAD, nicotinamide adenine dinucleotide; NETs, neutrophil extracellular traps; N; NLRP3, NOD-, LRR- and pyrin domain-containing protein 3; OA, oleic acid; oATP, oxidized ATP; OGD, oxygen-glucose deprivation; OVA, ovalbumin; P1R, P1 receptor; P2R, P2 receptor; PAMPs, pathogen-associated molecular patterns; PGE₂, prostaglandin E₂; PLA₂, phospholipase A₂; PLC, phospholipase C; PLD, phospholipase D; PMN, polymorphonuclear neutrophil; PNP, purine nucleotide phosphorylase; PPADS, pyridoxal phosphate-6-azophenyl-2',4'-disulfonic acid; RB-2, Reactive Blue-2; ROS, reactive oxygen species; rP2X7, rat P2X7; TACE, TNF-α converting enzyme; TM, transmembrane; TXB₂, thromboxane B₂; UDP, uridine diphosphate; UTP, uridine triphosphate; VASP, vasodilator-stimulated phosphoprotein; VRACs, volume-regulated anionic channels; WT, wildtype; zfP2X4, zebrafish P2X4; α,β-meATP, α,β-methylene adenosine triphosphate.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Journal of Leukocyte Biology* published by Wiley Periodicals LLC on behalf of Society for Leukocyte Biology.

availability of nucleotides in the extracellular environment. In this article, we review the characteristics of the main purinergic receptor subtypes present in the immune system, focusing on the P2 family. In addition, we describe the physiologic roles of the P2 receptors already identified in leukocytes and how they can positively or negatively modulate the development of infectious diseases, inflammation, and pain.

KEYWORDS

ATP, glia, immune cells, P2X, P2Y

1 | INTRODUCTION

Leukocytes have a critical role in immune surveillance and protection against viruses, bacteria, fungi, and others. These cells are mainly produced in the bone marrow and can be subdivided into several subtypes according to the proteins expressed on their cell membranes, including receptors that help them sense the environment.¹

Purinergic receptors are transmembrane proteins that respond to extracellular purines and pyrimidines, and are divided into 2 families: P1 (activated by adenosine [ADO]) and P2 (activated by nucleotides).² The physiologic agonists of P2 receptors include adenosine triphosphate (ATP), adenosine diphosphate (ADP), uridine triphosphate (UTP), uridine diphosphate (UDP), UDP-glucose, nicotinamide adenine dinucleotide (NAD), and nicotinic acid adenine dinucleotide phosphate (NAADP), among others.^{3–6} P2 receptors are further subdivided into 2 classes: P2X (ligand-gated ion channels) and P2Y (G-protein coupled receptors (GPCRs)).²

P2 receptors display a widespread distribution in mammalian tissues. Cells of the immune system express several P2 subtypes, which play important roles on chemotaxis, cytokine release and formation of proinflammatory mediators, among others.^{7,8} In this article, we review the main characteristics of the purinergic receptors in the immune system, focusing on the P2 family. In addition, we describe important discoveries on the physiologic roles of P2 receptors in leukocytes and how they can positively or negatively modulate the development of infectious diseases, inflammation, and pain.

1.1 | Historical background

In 1979, Cockroft and Gomperts reported that ATP⁴⁻, that is, ATP uncoupled to Mg²⁺, induced histamine release from mast cells, and also turned their plasma membrane permeable to large solutes.^{9,10} This observation followed the description, by Cohn and Parks, of the role of ATP in the formation of pinocytotic vesicles in cultured macrophages¹¹ and the descriptions by Burnstock of purinergic receptors in a variety of tissues,¹⁰ led Cockroft and Gomperts to propose that the effects of extracellular ATP on mast cells were mediated by purinergic receptors.¹² These observations were followed by the description of several physiological responses to ATP in a variety of tissues, including smooth muscle contraction and relaxation, neuron polarization, and membrane permeabilization.^{13–15}

The current classification of P2 receptors was conceived in the 1990s, with the advancement of molecular cloning studies of these proteins and their pharmacologic characterization with the patch-clamp electrophysiologic technique.^{16–18} In the current classification, the P2X class comprises the ionotropic receptors, whereas the P2Y class are GPCRs.

1.2 | P2 receptors and the current understanding of their general role in the immune system

The idea that extracellular nucleotides can play an important modulatory role in the immune system has long been proposed,^{19–21} and P1 and P2 receptors, as well as ectoenzymes that interact with their natural ligands, have been identified in many immune cell types, pointing to complex immune-neuro-endocrine interactions.^{22–27} Tissue injury situations, in particular, result in massive release of ATP and other nucleotides, leading to the activation of P2 receptors. The activation of these receptors induces proinflammatory responses, contributing to the development of acute inflammation. In turn, ectonucleotidases, by converting ATP to ADO, counteract the maintenance of acute inflammatory responses, as ADO activates P1 receptors, which have anti-inflammatory effects and are involved in the resolution of inflammation.²⁸ In this review, we will focus on the effects mediated by P2 receptors on leukocytes.

2 | P2 RECEPTOR

2.1 | P2X receptors

The sequencing of P2X receptors revealed a new class of receptors, with no significant homology to other purinergic receptors based on sequence and mutagenesis analysis.^{29,30} The P2X receptor class comprises 7 members, named P2X1–7. The sequence homologies between them ranges from 35 to 48%.³¹

Each P2X receptor subunit displays 2 hydrophobic transmembrane segments, with 1 extracellular loop and intracellular N- and C-terminals.^{30,32–34} The C-terminal domain has binding sites to interact with protein kinases. The two transmembrane (TM) domains, TM1 and TM2, are involved in the ion channel and pore formation, respectively. The extracellular loop has cysteine residues that form disulfide bridges.³⁵

TABLE 1 Homomeric and heteromeric P2X receptors

P2X receptor	Homomultimer	Heteromultimer(s)
P2X1	Yes	P2X1/2
		P2X1/4
		P2X1/5
P2X2	Yes	P2X1/2
		P2X2/3
		P2X2/6
P2X3	Yes	P2X2/3
P2X4	Yes	P2X1/4
		P2X4/6
P2X5	Yes	P2X1/5
P2X6	No	P2X2/6
		P2X4/6
P2X7	Yes	No

References 40–43.

All P2X receptors described so far, except for P2X6, can form homomultimer channels *in vitro*, since purinergic analogs activate currents when each subtype is expressed in oocytes or cell lines (P2X6, in contrast, only works as a heteromultimer).^{32,33} However, *in vivo* more than one P2X subtype may be found in the same cell type, including neurons and muscle cells, and these receptors also appear to form heteromultimers.^{36–38} Indeed, the coexpression of P2X2 and P2X3 results in a channel with characteristics that differ from an exogenous expression of P2X2 or P2X3 alone; and the same can be observed for P2X1 and P2X5.³⁹ Other heteromeric receptors described so far were P2X4/6⁴⁰ and P2X2/6⁴¹ in the CNS. Table 1 summarizes the homomultimers and heteromultimers formed by P2X.

Regarding the exact number of subunits that form the cation channel, Nicke and colleagues⁴² showed that P2X receptors present a trimeric arrangement, based on chemical cross-linking and blue native PAGE experiments. The trimeric assembly of these receptors was confirmed in 2009 by Kawate and colleagues,⁴⁴ who resolved the crystal structure of the zebrafish P2X4 receptor in its closed state.

Several studies have focused on clarifying the structural motifs responsible for P2X channel formation. These studies revealed that a 25 amino acid sequence, including the second transmembrane domain, is determinant for subunit co-assembly, whereas the extracellular region by itself is not sufficient for the assembly of the full-sized subunits.⁴¹ Additionally, mutations in the transmembrane domains of P2X receptors can alter pore dilation.^{45,46}

The P2X4 subunits resemble the shape of a dolphin: in this analogy, the extracellular domain represents the upper body, and the transmembrane domains represent the fluke (for a detailed explanation of the dolphin's shape, see Ref. 44). Additionally, there are some lateral fenestrations. The N- and C-terminals were not present in the crystalized receptor, preventing a more detailed study of the pore structure. Interestingly, the subunits are connected only by the upper portion of the extracellular domain, and the region immediately above the membrane is not connected. This conformation allows the transmembrane

domains, which are in contact, to move in the gating (opening and closing) of the channel. Moreover, the residues that form the body domain are conserved, which suggests that body-to-body interactions are a common feature among all P2X receptors.⁴⁴

The crystallographic data of the zebrafish P2X4 (zfP2X4) receptor complexed with ATP (in the open state) demonstrated that the three ATP-binding pockets are located in conserved residues distant 40 Å above the membrane border. These binding pockets are formed by the intersection of the three subunits and have several positively charged residues.⁴⁷ The transition from a closed state to an open state occurs when ATP binds the agonist site. In this situation, a small fold forms in the lower part of the extracellular loop and the transmembrane domains enlarge, causing pore dilation. The hydrated ions flow into the pore through the lateral fenestrations.⁴⁷

The crystal structure of human P2X3 (hP2X3) revealed that this receptor also presents a dolphin-like architecture. The transmembrane domains in hP2X3 are longer than those in zfP2X4, and the antagonists of the P2X3 receptor seem to fasten to the orthosteric binding pocket. hP2X3 has an intracellular motif, named “cytoplasmic cap,” which is formed by a network of three β -sheets that cap the cytoplasmic surface of the pore. This structure is only observed in the ATP-bound (open state) and generates lateral fenestrations for water and ion flux. So far, this “cytoplasmic cap” was only observed in P2X3 receptors.⁴⁸

More recently, the P2X7 structure was elucidated by X-ray crystallography. The crystal structure of chicken P2X7 (cP2X7) in complex with the antagonist TNP-ATP again revealed a dolphin-like architecture, with the Asp319 to Leu344 residues in the TM2 helices lining the pore. The binding site for TNP-ATP was the same as for the ATP molecule, allowing interaction with some parts of the structure of adjacent subunits.^{49,50} In contrast, the crystal structure of the P2X7 receptor from giant panda (gpP2X7) exposed the allosteric site, distinct from the ATP-binding site. This allosteric site was formed between two adjacent subunits and was able to accommodate different small molecules through hydrophobic interactions.⁵¹ Finally, the elucidation of the rat P2X7 (rP2X7) structure by cryo-electron microscopy, uncovered characteristics that distinguishes this receptor from other P2X receptors. rP2X7 contains a cytoplasmic segment with a distinctive folding with guanosine and zinc-binding sites as well as a C-cys linked to the membrane with some palmitoyl groups, which explains the reason why this receptor does not desensitize.⁵²

In terms of pharmacology, P2X receptors are physiologically activated by ATP with an EC₅₀ between 0.5 and 12 μ M, with the exception of P2X7, which has an EC₅₀ value greater than 100 μ M. Other general agonists for P2X receptors include 2-methylthio-adenosine triphosphate (2-MeSATP), adenosine-5'-o-(3-thio-triphosphate) (ATP γ S), α,β -methylene adenosine triphosphate (α,β -MeATP), and 2'-(3')-o-(4-benzoylbenzoyl) adenosine triphosphate (BzATP). Regarding antagonists, Suramin, pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid (PPADS), Reactive Blue-2 (RB-2), and TNP-ATP are considered nonselective antagonists. Selective antagonists have been identified for all P2X subtypes except P2X5 and P2X6, but those molecules may act on other subtypes at higher concentrations. Gefapixant, a P2X3 receptor antagonist, has recently completed phase

TABLE 2 P2Y-associated main effector systems

P2Y receptor	G-protein coupled	Main effector systems
P2Y ₁	G _q	PLC, Rac, and Rho activation
P2Y ₂	G _q ; G _i ; G _{12/13}	PLC, Rac, and Rho activation
P2Y ₄	G _q ; G _i	PLC activation
P2Y ₆	G _q ; G _{12/13}	PLC and Rho activation
P2Y ₁₁	G _q ; G _s	PLC and AC activation
P2Y ₁₂	G _i	PLC and Rho activation; AC inhibition
P2Y ₁₃	G _i	PLC and Rho activation; AC inhibition
P2Y ₁₄	G _i	PLC activation; AC inhibition

References 66 and 67.

3 in clinical trials, and the P2X7 receptor antagonists AZD9056 and CE-224,535, have also advanced to the stage of clinical trials, aiming at the treatment of rheumatoid arthritis; however, they did not show superior efficacy to the conventional treatment.^{53,54} Meanwhile, the P2X7 antagonist JNJ-54175446 is currently being tested for depression treatment.

In addition, knockout (KO) models have already been developed for *in vivo* studies with P2X receptors (for more updates on the pharmacology of these receptors, see Ref. 55).

2.2 | P2Y receptors

P2Y receptors belong to the GPCR superfamily. Since the initial cloning in the 1990s, eight subtypes of P2Y have been characterized, named P2Y₁, P2Y₂, P2Y₄, P2Y₆, and P2Y₁₁₋₁₄. Missing numbers consist of subtypes incorrectly classified as P2Y receptors or nonmammalian orthologues.^{17,56,57} As other GPCRs, these receptors present the following putative structure: seven transmembrane domains known as TM1-7, three intracellular loops, three extracellular loops, an intracellular C-terminal, and an extracellular N-terminal.³⁵ The agonist binding site is located between TM6-7 domains near the extracellular face of the plasma membrane.⁵⁸⁻⁶¹

Until now, only two P2Y receptors have had their structures elucidated by X-ray crystallography. The first was the P2Y₁₂ receptor, whose structure revealed the existence of two possible sub-pockets for ligands, and that the agonist binding promotes conformational changes in helices 6 and 7.^{62,63} Similar to P2Y₁₂, the crystal structure of P2Y₁ receptor also has two distinct ligand binding sites that differ in terms of location and shape, and the nucleotide-binding site is situated in the extracellular loop region.^{64,65}

Regarding the associated G protein, P2Y receptors may be coupled to G_q protein (P2Y₁, P2Y₂, P2Y₄, P2Y₆, and P2Y₁₁), whose activation results in the mobilization of intracellular calcium via phospholipase C (PLC)/inositol triphosphate (IP₃), or G_i protein (P2Y₂, P2Y₄, and P2Y₁₂₋₁₄), inhibiting the adenylate cyclase (AC) enzyme. The P2Y₁₁ receptor can couple to both the G_q protein as well as the G_s protein, which activates AC.⁶⁶ Table 2 summarizes the major G proteins associated with P2Y receptors and their main effector systems.

Regarding their pharmacology, human P2Y receptors are activated by different nucleotides, including ADP (P2Y₁, P2Y₁₂, and P2Y₁₃), ATP (P2Y₂ and P2Y₁₁), UDP (P2Y₆ and P2Y₁₄), UTP (P2Y₂ and P2Y₄), UDP-glucose (P2Y₁₄), diadenosine tetraphosphate (P2Y₂, P2Y₄, P2Y₁₂ and P2Y₁₃), and diuridine tetraphosphate (P2Y₂, P2Y₄, and P2Y₆). P2Y receptors have a number of selective antagonists for each of their subtypes (for more information, see Ref. 66). The development of selective antagonists allowed the replacement of nonselective antagonists such as Suramin, PPADS, and RB-2 in experimental tests that seek to understand the physiologic roles of these receptors.⁶⁸ The discovery of new antagonists has been boosted with the advancement of elucidation of the crystallographic structure of P2Y receptors and the use of cheminformatics techniques by research groups that study these receptors. Finally, it is important to emphasize that P2Y₁₂ receptor antagonists have been in the pharmaceutical market for over 20 years, being used to prevent platelet aggregation and acting as antithrombotic agents. These drugs include clopidogrel, prasugrel, ticagrelor, and cangrelor.⁶⁶

3 | PHYSIOLOGIC ACTIVATION OF P2 RECEPTORS IN IMMUNE CELLS

3.1 | Sources of extracellular nucleotides and nucleosides

A central question in the study of purinergic receptors in the immune system is how endogenously released nucleosides and nucleotides might reach concentrations sufficient to activate these receptors. Indeed, 20 years ago there was some skepticism that ATP and ADO could play a decisive role in immune-cell-specific phenomena, due to the lack of quantitative data about nucleotide and nucleoside species, sources, and concentrations in situations that might be relevant for the development of immune response. However, over the years, evidence has accumulated that these molecules can be selectively released from almost all cells studied and can indeed reach sufficient concentrations in many immunologically relevant circumstances. Another source of skepticism regarding the role of purinergic receptors in the immune system may be attributed to the formerly widespread idea

that the immune system is self-contained and exploits only “specific” molecules generated within itself, such as cytokines and chemokines, as its main soluble intercellular mediators. This scenario has changed as multiple and complex immuno-neuro-endocrine interactions have been the object of intense research over the last 20 years.

Currently, the main sources of extracellular nucleotides and nucleosides are believed to be: (a) corelease of nucleotides with other neurotransmitters by sympathetic and parasympathetic nerves; (b) exocytotic release from intracellular vesicles in non-neural cells such as platelets, astrocytes, and suprarenal cells; (c) release mediated by membrane transporters, such as the ATP-binding cassette protein superfamily; (d) electrochemically driven efflux through anion channels and transmembrane pores, such as connexin hemichannels and pannexin channels; (e) lytic release from injured/damaged cells, as might be the case in trauma, in the site of acute infection, and due to lysis by effector cells. In each of these cases, much attention has been given to the release of ATP. However, UTP, ADP, UDP, ADO, and other nucleotides and nucleosides are also released in many situations. The immune system might be potentially affected by all of these sources, as described below.^{69–73}

3.1.1 | Sympathetic and parasympathetic nerve release

Several research groups have demonstrated innervation by sympathetic and parasympathetic nerve fibers of primary and secondary lymphoid organs, including the thymus, bone marrow, spleen, lymph nodes, and gut-associated lymphoid tissue (GALT).^{74,75} In these organs, the innervation predominantly follows the vasculature, but intense nerve fiber ramification is also seen in lymphoid parenchyma—including the thymic cortex, splenic white pulp, the periaxillary lymphatic sheath, lymph node paracortical regions, and T-dependent regions of GALT. In some cases, direct contact between the nerve terminal and immune cells such as lymphocytes and macrophages was evident.⁷⁶ In all these sites, a possible release of ATP conjointly with classical neurotransmitters could be expected as demonstrated in other targets of sympathetic innervation.^{77,78} Similar to what occurs in other organs, the release of vesicular contents from sympathetic nerve terminals in the thymus is regulated at the presynaptic terminal by P1 receptors, as well as by other mediators.⁷⁹

3.1.2 | Exocytotic release

The release of nucleotides stored within intracellular vesicles by exocytosis also occurs in cells of non-neural origin. Those include platelets, mast cells, and suprarenal cells. After an appropriate stimulus, these cells release their vesicular nucleotide contents, modulating neighboring cells in a paracrine fashion.⁸ Evidence of nonlytic release of ATP was also reported for lymphocytes, macrophages, and microglial cells.^{36,80–82} Some data suggest that ATP can be released by both cytotoxic lymphocytes and target cells during the cytolytic effector function. This phenomenon was first reported in 1973 by Heney,

who recognized that the total amount of ATP released during a cytolytic reaction could be larger than what could be contained in the target cell pool, suggesting the release of ATP from both the effector and target cells.⁸³ Whereas target cells probably release ATP as a consequence of pores formed by perforin, the effector cells are engaged in an active process possibly involving the release of lytic granules.^{80,84}

3.1.3 | Release mediated by membrane transporters

Nonvesicular release of cytosolic nucleotides has also been proposed, as the concentration of cytoplasmic ATP in healthy eukaryotic cells ranges from 3 to 5 mM. As a strong anion, ATP cannot diffuse through the plasma membrane lipid bilayer; accordingly, membrane transporters have been proposed as mediators of nonvesicular release in healthy cells. Two possible candidates have been described, the P-glycoprotein and the cystic fibrosis transmembrane conductance regulator (CFTR), both belonging to the ATP-binding cassette protein superfamily.^{85,86} Although the ATP-transporter nature of CFTR is highly controversial, its possible indirect importance in ATP release could not be excluded.^{85,87–92}

3.1.4 | Release mediated by anion channels and transmembrane pores

Several pore-forming proteins have a pore diameter larger than the ATP molecule in cross-section (1.14–1.22 nm). These include connexins, pannexins, calcium homeostasis modulating channel 1 (CALHM1), volume-regulated anionic channels (VRACs), and maxi-anion channels (MACs).^{93,94}

Connexins are a family of hemichannels that form gap junctions, whose diameter of the narrowest pore region is 1.4 nm. Among all connexin family members, connexin 43 (Cx43) was observed to play essential role in ATP release in leukocytes.^{95,96} For example, Cx43 mediates ATP release from macrophages during sepsis.⁹⁶

Another family of hemichannels, pannexins, does not form gap junctions but can allow molecules up to 1 kDa to permeate through them. Pannexins are a pathway of ATP release in various conditions, such as shear stress in leukocytes, immunogenic cell death with chemotherapeutics, and others.^{97–100}

CALHM1, a protein belonging to the CALHM family, is the only family member that forms a functional ion channel. This voltage- and extracellular calcium-sensitive and calcium-permeable ion channel regulates intracellular calcium homeostasis. It was also shown to allow the flow of ATP to the extracellular medium and to have a diameter at the narrowest region of its pore of 1.42 nm. Although its role in ATP release in airway epithelial cells, taste bud cells, and bladder has been observed, no role for CALHM1 in ATP release from immune cells has been described yet.^{93,94,101–104}

VRACs carry negatively charged organic and inorganic osmolytes to the extracellular medium. ATP, besides flowing through these channels, binds to them directly to trigger their activities. VRACs have a pore size

of 1.2–1.4 nm and are expressed in RAW 264.7 murine macrophages, which were observed to release ATP.^{93,94,105}

MACs are high-conductance, ATP-permeable, voltage-dependent channels that are expressed in all cells investigated so far, including T and B cells. These channels regulate cell volume and fluid secretion through the transport of chloride, anions, and ATP through their pores, whose diameter is estimated to range from 1.1 to 1.5 nm. These channels are expressed in cardiomyocytes, for example, where they mediate the release of ATP to the extracellular medium in a physiologically significant manner (at micromolar level). On the other hand, millimolar concentrations of extracellular ATP block these channels, in a kind of negative feedback.^{93,94,104,106–108}

Finally, as ATP can induce the opening of large P2X7-associated pores, it is also conceivable that ATP, as well as other nucleotides can leak out of the cells through these channels. Such a mechanism could create a positive-feedback mechanism in which the release of intracellular ATP could trigger the release of even more ATP from the cytosol of the same cell and activate a similar process in neighboring cells. This process may be involved in the propagation of intracellular calcium waves observed in microglia and mast cells.¹⁰⁹ However, this positive feedback mechanism could also lead to the sustained opening of the P2X7 pore and cell death.

3.1.5 | Lytic release

Another important source of extracellular nucleotides comes from injured cells that undergo irreversible cell membrane damage. This may occur in different types of tissue trauma and may represent an important stimulus triggering the activation of P2, and P1, receptors in endothelial cells, platelets, polymorphonuclear cells, and other leukocytes. Extracellular nucleotides constitute a “find me” signal, via activation of purinergic receptors (e.g., P2Y₂), to attract phagocytes (monocytes, macrophages, dendritic cells [DCs], and neutrophils) and promote the clearance of necrotic or apoptotic cell bodies.^{72,110}

3.1.6 | Other sources

Other nonlytic ATP release mechanisms, independent of P-glycoprotein and CFTR, have been described,¹¹¹ including some triggered by cell volume alterations and mechanical stimuli.^{112,113} An example is the release of ATP from blood vessel endothelial cells by shear stress. The autocrine action of ATP leads to NO release and consequent smooth muscle relaxation. Another example, described by Lazarowski et al.,¹¹⁴ is a CFTR-independent, mechanically induced release of cytoplasmic UTP in the human 1321N1 astrocytoma. In both cases, the mechanism of ATP and UTP release is still unknown.

3.2 | Physiologic removal of purinergic agonists

Regardless of the primary source of nucleotides and nucleosides, ectoenzymes quickly alter the final composition of the extracellular

milieu. Therefore, the final physiologic effect of nucleotide release depends on the interplay between the agonists, these enzymes, and the subtypes of P2 receptors expressed on the surface of the cells. Figure 1 summarizes the main nucleotide release pathways and the action of these molecules on purinergic receptors.

There are at least four families of ectonucleotidases: (a) ectonucleoside triphosphate phosphohydrolase (E-NTPDase), which hydrolyze tri and diphosphate nucleotides such as ATP and ADP into adenosine monophosphate (AMP); (b) ectonucleotide pyrophosphate/phosphodiesterase (E-NPP), which perform hydrolysis of pyrophosphate and phosphodiester bonds, thereby converting ATP or ADP to AMP; (c) alkaline phosphatases, which cleave tri, di, or monophosphate nucleotides into nucleoside and phosphate molecule; and (d) ecto-5'-nucleotidases, which convert monophosphate nucleotides (e.g., AMP) to ADO.^{115,116}

In addition, the resulting ADO molecules can be further deaminated into inosine and, subsequently, hypoxanthine, by reactions mediated by adenosine deaminase (ADA) and purine nucleotide phosphorylase (PNP), respectively.^{115,116} Table 3 summarizes the main enzymes responsible for the cleavage of extracellular nucleotides and their degradation products.

4 | P2 RECEPTORS IN DIFFERENT COMPONENTS OF THE IMMUNE SYSTEM

A specific set of P2 subtypes are expressed in leukocytes, which sometimes act synergistically, inducing the secretion of inflammatory mediators, stimulating chemotaxis, and activating signaling pathways such as the MAPK. In the following sections, we will describe the main roles of P2 receptors in these cells.

4.1 | Mononuclear phagocyte system (MPS) cells

4.1.1 | Monocytes

Monocytes are derived from the bone marrow and are present in blood. These cells recognize damage-associated molecular patterns (DAMPs) and pathogen-associated molecular patterns (PAMPs), triggering a series of responses that include phagocytosis, antigen presentation, chemokine secretion, and migration to tissues, where they differentiate into macrophages and DCs.¹¹⁷

ATP acts as an immunomodulating agent in monocytes. ATP activation inhibits human leukocyte antigen G (HLA-G) and IL-10 secretion in lipopolysaccharide (LPS)-activated human monocytes in vitro.¹¹⁸ On the other hand, ATP stimulation after LPS treatment induces IL-1 β release from monocytes and potentiates cytokine release from cells stimulated with different PAMPs and DAMPs. Additionally, several PAMPs and DAMPs stimulate ATP release with autocrine effects.¹¹⁹ UDP, another nucleotide, stimulates macrophage inflammatory protein-3 α (MIP-3 α) release from human monocytes from peripheral blood.¹²⁰




FIGURE 1 ATP signaling in the immune system cells. ATP is stored in cells within intracellular vesicles, and thus may be released via exocytosis, through membrane pores such as connexin and pannexin-1, and by dying cells. The endogenously released ATP provides autocrine and paracrine signaling, acting upon nearby P2 receptors (P2R). ATP is also degraded by soluble and membrane-associated ectonucleotidases such as ecto-apyrases (CD39) and ecto-5'-nucleotidases (CD73), originating ADP/AMP and ADO, respectively. ADO, in turn, acts upon P1 receptors (P1R)

TABLE 3 Main ectonucleotidases and their products

Enzyme	The final product(s)
E-NTPDase	ATP → ADP + Pi ADP → AMP + Pi
E-NPP	ATP → AMP + PPi
Alkaline phosphatase	ATP → ADP + Pi ADP → AMP + Pi AMP → ADO + Pi
Ecto-5'-nucleotidase	AMP → ADO + Pi

References 115 and 116.

P2 receptors expressed in human monocytes are involved in intracellular calcium mobilization,^{121,122} and P2X7 activation induces the formation of a membrane pore permeable to large solutes. However, in contrast to what is observed in macrophages, this pore only allows dye uptake when the extracellular medium contains low sodium and chloride levels.¹²³

P2X7 involvement has been demonstrated in several immunologic responses mediated by monocytes. P2X7 activation induces surface CD86 expression in human monocytes, whereas P2X7 blockade diminishes its expression.¹²⁴ In patients with sepsis, an increase in P2X7 expression in monocytes and its activation could be implicated in mitochondrial depolarization.¹²⁵ Patients with Behçet's disease (BD),

an immune-inflammatory syndrome, present higher levels of P2X7 on the surface of monocytes, despite no difference in P2X7 mRNA expression, compared with controls. Moreover, in monocytes from BD patients, ATP triggers augmented calcium influx, pore formation, and IL-1 β release.¹²⁶ In contrast, monocytes in primary Sjögren's syndrome patients, also express significantly higher levels of P2X7 than those from control individuals, but, in this case, higher levels of P2X7 mRNA are also observed.¹²⁷

A summary of the P2 receptors expressed in monocytes and their functions is shown in Table 4.

4.1.2 | Macrophages

Macrophages are phagocytic cells that are present in tissues and body cavities even in the absence of inflammation, where they contribute to surveillance and protection against infection. Their functions include phagocytosis of pathogens, toxins, and dead cells; cytokine and chemokine secretion; activation and recruitment of leukocytes to injured tissue; antigen presentation to T cells; and oxygen and nitrogen reactive species formation.¹¹⁷

Most of the information about the effects of ATP in immune system cells was obtained in macrophages. In 1985, Sung et al.¹³⁸ demonstrated that exogenous ATP interferes with transmembrane ion flux and inhibits phagocytosis in mouse macrophages. Afterward,

TABLE 4 P2 receptors in monocytes

Cell source	P2 receptor	Effects	References
Human monocytes	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2X ₁ , P2X ₄ , and P2X ₇	Calcium mobilization, ↑CD86, and cyclooxygenase-2 (COX-2) expression, CCL20, HLA-G, IL-1β, and IL-10 release, MAPK and NF-κB activation, and pore formation	118,120-122,124,128-131
THP-1 cell line	P2X ₁ , P2X ₂ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X _{7a}	Calcium mobilization, chemotaxis, dye uptake, IL-1β, and IL-8 release, and ROS formation	119,132-135
U937 cell line	Some P2X and P2Y receptors	Calcium mobilization, ERK1/2 phosphorylation, and IL-8 release	136,137

^aOnly the expression of P2X receptors was investigated.

several research groups showed that ATP induces plasma membrane permeabilization to fluorescent dyes up to 900 Da as well as a non-selective plasma membrane conductance in macrophages, which can be observed through the patch-clamp technique in whole-cell configuration.¹³⁹⁻¹⁴² Both these effects were concentration and temperature dependent and were modulated by Mg²⁺ and blocked by P2X₇ antagonists.¹⁴³⁻¹⁴⁵ Due to its role in ATP-triggered permeabilization, P2X₇ is the most studied purinergic receptor in macrophages. This receptor was first named P2Z, but in the 1990s, it was named P2X₇ according to cloning and homology data.²⁹ A number of effects of ATP on macrophages have been attributed to P2X₇ activation, either in studies using specific antagonists or KO mice. Despite the importance of P2X₇ KO mice in discovering new functions associated with this receptor, some of these models display incomplete inactivation of the P2X₇ gene. As a result, the Glaxo P2X₇^{-/-} mouse lines display a functional P2X₇ receptor in T cells; and the Pfizer P2X₇^{-/-} mice express a P2X₇-like protein in the brain (reviewed in¹⁴⁶).

ATP and BzATP activate caspase-1 and MEK1/2, and promote IFN-β expression in macrophages. ATP-activated macrophages secrete several cytokines, including TNF-α, keratinocyte chemoattractant (KC), IL-1β, and macrophage inflammatory protein 2 (MIP-2), as well as eicosanoids, CD14, and cathepsin.¹⁴⁷⁻¹⁵⁴ ATP also promotes reactive oxygen species (ROS) production, cell migration, and the shedding of microvesicles containing IL-18.^{150,152,155} All of those effects are significantly inhibited by P2X₇ blockage or KO.^{147-150,152,156,157}

ATP exposure can induce lytic cell death in human macrophages, and this effect is enhanced by IFN-γ treatment. On the other hand, ATP-mediated cytotoxicity is reverted by granulocyte-macrophage colony-stimulating factor (GM-CSF) and P2X₇ antagonists.^{80,158,159}

Murine peritoneal macrophages also form multinucleated cells upon GM-CSF cytokine stimulation. This effect can be inhibited by the P2X₇ antagonists such as oxidized ATP (oATP) and A740003, by P2X₇ KO, or by pannexin-1 blockage, suggesting the participation of P2X₇ and pannexin-1 in cell fusion.¹⁶⁰

P2X₄ is another receptor that plays important role in macrophages. P2X₄ activation was connected to macrophage recruitment to the peritoneal cavity and inflammatory response, since it induces cytosolic phospholipase A₂ (PLA₂) activation and prostaglandin E₂ (PGE₂) release.^{161,162} P2X₄ seems to act together with P2X₇ in regulating macrophages, through facilitating cytokine release and suppressing

autophagy during inflammation.^{163,164} Recently, it was demonstrated that both apyrase and P2X₄/P2X₇ antagonists significantly diminish macrophage phagocytic activity, in an extracellular calcium-dependent manner, suggesting their role in paracrine and autocrine stimulation of macrophage phagocytosis.¹⁶⁵

Macrophages also express P2Y receptors, which mediate LPS-induced expression of iNOS and cytokine release such as IL-6, MIP-2, and TNF-α in murine macrophages.¹⁶⁶⁻¹⁶⁹ P2Y receptors also augment phagocytosis and endocytosis, chemotaxis, and induce PGE₂ and leukotriene C₄ (LTC₄) synthesis.¹⁷⁰⁻¹⁷³ In a peritonitis murine model evoked by *Escherichia coli*, P2Y₆ activation resulted in monocyte chemoattractant protein-1 (MCP-1)-induced chemotaxis of both monocytes and macrophages, and in clearance of bacteria.¹⁷³

A summary of the P2 receptors expressed in various types of macrophages and their functions is shown in Table 5. Osteoclasts, a resident bone cell involved in bone resorption, are also from myeloid origin and can be derived from macrophages; therefore, they are also included in this summary.¹⁷⁴

4.1.3 | Microglia

Microglia represent the main immune effector cell population of the CNS.²⁰¹ Microglia are known to release various cytokines, NO, and ROS.²⁰²

ATP induces different effects on microglial cells, including ion channel opening, intracellular calcium mobilization, apoptosis, phagocytosis, NF-κB activation, ATP release, cytokine release, and chemokine secretion (e.g., CXCL2).²⁰³⁻²¹² Many of these effects have been attributed to P2X₇ activation.²¹³

P2X₇ activation triggers CXCL2 production, NFAT and MAPK pathway activation, cytokine release, and ROS production.²¹⁴⁻²¹⁶ LPS augments P2X₇ expression in microglia and contributes to its activation in rats.^{217,218}

P2X₇ was overexpressed in microglia of Alzheimer's disease (AD) patients. Amyloid β (Aβ)₁₋₄₂ peptide stimulates P2X₇ expression in fetal human microglia, suggesting an important role for this receptor in the development of AD.²¹⁹ P2X₇ silencing enhanced Aβ₁₋₄₂ accumulation at extracellular medium, by diminishing microglia phagocytosis in vitro, suggesting a protective role of P2X₇ in AD. P2X₇ blockage

TABLE 5 P2 receptors in macrophages

Cell source	P2 receptor	Effects	References
Human macrophages	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2X ₁ , P2X ₄ , P2X ₅ , and P2X ₇	Calcium mobilization, cell death, cathepsin release, chemokine and cytokine (CCL2, CCL4, CXCL5, IFN- γ , IL-1 β , IL-6, IL-23, TNF- α) release, microvesicle shedding, mycobacterium killing, and \downarrow leishmanial infection	151,153–155,158,175–179
Human alveolar macrophages	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₄ , P2X ₁ , P2X ₄ , P2X ₅ , and P2X ₇	Calcium mobilization, induction of membrane currents and cytokine (IL-1 β , IL-6, and TNF- α) secretion	170,180
Human monocyte-derived macrophages	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₃ , P2X ₁ , P2X ₄ , P2X ₅ , and P2X ₇	Calcium mobilization and CXCL5 release	154
Human monocyte-derived osteoclasts	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2X ₁ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X ₇	Multinucleated cells formation and \uparrow osteoclast resorption	181,182
Murine peritoneal macrophages	P2Y ₁ , P2Y ₂ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₄ , P2X ₁ , P2X ₄ , and P2X ₇	Calcium mobilization, cell death, chemotaxis, cytokine (IL-1 α , IL-1 β , IL-6, KC, TNF- α , MIP-2) release, induction of membrane currents, membrane permeabilization, microbicide actions, multinucleated cells formation, oleic acid (OA) and arachidonic acid (AA) release, phagocytosis and endocytosis, PGE ₂ and LTC ₄ synthesis, ROS production, and \downarrow leishmanial infection	150,152,156,157,160, 161,172,173,180,183–186
Murine bone marrow derived-macrophages (BMDMs)	P2Y ₁ , P2Y ₂ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₄ , and P2X ₇	Caspase-1 and MEK1/2 activation, cathepsin release, IFN- β expression, inflammatory mediators (IL-1 β , leukotriene B ₄ (LTB ₄), PGE ₂ , and thromboxane B ₂ (TXB ₂)) release, phagocytosis, and \uparrow TNF- α converting enzyme (TACE) activity	147–151,165,187
Mouse alveolar macrophages	P2X ₁ , P2X ₃ , P2X ₄ , P2X ₅ , and P2X ₇ ^a	Induction of membrane currents	188
Murine osteoclasts	P2Y ₁ , P2Y ₂ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₃ , P2X ₄ , P2X ₅ , and P2X ₇	\uparrow area resorbed per osteoclast and \uparrow osteoclast number	189
THP-1 derived macrophages cell line	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X ₇	IL-1 β release	190,191
RAW 264.7 cell line	P2Y ₁ , P2Y ₂ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₄ , P2X ₂ , P2X ₄ , and P2X ₇	Actin reorganization, cell death, chemotaxis, expression of proinflammatory mediators (COX-2, GM-CSF, high mobility group box 1 (HMGB1), IFN- β , IL-1 α , IL-1 β , iNOS, MCP-1, NO, PGE ₂ , and TNF- α), NF- κ B, MAPK, caspase-3, and Ras activation, NO and ROS production, membrane blebbing, and membrane permeabilization	149,165,173,192–196
J774 cell line	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2X ₁ , P2X ₂ , and P2X ₇	Calcium mobilization, induction of membrane currents, IL-1 β release, membrane permeabilization, and \downarrow bacilli Calmette-Guerin (BCG) viability	139–142,195,197,198
KUP5 cell line	P2Y ₂ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , P2X ₄ , and P2X ₇	IL-1 β and IL-6 release	199,200

^aOnly the expression of P2X receptors was investigated.

or KO also prevents IL-1 β and TNF- α released by microglia.^{220,221} A β_{1-42} aggregates stimulate murine primary microglia to release ATP and augment P2Y₂ expression in these cells. Both ATP and UTP induce cell migration and A β_{1-42} uptake and degradation, which were not observed in P2Y₂^{-/-} cells, suggesting an important role for this receptor in the modulation of AD.²²²

Uridine nucleotides induce important physiologic effects in microglia. UDP stimulation promotes actin polymerization in primary rat microglia and vasodilator-stimulated phosphoprotein (VASP) phosphorylation, suggesting a possible role of VASP in UDP-induced actin aggregation.²²³ P2Y₆ activation induces MCP-1 and macrophage inflammatory protein-1 α (MIP-1 α) secretion by rat primary microglia in a calcium-dependent manner.²²⁴ UTP is also involved in microglia phagocytosis, CCL2 expression and ERK1/2 phosphorylation in rat spinal microglia.^{207,225} P2Y₂ and P2Y₄ receptors also mediate particle pinocytosis in the presence of ATP and UTP.²²⁶

Other P2Y receptors exert important functions in microglia. P2Y₁₃ activation causes intracellular calcium mobilization in dorsal spinal cord microglia in a concentration-dependent manner.²²⁷ P2Y₁₂ and P2Y₁₃ augment IL-1 β , IL-6, and TNF- α release in microglia from the rat dorsal spinal cord.²²⁸ ADP and ATP stimulate process extension and adhesion in microglial cells from rats.^{229,230} Loss of P2Y₁₂ receptor diminishes the number of microglial projections and could be responsible for the conversion from ramified to amoeboid cell state. ADP also stimulates membrane ruffling and chemotaxis of rat and murine microglia. However, P2Y₁₂ expression in LPS-activated microglia in vivo was not observed.²³¹⁻²³⁴ Microglial P2Y₁₂ KO prevented neuronal cell death and reduced cell migration and NF- κ B expression in oxygen-glucose deprivation (OGD), an in vitro model of ischemia.²³⁵

A summary of the P2 receptors expressed in microglia and their functions are shown in Table 6.

In summary, the activation of P2 receptors in MPS cells results in proinflammatory responses. Both P2X and P2Y receptors induce the release of cytokines, chemokines, and the formation of inflammatory mediators, but many of these effects are attributed to P2X7 activation. Its activation further results in the formation of multinucleated giant cells (MGCs) by macrophages. On the other hand, migration and chemotaxis activities, as well as phagocytosis, endocytosis, and pinocytosis, seem to be more related to the activation of P2Y receptors, although P2X receptors can also mediate these effects.

4.2 | Dendritic cells

DCs act as sentinels of the immune system, and they are the most important APCs of the body, being responsible for the initiation of adaptive immune responses, inducing the activation of Th1, Th2, or Th17 profiles.²⁵¹

P2 receptors are implicated in the maturation of DCs through ATP stimulation.^{252,253} Human immature and mature DCs respond to ATP, UTP, and ADP through calcium mobilization. However, mature DCs seem to be less sensitive to UTP and ADP. All of these nucleotides induce actin polymerization and chemotaxis in immature DCs, but fail

to do so in mature cells, suggesting that these nucleotides have chemotactic activity on immature cells, attracting them to inflammatory sites.^{254,255}

Nucleotides increase the production of cytokines such as IL-6, IL-10, and IL-12.^{256,257} Moreover, P2 activation increases antigen presentation by DCs.²⁵⁸ Taken together, these data suggest that ATP analogs might be useful in the treatment of some infectious diseases. Supporting this idea, it was demonstrated that an ATP analog protects mice infected with gram-negative bacteria.²⁵⁹

The P2X7 receptor plays several roles in DCs, including intracellular calcium mobilization, endocytosis and apoptosis.²⁶⁰ This receptor is also involved in antigen presentation and IL-1 β release.^{258,261} In addition, P2X7 displays a crucial role in DC-mediated cancer control. Dying tumor cells release ATP, which acts on the P2X7 receptor in DCs and triggers the NOD-, LRR-, and pyrin domain-containing protein 3 (NLRP3) inflammasome, allowing the secretion of IL-1 β cytokine, which is important to control cancer growth and spread.²⁶²

Other P2 receptors are expressed in DCs and may be associated with different functions. P2Y receptors induce intracellular calcium mobilization and modulate cytokine expression and secretion.^{256,263,264} P2Y receptors could be implicated in human DC migration in peripheral blood, as well as in pinocytosis.²⁶⁵⁻²⁶⁷ P2Y activation also favors Th2 and Th17 responses.^{253,268,269}

A summary of the P2 receptors expressed in DCs and their functions are shown in Table 7.

In summary, the role of P2 receptors on DCs is associated with chemotaxis to inflamed tissues for the capture of antigens and their presentation to lymphocytes, initiating adaptive immune responses. Furthermore, high concentrations of ATP seem to decrease the release of proinflammatory cytokines and activation of P2Y receptors seems to modulate the Th2 or Th17 response profile.

4.3 | Lymphocytes

4.3.1 | T cells

T cells are a subpopulation of lymphocytes whose precursors originate in the bone marrow and differentiate in the thymus. These cells are responsible for antigen recognition after its presentation through sentinel cells via the HLA. Once activated (with the help of costimulatory signals), T cells stimulate IL-2 secretion, clonal expansion, and can differentiate into effector and memory cells. According to their subset, they have different roles in immunity, such as cytokine production, aiding B cells in antibody production (CD4⁺ cells), and destroying malignant or virus-infected cells (CD8⁺ cells).²⁷⁶

Nucleotides have important effects on T cells. It has long been known that ATP may induce the blastogenesis of murine medullar thymocytes.²⁷⁷ ATP also induces intracellular calcium mobilization, sustained depolarization of the plasma membrane, and membrane permeabilization to low molecular weight molecules (~300 Da). In contrast, cytotoxic T cells are resistant to the permeabilizing effects of ATP.²⁷⁸⁻²⁸⁰ These findings opened the possibility that ATP could be an

TABLE 6 P2 receptors in microglia

Cell source	P2 receptor	Effects	References
Human	P2Y ₁ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , P2X ₄ , and P2X ₇	Calcium mobilization and COX-2, IL-6, IL-1 β , IL-12, MCP-1, and TNF- α expression	236
Rhesus macaque	P2Y ₁ , P2Y ₂ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2X ₁ , P2X ₄ , P2X ₅ , and P2X ₇	Not determined	237
Rat	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , and P2Y ₁₄ ^a	Actin polymerization, adhesion, CCL2, CCL3, IL-1 β , IL-6 and TNF- α expression and release, chemotaxis, ERK1/2 and VASP phosphorylation, H ₂ O ₂ and nitrite production, membrane ruffling, \uparrow or \downarrow phagocytosis, and process extension	207,215,216,223,225,228,230-234,238-240
Mouse	P2Y ₁ , P2Y ₂ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₄ , and P2X ₇	2-Arachidonoylglycerol (2-AG) production, ATP, IL-1 β and TNF- α release, chemotaxis, membrane ruffling, and phagocytosis	222,231-234,236,241-243
BV-2 cell line	P2Y ₂ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , and P2Y _{14a}	Chemotaxis and CXCL2 production	214,234,244
C8B4 cell line	P2Y ₁ , P2Y ₂ , P2Y ₄ , and P2Y ₆ ^a	Calcium mobilization	245
N9 cell line	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₃ , P2X ₄ , P2X ₆ and P2X ₇	Not determined	246-248
GL261 glioma-derived CD11b ⁺ cells	P2Y ₁ , P2Y ₂ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₃ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X ₇	\downarrow cell proliferation and \downarrow IL-6 release	246
N13 cell line	P2X ₇ ^a	ATP and IL-1 β release, cell death, cell swelling, and dye uptake	249
MG-5 cell line	P2X ₇ ^b	CCL3 release	238
EOC13 cell line	P2X ₇ ^c	Cell death, pore formation, and ROS and NO production	250

^aOnly the expression of P2Y receptors was investigated. ^bReceptor expression was evaluated by functional assays. ^cOnly the expression of P2X₇ receptor was evaluated.

additional cytotoxic molecule used by T cells in lytic events,^{82,84,281} but this was refuted by others.²⁸²

ATP may induce differentiation, growth, or death of CD4⁺ T cells. It has been shown that whereas ATP concentrations below 50 nM in the extracellular milieu does not alter CD4⁺ T cells activity, 250 nM can trigger proliferation, and 1 mM can induce cell death (via P2X₄ and P2X₇ receptors). However, exposure to 1 mM of ATP can also increase proliferation and immunoregulatory effects of T regulatory cells.²⁸³ ATP may also control the migration of T cells within lymph nodes in homeostatic situations.²⁸⁴

Some ATP-mediated effects on T cells have been attributed to the expression of P2X₇ receptor.²⁸⁵ P2X₇ receptor was initially characterized in T cells by pharmacologic and physiologic means, as a P2X-activated channel was implicated in the mitogenic stimulation and cell death of human T cells.²⁸⁶⁻²⁹⁰ Later, it was demonstrated that CD4/CD8-defined single positive thymocytes express more P2X₇ receptor than double negative or double-positive cells.^{290,291}

FasL, an apoptosis inducer, stimulates ATP release via pannexin-1 hemichannels, and can induce P2X₇-mediated apoptosis.²⁹² Pannexin hemichannels also contribute to the activation of P2 receptors in T cells. Both murine CD4⁺ and CD8⁺ T cells express pannexins 1 and 2 in the plasma membrane, and after TCR activation, ATP is released from T cell via pannexin channels and may activate this cell autocrinally.^{293,294}

Besides cell death, P2X₇ has also been implicated in CD62L and CD27 shedding, lymphocyte proliferation, caspase-1 activation, and IL-1 β release.²⁹⁵⁻²⁹⁷ P2X₇ stimulation regulates T cell activation, initiating effector immunity.²⁹⁶ Also, the participation of P2X₇ and P2Y₆ receptors in T cell activation via TCR has been observed.²⁹⁸ Finally, studies with P2X₇ KO mice have revealed that P2X₇ activation contributes to $\gamma\delta$ T cell lineage commitment and the development of peripheral $\gamma\delta$ T cells.²⁹⁹

P2X₇ activation also seems to inhibit Treg function. P2X₇^{-/-} Tregs display lower levels of released ATP and phosphorylated ERK than

TABLE 7 P2 receptors in dendritic cells

Cell source	P2 receptor	Effects	References
Human from peripheral blood	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2X ₁ , P2X ₄ , and P2X ₇	Actin polymerization, chemotaxis, CCL20 release, ↓IL-12, ↓IL-23, ↑IL-1β, ↑IL-27, intracellular calcium mobilization, and NLRP3 inflammasome activation	120,254,260–262,265,270,271
Human monocyte-derived dendritic cells	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₄ , P2X ₁ , P2X ₄ , P2X ₅ , and P2X ₇	Intracellular calcium mobilization, dye uptake and CD23 shedding	264,272
Human plasmacytoid dendritic cells	P2Y ₄ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , and P2Y ₁₄ ^a	Intracellular calcium mobilization and IFN-α release	263
Human Langerhans cells	P2X ₇ ^b	Resistance to lytic effects of ATP and dye uptake	273
Murine phagocytic cells of the thymic reticulum	P2X ₇ ^b	Intracellular calcium mobilization	141,274
Murine bone marrow dendritic cells	P2X ₇ ^b	↑IL-1β, ↑IL-23, NLRP3 inflammasome activation, silica nanoparticles uptake, and ROS production	269,275
Murine D2SC/1 cell line	P2X ₇ ^c	Cell permeabilization and cell death	258,260

^aOnly the expression of P2Y receptors was evaluated. ^bReceptor expression was evaluated by functional assays. ^cOnly the expression of P2X₇ receptor was evaluated.

controls.³⁰⁰ P2X₇ activation in murine macrophages diminishes the expression of MHC I, and ATP treatment diminishes antigen presentation, reducing CD8⁺ T cell activation.³⁰¹

P2X₄ is involved in Th17 activation and its inhibition impacts IL-17 release. Since Th17 is involved in the development of autoimmune arthritis, P2X₄ blockade might alleviate the severity of this disease.³⁰²

P2Y₆ expression was described in T cells in inflammatory bowel disease (IBD).³⁰³ In an experimental model of colitis, P2Y₆^{-/-} animals showed an increase of cytokines involved in Th17 differentiation, such as IL-1β, IL-6, TGF-β1, IL-17, and IL-23. An increase of Th17 cells in the gut was also observed, suggesting that P2Y₆ has a protective role in intestinal inflammation.³⁰⁴

Besides P2X₇ and P2Y₆, the role of other P2 receptors on T cells has also been demonstrated. ADP induces intracellular calcium mobilization in T cells, while UDP-glucose slightly inhibits T cell proliferation.³⁰⁵

A summary of the P2 receptors expressed in T cells and their functions is shown in Table 8.

4.3.2 | B cells

B cells are a subgroup of lymphocytes developed in bone marrow, which are responsible for antibody production. These cells can recognize different antigens through the B cell receptor. Once matured, these cells differentiate into several subsets, including follicular, marginal zone, and B-1 B cells.^{276,313}

Extracellular ATP exerts many effects on B cells. In human B cells isolated from patients with chronic lymphocytic leukemia (CLL), ATP induces cation permeability, intracellular calcium mobilization, and membrane pore opening.^{314–317} Human B cells express functional P2X₇ receptors, although, in B cells, this receptor promotes less dye

uptake than in monocytes and macrophages, and the membrane pore present a lower molecular weight cut-off (~300 Da).²⁸⁰ This permeability response is further reduced in some B-CLL patients, suggesting that this receptor could be nonfunctional because of mutations.^{318,319}

B cells may modulate the expression of P2 receptors depending on the site or the situation (naive or activated). P2X₇ activation in human leukemic lymphocytes has been associated with stimulation of phospholipase D (PLD) activity and modulation of L-selectin and the low-affinity IgE receptor (named CD23) by interfering with different proteases, suggesting some importance of this receptor in cellular adhesive processes.^{320–322} P2X₇ receptor is also important to IgM release, lymphocyte proliferation, cell migration and shedding of CD21, CD23, and CD62L.^{323–326}

A summary of the P2 receptors expressed in B cells and their functions is shown in Table 9.

In summary, ATP activation of T cells promotes activation, proliferation, and differentiation of CD4⁺ T cells with the participation of the P2X₇ receptor. The activation of P2Y receptors can inhibit the proliferation of these cells and induce differentiation to the Th17 subtype. Meanwhile, in B cells, activation of P2 receptors is associated with adhesion, migration and IgM secretion.

4.4 | NK cells

NK cells are cytotoxic lymphocytes that do not carry classical markers for B or T lymphocytes. They are pivotal cells of the innate immunity, acting in the initial immune response, especially in virus infections, exerting direct cytotoxicity against infected cells, and producing cytokines to activate other cells. NK cells also appear to be involved in tumor surveillance, since NK deficiency can lead to increased

TABLE 8 P2 receptors in T cells

Cell source	P2 receptor	Effects	References
Human from peripheral blood	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2X ₁ , P2X ₄ , and P2X7	Caspase-1 activation, cell motility reduction, differentiation, IL-1 β release, and proliferation	297,303,306–308
Human CD4 ⁺ T cells	P2Y ₂ , P2X ₁ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X7	Cell death, differentiation, and growth	283,309
Jurkat cell line	P2X ₁ , P2X ₄ , and P2X7	Cell death	292,309,310
Murine T cells	P2Y ₆ , P2Y ₁₄ , and P2X7	CD27 and CD62L shedding, CD25 expression, cell death, IL-2 production, and inhibition of cell proliferation	287,295,305
Murine thymocytes	P2Y ₁ , P2Y ₂ , P2X ₁ , P2X ₂ , P2X ₅ , P2X ₆ , and P2X7	Apoptosis, CD27 shedding, differentiation and T cell activation	287,288,290,296,298,308,311
Murine splenic CD4 ⁺ and CD8 ⁺ cells	P2X7 ^b	Cell death and differentiation	291
Rat thymocytes	P2Y ₂ , P2X ₁ , and P2X4	Not determined	312

^aOnly the expression of P2X receptors was evaluated. ^bOnly the expression of P2X7 receptor was evaluated.

TABLE 9 P2 receptors in B cells

Cell source	P2 receptor	Effects	References
Human B cell	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₃ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X7	CD21, CD23, and CD62L shedding, and IgM release	324–328
Epstein-Barr immortalized B cells	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X7	Not determined	328
Human leukemic B lymphocytes	P2X7 ^a	Bromide ethidium uptake, phospholipase D activation and modulation of L-selectin and CD23	314–316,318,320–322,329
Mouse and rat bone marrow B cells	P2X7 ^a	Not determined	37
Murine splenic B cells	P2X7 ^a	CD23 shedding	325

^a Only the expression of P2X7 receptor was evaluated.

susceptibility to neoplasia. They are also able to kill different tumor cells in vitro.³³⁰

Besides a possible role as a cytotoxin secreted by NK cells, ATP seems to be involved in the modulation of NK cell responses. The first evidence of the presence of purinergic receptors in NK cells was reported in 1983 by Henriksson, who showed the inhibition of natural killing activity by ADO ribonucleotides; these findings were later reproduced and confirmed.^{331–333} It was postulated that this ATP-related activity was due to post recognition signaling events mediated by P2 receptors.^{332,334}

ATP also inhibits the proliferation of NK cells, probably acting through a lineage-specific receptor.³³³ The fact that UTP failed to affect NK cells, associated with a lack of inhibition of the effects of adenine nucleotides by pertussis toxin, may indicate the involvement of a P2X receptor, but this hypothesis lacks experimental confirmation.³³³ The possible influence of phosphorylation reactions

taking place on NK cell surfaces in these phenomena should not be discarded.³³⁵

Surprisingly, ATP inhibited NK cell migration in response to CX3CL1 and abolished the CX3CL1-dependent NK killing of endothelial cells. These effects were attributed to P2Y₁₁ according to pharmacologic blockers used to elucidate the receptor activation.³³⁶ Recently, it was demonstrated that P2Y₆ receptor expression diminishes NK cell maturation and activation. On the other hand, P2Y₆ deficiency increases the NK cytotoxicity and antimetastatic activities, suggesting a new niche for cancer therapy.³³⁷

Taken together, the findings suggest that, in NK cells, the activation of P2 receptors generates inhibitory responses on the proliferation and chemotaxis of these cells, in contrast to what was observed in other cells of the immune system.

A summary of the P2 receptors expressed in NK cells and their functions is shown in Table 10.

TABLE 10 P2 receptors in NK cells

Cell source	P2 receptors	Effects	References
Human blood	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₄ , P2X ₅ , P2X ₆ , and P2X ₇	↓ cell proliferation, ↓ chemotaxis, and ↓ killing activity	332–334,336

4.5 | Granulocytes

4.5.1 | Neutrophils

Neutrophils, also called polymorphonuclear neutrophils (PMN), are myeloid cells with multilobulated nuclei. They normally represent 70% of the leukocyte population, being the most abundant granulocytes. Neutrophils are the first cells to arrive at an inflammatory site to deal with tissue damage and microorganisms. Marked neutropenia can lead to overwhelming infection.³³⁸

Important neutrophil functions are migration from the blood to the injured tissue, the release of granule contents, and oxidative burst, which results in the production of free radicals.³³⁹ Over the last decades, a growing body of evidence indicates the action of purine nucleotides on these functions and in neutrophils' intracellular signaling pathways.

Several studies published in the 1980s demonstrated the effect of extracellular ATP on neutrophils. In these works, the authors showed that extracellular ATP and its analogs were able to increase both intracellular calcium and the respiratory burst after stimulation with formylated chemotactic peptides.^{21,340–343}

P2Y₂ is predominantly expressed in neutrophils. The activation of this receptor is involved in elastase release, cell degranulation, ROS production, MAPK phosphorylation, and intracellular calcium mobilization.^{344,345} P2Y₂ is also associated with neutrophil migration toward the IL-8 gradient.^{346–348} Epithelial cells incubated with peptides from human neutrophils or with nucleotides (ATP, ADP, UTP, and UDP) secreted the chemokine IL-8 in a concentration-dependent manner, suggesting the participation of P2Y receptors in this process.³⁴⁹ P2Y₄ and P2Y₆ could also contribute to UTP and UDP-induced neutrophil migration *in vivo*.^{350,351} In an experimental model of colitis in mice, intrathecal administration of UDP recruited neutrophils to the bowel, aggravating the symptoms of inflammation such as stool consistency, blood in stool, and rectal bleeding.³⁵¹

Furthermore, P2Y₆ activation is associated with neutrophil extracellular traps (NETs) formation, IL-8 release, migration, phagocytosis, superoxide production, and calcium mobilization.³⁵²

P2Y₁ and P2Y₁₄ receptors are involved to chemotaxis *in vitro* and neutrophil migration *in vivo*.^{353–356} This latter receptor also mediates elastase release.³⁵⁷ Some recent evidence suggests that P2Y₁₂ antagonists used in thrombosis prevention, such as clopidogrel, prasugrel, and ticagrelor, could modulate neutrophil functions *in vitro* and *in vivo*. However, this role remains uncharacterized.^{358–360}

P2X₁ is another receptor involved in the chemotaxis of neutrophils. Genetic deletion of P2X₁ slightly reduced neutrophil migration *in vitro*, but significantly reduced the migration of these cells to the peritoneum

in a peritonitis experimental model.³⁶¹ In addition, P2X₁ plays a protective role in endotoxemia, as P2X₁^{-/-} animals survived less than wildtype (WT) animals, and produced higher levels of ROS and neutrophil infiltration.³⁶² P2X₁^{-/-} animals also present lower neutrophil and fibrin accumulation at the injury site.³⁶³

P2X₇ activation in neutrophils causes cell depolarization, membrane pore opening, intracellular calcium mobilization, and secretion of IL-1 β in an NLRP3 inflammasome-dependent manner.^{364,365} The P2X₇ receptor is up-regulated in both blood and bronchoalveolar lavage fluid (BALF) neutrophils after exposure to cigarette smoke. Mice exposed to cigarette smoke have increased neutrophil counts and higher cytokine levels on BALF neutrophils compared with control animals, which may indicate a role of purinergic signaling in neutrophils during lung inflammation.³⁶⁶ In contrast, Martel-Gallegos et al.³⁶⁷ demonstrated through different techniques (patch-clamp, dye uptake, ROS production, RT-PCR, Western blotting, and immunofluorescence) that human neutrophils do not express P2X₇ receptor. This fact may suggest that neutrophils respond to ATP via another P2 receptor.³⁶⁷ In this sense, ATP could activate purinergic P2 receptors in other cell types, such as macrophages or fibroblasts, which could release chemokines to recruit neutrophils to the injury site.^{152,368}

A summary of the P2 receptors expressed in neutrophils and their functions is shown in Table 11.

4.5.2 | Mast cells

Mast cells can be found in almost all vascularized tissues of the body, especially in mucosa and epithelia. They are characterized by the presence, in their cytoplasm, of many metachromatic granules rich in chemical mediators, including histamine and heparin. Upon activation, mast cells not only release granule contents, but also begin the synthesis of other chemical mediators, such as arachidonic acid (AA) metabolites and cytokines.³⁷¹ Another characteristic of mast cells is the presence of a high-affinity receptor for IgE on their surface. Mast cells are involved in inflammation, anaphylactic reactions, and in some parasite infections.³⁷²

Mast cells were one of the first cells of the immune system to have the presence of purinergic receptors characterized, with the description of the effect of ATP causing permeabilization of the plasma membrane to large solutes, through the activation of the P2X₇ receptor.^{373,374} Since then, extracellular nucleotides were found to have a large number of biologic effects on mast cells. These include the release of histamine by mast cells upon stimulation with ATP, as well as ADP and guanosine triphosphate (GTP). However, this effect was observed only on rat and mouse mast cells, and not in human or

TABLE 11 P2 receptors in neutrophils

Cell source	P2 receptor	Effects	References
Human	P2Y ₁ , P2Y ₂ , P2Y ₁₄ , P2X ₁ , P2X ₄ , P2X ₅ , and P2X ₇ ?	Elastase release, intracellular calcium mobilization, MAPK phosphorylation, migration, NET formation, phagocytosis, and ROS production	344,348,352,357,365,369,370
Mouse	P2Y ₁ , P2Y ₄ , P2Y ₆ , P2Y ₁₄ , and P2X ₁	Migration	350,351,353,361
Rat	P2Y ₂ , P2Y ₁₁ , P2X ₁ , P2X ₄ , P2X ₅ , and P2X ₇	Migration	356
HL-60 cell line	P2Y ₂ , P2Y ₁₄ , and P2X ₇	Chemotaxis and dye uptake	354,364

guinea pig cells.^{375,376} On the other hand, ATP, 2MeSATP, and UTP enhanced the anti-IgE-induced histamine release by human lung mast cells.³⁸⁵

P2X and some P2Y receptors are associated with an increase of intracellular calcium levels, suggesting a role for these receptors in calcium mobilization and mast cell degranulation.^{109,377–379} Conversely, histamine induces ATP release to the extracellular medium via pannexin-1, which could represent a positive feedback response.³⁸⁰

P2 receptors seem also to be involved in triggering other mast cell responses, such as differentiation to mucosal and serosal phenotypes and chemoattraction.^{375,381} Indeed, ADP, ATP, and UTP are effective chemoattractants, apparently through the activation of P2Y receptors.³⁸² These nucleotides were also reported to inhibit TNF- α generation dependent of TLR2 ligand, and abolish TNF- α , IL-8, and MIP-1 assembly in response to LTB₄.³⁸³ Finally, ATP induces mast cell apoptosis, cytokine production (IL-4, IL-6, IL-13, and TNF- α), and is involved in mast cell-induced inflammation.^{384,385}

A summary of the P2 receptors expressed in mast cells and their functions is shown in Table 12.

4.5.3 | Eosinophils

Eosinophils are granulocytes involved mainly in the inflammatory response to allergens and infection with helminthic parasites. They are characterized by cytoplasmic granules containing lysosomal hydro-lases, as well as cationic proteins.³⁹⁰

The first work that suggested the expression of P2 receptors in eosinophils was published by Dichmann et al. in 2000, describing ATP-induced ROS generation, CD11b up-regulation, calcium mobilization, and actin polymerization.³⁹¹ Since then, extracellular nucleotides, such as ATP, UTP, and UDP, were found to induce IL-8 and eosinophil cationic protein (ECP) release by eosinophils. The involvement of P2Y receptors in granule release is suggested by the fact that this effect was blocked by pertussis toxin. On the other hand, IL-8 release was blocked by KN-62, indicating P2X involvement.³⁹² Among P2 receptors, P2Y₂ was the main receptor linked to chemotaxis in vitro and migration of eosinophils in vivo.³⁹³

In an asthma animal model, treatment with KN-62, a P2X₇ receptor antagonist, significantly inhibited eosinophilia as well as lymphocytosis in bronchoalveolar lavage (BAL). On the other hand, the same response was observed using P2X₇^{-/-} animals. Considering the incomplete inactivation of the P2X₇ gene observed in some P2X₇ KO models (reviewed in¹⁴⁶), these apparently contradictory results do not allow a clear definition of the role, or lack of it, for P2X₇ in eosinophilia in asthma models. Indeed, an up-regulation of P2X₇ expression in asthmatic patients compared to healthy individuals has been reported, as well as a slight increase in oxygen radical production by these cells, although the number of individuals evaluated was small ($n = 8$).³⁹⁴ In contrast, some groups did not observe the expression of P2X₇ in asthmatic patients.^{369,395}

P2Y₁₄ is another P2 receptor overexpressed in eosinophils, especially in the asthma context, and is also involved in chemotaxis of eosinophils to airway.³⁹⁶

A summary of the P2 receptors expressed in eosinophils and their functions is shown in Table 13.

Despite their important effects on allergic diseases, there is still much to be studied regarding purinergic signaling in this cell type. In addition, there are no studies on the possible role of purinergic signaling in helminthic diseases, which may represent a new direction for research.

4.5.4 | Basophils

Basophils are the least abundant granulocyte found in peripheral blood. These cells contain histamine-rich cytoplasmic granules and are the circulating counterparts of mast cells, apparently having similar properties and functions.³⁹⁹

Despite their similarity to mast cells, research on basophils is still scarce, including in the field of purinergic receptors. Nevertheless, UDP stimulation has been reported to augment intracellular calcium mobilization and IgE-dependent degranulation in basophils.^{400,401}

A summary of the P2 receptors expressed in basophils and their functions is shown in Table 14.

TABLE 12 P2 receptors in mast cells

Cell source	P2 receptors	Effects	References
Human lung	P2Y ₁ , P2Y ₂ , P2X ₁ , P2X ₂ , and P2X ₇	Enhance the anti-IgE-induced histamine	386,387
Cord blood derived-human mast cell	P2Y ₁ , P2Y ₂ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2X ₁ , and P2X ₄	Reduce cytokine production (TNF- α , IL-8, and macrophage inflammatory protein-1 β (MIP-1 β))	383
Murine peritoneal mast cells	P2X ₁ , P2X ₃ , P2X ₄ , and P2X ₇ ^a	Intracellular calcium mobilization, induction of membrane currents, histamine, and chemokine release	388
Bone marrow-derived mast cells (BMMC)	P2Y ₁ , P2Y ₄ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₃ , P2X ₄ , P2X ₆ , and P2X ₇	Apoptosis, calcium influx, cell permeabilization, and cytokine production (IL-4, IL-6, IL-13, and TNF- α)	384,389
LAD2	P2Y ₁ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , and P2X ₇	Cell degranulation	109,379
MC/9	P2Y ₁ , P2Y ₆ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₄ , P2X ₆ , and P2X ₇	Apoptosis, calcium influx, cell permeabilization, and cytokine production (IL-4, IL-6, IL-13, and TNF- α)	384
P815	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , P2Y ₁₄ , P2X ₁ , P2X ₂ , P2X ₃ , P2X ₄ , P2X ₅ , P2X ₆ and P2X ₇	Apoptosis, calcium influx, and cell permeabilization	384
RBL-2H3	P2Y ₁ , P2Y ₂ , P2Y ₁₃ , and P2Y ₁₄ ^b	Cell degranulation	377,378

^aOnly the expression of P2X receptors was investigated.

^bOnly the expression of P2Y receptors was investigated.

TABLE 13 P2 receptors in eosinophils

Cell Source	P2 receptors	Effects	References
Human	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₁ , P2X ₁ , P2X ₄ , and P2X ₇	Actin polymerization, CD11b up-regulation, chemotaxis, IL-8 and ECP release, and ROS generation	369,391,392,394,395,397,398
Rat	P2Y ₁ , P2Y ₂ , P2Y ₄ , P2X ₁ , P2X ₂ , P2X ₄ , and P2X ₇	Chemotaxis and cell migration	393

TABLE 14 P2 receptors in basophils

Cell Source	P2 receptors	Effects	References
Human	P2Y ₂ , P2Y ₄ , P2Y ₆ , P2Y ₁₂ , P2Y ₁₃ , and P2Y ₁₄ ^a	Intracellular calcium mobilization and IgE-dependent degranulation	401

^aOnly the expression of P2Y receptors was investigated.

5 | ROLE OF P2 RECEPTORS IN DISEASES AND THERAPEUTIC PERSPECTIVES

The data discussed here indicates that P2 receptors are widely expressed in the cells of the immune system, where they play important roles in innate immune response, as well as in the initiation of adaptive immune response. However, the exacerbation of proinflammatory responses promoted by P2 receptors also seems to contribute to the development of inflammatory diseases and pain. In this scenario, the use of selective agonists and antagonists of P2 receptors can significantly contribute to the management of these conditions.

5.1 | Infectious diseases

A potential therapeutic use of P2 agonists in the treatment of some infectious diseases has emerged from the studies of their effects on the immune system. Macrophages are capable of killing intracellular *Mycobacterium tuberculosis* or BCG through a mechanism dependent on nitrogen and oxygen free radicals activated by P2 receptors.^{81,402,403} ATP induces apoptosis and autophagy of mycobacterium-infected macrophages, reducing BCG viability.^{176,177,404} Another ATP-activated pathway that reduces BCG viability is associated with PLD activity and phagosome-lysosome fusion.¹⁹⁷




FIGURE 2 Main roles of P2 receptors expressed on leukocytes in infection. At the site of infection, cells release higher ATP concentrations, which functions as a “find me” signal to activate leukocytes. Extracellular ATP activates P2 receptors expressed on leukocytes, promoting cytokines production such as IFN- γ and TNF- α , phagocytosis, and the production of NO and ROS. Those, together with cell apoptosis caused by P2X7 activation, mediate both extracellular and intracellular pathogen killing. The activation of P2 receptors, especially P2Y, stimulates chemokine release, production of adhesion molecules and augments vascular permeability, favoring chemotaxis of leukocytes to the site of infection.

P2X7 KO increases mycobacterial viability, suggesting that P2X7 is important to protect cells from mycobacterial infection.¹⁹⁷ P2X7 receptor could also play a dual role in tuberculosis depending on the mycobacterium strain. P2X7 activation contributes to the resolution of tuberculosis in mice infected with laboratory strain, but it could aggravate the disease in mice infected with a hypervirulent strain.^{405,406}

Polymorphisms in the P2X7 genes have been associated with loss of macrophage activity and the development of tuberculosis.⁴⁶⁷⁻⁴⁶⁹ Individuals presenting the rs3751143 polymorphism, in which Glu496 is exchanged to Ala, have reduced P2X7 expression and decreased apoptosis when infected with BCG, favoring mycobacteria survival.⁴⁰⁷ The same effect is observed in individuals presenting heterozygosity in polymorphism Thr357 to Ser.⁴⁰⁸

Inflammasome activation is another strategy to fight pathogens. The inflammasome is a multiprotein complex associated with inflammation and cell death. It is composed of three proteins: a molecular pattern sensing protein, (generally NLRP3), the ASC adapter protein, and the pro-caspase-1 enzyme, which is cleaved and activated upon integration into the inflammasome complex. Active caspase-1 promotes the activation of proinflammatory cytokines IL-1 β and IL-18, and may also mediate pyroptosis cell death, contributing to pathogen killing.⁴⁰⁹⁻⁴¹¹

Some studies have demonstrated that infected macrophages and DCs use this strategy in infection control.⁴¹²⁻⁴¹⁴

In chlamydia infection, P2X7-induced apoptosis, as well as membrane permeabilization and calcium mobilization, is inhibited in macrophages. In turn, the treatment of macrophages with ATP reduces infection levels, via PLD activation and vesicle fusion.⁴¹⁵

Similarly to the P2X7 receptor, P2Y₂ activation plays a role in microbial elimination. P2Y₂ activation augments the acidification of mycobacterial phagosomes in a calcium-dependent manner, killing the mycobacteria without causing macrophage death. This effect was mediated by the P2Y₂ receptor even in the absence of P2X7.⁴¹⁶

ATP treatment significantly reduces infection with *Toxoplasma gondii* in macrophages from swiss, C57Bl/6, and Balb/c mice strains. Brilliant blue G treatment or P2X7 KO abolished this protection, suggesting a role for P2X7 in *T. gondii* elimination.¹⁸⁶ UTP and UDP treatment, in addition to ATP treatment, in peritoneal macrophages infected with *T. gondii* also reduced the percentage of infected cells and the number of parasites per cell in a concentration-dependent manner, suggesting the participation of P2Y receptors in this effect.⁴¹⁷

Several P2 receptors are involved in the reduction of *Leishmania amazonensis* infection of macrophages by nucleotide treatment. The microbicidal actions of P2X7 on macrophages against *L. amazonensis*




FIGURE 3 Schematic illustration of the P2 receptors in pain. At inflamed tissue, immune cells, including macrophages, neutrophils, mast cells, and T cells, are activated by extracellular nucleotides. Once activated, these cells release several cytokines, including IL- 1β , IL-6, and TNF- α , as well as prostaglandins, histamine, and other inflammatory mediators, contributing to sensitization of nociceptor sensory neurons via ion channels, cytokine receptors and GPCRs, and promoting pain sensation

involve the production of lipid mediators and LTB₄ secretion.^{418–420} Among P2Y receptors, P2Y₂ may be one of the most important to control infection, since inhibition of this receptor augments it. UTP stimulation induces ATP and LTB₄ release, contributing to autocrine signaling and parasite death, respectively.^{184,185,421} Figure 2 illustrates the role of P2 receptors expressed on leukocytes in infectious diseases.

Recently, several studies have demonstrated the benefits of blocking P2 receptors in infectious diseases such as acquired immunodeficiency syndrome. Hazleton et al.⁴²² demonstrated that oATP and Suramin inhibited human immunodeficiency virus (HIV) p24 production and release as well as the percentage of infected cells. These authors suggested the participation of at least three receptors in these processes: P2Y₁, P2X₁, and P2X₇.⁴²² Giroud et al.⁴²³ also observed that P2X₁ antagonists blocked virus fusion.

5.2 | Inflammatory diseases

Nucleotides act as a “find me” signal to leukocytes, stimulating their migration to inflammatory sites. P2 receptors are activated as a

result of the accumulation of extracellular nucleotides at these sites and promote the formation and release of inflammatory mediators, such as cytokines, chemokines, ROS, among others, contributing to local inflammation, tissue damage, and pain sensation. In this scenario, P2 receptors emerged as a promising target for therapeutics in inflammatory diseases.^{110,424–426}

The participation of the P2X₇ in inflammatory diseases is related to its ability to activate the inflammasome, a target in the treatment of inflammatory disorders.⁴²⁶ Despite many studies, the exact mechanism by which P2X₇ promotes inflammasome activation is still unknown. Hypotheses proposed include a decrease in intracellular K⁺ concentration, ROS production, and the destabilization of lysosomes. This will likely be an important field of study in the coming years, as a strategy to identify new anti-inflammatory agents.^{409–411}

In animal models of arthritis, P2X₇ antagonists alleviate joint inflammation, including type II collagen-induced joint damage.^{306,427,428} P2X₇ KO mice developed fewer joint cartilage lesions than controls, and lower proteoglycan content and collagen degradation, reducing the severity of the disease.⁴²⁹ P2X₇ antagonists also inhibit the release of cathepsins (lysosomal proteases), suggesting a possible role in chronic inflammation.¹⁵¹

The P2Y₂ receptor role in cell recruitment to the inflammation site is well recognized.⁴³⁰ This receptor senses the nucleotide gradient, promoting neutrophil migration to the site of inflammation.^{431,432} Impairment of P2Y₂ activity could be beneficial in certain models of lung inflammatory disease. However, one exception is infection with the pneumonia virus of mice, as P2Y₂^{-/-} mice had lower counts of leukocytes in BALF, and had higher mortality than WT animals.⁴³³

P2Y₁ and P2Y₆ play important roles in vascular inflammation. P2Y₁ participates in vascular inflammation by recruiting leukocytes. P2Y₁^{-/-} mice present lower levels of adhesion molecules and of leukocytes. P2Y₁ blockage also reduced the number of rolling leukocytes in vivo.⁴³⁴ P2Y₆ overexpression was accompanied by the increased expression of VCAM-1 in LPS-injected animals.⁴³⁵ In ovalbumin (OVA) and house dust mite (HDM) models of allergic inflammation, P2Y₆ is up-regulated and modulates the classical allergic features such as eosinophilia, airway remodeling, Th2 cytokine secretion, and bronchial hyperresponsiveness, which was alleviated by the use of antagonists or in P2Y₆ receptor KO animals.^{436,437}

Regarding pain, P2X receptors seem to play a more important role (or their participation in pain processes is more studied) than P2Y receptors. P2X receptors promote the release of several inflammatory mediators, such as IL-1 β and prostaglandins, which are involved in inflammatory hyperalgesia.⁴³⁸

P2X_{2/3} activation is associated with inflammatory hyperalgesia induced by carrageenan, bradykinin, prostaglandins, and sympathomimetic amines, which could be reverted with the use of their specific antagonists.^{439–442} Moreover, P2X₃ and the heteromeric P2X_{2/3} were shown, by functional assays, to be key receptors involved in pain in primary sensory neurons, via PLA₂ activation.^{443–445} Recently, it was demonstrated that TNF- α released from macrophages induce P2X₃ expression in neurons, modulating allodynia.⁴⁴⁶ In turn, P2X₄ is related to mechanical allodynia after nerve injury.^{447–449} Additionally, P2X₄ is overexpressed in microglia after nerve injury and P2X₄^{-/-} mice have abolished tactile allodynia.^{447,448}

Selective blockers of the P2X₇ receptor diminished allodynia in different models of rat neuropathic pain.⁴⁵⁰ P2X₇ blockage relieved inflammatory pain and thermal hyperalgesia.^{451,452} KO of IL-1 $\alpha\beta$ in mice abolished the hyperalgesic effects of the complete Freund's adjuvant (CFA) model, indicating that hyperalgesia relief by the blockage of the P2X₇ is due to the blockage of IL-1 β .⁴⁵³

P2Y receptors also participate in neuropathic pain. P2Y₁₂ receptor inactivation, both pharmacologically and through receptor KO, abolished the pain after nerve injury.^{454,455} In contrast, other pharmacologic P2Y₁₂ inhibitors failed to block pain, indicating that the role of P2Y₁₂ in pain must be better explored. Figure 3 illustrates the role of P2 receptors activation in pain.

6 | CONCLUDING REMARKS

This review compiles broad evidence of the wide distribution of P2 receptors in cells of the immune system. In leukocytes, these receptors play important roles in the innate immune response and in the

initiation of the adaptive immune response, contributing significantly to pathogen death and infection control. However, the activation of these receptors is also related to the development of inflammatory diseases and pain, making them a promising target for the development of anti-inflammatory drugs and analgesics.

Purinergic therapy is still in its infancy, but it has several fields of application in which it can mature significantly. In this review we show that, over the approximately 50 years of the purinergic field, much has been discovered regarding the presence of P2 receptors and their roles in the organism; still, much more remains to be explored, especially concerning the immune system and CNS. Yet, the work on P2 receptor role in the immune system strongly suggests that they are a promising target for drug development over the next decades.

ACKNOWLEDGMENTS

This work was supported by Instituto Oswaldo Cruz (Fiocruz), Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). We thank Evelyne Araujo Dias and Filipe Faria Berçot by the design of the images. The authors dedicate this paper in honor of Prof. Geoffrey Burnstock, who pioneered the studies on purinergic receptors, and Prof. Pedro Persechini, who made great contributions to purinergic signaling in Brazil.

AUTHORSHIP

A. V. P. A., N. C. S. F., C. A., and L. A. A. conceived the idea. A. V. P. A., N. C. S. F., A. G. C. B., O. K. N., F. P. F., R. C. B., C. A., W. S., R. C. S., P. M. P., and L. A. A. wrote or contributed to the writing of the manuscript. A. V. P. A. and N. C. S. F. contributed equally to this work.

DISCLOSURE

The authors declare no conflict of interest.

REFERENCES

- Nicholson LB. The immune system. *Essays Biochem.* 2016;60:275–301.
- Fredholm BB, Abbracchio MP, Burnstock G, et al. Nomenclature and classification of purinoceptors. *Pharmacol Rev.* 1994;46:143–156.
- Haag F, Adriouch S, Braß A, et al. Extracellular NAD and ATP: partners in immune cell modulation. *Purinergic Signal.* 2007;3:71–81.
- Coddou C, Yan Z, Obsil T, et al. Activation and regulation of purinergic P2X receptor channels. *Pharmacol Rev.* 2011;63:641–683.
- Gallego D, Gil V, Aleu J, et al. Pharmacological characterization of purinergic inhibitory neuromuscular transmission in the human colon. *Neurogastroenterol Motil.* 2011;23:792–e338.
- Abbracchio MP, Burnstock G, Boeynaems J-M, et al. International Union of Pharmacology LVIII: update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy. *Pharmacol Rev.* 2006;58:281–341.
- Dubyak GR, El-Moatassim C. Signal transduction via P2-purinergic receptors for extracellular ATP and other nucleotides. *Am J Physiol Physiol.* 1993;265:C577–606.
- Ralevic V, Burnstock G. Receptors for purines and pyrimidines. *Pharmacol Rev.* 1998;50:413–492.
- Keller R. Tissue mast cells in immune reactions. *Monogr Allergy.* 1966;2:1–144.

10. Burnstock G. Purinergic nerves. *Pharmacol Rev.* 1972;24:509-581.
11. Cockcroft S, Gomperts BD. ATP induces nucleotide permeability in rat mast cells. *Nature.* 1979;279:541-542.
12. Cohn ZA, Parks E. The regulation of pinocytosis in mouse macrophages: II. Factors inducing vesicle formation. *J Exp Med.* 1967;125:213-232.
13. Burnstock G. A basis for distinguishing two types of purinergic receptor. In: Straub RW, Bolis L, eds. *Cell membrane receptors for drugs and hormones: A multidisciplinary approach.* Raven Press; 1978: 107-118.
14. Gordon JL. Extracellular ATP: effects, sources and fate. *Biochem J.* 1986;233:309-319.
15. O'Connor SE, Dainty IA, Leff P. Further subclassification of ATP receptors based on agonist studies. *Trends Pharmacol Sci.* 1991;12:P137-141.
16. Burnstock G. The past, present and future of purine nucleotides as signalling molecules. *Neuropharmacology.* 1997;36:1127-1139.
17. Burnstock G. Purine and purinergic receptors. *Brain Neurosci Adv.* 2018;2:239821281881749.
18. Burnstock G. P2X ion channel receptors and inflammation. *Purinergic Signal.* 2016;12:59-67.
19. Cowen DS, Lazarus HM, Shurin SB, et al. Extracellular adenosine triphosphate activates calcium mobilization in human phagocytic leukocytes and neutrophil/monocyte progenitor cells. *J Clin Invest.* 1989;83:1651-1660.
20. Dubyak GR, Cowen DS, Lazarus HM. Activation of the inositol phospholipid signaling system by receptors for extracellular ATP in human neutrophils, monocytes, and neutrophil/monocyte progenitor cells. *Ann N Y Acad Sci.* 1988;551: 218-237. discussion 237-8.
21. Dubyak GR, Cowen DS, Mueller LM. Activation of inositol phospholipid breakdown in HL60 cells by P2-purinergic receptors for extracellular ATP. Evidence for mediation by both pertussis toxin-sensitive and pertussis toxin-insensitive mechanisms. *J Biol Chem.* 1988;263:18108-18117.
22. Deaglio S, Robson SC. Ectonucleotidases as regulators of purinergic signaling in thrombosis, inflammation, and immunity. *Adv Pharmacol.* 2011;61:301-332.
23. Kukulski F, Lévesque SA, Sévigny J. Chapter 9 - Impact of ectoenzymes on P2 and P1 receptor signaling. In: Jacobson KA, Linden J, eds. *Advances in Pharmacology.* Academic Press; 2011:263-299.
24. Longhi MS, Robson SC, Bernstein SH, et al. Biological functions of ecto-enzymes in regulating extracellular adenosine levels in neoplastic and inflammatory disease states. *J Mol Med.* 2013;91:165-172.
25. Slakey LL, Gordon EL, Pearson JD. A comparison of ectonucleotidase activities on vascular endothelial and smooth muscle cells. *Ann N Y Acad Sci.* 1990;603:366-378. discussion 378-379.
26. Zimmermann H. Extracellular metabolism of ATP and other nucleotides. *Naunyn Schmiedeberg's Arch Pharmacol.* 2000;362: 299-309.
27. Zimmermann H. Extracellular purine metabolism. *Drug Dev Res.* 1996;39:337-352.
28. Cekic C, Linden J. Purinergic regulation of the immune system. *Nat Rev Immunol.* 2016;16:177-192.
29. Surprenant A, Rassendren F, Kawashima E, et al. The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X7). *Science.* 1996;272:735-738.
30. North RA. P2X receptors: a third major class of ligand-gated ion channels. *Ciba Found Symp.* 1996;198:91-105.
31. North RA. Families of ion channels with two hydrophobic segments. *Curr Opin Cell Biol.* 1996;8:474-483.
32. Brake AJ, Wagenbach MJ, Julius D. New structural motif for ligand-gated ion channels defined by an ionotropic ATP receptor. *Nature.* 1994;371:519-523.
33. Valera S, Hussy N, Evans RJ, et al. A new class of ligand-gated ion channel defined by P2x receptor for extracellular ATP. *Nature.* 1994;371:516-519.
34. Surprenant A, Buell G, Alan North R. P2X receptors bring new structure to ligand-gated ion channels. *Trends Neurosci.* 1995;18:224-229.
35. Burnstock G. Purine and pyrimidine receptors. *Cell Mol Life Sci.* 2007;64:1471-1483.
36. Buell G, Collo G, Rassendren F. P2X receptors: an emerging channel family. *Eur J Neurosci.* 1996;8:2221-2228.
37. Collo G, Neidhart S, Kawashima E, et al. Tissue distribution of the P2x7 receptor. *Neuropharmacology.* 1997;36:1277-1283.
38. Collo G, North RA, Kawashima E, et al. Cloning of P2x5 and P2x6 receptors and the distribution and properties of an extended family of ATP-gated ion channels. *J Neurosci.* 1996;16:2495-2507.
39. Lê KT, Boué-Grabot EÉ, Archambault V, et al. Functional and biochemical evidence for heteromeric ATP-gated channels composed of P2x1 and P2x5 subunits. *J Biol Chem.* 1999;274:15415-15419.
40. Royle SJ, Bobanović LK, Murrell-Lagnado RD. Identification of a non-canonical tyrosine-based endocytic motif in an ionotropic receptor. *J Biol Chem.* 2002;277:35378-35385.
41. Torres GE, Egan TM, Voigt MM. Hetero-oligomeric assembly of P2X receptor subunits. Specificities exist with regard to possible partners. *J Biol Chem.* 1999;274:6653-6659.
42. Nicke A, Kerschensteiner D, Soto F. Biochemical and functional evidence for heteromeric assembly of P2x1 and P2x4 subunits. *J Neurochem.* 2005;92:925-933.
43. Coddou C, Yan Z, Obsil T, et al. Activation and regulation of purinergic P2X receptor channels. *Pharmacol Rev.* 2011;63:641-683.
44. Kawate T, Michel JC, Birdsong WT, et al. Crystal structure of the ATP-gated P2x4 ion channel in the closed state. *Nature.* 2009;460:592-598.
45. Khakh BS, Bao XR, Labarca C, et al. Neuronal P2X transmitter-gated cation channels change their ion selectivity in seconds. *Nat Neurosci.* 1999 24. 1999;2:322-330.
46. Virginio C, MacKenzie A, Rassendren FA, et al. Pore dilation of neuronal P2X receptor channels. *Nat Neurosci.* 1999;2:315-321.
47. Hattori M, Gouaux E. Molecular mechanism of ATP binding and ion channel activation in P2X receptors. *Nature.* 2012;485:207-212.
48. Mansoor SE, Lü W, Oosterheert W, et al. X-ray structures define human P2x3 receptor gating cycle and antagonist action. *Nature.* 2016;538:66-71.
49. Kasuya G, Yamaura T, Ma X-B, et al. Structural insights into the competitive inhibition of the ATP-gated P2X receptor channel. *Nat Commun.* 2017;8:876.
50. Kasuya G, Fujiwara Y, Tsukamoto H, et al. Structural insights into the nucleotide base specificity of P2X receptors. *Sci Rep.* 2017;7:45208.
51. Karasawa A, Kawate T. Structural basis for subtype-specific inhibition of the P2x7 receptor. *Elife.* 2016;5:e22153.
52. McCarthy AE, Yoshioka C, Mansoor SE. Full-length P2x7 structures reveal how palmitoylation prevents channel desensitization. *Cell.* 2019;179:659-670.e13.
53. Keystone EC, Wang MM, Layton M, et al. Clinical evaluation of the efficacy of the P2x7 purinergic receptor antagonist AZD9056 on the signs and symptoms of rheumatoid arthritis in patients with active disease despite treatment with methotrexate or sulphasalazine. *Ann Rheum Dis.* 2012;71:1630-1635.
54. Stock TC, Bloom BJ, Wei N, et al. Efficacy and safety of CE-224,535, an antagonist of P2x7 receptor, in treatment of patients with rheumatoid arthritis inadequately controlled by methotrexate. *J Rheumatol.* 2012;39:720-727.
55. Illes P, Müller CE, Jacobson KA, et al. Update of P2X receptor properties and their pharmacology: iUPHAR Review 30. *Br J Pharmacol.* 2021;178:489-514.
56. Burnstock G. Purinergic signalling: past, present and future. *Brazilian J Med Biol Res.* 2009;42:3-8.

57. Burnstock G. Purinergic signalling: from discovery to current developments. *Exp Physiol*. 2014;99:16-34.
58. Erb L, Liao Z, Seye CI, et al. P2 receptors: intracellular signaling. *Pflugers Arch*. 2006;452:552-62.
59. Erb L, Garrad R, Wang Y, et al. Site-directed mutagenesis of P2U purinoceptors. Positively charged amino acids in transmembrane helices 6 and 7 affect agonist potency and specificity. *J Biol Chem*. 1995;270:4185-4188.
60. Jiang Q, Guo D, Lee BX, et al. A mutational analysis of residues essential for ligand recognition at the human P2Y1 receptor. *Mol Pharmacol*. 1997;52:499-507.
61. Hoffmann C, Moro S, Nicholas RA, et al. The role of amino acids in extracellular loops of the human P2Y1 receptor in surface expression and activation processes. *J Biol Chem*. 1999;274:14639-14647.
62. Zhang J, Zhang K, Gao ZG, et al. Agonist-bound structure of the human P2Y12 receptor. *Nature*. 2014;509:119-122.
63. Zhang K, Zhang J, Gao ZG, et al. Structure of the human P2Y12 receptor in complex with an antithrombotic drug. *Nature*. 2014;509:115-118.
64. Zhang D, Gao ZG, Zhang K, et al. Two disparate ligand-binding sites in the human P2Y1 receptor. *Nature*. 2015;520:317-321.
65. Jacobson KA, Paoletta S, Katritch V, et al. Nucleotides acting at P2Y receptors: connecting structure and function. *Mol Pharmacol*. 2015;88:220-230.
66. Jacobson KA, Delicado EG, Gachet C, et al. Update of P2Y receptor pharmacology: iUPHAR Review 27. *Br J Pharmacol*. 2020;177:2413-2433.
67. Erb L, Weisman GA. Coupling of P2Y receptors to G proteins and other signaling pathways. *Wiley Interdiscip Rev Membr Transp Signal*. 2012;1:789-803.
68. Jacobson KA, Boeynaems J-M. P2Y nucleotide receptors: promise of therapeutic applications. *Drug Discov Today*. 2010;15:570-578.
69. Bodin P, Burnstock G. Purinergic signalling: ATP release. *Neurochem Res*. 2001;26:959-969.
70. Fields RD. Nonsynaptic and nonvesicular ATP release from neurons and relevance to neuron-glia signaling. *Semin Cell Dev Biol*. 2011;22:214-219.
71. Lazarowski ER. Vesicular and conductive mechanisms of nucleotide release. *Purinergic Signal*. 2012;8:359-373.
72. Dosch M, Gerber J, Jebbawi F, et al. Mechanisms of ATP release by inflammatory cells. *Int J Mol Sci*. 2018;19:1-16.
73. Giuliani AL, Sarti AC, Di Virgilio F. Extracellular nucleotides and nucleosides as signalling molecules. *Immunol Lett*. 2019;205:16-24.
74. Felten DL, Felten SY, Carlson SL, et al. Noradrenergic and peptidergic innervation of lymphoid tissue. *J Immunol*. 1985;135:755s-765s.
75. Stevens-Felten SY, Bellingier DL. Noradrenergic and peptidergic innervation of lymphoid organs. *Chem Immunol*. 1997;69:99-131.
76. Phillips RJ, Powley TL. Macrophages associated with the intrinsic and extrinsic autonomic innervation of the rat gastrointestinal tract. *Auton Neurosci*. 2012;169:12-27.
77. von Kügelgen I, Starke K. Noradrenaline-ATP co-transmission in the sympathetic nervous system. *Trends Pharmacol Sci*. 1991;12:319-324.
78. Westfall DP, Sedaa KO, Shinozuka K, et al. ATP as a cotransmitter. *Ann N Y Acad Sci*. 1990;603:300-310.
79. Haskó G, Elenkov IJ, Vizi ES. Presynaptic receptors involved in the modulation of release of noradrenaline from the sympathetic nerve terminals of the rat thymus. *Immunol Lett*. 1995;47:133-137.
80. Blanchard DK, Hoffman SL, Djeu JY. Inhibition of extracellular ATP-mediated lysis of human macrophages by calmodulin antagonists. *J Cell Biochem*. 1995;57:452-464.
81. Ferrari D, Chiozzi P, Falzoni S, et al. Extracellular ATP triggers IL-1 beta release by activating the purinergic P2Z receptor of human macrophages. *J Immunol*. 1997;159:1451-1458.
82. Filippini A, Taffs RE, Sitkovsky MV. Extracellular ATP in T-lymphocyte activation: possible role in effector functions. *Proc Natl Acad Sci USA*. 1990;87:8267-8271.
83. Henney S. Studies on the mechanism of T lymphocyte-mediated cytotoxicity. *Int Arch Allergy Immunol*. 1973;45:281-284.
84. Apasov S, Koshiba M, Redegeld F, et al. Role of extracellular ATP and P1 and P2 classes of purinergic receptors in T-cell development and cytotoxic T lymphocyte effector functions. *Immunol Rev*. 1995;146:5-19.
85. Reising IL, Prats AG, Abraham EH, et al. The cystic fibrosis transmembrane conductance regulator is a dual ATP and chloride channel. *J Biol Chem*. 1994;269:20584-20591.
86. Abraham EH, Prat AG, Gerweck L, et al. The multidrug resistance (mdr1) gene product functions as an ATP channel. *Proc Natl Acad Sci USA*. 1993;90:312-316.
87. Sprague RS, Ellsworth ML, Stephenson AH, et al. Deformation-induced ATP release from red blood cells requires CFTR activity. *Am J Physiol*. 1998;275:H1726-1732.
88. Prat AG, Reising IL, Ausiello DA, et al. Cellular ATP release by the cystic fibrosis transmembrane conductance regulator. *Am J Physiol*. 1996;270:C538-545.
89. Schwiebert EM, Egan ME, Hwang TH, et al. CFTR regulates outwardly rectifying chloride channels through an autocrine mechanism involving ATP. *Cell*. 1995;81:1063-1073.
90. Watt WC, Lazarowski ER, Boucher RC. Cystic fibrosis transmembrane regulator-independent release of ATP. Its implications for the regulation of P2Y2 receptors in airway epithelia. *J Biol Chem*. 1998;273:14053-14058.
91. Jiang Q, Mak D, Devidas S, et al. Cystic fibrosis transmembrane conductance regulator-associated ATP release is controlled by a chloride sensor. *J Cell Biol*. 1998;143:645-657.
92. Reddy MM, Quinton PM, Haws C, et al. Failure of the cystic fibrosis transmembrane conductance regulator to conduct ATP. *Science (80-)*. 1996;271:1876-1879.
93. Taruno A. ATP release channels. *Int J Mol Sci*. 2018;19:808.
94. Sabirov RZ, Okada Y. ATP release via anion channels. *Purinergic Signal*. 2005;1:311-328.
95. Eltzhig HK, Eckle T, Mager A, et al. ATP release from activated neutrophils occurs via connexin 43 and modulates adenosine-dependent endothelial cell function. *Circ Res*. 2006;99:1100-1108.
96. Dosch M, Zindel J, Jebbawi F, et al. Connexin-43 dependent ATP release mediates macrophage activation during sepsis. *Elife*. 2019;8:e42670.
97. Barbe MT, Monyer H, Bruzzone R. Cell-cell communication beyond connexins: the pannexin channels. *Physiol*. 2006;21:103-114.
98. Bruzzone R, Hormuzdi SG, Barbe MT, et al. Pannexins, a family of gap junction proteins expressed in brain. *Proc Natl Acad Sci USA*. 2003;100:13644-13649.
99. Locovei S, Bao L, Dahl G. Pannexin 1 in erythrocytes: function without a gap. *Proc Natl Acad Sci USA*. 2006;103:7655-7659.
100. Bao L, Locovei S, Dahl G. Pannexin membrane channels are mechanosensitive conduits for ATP. *FEBS Lett*. 2004;572:65-68.
101. Taruno A, Vingtdoux V, Ohmoto M, et al. CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and umami tastes. *Nature*. 2013;495:223-226.
102. Sana-Ur-Rehman H, Markus I, Moore KH, et al. Expression and localization of pannexin-1 and CALHM1 in porcine bladder and their involvement in modulating ATP release. *Am J Physiol Regul Integr Comp Physiol*. 2017;312:R763-R772.
103. Workman AD, Carey RM, Chen B, et al. CALHM1-mediated ATP release and ciliary beat frequency modulation in nasal epithelial cells. *Sci Rep*. 2017;7:6687.
104. Sabirov RZ, Okada Y. The maxi-anion channel: a classical channel playing novel roles through an unidentified molecular entity. *J Physiol Sci*. 2009;59:3-21.

105. Burow P, Klapperstück M, Markwardt F. Activation of ATP secretion via volume-regulated anion channels by sphingosine-1-phosphate in RAW macrophages. *Pflügers Arch Eur J Physiol*. 2015;467:1215-1226.
106. Dutta AK, Sabirov RZ, Uramoto H, et al. Role of ATP-conductive anion channel in ATP release from neonatal rat cardiomyocytes in ischaemic or hypoxic conditions. *J Physiol*. 2004;559:799-812.
107. Bosma MM. Anion channels with multiple conductance levels in a mouse B lymphocyte cell line. *J Physiol*. 1989;410:67-90.
108. Schlichter LC, Grygorczyk R, Pahapill PA, et al. A large, multiple-conductance chloride channel in normal human T lymphocytes. *Pflügers Arch Eur J Physiol*. 1990;416:413-421.
109. Wareham KJ, Seward EP. P2×7 receptors induce degranulation in human mast cells. *Purinergic Signal*. 2016;12:235-246.
110. Elliott MR, Chekeni FB, Trampont PC, et al. Nucleotides released by apoptotic cells act as a find-me signal to promote phagocytic clearance. *Nature*. 2009;461:282-286.
111. Mitchell CH, Carré DA, Mcglinn AM, et al. A release mechanism for stored ATP in ocular ciliary epithelial cells. *Proc Natl Acad Sci USA*. 1998;95:7174-7178.
112. Grygorczyk R, Hanrahan JW. CFTR-independent ATP release from epithelial cells triggered by mechanical stimuli. *Am J Physiol*. 1997;272:C1058-1066.
113. Wang Y, Roman R, Lidofsky SD, et al. Autocrine signaling through ATP release represents a novel mechanism for cell volume regulation. *Proc Natl Acad Sci USA*. 1996;93:12020-12025.
114. Lazarowski ER, Homolya L, Boucher RC, et al. Direct demonstration of mechanically induced release of cellular UTP and its implication for uridine nucleotide receptor activation. *J Biol Chem*. 1997;272:24348-24354.
115. Zimmermann H. Ectonucleotidases: some recent developments and a note on nomenclature. *Drug Dev Res*. 2001;52:44-56.
116. Yegutkin GG. Nucleotide- and nucleoside-converting ectoenzymes: important modulators of purinergic signalling cascade. *Biochim Biophys Acta Mol Cell Res*. 2008;1783:673-694.
117. Chiu S, Bharat A. Role of monocytes and macrophages in regulating immune response following lung transplantation. *Cuurent Opin Organ Transplant*. 2016;21:239-245.
118. Rizzo R, Ferrari D, Melchiorri L, et al. Extracellular ATP acting at the P2×7 receptor inhibits secretion of soluble HLA-G from human monocytes. *J Immunol*. 2009;183:4302-4311.
119. Piccini A, Carta S, Tassi S, et al. ATP is released by monocytes stimulated with pathogen-sensing receptor ligands and induces IL-1β and IL-18 secretion in an autocrine way. *Proc Natl Acad Sci USA*. 2008;105:8067-8072.
120. Marcet B, Horckmans M, Libert F, et al. Extracellular nucleotides regulate CCL20 release from human primary airway epithelial cells, monocytes and monocyte-derived dendritic cells. *J Cell Physiol*. 2007;211:716-727.
121. Grahnert A, Klein C, Hauschildt S. Involvement of P2X receptors in the NAD⁺-induced rise in [Ca²⁺]_i in human monocytes. *Purinergic Signal*. 2009;5:309-319.
122. De Ita M, Vargas MH, Carbajal V, et al. ATP releases ATP or other nucleotides from human peripheral blood leukocytes through purinergic P2 receptors. *Life Sci*. 2016;145:85-92.
123. Gudipaty L, Humphreys BD, Buell G, et al. Regulation of P2×7 nucleotide receptor function in human monocytes by extracellular ions and receptor density. *Am J Physiol Cell Physiol*. 2001;280:C943-C953.
124. Lioi AB, Ferrari BM, Dubyak GR, et al. Human beta defensin-3 increases CD86 expression on monocytes by activating the ATP-gated channel P2×7. *J Immunol*. 2015;195:4438-4445.
125. Martínez-García JJ, Martínez-Banaclocha H, Angosto-Bazarra D, et al. P2×7 receptor induces mitochondrial failure in monocytes and compromises NLRP3 inflammasome activation during sepsis. *Nat Commun*. 2019;10:2711.
126. Castrichini M, Lazzerini PE, Gamberucci A, et al. The purinergic P2×7 receptor is expressed on monocytes in Behçet's disease and is modulated by TNF-α. *Eur J Immunol*. 2014;44:227-238.
127. Yu J, Chen Y, Li M, et al. Paeoniflorin down-regulates ATP-induced inflammatory cytokine production and P2×7R expression on peripheral blood mononuclear cells from patients with primary Sjögren's syndrome. *Int Immunopharmacol*. 2015;28:115-120.
128. López-López C, Jaramillo-Polanco J, Portales-Pérez DP, et al. Two P2×1 receptor transcripts able to form functional channels are present in most human monocytes. *Eur J Pharmacol*. 2016;793:82-88.
129. Wang L, Jacobsen SEW, Bengtsson A, et al. P2 receptor mRNA expression profiles in human lymphocytes, monocytes and CD34⁺ stem and progenitor cells. *BMC Immunol*. 2004;5:1-7.
130. Gudipaty L, Humphreys BD, Buell G, et al. Regulation of P2×7 nucleotide receptor function in human monocytes by extracellular ions and receptor density. *Am J Physiol Cell Physiol*. 2001;280:943-953.
131. Aga M, Johnson CJ, Hart AP, et al. Modulation of monocyte signaling and pore formation in response to agonists of the nucleotide receptor P2X(7). *J Leukoc Biol*. 2002;72:222-32.
132. Layhadi JA, Fountain SJ. P2×4 receptor-dependent Ca²⁺ influx in model human monocytes and macrophages. *Int J Mol Sci*. 2017;18:2261.
133. Hewinson J, Moore SF, Glover C, et al. A key role for redox signaling in rapid P2×7 receptor-induced IL-1β processing in human monocytes. *J Immunol*. 2008;180:8410-8420.
134. Warny M, Aboudola S, Robson SC, et al. P2Y(6) nucleotide receptor mediates monocyte interleukin-8 production in response to UDP or lipopolysaccharide. *J Biol Chem*. 2001;276:26051-26056.
135. Sivaramakrishnan V, Bidula S, Campwala H, et al. Constitutive lysosome exocytosis releases ATP and engages P2Y receptors in human monocytes. *J Cell Sci*. 2012;125:4567-4575.
136. Santiago-Pérez LI, Flores RV, Santos-Berríos C, et al. P2Y(2) nucleotide receptor signaling in human monocytic cells: activation, desensitization, and coupling to mitogen-activated protein kinases. *J Cell Physiol*. 2001;187:196-208.
137. Syeda F, Liu HY, Tullis E, et al. Differential signaling mechanisms of HNP-induced IL-8 production in human lung epithelial cells and monocytes. *J Cell Physiol*. 2008;214:820-827.
138. Sung SSS, Young JDE, Origlio AM, et al. Extracellular ATP perturbs transmembrane ion fluxes, elevates cytosolic [Ca²⁺]_i, and inhibits phagocytosis in mouse macrophages. *J Biol Chem*. 1985;260:13442-13449.
139. Buisman HP, Steinberg TH, Fischbarg J, et al. Extracellular ATP induces a large nonselective conductance in macrophage plasma membranes. *Proc Natl Acad Sci USA*. 1988;85:7988-7992.
140. Steinberg TH, Silverstein SC. Extracellular ATP⁴⁻ promotes cation fluxes in the J774 mouse macrophage cell line. *J Biol Chem*. 1987;262:3118-3122.
141. Coutinho-Silva R, Alves LA, Savino W, et al. A cation non-selective channel induced by extracellular ATP in macrophages and phagocytic cells of the thymic reticulum. *Biochim Biophys Acta Biomembr*. 1996;1278:125-130.
142. Eschke D, Wüst M, Hauschildt S, et al. Pharmacological characterization of the P2X(7) receptor on human macrophages using the patch-clamp technique. *Naunyn Schmiedebergs Arch Pharmacol*. 2002;365:168-171.
143. Steinberg TH, Newman AS, Swansonq JA, et al. ATP₄-permeabilizes the plasma membrane of mouse macrophages to fluorescent dyes. *J Biol Chem*. 1987;262:8884-8888.
144. Murgial M, Hanaug S, Pizzol P, et al. Oxidized ATP. An irreversible inhibitor of the macrophage purinergic P2Z receptor. *J Biol Chem*. 1993;268:8199-8203.

145. Nuttles LC, Dubyak GR. Differential activation of cation channels and non-selective pores by macrophage P2, purinergic receptors expressed in *Xenopus* oocytes. *J Biol Chem.* 1994;269:13988-13996.
146. Kaczmarek-Hájek K, Lőrinczi É, Hausmann R, et al. Molecular and functional properties of P2X receptors-recent progress and persisting challenges. *Purinergic Signal.* 2012;8:375-417.
147. Martin U, Scholler J, Gurgel J, et al. Externalization of the leaderless cytokine IL-1F6 occurs in response to lipopolysaccharide/ATP activation of transduced bone marrow macrophages. *J Immunol.* 2009;183:4021-4030.
148. Barberà-Cremades M, Baroja-Mazo A, Gomez AI, et al. P2×7 receptor-stimulation causes fever via PGE2 and IL-1β release. *FASEB J.* 2012;26:2951-2962.
149. Gavala ML, Liu Y-P, Lenertz LY, et al. Nucleotide receptor P2RX7 stimulation enhances LPS-induced interferon-β production in murine macrophages. *J Leukoc Biol.* 2013;94:759-768.
150. Barberà-Cremades M, Gómez AI, Baroja-Mazo A, et al. P2×7 receptor induces tumor necrosis factor-α converting enzyme activation and release to boost TNF-α production. *Front Immunol.* 2017;8:862.
151. Lopez-Castejon G, Theaker J, Pelegrin P, et al. P2×7 receptor-mediated release of cathepsins from macrophages is a cytokine-independent mechanism potentially involved in joint diseases. *J Immunol.* 2010;185:2611-2619.
152. Kawamura H, Kawamura T, Kanda Y, et al. Extracellular ATP-stimulated macrophages produce macrophage inflammatory protein-2 which is important for neutrophil migration. *Immunology.* 2012;136:448-458.
153. Gicquel T, Victoni T, Fautrel A, et al. Involvement of purinergic receptors and NOD-like receptor-family protein 3-inflammasome pathway in the adenosine triphosphate-induced cytokine release from macrophages. *Clin Exp Pharmacol Physiol.* 2014;41:279-286.
154. Layhadi JA, Turner J, Crossman D, et al. ATP evokes Ca(2+) responses and CXCL5 secretion via P2×4 receptor activation in human monocyte-derived macrophages. *J Immunol.* 2018;200:1159-1168.
155. Gulinelli S, Salaro E, Vuerich M, et al. IL-18 associates to microvesicles shed from human macrophages by a LPS/TLR-4 independent mechanism in response to P2X receptor stimulation. *Eur J Immunol.* 2012;42:3334-3345.
156. Solle M, Labasi J, Perregaux DG, et al. Altered cytokine production in mice lacking P2×7 receptors. *J Biol Chem.* 2001;276:125-132.
157. El Ouaaliti M, Seil M, Dehaye JP. Activation of calcium-insensitive phospholipase A2 (iPLA 2) by P2×7 receptors in murine peritoneal macrophages. *Prostaglandins Other Lipid Mediat.* 2012;99:116-123.
158. Blanchard DK, McMillen S, Djeu JY. IFN-gamma enhances sensitivity of human macrophages to extracellular ATP-mediated lysis. *J Immunol.* 1991;147:2579-2585.
159. Spranzi E, Djeu JY, Hoffman SL, et al. Lysis of human monocytic leukemia cells by extracellular adenosine triphosphate: mechanism and characterization of the adenosine triphosphate receptor. *Blood.* 1993;82:1578-1585.
160. Lemaire I, Falzoni S, Zhang B, et al. The P2×7 receptor and pannexin-1 are both required for the promotion of multinucleated macrophages by the inflammatory cytokine GM-CSF. *J Immunol.* 2011;187:3878-3887.
161. Brône B, Moechars D, Marrannes R, et al. P2X currents in peritoneal macrophages of wild type and P2×4^{-/-} mice. *Immunol Lett.* 2007;113:83-89.
162. Ulmann L, Hirbec H, Rassendren F. P2×4 receptors mediate PGE2 release by tissue-resident macrophages and initiate inflammatory pain. *EMBO J.* 2010;29:2290-300.
163. Kawano A, Tsukimoto M, Mori D, et al. Regulation of P2×7-dependent inflammatory functions by P2×4 receptor in mouse macrophages. *Biochem Biophys Res Commun.* 2012;420:102-107.
164. Kawano A, Tsukimoto M, Noguchi T, et al. Involvement of P2×4 receptor in P2×7 receptor-dependent cell death of mouse macrophages. *Biochem Biophys Res Commun.* 2012;419:374-380.
165. Zumerle S, Cali B, Munari F, et al. Intercellular calcium signaling induced by ATP potentiates macrophage phagocytosis. *Cell Rep.* 2019;27:1-10.
166. Tonetti M, Sturla L, Giovine M, et al. Extracellular ATP enhances mRNA levels of nitric oxide synthase and TNF-alpha in lipopolysaccharide-treated RAW 264.7 murine macrophages. *Biochem Biophys Res Commun.* 1995;214:125-130.
167. Tonetti M, Sturla L, Bistolfi T, et al. Extracellular ATP potentiates nitric oxide synthase expression induced by lipopolysaccharide in RAW 264.7 murine macrophages. *Biochem Biophys Res Commun.* 1994;203:430-435.
168. Denlinger LC, Fisette PL, Garis KA, et al. Regulation of inducible nitric oxide synthase expression by macrophage purinoreceptors and calcium. *J Biol Chem.* 1996;271:337-342.
169. Bar I, Guns PJ, Metallo J, et al. Knockout mice reveal a role for P2Y6 receptor in macrophages, endothelial cells, and vascular smooth muscle cells. *Mol Pharmacol.* 2008;74:777-784.
170. Myrtek D, Müller T, Geyer V, et al. Activation of human alveolar macrophages via P2 receptors: coupling to intracellular Ca2+ increases and cytokine secretion. *J Immunol.* 2008;181:2181-2188.
171. Zhang Z, Wang Z, Ren H, et al. P2Y6 agonist uridine 5'-diphosphate promotes host defense against bacterial infection via monocyte chemoattractant protein-1-mediated monocytes/macrophages recruitment. *J Immunol.* 2011;186:5376-5387.
172. Pfeilschifter J, Thüring B, Festa F. Extracellular ATP stimulates poly(inositol phospholipid) hydrolysis and eicosanoid synthesis in mouse peritoneal macrophages in culture. *Eur J Biochem.* 1989;186:509-513.
173. Zhang L, Li HY, Li H, et al. Lipopolysaccharide activated phosphatidylcholine-specific phospholipase C and induced IL-8 and MCP-1 production in vascular endothelial cells. *J Cell Physiol.* 2011;226:1694-1701.
174. Sun Y, Li J, Xie X, et al. Macrophage-osteoclast associations: origin, polarization, and subgroups. *Front Immunol.* 2021;12:778078.
175. Gicquel T, Robert S, Loyer P, et al. IL-1β production is dependent on the activation of purinergic receptors and NLRP3 pathway in human macrophages. *FASEB J.* 2015;29:4162-4173.
176. Biswas D, Qureshi OS, Lee W-Y, et al. ATP-induced autophagy is associated with rapid killing of intracellular mycobacteria within human monocytes/macrophages. *BMC Immunol.* 2008;9:35.
177. Placido R, Auricchio G, Falzoni S, et al. P2×7 purinergic receptors and extracellular ATP mediate apoptosis of human monocytes/macrophages infected with *Mycobacterium tuberculosis* reducing the intracellular bacterial viability. *Cell Immunol.* 2006;244:10-18.
178. Marques-da-Silva C, Chaves MM, Thorstenberg ML, et al. Intraleisional uridine-5'-triphosphate (UTP) treatment induced resistance to *Leishmania amazonensis* infection by boosting Th1 immune responses and reactive oxygen species production. *Purinergic Signal.* 2018;14:201-211.
179. Hanley PJ, Musset B, Renigunta V, et al. Extracellular ATP induces oscillations of intracellular Ca2+ and membrane potential and promotes transcription of IL-6 in macrophages. *Proc Natl Acad Sci USA.* 2004;101:9479-9484.
180. Stokes L, Surprenant A. Dynamic regulation of the P2×4 receptor in alveolar macrophages by phagocytosis and classical activation. *Eur J Immunol.* 2009;39:986-995.
181. Gartland A, Buckley KA, Bowler WB, et al. Blockade of the pore-forming P2×7 receptor inhibits formation of multinucleated human osteoclasts in vitro. *Calcif Tissue Int.* 2003;73:361-369.
182. Buckley KA, Hipkind RA, Gartland A, et al. Adenosine triphosphate stimulates human osteoclast activity via upregulation of

- osteoblast-expressed receptor activator of nuclear factor-kappa B ligand. *Bone*. 2002;31:582-590.
183. Bar I, Guns PJ, Metallo J, et al. Knockout mice reveal a role for P2Y6 receptor in macrophages, endothelial cells, and vascular smooth muscle cells. *Mol Pharmacol*. 2008;74:777-784.
 184. Marques-da-Silva C, Chaves MM, Chaves SP, et al. Infection with *Leishmania amazonensis* upregulates purinergic receptor expression and induces host-cell susceptibility to UTP-mediated apoptosis. *Cell Microbiol*. 2011;13:1410-1428.
 185. Thorstenberg ML, Ferreira MVR, Amorim N, et al. Purinergic cooperation between P2Y2 and P2x7 receptors promote cutaneous leishmaniasis control: involvement of pannexin-1 and leukotrienes. *Front Immunol*. 2018;9.
 186. Corrêa G, Marques da Silva C, de Abreu Moreira-Souza AC, et al. Activation of the P2x7 receptor triggers the elimination of *Toxoplasma gondii* tachyzoites from infected macrophages. *Microbes Infect*. 2010;12:497-504.
 187. Merz J, Nettesheim A, von Garlen S, et al. Pro- and anti-inflammatory macrophages express a sub-type specific purinergic receptor profile. *Purinergic Signal*. 2021;17:481-492.
 188. Kessler S, Clauss WG, Günther A, et al. Expression and functional characterization of P2X receptors in mouse alveolar macrophages. *Pflugers Arch Eur J Physiol*. 2011;462:419-430.
 189. Orriss IR, Wang N, Burnstock G, et al. The P2Y6 receptor stimulates bone resorption by osteoclasts. *Endocrinology*. 2011;152:3706-3716.
 190. Layhadi JA, Fountain SJ. P2x4 receptor-dependent Ca²⁺ influx in model human monocytes and macrophages. *Int J Mol Sci*. 2017;18:2261.
 191. Baron L, Gombault A, Fanny M, et al. The NLRP3 inflammasome is activated by nanoparticles through ATP, ADP and adenosine. *Cell Death Dis*. 2015;6:e1629-15.
 192. Aga M, Watters JJ, Pfeiffer ZA, et al. Evidence for nucleotide receptor modulation of cross talk between MAP kinase and NF- κ B signaling pathways in murine RAW 264.7 macrophages. *Am J Physiol Cell Physiol*. 2004;286:923-930.
 193. Pfeiffer ZA, Aga M, Prabhu U, et al. The nucleotide receptor P2x7 mediates actin reorganization and membrane blebbing in RAW 264.7 macrophages via p38 MAP kinase and Rho. *J Leukoc Biol*. 2004;75:1173-1182.
 194. Noguchi T, Ishii K, Fukutomi H, et al. Requirement of reactive oxygen species-dependent activation of ASK1-p38 MAPK pathway for extracellular ATP-induced apoptosis in macrophage. *J Biol Chem*. 2008;283:7657-7665.
 195. Pelegrin P, Barroso-Gutierrez C, Surprenant A. P2x7 receptor differentially couples to distinct release pathways for IL-1 β in mouse macrophage. *J Immunol*. 2008;180:7147-7157.
 196. Eun SY, Seo J, Park SW, et al. LPS potentiates nucleotide-induced inflammatory gene expression in macrophages via the upregulation of P2Y2 receptor. *Int Immunopharmacol*. 2014;18:270-276.
 197. Fairbairn IP, Stober CB, Kumararatne DS, et al. ATP-mediated killing of intracellular mycobacteria by macrophages is a P2x7-dependent process inducing bacterial death by phagosome-lysosome fusion. *J Immunol*. 2001;167:3300-3307.
 198. Ferreira NC da S, Soares-Bezerra RJ, da Silveira RFC, et al. New insights in purinergic therapy: novel antagonists for uridine 5'-triphosphate-activated P2Y receptors from Brazilian flora. *J Med Food*. 2019;22:211-224.
 199. Kojima S, Negishi Y, Tsukimoto M, et al. Purinergic signaling via P2x7 receptor mediates IL-1 β production in Kupffer cells exposed to silica nanoparticle. *Toxicology*. 2014;321:13-20.
 200. Ishimaru M, Yusuke N, Tsukimoto M, et al. Purinergic signaling via P2Y receptors up-mediates IL-6 production by liver macrophages/Kupffer cells. *J Toxicol Sci*. 2014;39:413-423.
 201. Eric Thomas W. Brain macrophages: evaluation of microglia and their functions. *Brain Res Brain Res Rev*. 1992;17:61-74.
 202. Smith JA, Das A, Ray SK, et al. Role of pro-inflammatory cytokines released from microglia in neurodegenerative diseases. *Brain Res Bull*. 2012;87:10-20.
 203. Inoue K, Nakajima K, Morimoto T, et al. ATP stimulation of Ca²⁺-dependent plasminogen release from cultured microglia. *Br J Pharmacol*. 1998;123:1304-1310.
 204. Ferrari D, Villalba M, Chiozzi P, et al. Mouse microglial cells express a plasma membrane pore gated by extracellular ATP. *J Immunol*. 1996;156:1531-1539.
 205. Illes P, Norenberg W, Gebicke-Haerter PJ. Molecular mechanisms of microglial activation. B. Voltage- and purinoceptor-operated channels in microglia. *Neurochem Int*. 1996;29:13-24.
 206. Light AR, Wu Y, Hughen RW, et al. Purinergic receptors activating rapid intracellular Ca²⁺ increases in microglia. *Neuron Glia Biol*. 2006;2:125-138.
 207. Koizumi S, Shigemoto-Mogami Y, Nasu-Tada K, et al. UDP acting at P2Y6 receptors is a mediator of microglial phagocytosis. *Nature*. 2007;446:1091-1095.
 208. Wu LJ, Vadakkan KI, Zhuo M. ATP-induced chemotaxis of microglial processes requires P2Y receptor-activated initiation of outward potassium currents. *Glia*. 2007;55:810-821.
 209. Takenouchi T, Nakai M, Iwamaru Y, et al. The activation of P2x7 receptor impairs lysosomal functions and stimulates the release of autophagolysosomes in microglial cells. *J Immunol*. 2009;182:2051-2062.
 210. Lambert C, Ase AR, Séguéla P, et al. Distinct migratory and cytokine responses of human microglia and macrophages to ATP. *Brain Behav Immun*. 2010;24:1241-1248.
 211. Inoue K. Purinergic systems in microglia. *Cell Mol Life Sci*. 2008;65:3074-3080.
 212. Fang KM, Wang YL, Huang MC, et al. Expression of macrophage inflammatory protein-1 α and monocyte chemoattractant protein-1 in glioma-infiltrating microglia: involvement of ATP and P2x7 receptor. *J Neurosci Res*. 2011;89:199-211.
 213. Di Virgilio F. Dr. Jekyll/Mr. Hyde: the dual role of extracellular ATP. *J Auton Nerv Syst*. 2000;81:59-63.
 214. Shiratori M, Tozaki-Saitoh H, Yoshitake M, et al. P2x7 receptor activation induces CXCL2 production in microglia through NFAT and PKC/MAPK pathways. *J Neurochem*. 2010;114:810-819.
 215. Hide I, Tanaka M, Inoue A, et al. Extracellular ATP triggers tumor necrosis factor-alpha release from rat microglia. *J Neurochem*. 2000;75:965-972.
 216. Parvathenani LK, Tertysnikova S, Greco CR, et al. P2x7 mediates superoxide production in primary microglia and is up-regulated in a transgenic mouse model of Alzheimer's disease. *J Biol Chem*. 2003;278:13309-13317.
 217. Choi HB, Ryu JK, Kim SU, et al. Modulation of the purinergic P2x7 receptor attenuates lipopolysaccharide-mediated microglial activation and neuronal damage in inflamed brain. *J Neurosci*. 2007;27:4957-4968.
 218. Shieh CH, Heinrich A, Serchov T, et al. P2x7-dependent, but differentially regulated release of IL-6, CCL2, and TNF- α in cultured mouse microglia. *Glia*. 2014;62:592-607.
 219. McLarnon JG, Ryu JK, Walker DG, et al. Upregulated expression of purinergic P2x7 receptor in Alzheimer disease and amyloid- β peptide-treated microglia and in peptide-injected rat hippocampus. *J Neuropathol Exp Neurol*. 2006;65:1090-1097.
 220. Ni J, Wang P, Zhang J, et al. Silencing of the P2x7 receptor enhances amyloid- β phagocytosis by microglia. *Biochem Biophys Res Commun*. 2013;434:363-369.
 221. Facci L, Barbierato M, Zusso M, et al. Serum amyloid A primes microglia for ATP-dependent interleukin-1 β release. *J Neuroinflammation*. 2018;15:1-11.

222. Kim HJ, Ajit D, Peterson TS, et al. Nucleotides released from A β 1-42-treated microglial cells increase cell migration and A β 1-42 uptake through P2Y2 receptor activation. *J Neurochem*. 2012;121:228-238.
223. Kataoka A, Koga Y, Uesugi A, et al. Involvement of vasodilator-stimulated phosphoprotein in UDP-induced microglial actin aggregation via PKC- and Rho-dependent pathways. *Purinergic Signal*. 2011;7:403-411.
224. Kim B, Jeong H, Kim J, et al. Uridine 5'-diphosphate induces chemokine expression in microglia and astrocytes through activation of the P2Y6 receptor. *J Immunol*. 2011;186:3701-3709.
225. Morioka N, Tokuhara M, Harano S, et al. The activation of P2Y6 receptor in cultured spinal microglia induces the production of CCL2 through the MAP kinases-NF- κ B pathway. *Neuropharmacology*. 2013;75:116-125.
226. Li H-q, Chen C, Dou Y, et al. P2Y4 receptor-mediated pinocytosis contributes to amyloid beta-induced self-uptake by microglia. *Mol Cell Biol*. 2013;33:4282-4293.
227. Zeng J, Wang G, Liu X, et al. P2Y13 receptor-mediated rapid increase in intracellular calcium induced by ADP in cultured dorsal spinal cord microglia. *Neurochem Res*. 2014;39:2240-2250.
228. Liu PW, Yue MX, Zhou R, et al. P2Y12 and P2Y13 receptors involved in ADP β s induced the release of IL-1 β , IL-6 and TNF- α from cultured dorsal horn microglia. *J Pain Res*. 2017;10:1755-1767.
229. Ohsawa K, Irino Y, Nakamura Y, et al. Involvement of P2 \times 4 and P2Y12 receptors in ATP-induced microglial chemotaxis. *Glia*. 2007;55:604-616.
230. Ohsawa K, Irino Y, Sanagi T, et al. P2Y12 receptor-mediated integrin-beta1 activation regulates microglial process extension induced by ATP. *Glia*. 2010;58:790-801.
231. Honda S, Sasaki Y, Ohsawa K, et al. Extracellular ATP or ADP induce chemotaxis of cultured microglia through Gi/o-coupled P2Y receptors. *J Neurosci*. 2001;21:1975-1982.
232. Haynes SE, Hollopeter G, Yang G, et al. The P2Y12 receptor regulates microglial activation by extracellular nucleotides. *Nat Neurosci*. 2006;9:1512-1519.
233. De Simone R, Niturad CE, De Nuccio C, et al. TGF- β and LPS modulate ADP-induced migration of microglial cells through P2Y1 and P2Y12 receptor expression. *J Neurochem*. 2010;115:450-459.
234. Lee R, Williams JC, Mackman N. P2 \times 7 regulation of macrophage tissue factor activity and microparticle generation. *J Thromb Haemost*. 2012;10:1965-1967.
235. Webster CM, Hokari M, McManus A, et al. Microglial P2Y12 deficiency/inhibition protects against brain ischemia. *PLoS One*. 2013;8:e70927.
236. He Y, Taylor N, Fourceaud L, et al. The role of microglial P2 \times 7: modulation of cell death and cytokine release. *J Neuroinflammation*. 2017;14:1-13.
237. Burm SM, Zuiderwijk-Sick EA, Weert PM, et al. ATP-induced IL-1 β secretion is selectively impaired in microglia as compared to hematopoietic macrophages. *Glia*. 2016;64:2231-2246.
238. Kataoka A, Tozaki-Saitoh H, Koga Y, et al. Activation of P2 \times 7 receptors induces CCL3 production in microglial cells through transcription factor NFAT. *J Neurochem*. 2009;108:115-125.
239. Fang KM, Yang CS, Sun SH, et al. Microglial phagocytosis attenuated by short-term exposure to exogenous ATP through P2 \times 7 receptor action. *J Neurochem*. 2009;111:1225-1237.
240. Kim B, Jeong H, Kim J, et al. Uridine 5'-diphosphate induces chemokine expression in microglia and astrocytes through activation of the P2Y6 receptor. *J Immunol*. 2011;186:3701-3709.
241. Witting A, Walter L, Wacker J, et al. P2 \times 7 receptors control 2-arachidonoylglycerol production by microglial cells. *Proc Natl Acad Sci USA*. 2004;101:3214-3219.
242. Ni J, Wang P, Zhang J, et al. Silencing of the P2 \times 7 receptor enhances amyloid-B phagocytosis by microglia. *Biochem Biophys Res Commun*. 2013;434:363-369.
243. Crain JM. Microglial P2 purinergic receptor and immunomodulatory gene transcripts vary by region, sex, and age in the healthy mouse CNS. *Transcr Open Access*. 2015;3:124.
244. Jiang P, Xing F, Guo B, et al. Nucleotide transmitters ATP and ADP mediate intercellular calcium wave communication via P2Y12/13 receptors among BV-2 microglia. *PLoS One*. 2017;12:e0183114.
245. Langfelder A, Okonji E, Deca D, et al. Extracellular acidosis impairs P2Y receptor-mediated Ca $^{2+}$ signalling and migration of microglia. *Cell Calcium*. 2015;57:247-256.
246. Curet MA, Watters JJ. P2Y14 receptor activation decreases interleukin-6 production and glioma GL261 cell proliferation in microglial transwell cultures. *J Neurooncology*. 2018;137:23-31.
247. Xiang Z, Burnstock G. Expression of P2X receptors on rat microglial cells during early development. *Glia*. 2005;52:119-126.
248. Bianco F, Fumagalli M, Pravettoni E, et al. Pathophysiological roles of extracellular nucleotides in glial cells: differential expression of purinergic receptors in resting and activated microglia. *Brain Res Rev*. 2005;48:144-156.
249. Sanz JM, Chiozzi P, Ferrari D, et al. Activation of microglia by amyloid B requires P2 \times 7 receptor expression. *J Immunol*. 2009;182:4378-4385.
250. Bartlett R, Yerbury JJ, Sluyter R. P2 \times 7 receptor activation induces reactive oxygen species formation and cell death in murine eoc13 microglia. *Mediators Inflamm*. 2013;2013.
251. Castell-Rodríguez A, Piñón-Zárate G, Herrera-, et al. Dendritic cells: location, function, and clinical implications. *Biology of Myelomonocytic Cells*. 2017:21-50.
252. Wilkin F, Duhant X, Bruyns C, et al. The P2Y11 receptor mediates the ATP-induced maturation of human monocyte-derived dendritic cells. *J Immunol*. 2001;166:7172-7177.
253. la Sala A, Ferrari D, Corinti S, et al. Extracellular ATP induces a distorted maturation of dendritic cells and inhibits their capacity to initiate Th1 responses. *J Immunol*. 2001;166:1611-1617.
254. Idzko M, Dichmann S, Ferrari D, et al. Nucleotides induce chemotaxis and actin polymerization in immature but not mature human dendritic cells via activation of pertussis toxin-sensitive P2Y receptors. *Blood*. 2002;100:925-932.
255. Horckmans M, Marcet B, Marteau F, et al. Extracellular adenine nucleotides inhibit the release of major monocyte recruiters by human monocyte-derived dendritic cells. *FEBS Lett*. 2006;580:747-754.
256. Marriott I, Insko EW, Bost KL. Extracellular uridine nucleotides initiate cytokine production by murine dendritic cells. *Cell Immunol*. 1999;195:147-156.
257. Marteau F, Communi D, Boeynaems J-M, et al. Involvement of multiple P2Y receptors and signaling pathways in the action of adenine nucleotides diphosphates on human monocyte-derived dendritic cells. *J Leukoc Biol*. 2004;76:796-803.
258. Mutini C, Falzoni S, Ferrari D, et al. Mouse dendritic cells express the P2 \times 7 purinergic receptor: characterization and possible participation in antigen presentation. *J Immunol*. 1999;163:1958-1965.
259. Proctor RA, Denlinger LC, Leventhal PS, et al. Protection of mice from endotoxic death by 2-methylthio-ATP. *Proc Natl Acad Sci USA*. 1994;91:6017-6020.
260. Coutinho-Silva R, Persechini PM, Da Cunha Bisaggio R, et al. P2Z/P2 \times 7 receptor-dependent apoptosis of dendritic cells. *Am J Physiol*. 1999;276:C1139-1147.
261. Pizzirani C, Ferrari D, Chiozzi P, et al. Stimulation of P2 receptors causes release of IL-1beta-loaded microvesicles from human dendritic cells. *Blood*. 2007;109:3856-3864.
262. Ghiringhelli F, Apetoh L, Tesniere A, et al. Activation of the NLRP3 inflammasome in dendritic cells induces IL-1beta-dependent adaptive immunity against tumors. *Nat Med*. 2009;15:1170-1178.

263. Shin A, Toy T, Rothenfusser S, et al. P2Y receptor signaling regulates phenotype and IFN- α secretion of human plasmacytoid dendritic cells. *Blood*. 2008;111:3062-3069.
264. Skelton L, Cooper M, Murphy M, et al. Human immature monocyte-derived dendritic cells express the G protein-coupled receptor GPR105 (KIAA0001, P2Y14) and increase intracellular calcium in response to its agonist, uridine diphosphoglucose. *J Immunol*. 2003;171:1941-1949.
265. Liu QH, Bohlen H, Titzer S, et al. Expression and a role of functionally coupled P2Y receptors in human dendritic cells. *FEBS Lett*. 1999;445:402-408.
266. Ferrari D, Gorini S, Callegari G, et al. Shaping immune responses through the activation of dendritic cells' P2 receptors. *Purinergic Signal*. 2007;3:99-107.
267. Ben Addi A, Cammarata D, Conley PB, et al. Role of the P2Y12 receptor in the modulation of murine dendritic cell function by ADP. *J Immunol*. 2010;185:5900-5906.
268. Wilkin F, Stordeur P, Goldman M, et al. Extracellular adenine nucleotides modulate cytokine production by human monocyte-derived dendritic cells: dual effect on IL-12 and stimulation of IL-10. *Eur J Immunol*. 2002;32:2409-2417.
269. Zhang J, Li Z, Hu X, et al. Knockout of P2Y12 aggravates experimental autoimmune encephalomyelitis in mice via increasing of IL-23 production and Th17 cell differentiation by dendritic cells. *Brain Behav Immun*. 2017;62:245-255.
270. Schnurr M, Toy T, Shin A, et al. Extracellular nucleotide signaling by P2 receptors inhibits IL-12 and enhances IL-23 expression in human dendritic cells: a novel role for the cAMP pathway. *Blood*. 2005;105:1582-1589.
271. Ferrari D, La Sala A, Chiozzi P, et al. The P2 purinergic receptors of human dendritic cells: identification and coupling to cytokine release. *FASEB J*. 2000;14:2466-2476.
272. Berchtold S, Ogilvie ALJ, Bogdan C, et al. Human monocyte derived dendritic cells express functional P2X and P2Y receptors as well as ecto-nucleotidases. *FEBS Lett*. 1999;458:424-428.
273. Girolomoni G, Santantonio ML, Pastore S, et al. Epidermal Langerhans cells are resistant to the permeabilizing effects of extracellular ATP: in vitro evidence supporting a protective role of membrane ATPase. *J Invest Dermatol*. 1993;100:282-287.
274. Coutinho-Silva R, Alves LA, De Carvalho ACC, et al. Characterization of P2Z purinergic receptors on phagocytic cells of the thymic reticulum in culture. *Biochim Biophys Acta Biomembr*. 1996;1280:217-222.
275. Nakanishi K, Tsukimoto M, Tanuma S-I, et al. Silica nanoparticles activate purinergic signaling via P2 \times 7 receptor in dendritic cells, leading to production of pro-inflammatory cytokines. *Toxicol Vitro*. 2016;35:202-211.
276. Cano LE, Lopera HDE. Introduction to T and B lymphocytes. In: Anaya J-M, Shoenfeld Y, Rojas-Villarraga A, eds. *Autoimmunity: From bench to bedside*. El Rosario University Press; 2013:77-96.
277. El-Moatassim C, Dornand J, Mani JC. Extracellular ATP increases cytosolic free calcium in thymocytes and initiates the blastogenesis of the phorbol 12-myristate 13-acetate-treated medullary population. *Biochim Biophys Acta*. 1987;927:437-444.
278. El-Moatassim C, Bernad N, Mani JC, et al. Extracellular ATP induces a nonspecific permeability of thymocyte plasma membranes. *Biochem Cell Biol*. 1989;67:495-502.
279. Di Virgilio F, Bronte V, Collavo D, et al. Responses of mouse lymphocytes to extracellular adenosine 5'-triphosphate (ATP). Lymphocytes with cytotoxic activity are resistant to the permeabilizing effects of ATP. *J Immunol*. 1989;143:1955-1960.
280. Adinolfi E, Pizzirani C, Idzko M, et al. P2 \times 7 receptor: death or life? *Purinergic Signal*. 2005;1:219-227.
281. Redegeld F, Filippini A, Trenn G. Possible role of extracellular ATP in cell-cell interactions leading to CTL-mediated cytotoxicity. In: Sitkovsky M V, Henkart PA, eds. *Cytotoxic cells: Recognition, effector function, generation, and methods*. Birkhäuser Boston; 1993:307-313.
282. Zamboni A, Bronte V, Di Virgilio F, et al. Role of extracellular ATP in cell-mediated cytotoxicity: a study with ATP-sensitive and ATP-resistant macrophages. *Cell Immunol*. 1994;156:458-467.
283. Trabanelli S, Očadlíková D, Gulinelli S, et al. Extracellular ATP exerts opposite effects on activated and regulatory CD4 $^{+}$ T cells via purinergic P2 receptor activation. *J Immunol*. 2012;189:1303-1310.
284. Kobayashi D, Sugiura Y, Umemoto E, et al. Extracellular ATP limits homeostatic T cell migration within lymph nodes. *Front Immunol*. 2021;12:786595.
285. Alves LA, Coutinho-Silva R, Savino W. Extracellular ATP: a further modulator in neuroendocrine control of the thymus. *Neuroimmunomodulation*. 1999;6:81-89.
286. Baricordi OR, Ferrari D, Melchiorri L, et al. An ATP-activated channel is involved in mitogenic stimulation of human T lymphocytes. *Blood*. 1996;87:682-690.
287. Freedman BD, Liu QH, Gaulton G, et al. ATP-evoked Ca $^{2+}$ transients and currents in murine thymocytes: possible role for P2X receptors in death by neglect. *Eur J Immunol*. 1999;29:1635-1646.
288. Chvatchko Y, Valera S, Aubry JP, et al. The involvement of an ATP-gated ion channel, P2 \times 1, in thymocyte apoptosis. *Immunity*. 1996;5:275-283.
289. Nagy P, Panyi G, Jenei A, et al. Ion-channel activities regulate transmembrane signaling in thymocyte apoptosis and T-cell activation. *Immunol Lett*. 1995;44:91-95.
290. Apasov SG, Koshiba M, Chused TM, et al. Effects of extracellular ATP and adenosine on different thymocyte subsets: possible role of ATP-gated channels and G protein-coupled purinergic receptor. *J Immunol*. 1997;158:5095-5105.
291. Chused TM, Apasov S, Sitkovsky M. Murine T lymphocytes modulate activity of an ATP-activated P2Z-type purinoceptor during differentiation. *J Immunol*. 1996;157:1371-1380.
292. Aguirre A, Shoji KF, Sáez JC, et al. FasL-triggered death of Jurkat cells requires caspase 8-induced, ATP-dependent cross-talk between Fas and the purinergic receptor P2 \times 7. *J Cell Physiol*. 2013;228:485-493.
293. Shoji KF, Sáez PJ, Harcha PA, et al. Pannexin1 channels act downstream of P2 \times 7 receptors in ATP-induced murine T-cell death. *Channels*. 2014;8:142-156.
294. Schenk U, Westendorf AM, Radaelli E, et al. Purinergic control of T cell activation by ATP released through pannexin-1 hemichannels. *Sci Signal*. 2008;1.
295. Elliott JI, Higgins CF. Major histocompatibility complex class I shedding and programmed cell death stimulated through the proinflammatory P2 \times 7 receptor: a candidate susceptibility gene for NOD diabetes. *Diabetes*. 2004;53:2012-2017.
296. Moon H, Na HY, Chong KH, et al. P2 \times 7 receptor-dependent ATP-induced shedding of CD27 in mouse lymphocytes. *Immunol Lett*. 2006;102:98-105.
297. Eleftheriadis T, Pissas G, Karioti A, et al. Uric acid induces caspase-1 activation, IL-1 β secretion and P2 \times 7 receptor dependent proliferation in primary human lymphocytes. *Hippokratia*. 2013;17:141-145.
298. Tsukimoto M, Tokunaga A, Harada H, et al. Blockade of murine T cell activation by antagonists of P2Y6 and P2 \times 7 receptors. *Biochem Biophys Res Commun*. 2009;384:512-518.
299. Frascoli M, Marcandalli J, Schenk U, et al. Purinergic P2 \times 7 receptor drives T cell lineage choice and shapes peripheral $\gamma\delta$ cells. *J Immunol*. 2012;189:174-180.
300. Schenk U, Frascoli M, Proietti M, et al. ATP inhibits the generation and function of regulatory T cells through the activation of purinergic P2X receptors. *Sci Signal*. 2011;4:1-12.
301. Baroja-Mazo A, Barberà-Cremades M, Pelegrín P. P2 \times 7 receptor activation impairs exogenous MHC class I oligopeptides presentation in antigen presenting cells. *PLoS One*. 2013;8:e70577.

302. Hamoudi C, Zhao C, Abderrazak A, et al. The purinergic receptor P2x4 promotes Th17 activation and the development of arthritis. *J Immunol.* 2022;208:1115-1127.
303. Somers GR, Hammet FM, Trute L, et al. Expression of the P2Y6 purinergic receptor in human T cells infiltrating inflammatory bowel disease. *Lab Invest.* 1998;78:1375-1383.
304. Salem M, El Azreq MA, Pelletier J, et al. Exacerbated intestinal inflammation in P2Y6 deficient mice is associated with Th17 activation. *Biochim Biophys Acta Mol Basis Dis.* 2019;1865:2595-2605.
305. Scrivens M, Dickenson JM. Functional expression of the P2Y14 receptor in murine T-lymphocytes. *Br J Pharmacol.* 2005;146:435-444.
306. Fan ZD, Zhang YY, Guo YH, et al. Involvement of P2x7 receptor signaling on regulating the differentiation of Th17 cells and type II collagen-induced arthritis in mice. *Sci Rep.* 2016;6:1-12.
307. Jin J, Dasari VR, Sistare FD, et al. Distribution of P2Y receptor subtypes on haematopoietic cells. *Br J Pharmacol.* 1998;123:789-794.
308. Lê KT, Paquet M, Nouel D, et al. Primary structure and expression of a naturally truncated human P2X ATP receptor subunit from brain and immune system. *FEBS Lett.* 1997;418:195-199.
309. Wang CM, Ploia C, Anselmi F, et al. Adenosine triphosphate acts as a paracrine signaling molecule to reduce the motility of T cells. *EMBO J.* 2014;33:1354-1364.
310. Woehrle T, Yip L, Elkhali A, et al. Pannexin-1 hemichannel-mediated ATP release together with P2x1 and P2x4 receptors regulate T-cell activation at the immune synapse. *Blood.* 2010;116:3475-3484.
311. Koshiha M, Apasov S, Sverdlov V, et al. Transient up-regulation of P2Y2 nucleotide receptor mRNA expression is an immediate early gene response in activated thymocytes. *Proc Natl Acad Sci USA.* 1997;94:831-836.
312. Glass R, Townsend-Nicholson A, Burnstock G. P2 receptors in the thymus: expression of P2X and P2Y receptors in adult rats, an immunohistochemical and in situ hybridisation study. *Cell Tissue Res.* 2000;300:295-306.
313. Lebien TW, Tedder TF. B lymphocytes: how they develop and function. *Blood.* 2008;112:1570-1580.
314. Wiley JS, Chen R, Jamieson GP. The ATP4+ receptor-operated channel (P2Z class) of human lymphocytes allows Ba2+ and ethidium+ uptake: inhibition of fluxes by Suramin. *Arch Biochem Biophys.* 1993;305:54-60.
315. Wiley JS, Chen R, Wiley MJ, et al. The ATP4- receptor-operated ion channel of human lymphocytes: inhibition of ion fluxes by amiloride analogs and by extracellular sodium ions. *Arch Biochem Biophys.* 1992;292:411-418.
316. Wiley JS, Chen JR, Snook MB, et al. The P2Z-purinoceptor of human lymphocytes: actions of nucleotide agonists and irreversible inhibition by oxidized ATP. *Br J Pharmacol.* 1994;112:946-950.
317. Padeh S, Cohen A, Roifman CM. ATP-induced activation of human B lymphocytes via P2-purinoceptors. *J Immunol.* 1991;146:1626-1632.
318. Gu BJ, Zhang WY, Bendall LJ, et al. Expression of P2x7 purinoceptors on human lymphocytes and monocytes: evidence for nonfunctional P2x7 receptors. *Am J Physiol Cell Physiol.* 2000;279:1189-1197.
319. Gu BJ, Zhang W, Worthington RA, et al. A Glu-496 to Ala polymorphism leads to loss of function of the human P2x7 receptor. *J Biol Chem.* 2001;276:11135-11142.
320. Gu B, Bendall LJ, Wiley JS. Adenosine triphosphate-induced shedding of CD23 and L-selectin (CD62L) from lymphocytes is mediated by the same receptor but different metalloproteases. *Blood.* 1998;92:946-951.
321. Jamieson GP, Snook MB, Thurlow PJ, et al. Extracellular ATP causes loss of L-selectin from human lymphocytes via occupancy of P2Z purinoceptors. *J Cell Physiol.* 1996;166:637-642.
322. Gargett CE, Cornish EJ, Wiley JS. Phospholipase D activation by P2Z-purinoceptor agonists in human lymphocytes is dependent on bivalent cation influx. *Biochem J.* 1996;313:529-535.
323. Baricordi OR, Melchiorri L, Adinolfi E, et al. Increased proliferation rate of lymphoid cells transfected with the P2X(7) ATP receptor. *J Biol Chem.* 1999;274:33206-33208.
324. Sengstake S, Boneberg EM, Illges H. CD21 and CD62L shedding are both inducible via P2x7Rs. *Int Immunol.* 2006;18:1171-1178.
325. Pupovac A, Geraghty NJ, Watson D, et al. Activation of the P2x7 receptor induces the rapid shedding of CD23 from human and murine B cells. *Immunol Cell Biol.* 2015;93:77-85.
326. Sakowicz-Burkiewicz M, Kocbuch K, Grden M, et al. High glucose concentration impairs ATP outflow and immunoglobulin production by human peripheral B lymphocytes: involvement of P2x7 receptor. *Immunobiology.* 2013;218:591-601.
327. Sakowicz-Burkiewicz M, Grden M, Maciejewska I, et al. High glucose impairs ATP formation on the surface of human peripheral blood B lymphocytes. *Int J Biochem Cell Biol.* 2013;45:1246-1254.
328. Lee DH, Park KS, Kong ID, et al. Expression of P2 receptors in human B cells and Epstein-Barr virus-transformed lymphoblastoid cell lines. *BMC Immunol.* 2006;7:1-11.
329. Wiley JS, DUBYAK GR. Extracellular adenosine triphosphate increases cation permeability of chronic lymphocytic leukemic lymphocytes. *Blood.* 1989;73:1316-1323.
330. Abel AM, Yang C, Thakar MS, et al. Natural killer cells: development, maturation, and clinical utilization. *Front Immunol.* 2018;9:1869.
331. Henriksson T. Inhibition of natural killing by adenosine ribonucleotides. *Immunol Lett.* 1983;7:171-176.
332. Schmidt A, Ortaldo JR, Herberman RB. Inhibition of human natural killer cell reactivity by exogenous adenosine 5'-triphosphate. *J Immunol.* 1984;132:146-150.
333. Miller JS, Cervenka T, Lund J, et al. Purine metabolites suppress proliferation of human NK cells through a lineage-specific purine receptor. *J Immunol.* 1999;162:7376-82.
334. Krishnaraj R. Negative modulation of human NK cell activity by purinoceptors. 1. Effect of exogenous adenosine triphosphate. *Cell Immunol.* 1992;141:306-322.
335. Myers LK, Kang ES. Modulation of natural killer cell activity by surface phosphorylation reactions. *Exp Cell Res.* 1990;187:270-276.
336. Gorini S, Callegari G, Romagnoli G, et al. ATP secreted by endothelial cells blocks CX3CL1-elicited natural killer cell chemotaxis and cytotoxicity via P2Y11 receptor activation. *Blood.* 2010;116:4492-4500.
337. Li Z, Gao Y, He C, et al. Purinergic receptor P2Y6 is a negative regulator of NK cell maturation and function. *J Immunol.* 2021;207:1555-1565.
338. Rosales C. Neutrophil: a cell with many roles in inflammation or several cell types? *Front Physiol.* 2018;9:113.
339. El-Benna J, Hurtado-Nedelec M, Marzaioli V, et al. Priming of the neutrophil respiratory burst: role in host defense and inflammation. *Immunol Rev.* 2016;273:180-193.
340. Ward PA, Cunningham TW, McCulloch KK, et al. Regulatory effects of adenosine and adenine nucleotides on oxygen radical responses of neutrophils. *Lab Invest.* 1988;58:438-447.
341. Ward PA, Cunningham TW, McCulloch KK, et al. Platelet enhancement of O2- responses in stimulated human neutrophils. Identification of platelet factor as adenine nucleotide. *Lab Invest.* 1988;58:37-47.
342. Ward PA, Cunningham TW, Walker BAM, et al. Differing calcium requirements for regulatory effects of ATP, ATPγS and adenosine on O2• responses of human neutrophils. *Biochem Biophys Res Commun.* 1988;154:746-751.
343. Kuhns DB, Wright DG, Nath J, et al. ATP induces transient elevations of [Ca2+]i in human neutrophils and primes these cells for enhanced O2- generation. *Lab Invest.* 1988;58:448-453.
344. Meshki J, Tuluc F, Bredeteau O, et al. Molecular mechanism of nucleotide-induced primary granule release in human neutrophils: role for the P2Y2 receptor. *Am J Physiol Cell Physiol.* 2004;286:264-271.

345. Tuluc F, Bredeteau O, Brailoiu E, et al. The priming effect of extracellular UTP on human neutrophils: role of calcium released from thapsigargin-sensitive intracellular stores. *Purinergic Signal*. 2005;1:359-368.
346. Kukulski F, Ben Yebdri F, Lecka J, et al. Extracellular ATP and P2 receptors are required for IL-8 to induce neutrophil migration. *Cytokine*. 2009;46:166-170.
347. Yebdri FB, Kukulski F, Tremblay A, et al. Concomitant activation of P2Y2 and P2Y6 receptors on monocytes is required for TLR1/2-induced neutrophil migration by regulating IL-8 secretion. *Eur J Immunol*. 2009;39:2885-2894.
348. Cicko S, Lucattelli M, Müller T, et al. Purinergic receptor inhibition prevents the development of smoke-induced lung injury and emphysema. *J Immunol*. 2010;185:688-697.
349. Khine AA, Del Sorbo L, Vaschetto R, et al. Human neutrophil peptides induce interleukin-8 production through the P2Y6 signaling pathway. *Blood*. 2006;107:2936-2942.
350. Kukulski F, Yebdri FB, Bahrami F, et al. The P2 receptor antagonist PPADS abrogates LPS-induced neutrophil migration in the murine air pouch via inhibition of MIP-2 and KC production. *Mol Immunol*. 2010;47:833-839.
351. Grbic DM, Degagné É, Larrivière J-F, et al. P2Y6 receptor contributes to neutrophil recruitment to inflamed intestinal mucosa by increasing CXC chemokine ligand 8 expression in an AP-1-dependent manner in epithelial cells. *Inflamm Bowel Dis*. 2012;18:1456-69.
352. Sil P, Hayes CP, Reaves BJ, et al. P2Y6 receptor antagonist MRS2578 inhibits neutrophil activation and aggregated neutrophil extracellular trap formation induced by gout-associated monosodium urate crystals. *J Immunol*. 2017;198:428-442.
353. Amison RT, Arnold S, O'Shaughnessy BG, et al. Lipopolysaccharide (LPS) induced pulmonary neutrophil recruitment and platelet activation is mediated via the P2Y1 and P2Y14 receptors in mice. *Pulm Pharmacol Ther*. 2017;45:62-68.
354. Barrett MO, Sesma JI, Ball CB, et al. A selective high-affinity antagonist of the P2Y14 receptor inhibits UDP-glucose-stimulated chemotaxis of human neutrophils. *Mol Pharmacol*. 2013;84:41-9.
355. Lintzmaier Petiz L, Glaser T, Scharfstein J, et al. P2Y14 receptor as a target for neutrophilia attenuation in severe COVID-19 cases: from hematopoietic stem cell recruitment and chemotaxis to thrombo-inflammation. *Stem Cell Rev Rep*. 2021;17:241-252.
356. Alkayed F, Kashimata M, Koyama N, et al. P2Y11 purinoceptor mediates the ATP-enhanced chemotactic response of rat neutrophils. *J Pharmacol Sci*. 2012;120:288-295.
357. Scrivens M, Dickenson JM. Functional expression of the P2Y14 receptor in human neutrophils. *Eur J Pharmacol*. 2006;543:166-173.
358. O'Donoghue M, Morrow DA, Cannon CP, et al. Association between baseline neutrophil count, clopidogrel therapy, and clinical and angiographic outcomes in patients with ST-elevation myocardial infarction receiving fibrinolytic therapy. *Eur Heart J*. 2008;29:984-991.
359. Liverani E, Rico MC, Garcia AE, et al. Prasugrel metabolites inhibit neutrophil functions. *J Pharmacol Exp Ther*. 2013;344:231-243.
360. Rahman M, Gustafsson D, Wang Y, et al. Ticagrelor reduces neutrophil recruitment and lung damage in abdominal sepsis. *Platelets*. 2014;25:257-263.
361. Lecut C, Frederix K, Johnson DM, et al. P2x1 ion channels promote neutrophil chemotaxis through Rho kinase activation. *J Immunol*. 2009;183:2801-2809.
362. Lecut C, Faccinnetto C, Delierneux C, et al. ATP-gated P2x1 ion channels protect against endotoxemia by dampening neutrophil activation. *J Thromb Haemost*. 2012;10:453-65.
363. Darbousset R, Delierneux C, Mezouar S, et al. P2x1 expressed on polymorphonuclear neutrophils and platelets is required for thrombosis in mice. *Blood*. 2014;124:2575-85.
364. Suh B-C, Kim J-S, Namgung U, et al. P2x7 nucleotide receptor mediation of membrane pore formation and superoxide generation in human promyelocytes and neutrophils. *J Immunol*. 2001;166:6754-6763.
365. Karmakar M, Katsnelson MA, Dubyak GR, et al. Neutrophil P2x7 receptors mediate NLRP3 inflammasome-dependent IL-1 β secretion in response to ATP. *Nat Commun*. 2016;7:1-13.
366. Lucattelli M, Cicko S, Müller T, et al. P2x7 receptor signaling in the pathogenesis of smoke-induced lung inflammation and emphysema. *Am J Respir Cell Mol Biol*. 2011;44:423-429.
367. Martel-Gallegos G, Casas-Pruneda G, Ortega-Ortega F, et al. Oxidative stress induced by P2x7 receptor stimulation in murine macrophages is mediated by c-Src/Pyk2 and ERK1/2. *Biochim Biophys Acta*. 2013;1830:4650-4659.
368. Ding Q, Quah SY, Tan KS. Secreted adenosine triphosphate from *Aggregatibacter actinomycetemcomitans* triggers chemokine response. *Mol Oral Microbiol*. 2016;31:423-434.
369. Mohanty JG, Raible DG, McDermott LJ, et al. Effects of purine and pyrimidine nucleotides on intracellular Ca²⁺ in human eosinophils: activation of purinergic P2Y receptors. *J Allergy Clin Immunol*. 2001;107:849-855.
370. Kudo F, Nishiguchi N, Mizuike R, et al. Neutrophil phagocytosis is down-regulated by nucleotides until encounter with pathogens. *Immunol Lett*. 2012;144:24-32.
371. Bulanova E, Bulfone-Paus S. P2 receptor-mediated signaling in mast cell biology. *Purinergic Signal*. 2010;6:3-17.
372. Krystel-Whittemore M, Dileepan KN, Wood JG. Mast cell: a multi-functional master cell. *Front Immunol*. 2016;6:620.
373. Bennett JP, Cockcroft S, Gomperts BD. Rat mast cells permeabilized with ATP secrete histamine in response to calcium ions buffered in the micromolar range. *J Physiol*. 1981;317:335-345.
374. Tatham PER, Cusack NJ, Gomperts BD. Characterisation of the ATP4-receptor that mediates permeabilisation or rat mast cells. *Eur J Pharmacol*. 1988;147:13-21.
375. McCloskey MA, Qian YX. Selective expression of potassium channels during mast cell differentiation. *J Biol Chem*. 1994;269:14813-14819.
376. Jaffar ZH, Pearce FL. Histamine secretion from mast cells stimulated with ATP. *Agents Actions*. 1990;30:64-66.
377. Gao ZG, Ding Y, Jacobson KA. UDP-glucose acting at P2Y14 receptors is a mediator of mast cell degranulation. *Biochem Pharmacol*. 2010;79:873-879.
378. Zhan-Guo GG, Ding Y, Jacobson KA. P2Y13 receptor is responsible for ADP-mediated degranulation in RBL-2H3 rat mast cells. *Pharmacol Res*. 2010;62:500-505.
379. Gao ZG, Wei Q, Jayasekara MPS, et al. The role of P2Y14 and other P2Y receptors in degranulation of human LAD2 mast cells. *Purinergic Signal*. 2013;9:31-40.
380. Pinheiro AR, Paramos-De-carvalho D, Certal M, et al. Histamine induces ATP release from human subcutaneous fibroblasts, via pannexin-1 hemichannels, leading to Ca²⁺ mobilization and cell proliferation. *J Biol Chem*. 2013;288:27571-27583.
381. Rossi GL, Yen A, Barrett KE. IgE and adenosine 5' triphosphate receptors on immature murine mast cells are functionally linked to signal transduction mechanisms. *J Allergy Clin Immunol*. 1992;90:765-771.
382. McCloskey MA, Fan Y, Luther S. Chemotaxis of rat mast cells toward adenine nucleotides. *J Immunol*. 1999;163:970-7.
383. Feng C, Mery AG, Beller EM, et al. Adenine nucleotides inhibit cytokine generation by human mast cells through a Gs-coupled receptor. *J Immunol*. 2004;173:7539-7547.
384. Bulanova E, Budagian V, Orinska Z, et al. Extracellular ATP induces cytokine expression and apoptosis through P2x7 receptor in murine mast cells. *J Immunol*. 2005;174:3880-3890.
385. Kurashima Y, Amiya T, Nochi T, et al. Extracellular ATP mediates mast cell-dependent intestinal inflammation through P2x7 purinoceptors. *Nat Commun*. 2012;3:1034.

386. Schulman ES, Glaum MC, Post T, et al. ATP modulates anti-IgE-induced release of histamine from human lungs mast cells. *Am J Respir Cell Mol Biol.* 1999;20:530-537.
387. Wareham K, Vial C, Wykes RCE, et al. Functional evidence for the expression of P2_{x1}, P2_{x4} and P2_{x7} receptors in human lung mast cells. *Br J Pharmacol.* 2009;157:1215-1224.
388. Jiang Y, Ye F, Du Y, et al. P2_{x7R} in mast cells is a potential target for salicylic acid and aspirin in treatment of inflammatory pain. *J Inflamm Res.* 2021;14:2913-2931.
389. Yoshida K, Ito M, Matsuoka I. Divergent regulatory roles of extracellular ATP in the degranulation response of mouse bone marrow-derived mast cells. *Int Immunopharmacol.* 2017;43:99-107.
390. Muniz VS, Weller PF, Neves JS. Eosinophil crystalloid granules: structure, function, and beyond. *J Leukoc Biol.* 2012;92:281-288.
391. Dichmann S, Idzko M, Zimpfer U, et al. Adenosine triphosphate-induced oxygen radical production and CD11b up-regulation: ca(++) mobilization and actin reorganization in human eosinophils. *Blood.* 2000;95:973-978.
392. Idzko M, Panther E, Bremer HC, et al. Stimulation of P2 purinergic receptors induces the release of eosinophil cationic protein and interleukin-8 from human eosinophils. *Br J Pharmacol.* 2003;138:1244-1250.
393. Alberto AVP, Faria RX, de Menezes JRL, et al. Role of P2 receptors as modulators of rat eosinophil recruitment in allergic inflammation. *PLoS One.* 2016;11:e0145392.
394. Müller T, Vieira RP, Grimm M, et al. A potential role for P2_{x7R} in allergic airway inflammation in mice and humans. *Am J Respir Cell Mol Biol.* 2011;44:456-464.
395. Wright A, Mahaut-Smith M, Symon F, et al. Impaired P2_{x1} receptor-mediated adhesion in eosinophils from asthmatic patients. *J Immunol.* 2016;196:4877-4884.
396. Karcz TP, Whitehead GS, Nakano K, et al. UDP-glucose and P2Y₁₄ receptor amplify allergen-induced airway eosinophilia. *J Clin Invest.* 2021;131:e140709.
397. Idzko M, Dichmann S, Panther E, et al. Functional characterization of P2Y and P2X receptors in human eosinophils. *J Cell Physiol.* 2001;188:329-336.
398. Ferrari D, Idzko M, Dichmann S, et al. P2 purinergic receptors of human eosinophils: characterization and coupling to oxygen radical production. *FEBS Lett.* 2000;486:217-224.
399. Chirumbolo S. State-of-the-art review about basophil research in immunology and allergy: is the time right to treat these cells with the respect they deserve? *Blood Transfus.* 2012;10:148-168.
400. Siracusa MC, Kim BS, Spergel JM, et al. Basophils and allergic inflammation. *J Allergy Clin Immunol.* 2013;132:789-801.
401. Nakano M, Ito K, Yuno T, et al. UDP/P2Y₆ receptor signaling regulates IgE-dependent degranulation in human basophils. *Allergol Int.* 2017;66:574-580.
402. Lammas DA, Stober C, Harvey CJ, et al. ATP-induced killing of mycobacteria by human macrophages is mediated by purinergic P2Z(P2_{x7}) receptors. *Immunity.* 1997;7:433-444.
403. Sikora A, Liu J, Brosnan C, et al. Cutting edge: purinergic signaling regulates radical-mediated bacterial killing mechanisms in macrophages through a P2_{x7}-independent mechanism. *J Immunol.* 1999;163:558-561.
404. Molloy A, Laochumroonvorapong P, Kaplan G. Apoptosis, but not necrosis, of infected monocytes is coupled with killing of intracellular bacillus Calmette-Guérin. *J Exp Med.* 1994;180:1499-1509.
405. Santos AA, Rodrigues-Junior V, Zanin RF, et al. Implication of purinergic P2_{x7} receptor in M. tuberculosis infection and host interaction mechanisms: a mouse model study. *Immunobiology.* 2013;218:1104-1112.
406. Amaral EP, Ribeiro SCM, Lanes VR, et al. Pulmonary infection with hypervirulent mycobacteria reveals a crucial role for the P2_{x7} receptor in aggressive forms of tuberculosis. *PLoS Pathog.* 2014;10.
407. Saunders BM, Fernando SL, Sluyter R, et al. A loss-of-function polymorphism in the human P2_{x7} receptor abolishes ATP-mediated killing of mycobacteria. *J Immunol.* 2003;171:5442-5446.
408. Shemon AN, Sluyter R, Fernando SL, et al. A Thr357 to Ser polymorphism in homozygous and compound heterozygous subjects causes absent or reduced P2_{x7} function and impairs ATP-induced mycobacterial killing by macrophages. *J Biol Chem.* 2006;281:2079-2086.
409. Pelegrin P. P2_{x7} receptor and the NLRP3 inflammasome: partners in crime. *Biochem Pharmacol.* 2021;187:114385.
410. Yang Y, Wang H, Kouadir M, et al. Recent advances in the mechanisms of NLRP3 inflammasome activation and its inhibitors. *Cell Death Dis.* 2019;10:128.
411. Abderrazak A, Syrovets T, Couchie D, et al. NLRP3 inflammasome: from a danger signal sensor to a regulatory node of oxidative stress and inflammatory diseases. *Redox Biol.* 2015;4:296-307.
412. Jang HM, Park JY, Lee YJ, et al. TLR2 and the NLRP3 inflammasome mediate IL-1 β production in *Prevotella nigrescens*-infected dendritic cells. *Int J Med Sci.* 2021;18:432-440.
413. Guan F, Jiang W, Bai Y, et al. Purinergic P2_{x7} receptor mediates the elimination of *Trichinella spiralis* by activating NF- κ B/NLRP3/IL-1 β pathway in macrophages. *Infect Immun.* 2021;89:e00683-20.
414. Li LH, Chen TL, Chiu HW, et al. Critical role for the NLRP3 inflammasome in mediating IL-1 β production in *Shigella sonnei*-infected macrophages. *Front Immunol.* 2020;11:1115.
415. Coutinho-Silva R, Perfettini JL, Persechini PM, et al. Modulation of P2Z/P2X(7) receptor activity in macrophages infected with *Chlamydia psittaci*. *Am J Physiol Cell Physiol.* 2001;280:C81-9.
416. Stober CB, Lammas DA, Li CM, et al. ATP-mediated killing of *Mycobacterium bovis* bacille Calmette-Guérin within human macrophages is calcium dependent and associated with the acidification of mycobacteria-containing phagosomes. *J Immunol.* 2001;166:6276-6286.
417. Moreira-Souza ACA, Marinho Y, Correa G, et al. Pyrimidinergic receptor activation controls *Toxoplasma gondii* infection in macrophages. *PLoS One.* 2015;10:1-23.
418. Marques-da-Silva C, Chaves MM, Castro NG, et al. Colchicine inhibits cationic dye uptake induced by ATP in P2_{x2} and P2_{x7} receptor-expressing cells: implications for its therapeutic action. *Br J Pharmacol.* 2011;163:912-26.
419. Purines SP, Torres-Santos EC, Marques C, et al. Modulation of P2_{x7} purinergic receptor in macrophages by *Leishmania amazonensis* and its role in parasite elimination. *Microbes Infect.* 2009;11:842-849.
420. Chaves MM, Marques-da-Silva C, Monteiro APT, et al. Leukotriene B₄ modulates P2_{x7} receptor-mediated *Leishmania amazonensis* elimination in murine macrophages. *J Immunol.* 2014;192:4765-4773.
421. Marques-da-Silva C, Chaves MM, Thorstenberg ML, et al. Intraleisional uridine-5'-triphosphate (UTP) treatment induced resistance to *Leishmania amazonensis* infection by boosting Th1 immune responses and reactive oxygen species production. *Purinergic Signal.* 2018;14:201-211.
422. Hazleton JE, Berman JW, Eugenin EA. Purinergic receptors are required for HIV-1 infection of primary human macrophages. *J Immunol.* 2012;188:4488-4495.
423. Giroud C, Marin M, Hammonds J, et al. P2_{x1} receptor antagonists inhibit HIV-1 fusion by blocking virus-coreceptor interactions. *J Virol.* 2015;89:9368-9382.
424. Trautmann A. Extracellular ATP in the immune system: more than just a "danger signal. *Sci Signal.* 2009;2:pe6.
425. Riteau N, Gasse P, Fauconnier L, et al. Extracellular ATP is a danger signal activating P2_{x7} receptor in lung inflammation and fibrosis. *Am J Respir Crit Care Med.* 2010;182:774-83.
426. Moll M, Kuemmerle-Deschner JB. Inflammasome and cytokine blocking strategies in autoinflammatory disorders. *Clin Immunol.* 2013;147:242-275.

427. Dell'Antonio G, Quattrini A, Dal Cin E, et al. Relief of inflammatory pain in rats by local use of the selective P2_x7 ATP receptor inhibitor, oxidized ATP. *Arthritis Rheum*. 2002;46:3378-3385.
428. Dell'Antonio G, Quattrini A, Dal Cin E, et al. Antinociceptive effect of a new P(2Z)/P2_x7 antagonist, oxidized ATP, in arthritic rats. *Neurosci Lett*. 2002;327:87-90.
429. Labasi JM, Petrushova N, Donovan C, et al. Absence of the P2_x7 receptor alters leukocyte function and attenuates an inflammatory response. *J Immunol*. 2002;168:6436-6445.
430. Burnstock G, Brouns I, Adriaensen D, et al. Purinergic signaling in the airways. *Pharmacol Rev*. 2012;64:834-868.
431. Chen Y, Corriden R, Inoue Y, et al. ATP release guides neutrophil chemotaxis via P2Y2 and A3 receptors. *Science (80-)*. 2006;314:1792-1795.
432. Inoue Y, Chen Y, Hirsh MI, et al. A3 and P2Y2 receptors control the recruitment of neutrophils to the lungs in a mouse model of sepsis. *Shock*. 2008;30:1.
433. Vanderstocken G, van de Paar E, Robaye B, et al. Protective role of P2Y2 receptor against lung infection induced by pneumonia virus of mice. *PLoS One*. 2012;7:1-6.
434. Zerr M, Hechler B, Freund M, et al. Major contribution of the P2Y1 receptor in purinergic regulation of TNF α -induced vascular inflammation. *Circulation*. 2011;123:2404-2413.
435. Riegel AK, Faigle M, Zug S, et al. Selective induction of endothelial P2Y6 nucleotide receptor promotes vascular inflammation. *Blood*. 2011;117:2548-2555.
436. Vieira RP, Müller T, Grimm M, et al. Purinergic receptor type 6 contributes to airway inflammation and remodeling in experimental allergic airway inflammation. *Am J Respir Crit Care Med*. 2011;184:215-223.
437. Giannattasio G, Ohta S, Boyce JR, et al. The purinergic G protein-coupled receptor 6 inhibits effector T cell activation in allergic pulmonary inflammation. *J Immunol*. 2011;187:1486-1495.
438. Pinho-Ribeiro FA, Verri WA, Chiu IM. Nociceptor sensory neuron-immune interactions in pain and inflammation. *Trends Immunol*. 2017;38:5-19.
439. Oliveira MCG, Pelegrini-da-Silva A, Tambeli CH, et al. Peripheral mechanisms underlying the essential role of P2_x3,2/3 receptors in the development of inflammatory hyperalgesia. *Pain*. 2009;141:127-134.
440. Teixeira JM, Oliveira MCG, Nociti FH, et al. Involvement of temporomandibular joint P2_x3 and P2_x2/3 receptors in carrageenan-induced inflammatory hyperalgesia in rats. *Eur J Pharmacol*. 2010;645:79-85.
441. de Oliveira Fusaro MCG, Pelegrini-da-Silva A, Araldi D, et al. P2_x3 and P2_x2/3 receptors mediate mechanical hyperalgesia induced by bradykinin, but not by pro-inflammatory cytokines, PGE2 or dopamine. *Eur J Pharmacol*. 2010;649:177-182.
442. Prado FC, Araldi D, Vieira AS, et al. Neuronal P2_x3 receptor activation is essential to the hyperalgesia induced by prostaglandins and sympathomimetic amines released during inflammation. *Neuropharmacology*. 2013;67:252-258.
443. Rae MG, Rowan EG, Kennedy C. Pharmacological properties of P2_x3-receptors present in neurones of the rat dorsal root ganglia. *Br J Pharmacol*. 1998;124:176-180.
444. Ueno S, Tsuda M, Iwanaga T, et al. Cell type-specific ATP-activated responses in rat dorsal root ganglion neurons. *Br J Pharmacol*. 1999;126:429-436.
445. Hasegawa S, Kohro Y, Tsuda M, et al. Activation of cytosolic phospholipase A2 in dorsal root ganglion neurons by Ca²⁺/calmodulin-dependent protein kinase II after peripheral nerve injury. *Mol Pain*. 2009;5:22.
446. Koizumi M, Asano S, Furukawa A, et al. P2_x3 receptor upregulation in trigeminal ganglion neurons through TNF α production in macrophages contributes to trigeminal neuropathic pain in rats. *J Headache Pain*. 2021;22:31.
447. Tsuda M, Shigemoto-Mogami Y, Koizumi S, et al. P2_x4 receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature*. 2003;424:778-83.
448. Ulmann L, Hatcher JP, Hughes JP, et al. Up-regulation of P2_x4 receptors in spinal microglia after peripheral nerve injury mediates BDNF release and neuropathic pain. *J Neurosci*. 2008;28:11263-11268.
449. Tsuda M, Kuboyama K, Inoue T, et al. Behavioral phenotypes of mice lacking purinergic P2_x4 receptors in acute and chronic pain assays. *Mol Pain*. 2009;5:28.
450. Honore P, Donnelly-Roberts D, Namovic MT, et al. A-740003 [N-(1-[[[cyanoimino](5-quinolinylamino) methyl]amino]-2,2-dimethylpropyl)-2-(3,4-dimethoxyphenyl)acetamide], a novel and selective P2_x7 receptor antagonist, dose-dependently reduces neuropathic pain in the rat. *J Pharmacol Exp Ther*. 2006;319:1376-85.
451. Fulgenzi A, Dell'Antonio G, Foglieni C, et al. Inhibition of chemokine expression in rat inflamed paws by systemic use of the antihyperalgesic oxidized ATP. *BMC Immunol*. 2005;6:1-11.
452. Teixeira JM, Oliveira MCG, Parada CA, et al. Peripheral mechanisms underlying the essential role of P2_x7 receptors in the development of inflammatory hyperalgesia. *Eur J Pharmacol*. 2010;644:55-60.
453. Honore P, Donnelly-Roberts D, Namovic M, et al. The antihyperalgesic activity of a selective P2_x7 receptor antagonist, A-839977, is lost in IL-1 α knockout mice. *Behav Brain Res*. 2009;204:77-81.
454. Kobayashi K, Yamanaka H, Yanamoto F, et al. Multiple P2Y subtypes in spinal microglia are involved in neuropathic pain after peripheral nerve injury. *Glia*. 2012;60:1529-1539.
455. Tozaki-Saitoh H, Tsuda M, Miyata H, et al. P2Y12 receptors in spinal microglia are required for neuropathic pain after peripheral nerve injury. *J Neurosci*. 2008;28:4949-4956.

How to cite this article: Alberto AVP, Ferreira NCS, Bonavita AGC, et al. Physiologic roles of P2 receptors in leukocytes. *J Leukoc Biol*. 2022;112:983-1012.
<https://doi.org/10.1002/JLB.2RU0421-226RR>