

# BMJ Open Comparison of common interventions for the treatment of infantile colic: a systematic review of reviews and guidelines

Julie Ellwood,<sup>1</sup> Jerry Draper-Rodi,<sup>1</sup> Dawn Carnes <sup>1,2</sup>

**To cite:** Ellwood J, Draper-Rodi J, Carnes D. Comparison of common interventions for the treatment of infantile colic: a systematic review of reviews and guidelines. *BMJ Open* 2020;**10**:e035405. doi:10.1136/bmjopen-2019-035405

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-035405>).

Received 04 November 2019  
Revised 31 January 2020  
Accepted 03 February 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Research Department, University College of Osteopathy, London, UK

<sup>2</sup>Faculty of Health, University of Applied Sciences Western Switzerland, Fribourg, Switzerland

## Correspondence to

Dr Dawn Carnes;  
dawn.carnes@uco.ac.uk

## ABSTRACT

**Objective** To conduct a systematic review of systematic reviews and national guidelines to assess the effectiveness of four treatment approaches (manual therapy, probiotics, proton pump inhibitors and simethicone) on colic symptoms including infant crying time, sleep distress and adverse events.

**Methods** We searched PubMed, Embase, Cochrane and Mantis for studies published between 2009 and 2019. Inclusion criteria were systematic reviews and guidelines that used evidence and expert panel opinion. Three reviewers independently selected articles by title, abstract and full paper review. Data were extracted by one reviewer and checked by a second. Selected studies were assessed for quality using modified standardised checklists by two authors. Meta-analysed data for our outcomes of interest were extracted and narrative conclusions were assessed.

**Results** Thirty-two studies were selected. High-level evidence showed that probiotics were most effective for reducing crying time in breastfed infants (range –25 min to –65 min over 24 hours). Manual therapies had moderate to low-quality evidence showing reduced crying time (range –33 min to –76 min per 24 hours). Simethicone had moderate to low evidence showing no benefit or negative effect. One meta-analysis did not support the use of proton pump inhibitors for reducing crying time and fussing. Three national guidelines unanimously recommended the use of education, parental reassurance, advice and guidance and clinical evaluation of mother and baby. Consensus on other advice and treatments did not exist.

**Conclusions** The strongest evidence for the treatment of colic was probiotics for breastfed infants, followed by weaker but favourable evidence for manual therapy indicated by crying time. Both forms of treatment carried a low risk of serious adverse events. The guidance reviewed did not reflect these findings.

**PROSPERO registration number** CRD42019139074.

## INTRODUCTION

Infantile colic, which is defined as excessive crying in the first few months of life, is a common but poorly understood and often frustrating problem for parents and carers. Infantile colic affects somewhere between 3% and 40% of infants worldwide, depending on

## Strengths and limitations of this study

- This study compares meta-data from different treatments for infantile colic on common outcomes.
- Guidance and evidence are compared.
- Distinguishing superiority of treatments is difficult with multiple subpopulations and outcomes.
- Where aetiology of a condition is uncertain, rationale for treatments are difficult to justify.
- Effectiveness studies and efficacy studies are needed in this field.

geography and definitions used.<sup>1</sup> It is estimated that around one in six families (17%) with children consult a health professional about symptoms associated with infantile colic, and these include excessive crying, fussing and distress.<sup>2</sup>

Infants cry for various reasons to express discomfort caused by conditions ranging from benign disorders to life-threatening illness. A meta-analysis published in 2017, which included 28 diary studies covering 8690 infants, reported a mean daily fuss and cry duration of 117 min to 133 min in the first 6 weeks of life, followed by a decline in crying time to a mean of 68 min per day by 10 to 12 weeks of age.<sup>3</sup> It is suggested that less than 5% of distressed infants have identifiable medical explanations for their crying.<sup>4</sup>

There is confusion around the terminology and diagnosis of infantile colic with other diagnostic terms such as silent reflux, functional gastrointestinal disorder and sometimes infantile headache to explain the symptoms of colic. A systematic review of definitions and outcome measures in trials of infantile colic reported the current variability in defining infantile colic, which parallels the non-uniformity of measuring the condition.<sup>5</sup> Most definitions are based on Wessel's criteria, also known as the 'rule of threes', which defines colic as paroxysms of irritability, fussing or

crying lasting  $\geq 3$  hours per day on  $\geq 3$  days per week for  $> 3$  weeks<sup>6</sup> in an otherwise healthy baby aged 2 weeks to 4 months. However, these criteria have been found to be out of date and impractical to use.<sup>7</sup> The most recent diagnostic criteria, formulated by the Rome IV committee, are recurrent and prolonged periods of crying without an obvious cause or evidence of failure to thrive or illness in infants younger than 5 months.<sup>8</sup>

The natural history of infantile colic is favourable with symptoms gradually disappearing by the time the infant is 4 months old.<sup>9</sup> However, the impact of excessive infant crying on healthcare services is the most common reason for paediatric consultations and hospital emergency department visits in the first weeks of life.<sup>10</sup> The consequences of having an excessively crying infant in the family are harmful to relationships and health.<sup>11</sup> Excessive infant crying is associated with maternal issues such as depression, anxiety and loss of parenting confidence.<sup>12 13</sup> It is also a common cause of early breastfeeding cessation<sup>14</sup> and has been associated with severe infant injury or death as a result of abuse.<sup>15</sup>

Recommended management strategies usually centre around parental support and reassurance that the infant is otherwise healthy. However, parents are often in a state of crisis and feel that they want to take action. A number of treatment options exist, which include pharmacological treatments (eg, dicyclomine hydrochloride, cimetropium bromide, simethicone and proton pump inhibitors), probiotics, complementary therapies (including herbal agents and sucrose), manual therapies (for example chiropractic, osteopathy and physiotherapy), dietary interventions and parental behavioural interventions.

Reviews to date have assessed the effectiveness of manipulative therapies,<sup>16</sup> probiotics,<sup>17</sup> dietary modification,<sup>18</sup> complementary and alternative therapies (herbal formulations, sucrose or glucose)<sup>19</sup> and pain-relieving agents.<sup>19</sup> However, there is limited research which compares these treatments to inform the management of infants experiencing colic and parental decision-making.<sup>9</sup>

## AIM

The aim of this study was to review and compare the effectiveness of manual therapy to three of the most common interventions (probiotics, simethicone and proton pump inhibitors) on colic symptoms in infants, including crying time, sleep and infant distress and adverse events.

## METHODS

We conducted a pragmatic narrative systematic review of systematic reviews (SRs) and clinical guidelines for the most common treatments of infantile colic. We employed a review approach using existing analyses of information whether it was based on compiled narrative analyses, meta-analysis or guideline consensus review. A network meta-analysis for comparing outcomes across treatments was not considered as we were aware from

the pre-scoping stage of this review that the studies were too heterogeneous in terms of quality between treatment type. The review protocol was registered on the International prospective register of systematic reviews, and this report follows the guidance in the PRISMA statement<sup>20</sup> for reporting reviews.

## Eligibility criteria

Our population of interest were infants (under 6 months old) with colic as defined by either Wessel's or the ROME III or IV criteria.<sup>6 7</sup> We were not considering infants with diagnosed gastro-oesophageal reflux disease (GORD), as this is considered a pathological condition and differs from colic in that the infants are generally not thriving.

Only children who were otherwise healthy and thriving were included in this study, that is, no other dominant or serious comorbidity requiring medical care in a non-primary care setting.

We considered four treatments for infantile colic: manual therapy, simethicone, proton pump inhibitors and probiotics as these were considered most widely recognised and of interest to the funder. We included the use of proton pump inhibitors because, while it is not recommended for use in infants with colic, it is increasingly being used with these infants to 'rule out' reflux and or GORD, as symptoms of colic are thought to be secondary to GORD.<sup>21</sup> Dicyclomine treatment for infantile colic (anti-spasmodic drug) was excluded in this study as it is not recommended for infants under 6 months due to contraindications, mainly respiratory symptoms varying in severity and consequence.<sup>22</sup>

We defined manual therapy as any predominantly (more than 75%) touch-based therapy administered by a trained and registered manual therapist, such as a chiropractor, osteopath, osteopathic physician, physical therapist or physiotherapist.

The outcomes of interest were crying time, sleep and parental distress, and in addition, where data existed, adverse events data for each treatment were reviewed.

We limited the type of literature we reviewed to systematic reviews of effectiveness and national clinical guidelines.

We included reviews that reported a systematic review methodology and which had more than one researcher indicated in the review process. Narrative literature reviews and editorials were not included.

We included clinical guidelines where clear methodological procedures for development were reported and included systematically designed evidence reviews and expert panel consensus. The guidance had to be intended for broad use at a national level rather than intended as guidance for a single clinic or hospital or a specific setting. We excluded guidance targeted at parents.

We limited the literature to that published in the last decade between 2009 and 2019 and written in English.

## Information sources

For systematic reviews, we searched PubMed, MANTIS, EMBASE, Cochrane and for guidelines we searched those published in English by national organisations. The searches were conducted in June 2019.

## Search

Key search terms used were ‘systematic reviews’, ‘infant\*’, ‘colic’, ‘manual therapy’, ‘manipulative medicine’, ‘simethicone’, ‘proton pump inhibitors’, ‘PPIs’, ‘probiotics’, ‘probiotic agents’ (see online supplementary appendix 1).

We searched the central clearing guideline database and known national guideline centres for guidelines on the treatment, management and care of babies with unsettled or distressed behaviour, including infantile colic.

## Study selection

Results from searches on each database were downloaded into a Reference Management Software, EndNote (V.X4.0.2), and duplicates were removed; titles and abstracts were screened by two independent researchers. Inclusion/exclusion decisions were made by all three authors. Citation tracking was used to triangulate our searches and check for missing reviews and guidance, as well as to identify other articles that may have not been indexed in PubMed. Full-text papers were obtained for those that met the inclusion criteria and for those where it was unclear whether or not the abstract and title met the inclusion criteria. If a more recent or updated versions of reviews or guidelines were found, the more recent ones replaced those initially found.

## Quality appraisal

We appraised the quality of the systematic reviews using a modified version of the AMSTAR 2 critical appraisal tool.<sup>23</sup> Of the 15 quality categories assessed, 8 were selected by the authors for the final calculation of quality and we allocated a score of 1 for Yes, 0 for No and 0.5 for Partial Yes. The scale ranged from 0 to 8 (Highest quality). Each study was appraised by two independent reviewers and a third reviewer was used if mitigation was required (see online supplementary appendix 2).

We reported on the quality of the guidelines using a modified version of the AGREE II framework.<sup>24</sup> The AGREE II checklist has 23 items evaluating six domains. Where more than 75% of the 23 items (>17/23) were met, we rated these as high-quality guidance. In guidance, where there was insufficient information to record a verdict, we left these domains blank. Guidelines of 16 or less domains did not receive an evaluation of high quality. Each guideline was appraised by one reviewer (see online supplementary appendix 3).

## Data extraction and items

We summarised studies by type of intervention, number of studies included, number of participants and outcomes of interest and measure used (table 1). We extracted meta-analysed data on hours of crying time and sleeping

time between intervention and control groups and other active interventions. For parent distress and confidence, we proposed to compare effect sizes and/or change scores on similar outcomes. In the absence of meta-analysis, change scores or effect size data, we conducted a narrative synthesis of outcomes.

We also extracted adverse event incident data and compared risk ratios between treatments and controls where possible. One researcher extracted characteristics and data from the selected articles, and this was checked by a second researcher. A third reviewer’s opinion was sought in cases of disagreement in the extraction process.

## Level of evidence

We used reported levels of evidence as published in the reviews and guidance and analysed these to indicate overall level of publication consensus on effectiveness and safety. The strength of the overall evidence was determined by the reviews and guidance evaluated as *Favourable*—significant benefit to treatment compared with control; *Not favourable*—no significant benefit to treatment compared with control or no difference between treatment and controls; *Unfavourable*—results for treatment worse than control; *Inconclusive*—unable to draw conclusions from results.

## Patient and public involvement

No patient involvement.

## RESULTS

### Study selection

We found 201 studies for screening against title and abstract of which 126 were excluded and 75 were selected for full paper review. Of these, a further 43 were excluded and 32 references were finally included in this systematic review. See figure 1 flow chart illustrating the process and reasons for exclusion.

Eleven studies investigated the effectiveness of manual therapy for treating infantile colic, 12 for probiotics, 1 for proton pump inhibitors, 4 for simethicone and 4 papers which included a combination of the interventions of interest in this review.

The characteristics of the final selection of studies are shown in tables 1–5 ordered by interventions: manual therapy, simethicone, probiotics, proton pump inhibitors and mixed interventions.

The control arms varied for the studies included in the systematic reviews: for example, in the probiotic, proton pump inhibitors and simethicone studies, the controls were mainly placebos or usual feeding method (mainly breast feeding); and in the manual therapy studies, the controls were either no treatment, usual care, dimethicone and in one craniosacral decompression.


### Outcome measures

Crying time was the most reported outcome (26/32 studies). In three studies, crying episodes (as opposed

**Table 1** Characteristics of studies for manual therapy for infantile colic

Authors	Participants/n/age/gender	Timing of intervention	Number of studies: type of studies included in review	Method of data synthesis (narrative/meta-analysis)
Alcantara <i>et al</i> <sup>38</sup>	Infants/n=980 (where reported)/<18 years/gender not reported	<4 weeks/30 days	26: 3 clinical trials, 2 survey studies, 6 case reports, 2 case series, 4 cohort studies, 5 commentaries and 4 reviews of the literature	Narrative
Carnes <i>et al</i> <sup>43</sup>	Infants/n=866 (one study not reported)/gender not reported	8 days to 4 weeks FU 2–11 years	11: 5× RCTs, 1× retrospective review, 2× cohort studies, 3× cases series	Both
Clar <i>et al</i> <sup>39</sup>	Infants/n=not reported	Not reported	19: 10 SRs and 9 RCTs	Narrative
Dobson <i>et al</i> <sup>16</sup>	Infants/n=358/birth–12 weeks/gender percentages varied (where reported) between 34% and 93% male (mean 56%, median 47%)	8 days to 6 weeks	6 RCTs	Meta-analysis
Driehuis <i>et al</i> <sup>42</sup>	Infants/n=283/birth–10 weeks/gender not reported	8–14 days	4 RCTs	Both
Ernst <sup>41</sup>	Infants/n=198/2–10 weeks/gender not specified	8–15 days	3 RCTs	Narrative
Gleberzon <i>et al</i> <sup>36</sup>	Infants/n=1459/birth–3 years	Immediate—4 weeks	6: 3× RCTs, 1× prospective uncontrolled, 2× retrospective studies	Narrative
Parnell Prevost <i>et al</i> <sup>40</sup>	Infants/n=>4000/0–12 weeks	Not reported	5: 4 RCTs, 1 obs	Narrative
Posadzki and Ernst <sup>27</sup>	Infants/n=not reported	Not reported	2 SRs	Narrative
Posadzki <i>et al</i> <sup>37</sup>	Infants/n=28/1–12 weeks	28 days	1 RCT	Narrative
Salehi <i>et al</i> <sup>28</sup>	Infants/n=368/age=not reported	Not reported	11 SRs—2 SRs for colic	Narrative

FU, follow-up; RCT, randomised controlled trial; SR, systematic review.



**Figure 1** Flowchart of search process for the review. PPI, proton pump inhibitor.

to crying duration) were recorded as the outcome measure.<sup>18 25 26</sup> Three studies did not report outcome data.<sup>27–29</sup>

Fourteen of the 26 papers that reported crying time conducted a meta-analysis. One study<sup>30</sup> reported outcomes for all three interventions and another<sup>31</sup> reported meta-analysed data for probiotic use and narrative analysis for simethicone. There were 10 meta-analysed results for probiotics, 2 for simethicone and 4 for manual therapy. The data indicated that for breastfed infants, probiotics significantly decreased daily crying time (range –25 min to –65 min). The level of evidence for these findings was reported as high (as indicated by the systematic reviewer authors' appraisal of quality and/or risk of bias) with seven studies reporting high quality, one moderate and two low quality. There were four systematic reviews investigating effectiveness of manual therapy for crying, three showed favourable effects (from –33 min to –76 hours of crying time reduction in 24 hours) and one review was inconclusive. The quality of these randomised controlled trials (RCTs) ranged from moderate to low by the authors of the SRs.

The findings from two studies investigating the effects of simethicone were inconclusive or not favourable and were graded as moderate/low quality. [Table 6](#) shows the meta-analysed data for crying time.

There were three studies that reported the number of crying episodes; one was a meta-analysis on the effectiveness of probiotics<sup>25</sup> and the other two were narrative analyses, one for probiotics<sup>26</sup> and one where simethicone was used as the control arm compared with partially hydrolysed formula.<sup>18</sup> The results from these three studies did not support the use of probiotics or simethicone for the treatment of infantile colic and the quality of the studies were reported as low in all studies (see [table 7](#)).

## Narrative analysis

### Crying time

Thirteen of the 26 studies reporting on crying times conducted a narrative review. Two of the studies<sup>2 32</sup> reported on the effectiveness of both probiotics and manual therapy for treating infantile colic. Of the four studies that reported on probiotics, one reported a favourable outcome when compared with simethicone.<sup>33</sup> The other studies did not support the use of probiotics and were of low-grade quality.

Two reviews narratively reported on the effects on crying time using simethicone; the findings did not support its use.<sup>31 34</sup> The quality of the studies in these reviews were graded as low by the review authors.

One review reported on crying time, crying episodes, crying time post feeding and parent perception of change in fussing time between infants taking proton pump inhibitors and a placebo. They found no significant differences between groups on any outcome and concluded in the light of associated adverse events unfavourable evidence for the use of proton pump inhibitors.<sup>35</sup>

Eight studies reported on the effectiveness of manual therapy for infantile colic. Two studies reported a favourable outcome,<sup>36 37</sup> four were inconclusive,<sup>32 38–40</sup> one study showed no beneficial effects over the control arm<sup>41</sup> and one study reported an unfavourable outcome.<sup>2</sup> Most of the reviews for manual therapy graded the RCT studies as low quality.

The findings are summarised in [table 8](#).

Three further narrative studies, two for manual therapy<sup>27 28</sup> and one for simethicone,<sup>29</sup> did not report details on how the effectiveness of interventions was measured. The results of these studies were reported as unfavourable or inconclusive and the quality of the studies were low.

### Adverse events

When reported, there were no serious adverse events reported in the RCTs for either manual therapy, simethicone or probiotics. In one SR,<sup>42</sup> four serious adverse events were documented in case studies of manual therapy, including a death, temporary paralysis and a rib fracture, but because these were poorly described and missed underlying pathology could not be ruled out, the overall risk where reported was low. Risks of developing infections with probiotics were reported as extremely low.<sup>33</sup> The review on PPIs showed both arms had adverse events, one RCT showed a significant difference between lansoprazole and the placebo for totality of serious adverse effects, mainly lower respiratory tract infections (10/81 vs 2/81, risk ratio 5, 95% CI 1.3 to 20, number needed to harm 11, 95% CI 6 to 49)<sup>21</sup> (see online supplementary appendix 4).

### Guidelines for the treatment, care and management of infants with colic

We found three national clinical guidelines about the treatment and management of 'colicky' infants.

[Table 9](#) summarises the recommendations they made. The quality rating for the UK guideline development

**Table 2** Characteristics of studies for probiotics for infantile colic

Authors	Participants/n/age/gender	Treatment and timing of intervention	Number of studies: type of studies included in review	Method of data synthesis (narrative/meta-analysis)
Anabrees <i>et al</i> <sup>52</sup>	Breastfed or partly breastfed infants/n=209/<4 months/gender not reported	<i>L. reuteri</i> (strains—American Type Culture Collection Strain 55730 and DSM 17938) 21 days and 28 days	3: 1× Open prospective randomised study, 2× double-blind RCTs	Meta-analysis
Batchelor <i>et al</i> <sup>53</sup>	Infants/n=924/newborns <3 months (where reported)/gender not reported	<i>L. reuteri</i> . 21 days and 90 days (where reported)	5: 3× RCTs, 1× observational prospective and 1× meta-analysis	Narrative
Cruchet <i>et al</i> <sup>33</sup>	Infants/n=not reported/age not reported/gender not reported	<i>L. reuteri</i> , <i>B. lactis</i> and <i>Str. thermophiles</i> Not reported	9: 1× SR, 8× RCTs	Narrative
Dryl and Szajewska <sup>54</sup>	Infants/n=471/<6 months/gender not reported	<i>L. reuteri</i> . 21–30 days	7 RCTs	Meta-analysis
Mugambi <i>et al</i> <sup>25</sup>	Healthy infants	Mixed: synbiotics, probiotics, prebiotics (various strains used). Not reported	25 RCTs	Meta-analysis
Ong <i>et al</i> <sup>17</sup>	Infants/n=1886/<1 month	<i>L. reuteri</i> DSM, multistrain probiotics, <i>L. rhamnosus</i> , <i>L. paracasei</i> and <i>B. animalis</i> 4/52 before delivery to 6 months	6 RCTs	Meta-analysis
Schreck Bird <i>et al</i> <sup>55</sup>	Infants/n=444/31–52 days	<i>L. reuteri</i> DSM 17938 (10 <sup>8</sup> CFU) 21 and 28 days	5 RCTs	Narrative
Skórka <i>et al</i> <sup>26</sup>	Infants/4 weeks to 36 months	Various probiotic strains included. 4 weeks to 12 months	20 RCTs	Narrative
Sung <i>et al</i> <sup>66</sup>	Infants/n=1825/<3 months	Various strains probiotics. 2 weeks to 6 months	12 RCTs	Meta-analysis
Sung <i>et al</i> <sup>67</sup>	Infants/n=345/birth to 23 months	<i>L. reuteri</i> DSM 17938 21 days	4 RCTs	Meta-analysis

CFU, colony-forming unit; RCT, randomised controlled trial; SR, systematic review.

process and reporting met 18/23 criteria, the USA and the Irish guidance 11/23.

## DISCUSSION

### Summary of findings and context

We found 32 relevant systematic reviews and three examples of guidance. Many of the RCTs were repeated within the reviews and this is reflected by fairly consistent results but differing interpretations. Overall, the meta-analysed results showed that both probiotics in breastfed infants and manual therapy can reduce crying time. The daily reduction in crying is between 33 and 76 min with manual therapy and between 25 min and 65 min with probiotics in breastfed infants. The quality and strength of evidence was higher for probiotics than manual therapy. The evidence for probiotics centred on breastfed infants rather than formula-fed infants and there were a

number of different types of strains of probiotics. The manual therapy evidence was based on low to moderate quality RCTs and therefore larger blinded RCTs were recommended. In addition, crying time was reported as the primary outcome in most studies which was used as a proxy indicator of colic resolution or improvement.

There were no serious adverse events reported for either probiotics or manual therapy, indicating that both represent a low risk to infants, although we cannot conclude they are without any risk.<sup>27 42</sup> Two reviews<sup>16 43</sup> analysed the risks of adverse events with manual therapy, one showed 88% less risk of an adverse event in the manual therapy groups than in the control groups<sup>43</sup> and the other showed one in six parents reported non-serious adverse events.<sup>16</sup> Another study<sup>42</sup> reported data from non-RCTs which included four case studies reporting serious incidents of harm but there was some doubt over causality as a result

**Table 3** Characteristics of studies for simethicone for infantile colic

Authors	Participants/n/age/gender	Timing of intervention	Number and type of studies included in review	Method of data synthesis (narrative/meta-analysis)
Biagioli <i>et al</i> <sup>19</sup>	Infants/n=167/<5 weeks/males 45.5%, 44.4%, 49.4%	7 days	4 RCTs	Meta-analysis
Gordon <i>et al</i> <sup>18</sup>	Infants/n=1121/2–16 weeks/balanced numbers of boys and girls	4–21 days	1 RCT	Narrative
Hall <i>et al</i> <sup>34</sup>	Infants/n=309 (27/32 and 83/92 and 199/267)/2–8 weeks/gender not reported	Not reported	3 RCTs	Narrative
Salvatore <i>et al</i> <sup>29</sup>	1 consensus review, 1 review, 1 SR	NA	NA	Narrative
Xu <i>et al</i> <sup>58</sup>	Infants/n=423/3–6 months	<i>L. reuteri</i> . 21 and 28 days	6 RCTs	Meta-analysis
Urbańska and Szajewska <sup>59</sup>	Infants/n=838/birth to 26 weeks	<i>L. reuteri</i> . Up to 1 month for management group and up to 3 months for the prevention group	5 RCTs	Meta-analysis

NA, not available; RCT, randomised controlled trial; SR, systematic review.

of the treatments given. The risk with probiotic was very low.<sup>17</sup>

The data for simethicone and proton pump inhibitors were unfavourable with five reviews concluding either no difference or worsening of symptoms with the use of simethicone. One review concluded no significant differences in crying time or episodes with proton pump inhibitors compared with a placebo, but there was evidence of serious adverse events with the proton pump inhibitor group (one RCT).<sup>42</sup> Other older reviews have concluded the same.<sup>22</sup>

We found few systematic reviews assessing the effectiveness of PPIs for colic despite their increasing use in this population.<sup>21 44</sup> This is because most studies investigating the use of PPIs were for GORD with symptoms of colic. We are aware, however, of the practice of using PPIs to diagnose GORD by treatment.<sup>44</sup> PPIs are designed to suppress acid but have consistently been shown to be ineffective for irritability and fussing in infants with GORD, and they are associated with an increased risk of adverse effects such as infections, allergies and hospital admissions.<sup>9 21 35 45 46</sup> There is evidence from Australia to show that PPIs are being over-prescribed for infants with physiological reflux and symptoms of colic who may or do not have GORD. The reasons are complex, but the authors suggest they centre around inconsistent diagnostic criteria and

diagnostic labelling and ‘defensive’ medicine practice to substantiate the diagnosis with the expectation to medicate.<sup>47</sup> None of the guidance reviewed recommended the use of PPIs for the treatment of colic.

We found three nationally representative guidance, and the only other guidance we found was directed at parents with unclear sources of evidence and guidance justification. Clinical evaluation, information, advice, support and reassurance were the only guidance that was agreed in all four guidelines. Three of the four guidelines recommended to continue to breast feed and use physical contact and not to recommend simethicone and manual therapy, despite the difference in current evidence between them, that is, favourable moderate to low quality for manual therapy and unfavourable low quality for simethicone. Despite the stronger evidence for probiotics in breastfed infants, this was only recommended as a treatment to consider in the USA and Irish guidance. The Canadian Paediatric Association issued a position statement in 2012 which was updated in 2019 stating: “While there may be a role for probiotics in treating infantile colic, there is insufficient evidence to recommend for or against using probiotics to manage this condition”.<sup>48</sup>

Only the USA guidance specifically states not to use proton pump inhibitors for the treatment of colic. We do not know the level of proton pump inhibitors use in

**Table 4** Characteristics of studies for proton pump inhibitors for infantile colic

Authors	Participants/n/age/gender	Timing of intervention	Number and type of studies included in review
Gieruszczak-Białek <i>et al</i> <sup>35</sup>	Infants/n=404/<12 months	2–4 weeks	5 RCTs Narrative

RCT, randomised controlled trial.

**Table 5** Characteristics of studies for mixed intervention studies for infantile colic

Authors	Intervention	Participants/n/age/gender	Timing of intervention	Number and type of studies included in review	Method of data synthesis (narrative/meta-analysis)
Gutiérrez-Castrellón <i>et al</i> <sup>30</sup>	Probiotics ( <i>L. reuteri</i> )	Infants/n=347/4.1–7.2 weeks/56% female	21 and 28 days	5 RCTs	Meta-analysis
	Manual therapy	Infants/n=257/5.1–5.9 weeks/56% female	11–28 days	5 RCTs	Meta-analysis
	Simethicone	Infants/n=14/4.1 weeks/56% female	17 days	1 RCT	Meta-analysis
Harb <i>et al</i> <sup>31</sup>	Probiotics ( <i>L. reuteri</i> )	Infants less than 6 months old/n=472	21 and 30 days	7: 5 double-blind RCTs, 1 single blind RCT, 1 prospective RCT	Meta-analysis
	Simethicone	Infants 2 to 8 weeks old/n=193	Not reported	3 RCTs (simethicone was intervention in 1 and control in 2)	Narrative
Lucassen <sup>2</sup>	Manual therapy	Infants/n=not reported/1–9 weeks/gender not reported	10–14 days	1 SR	Narrative
	Probiotics ( <i>L. reuteri</i> )	Infants/n=126/2–16 weeks/gender not reported	Not reported	1 SR	Narrative
Perry <i>et al</i> <sup>32</sup>	Probiotics ( <i>L. reuteri</i> , <i>L. rhamnosus</i> and <i>P. freudenreichii</i> )	Infants/n=101/0–16 weeks/M=47, F=45	14 days and 28 days	2 RCTs	Narrative
	Manual therapy	Infants/n=not reported /0–16 weeks/gender not reported	Not reported	4 RCTs	Narrative

RCT, randomised controlled trial; SR, systematic review.



**Table 6** Summary of meta-analyses of treatments for infant colic measured by crying time

Author	Time point	24 hours crying time	Effect	Level of evidence	AMSTAR
<b>Probiotics*</b>					
Ong <i>et al</i> <sup>17</sup>	30–90 days	–32.57 min (95% CI –55.6 to –9.54)	Favourable	Low	7
Dryl and Szajewska <sup>54</sup>	21 days and 28 days	–49 min (95% CI –66 to –33) for <i>L. reuteri</i> only	Favourable	High	5.5
Sung <i>et al</i> <sup>57</sup>	21 days	–25.4 min (95% CI –47.3 to –3.5)	Favourable	High	6.5
Gutiérrez-Castrellón <i>et al</i> <sup>30</sup>	7–28 days	WMD –51.3 min (95% CI –30.5 to –72.2 min), p=0.0001	Favourable	High	7
Schreck Bird <i>et al</i> <sup>55</sup>	21 days 28 days	2.3 times more likely to have a ≥50% reduction (p=0.01)	Favourable	High	6
Harb <i>et al</i> <sup>31</sup>	21 and 30 days	–55.8 min (95% CI –64.4 to –47.3)	Favourable	High	8
Xu <i>et al</i> <sup>58</sup>	14 days 21 days	–42.89 min (95% CI –60.50 to –25.29; p=0.000) –45.83 min (95% CI –59.45 to –32.21; p=0.000)	Favourable	High	7
Urbańska and Szajewska <sup>59</sup>	21 days 1 month	–43 min/day (95% CI –68 to –19)	Favourable	High	4.5
Sung <i>et al</i> <sup>56</sup>	21 days	–65 min (95% CI –86 to –44)	Inconclusive	Low	8
Anabrees <i>et al</i> <sup>52</sup>	21 and 28 days	–56.03 min (95% CI –59.92 to –52.15), RR of 0.06 (95% CI 0.01 to 0.25) NNT 2	Favourable	Moderate	8
<b>Simethicone</b>					
Gutiérrez-Castrellón <i>et al</i> <sup>30</sup>	7–28 days	WMD –30.0 min (95% CI –20.8 to –39.0 min), p=0.001 for drugs including simethicone	Inconclusive	Moderate	7
Biagioli <i>et al</i> <sup>19</sup>	7 days	–0.13 hours (95% CI –1.40 to 1.14)	Not favourable	Low	8
<b>Manual therapy</b>					
Carnes <i>et al</i> <sup>43</sup>	1–4 weeks	–1.27 hours (95% CI –2.19 to –0.36)	Favourable	Moderate	8
Gutiérrez-Castrellón <i>et al</i> <sup>30</sup>	7–28 days	WMD –37.4 min (95% CI –21.5 to –67.0 min), p=0.001	Inconclusive	Moderate	7
Dobson <i>et al</i> <sup>16</sup>	8 days to 4 weeks	–1.2 hours (95% CI –1.89 to 0.51)	Favourable	Low	8
Driehuis <i>et al</i> <sup>42</sup>	8–10 days	–0.33 hours per day (95% CI –0.012 to 0.59)	Favourable	Low	8

\*Predominantly breastfed infants.

NNT, number needed to treat; RR, risk ratio; WMD, weighted mean difference.

the infantile colic population; further research is needed to understand the prescribing practice in this field and the impact on the infants. Conversely, UK NICE guidance development group for the diagnosis and management of GORD stated that: “it was not unreasonable to offer a trial of either an  $H_2RA$  or a PPI for suspected reflux oesophagitis in infants without endoscopic evidence and that the clinical presentation of reflux oesophagitis would usually be obvious” (NICE 2015).

Interestingly, the guidance and the evidence do not reflect each other; this may be due to the timing of published evidence and guideline development. The lack of consistent guidance available for parents, pharmacists

and clinicians compounds the uncertainty relating to the care of infants with the symptoms of colic.<sup>49</sup>

We did not find any evidence around cost-effectiveness analyses of the treatments and there was not enough data to analyse our other outcomes of interest, sleep and parental distress.

### Strengths, limitations and challenges

This review was limited to assessing four treatments. The two treatments, probiotics and manual therapy, indicated an effect on reducing crying time. We do not know, however, whether this reduction is meaningful to parents or whether it is sufficient to improve the

**Table 7** Summary of meta-analyses of treatments for infant colic measured by number of crying episodes

Author	Intervention	Time point	Number of crying episodes	Effect	Level of evidence	AMSTAR
Gordon <i>et al</i> <sup>18</sup>	Simethicone	14 days	3.32 more episodes of crying and fussing with simethicone compared with partially hydrolysed formula	Not favourable c/w partially hydrolysed formula	Very low	8
Mugambi <i>et al</i> <sup>25</sup>	Probiotics	Up to 7 months	MD 0.60 (95% CI 0.20 to 1.00) in favour of control for 1 out of 4 studies reported	Not favourable	Low	8
Skórka <i>et al</i> <sup>26</sup>	Probiotics	4 weeks to 36 months	Colic symptoms and crying	Inconclusive (not favourable)	Low	7

MD, mean difference.

parent/infant relationship and/or the well-being of the infants, their parents and siblings. We did not compare manual therapy against other therapies such as dietary modification, herbal remedies, white noise, winding, swaddling and baby massage as suggested in some of the guidance. These other remedies and approaches do not appear to have a strong evidence base but are perhaps deemed to be relatively risk free by the expert

consensus guidance panels. Although one recent editorial suggests that herbal mixtures and swaddling, while potentially effective, may be harmful because herbal mixtures may affect optimal milk consumption and swaddling may increase hip dysplasia.<sup>9</sup> Simethicone and proton pump inhibitors were not considered as risk free and were not recommended in the guidance either explicitly or implicitly.

**Table 8** Summary of narrative analyses of treatments for infant colic measured by crying time

Author	vs control	Time point	Effect	Level of evidence	AMSTAR
<b>Probiotics</b>					
Batchelor <i>et al</i> <sup>53</sup>	Not reported	21 days and 90 days	Not favourable	Not reported	2.5
Cruchet <i>et al</i> <sup>33</sup>	Simethicone	7, 14 and 21 days	Favourable	High prevention, Moderate treatment	4.5
Lucassen <sup>2</sup>	Placebo	21 days	Inconclusive	Very Low	2
Perry <i>et al</i> <sup>32</sup>	Placebo and Simethicone	14 and 28 days	Inconclusive	Low	8
<b>Simethicone</b>					
Harb <i>et al</i> <sup>31</sup>	Placebo, mint and probiotics	17–21 days	Not favourable	Not reported	8
Hall <i>et al</i> <sup>34</sup>	Placebo	2–8 weeks	Not favourable	Low	4
<b>Manual therapy</b>					
Alcantara <i>et al</i> <sup>38</sup>	Dimethicone, no treatment, occipitosacral decompression	14 days and 4 weeks where reported	Inconclusive	Not reported	5
Ernst <sup>41</sup>	Not reported	8 days to 2 weeks	Not favourable	Low	4.5
Lucassen <sup>2</sup>	No treatment	10–14 days	Unfavourable	Low	2
Perry <i>et al</i> <sup>32</sup>	Dimethicone or no treatment	8 days to 4 weeks	Inconclusive	Low	8
Clar <i>et al</i> <sup>39</sup>	Not reported	Not reported	Inconclusive (favourable)	Low to moderate	8
Gleberzon <i>et al</i> <sup>36</sup>	Not reported	<4 weeks	Favourable	Low	6
Parnell Prevost <i>et al</i> <sup>40</sup>	Not reported	Not reported	Inconclusive	Moderate to high	7
Posadzki <i>et al</i> <sup>37</sup>	Not reported	4 weeks	Favourable (1 RCT only)	Low	7
<b>Proton pump inhibitors</b>					
Gieruszczak-Białek <i>et al</i> <sup>35</sup>	Placebo	2–4 weeks	Not favourable	Not reported	3.5

RCT, randomised controlled trial.

**Table 9** National guideline recommendations infantile colic

Recommendation/suggestion	UK 2013 2017*	USA 2015†	Ireland 2014‡
Clinician evaluation of mother and baby	✓	✓	✓
Parenting information, advice, support and reassurance	✓	✓	✓
Continue breast feeding	✓	✓	
Maternal diet modification	×	✓	
Change formula if formula fed (+unless milk allergy identified)	×+	✓	
Probiotic supplements (++)breastfed-only infants)	×	✓++	✓
Simethicone (eg, infacol)	×	×	
Herbal supplements (eg, fennel)	×	×	
Proton pump inhibitors (eg, omeprazole, Losec)		×	
Lactase (eg, Co-lief drops)	×		
Aniticholinergic medication (including dicyclomine)		×	
Gripe water		×	
<b>Medicine generally</b>			
Infant massage			✓
Manual therapy (including spinal manipulation and cranial osteopathy)	×	×	
Physical contact (eg, holding, rocking)	✓		✓
White noise	✓		
Bathing	✓		
Winding	✓		
Swaddling		×	
Acupuncture		×	
Sleep routine			

Blank boxes indicate no recommendation or suggestion made not considered or reported. Ticks indicate recommended interventions; cross indicates non-recommended interventions. \*UK NICE [www.nice.org.uk](http://www.nice.org.uk) Clinical Knowledge Summary Infant Colic 2017 and *Postnatal care: routine postnatal care of women and their babies* (NICE Guideline, 2015). †USA 2015 American Academy of Family Physicians (<http://www.aafp.org>). ‡Ireland: Irish College of General Practitioners (<https://www.icgp.ie>).

In addition, it was not always clear what manual treatments were given and how many; therefore, we are unable to comment on type or quantity of manual treatment that may or may not have an effect.

We reviewed data published in the last decade to avoid over-duplication and we only reviewed articles published in English in peer-reviewed journals; this may have introduced some publication bias.

We suspect that the number and diversity of treatments that exist for infantile colic are borne out of desperation and lack of understanding about the aetiology of the colic and its persistence, although self-limiting in most cases. It has also been suggested that this has led to heterogeneous outcomes being measured,<sup>11</sup> but several reviews were able to meta-analyse reduction in crying time. None of the reviews measured the well-being or confidence of the parents in managing their infants, or relationships and bonding between the parents and the infants and/or their siblings.

The strength of this review is that we have compared at least two treatments—probiotics and manual therapy—for colic and compared the recommendations in the guidance with the evidence which highlighted the difference in certainty between the evidence and the recommendations. This review found favourable but low to moderate quality evidence for manual therapy. This is different to other reviews, which generally reject manual therapy as an option due to poor quality data. New studies have raised the quality level of data and our more favourable interpretation may, in part, be due to the background of the authors and their understanding that manual therapy is a multicomponent therapy consisting of more than touch alone. The role of the manual therapist to reassure, guide, advise and support parents through this particularly difficult time may also have a therapeutic role to play which may affect parenting and outcomes. More high-quality studies are needed to increase the level of certainty surrounding the findings about the effect of manual therapy for infantile colic.

Future studies should consider the effect of treatment on the parents to explore parenting confidence, parent/infant bonding and meaningful levels of change in crying and sleeping time.

Further research will probably change recommendations in the guidance as the evidence evolves and increases in quality. The guidance we reviewed did not reflect the emerging evidence. Overall, there is uncertainty about the management and care of infants with infantile colic in part due to the lack of consensus surrounding its aetiology and justification for treatments, although increasingly, newer research is providing plausible mechanistic explanations why probiotics may be beneficial due to the presence of gut inflammatory markers.<sup>50 51</sup> Infantile colic is self-limiting in nature and understanding what symptoms and the level of reduction of these symptoms that are important to the infants and their carers needs further investigating. It is understandable that a range of approaches to treatment, management and care exist. Preserving parent choice and balancing this with the limited evidence about effectiveness and safety remains difficult to determine.

## CONCLUSIONS

We found that the strongest evidence for the treatment of infantile colic was probiotics, particularly *Lactobacillus reuteri* for breastfed infants, followed by weaker but

favourable evidence for manual therapy indicated by crying time. Both forms of treatment carried a low risk of serious adverse events. Current guidelines will probably change over time in light of existing new and emerging evidence.

**Contributors** DC conceptualised and designed the study, contributed to the data selection, extraction and analysis, drafted the initial manuscript, reviewed and revised the manuscript. JE managed the data, contributed to the data selection, extraction and analysis, reviewed and revised the manuscript. JD-R contributed to the data selection, extraction and analysis, reviewed and revised the manuscript. All the authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

**Funding** This review was funded primarily by the Australian College of Chiropractic Paediatrics (ACCP) with part of the funding provided by the College of Chiropractic Paediatrics (CCP).

**Disclaimer** ACCP and CCP were not involved in the literature search, extraction of data, analysis and synthesis.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Full datasets, analyses and full searches are available on request from the corresponding author (Dawn.Carnes@uco.ac.uk). No individual patient-level data were used in this study.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Dawn Carnes <http://orcid.org/0000-0002-3152-3133>

## REFERENCES

- Lucassen PL, Assendelft WJ. Systematic review of treatments for infant colic. *Pediatrics* 2001;108:1047–8.
- Lucassen P. Colic in infants. *BMJ Clin Evid* 2015;2015. [Epub ahead of print: 11 Aug 2015].
- Wolke D, Bilgin A, Samara M. Systematic review and meta-analysis: fussing and crying durations and prevalence of colic in infants. *J Pediatr* 2017;185:55–61.
- Heine RG. Gastroesophageal reflux disease, colic and constipation in infants with food allergy. *Curr Opin Allergy Clin Immunol* 2006;6:220–5.
- Steutel NF, Benninga MA, Langendam MW, et al. Reporting outcome measures in trials of infant colic. *J Pediatr Gastroenterol Nutr* 2014;59:341–6.
- Wessel MA, Cobb JC, Jackson EB, et al. Paroxysmal fussing in infancy, sometimes called colic. *Pediatrics* 1954;14:421–35.
- Zeevenhooven J, Browne PD, L'Hoir MP, et al. Infant colic: mechanisms and management. *Nat Rev Gastroenterol Hepatol* 2018;15:479–96.
- Benninga MA, Faure C, Hyman PE, et al. Childhood functional gastrointestinal disorders: neonate/toddler. *Gastroenterology* 2016;145:5.
- Sung V. Infantile colic. *Aust Prescr* 2018;41:105–10.
- Halpern R, Coelho R. Excessive crying in infants. *J Pediatr* 2016;92:S40–5.
- Steutel NF, Benninga MA, Langendam MW, et al. Developing a core outcome set for infant colic for primary, secondary and tertiary care settings: a prospective study. *BMJ Open* 2017;7:e015418.
- Johnson JD, Cocker K, Chang E. Infantile colic: recognition and treatment. *Am Fam Physician* 2015;92:577–82.
- Kurth E, Spichiger E, Cignacco E, et al. Predictors of crying problems in the early postpartum period. *J Obstet Gynecol Neonatal Nurs* 2010;39:250–62.
- Howard CR, Lanphear N, Lanphear BP, et al. Parental responses to infant crying and colic: the effect on breastfeeding duration. *Breastfeed Med* 2006;1:146–55.
- Overpeck MD, Brenner RA, Trumble AC, et al. Risk factors for infant homicide in the United States. *N Engl J Med* 1998;339:1211–6.
- Dobson D, Lucassen PLBJ, Miller JJ, et al. Manipulative therapies for infantile colic. *Cochrane Database Syst Rev* 2012;12:CD004796.
- Ong TG, Gordon M, Banks SS, et al. Probiotics to prevent infantile colic. *Cochrane Database Syst Rev* 2019;3:CD012473.
- Gordon M, Biagioli E, Sorrenti M, et al. Dietary modifications for infantile colic. *Cochrane Database Syst Rev* 2018;10:CD011029.
- Biagioli E, Tarasco V, Lingua C, et al. Pain-relieving agents for infantile colic. *Cochrane Database Syst Rev* 2016;9:CD009999.
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700.
- Safe M, Chan WH, Leach ST, et al. Widespread use of gastric acid inhibitors in infants: are they needed? Are they safe? *World J Gastrointest Pharmacol Ther* 2016;7:531–9.
- Garrison MM, Christakis DA. A systematic review of treatments for infant colic. *Pediatrics* 2000;106:184–90.
- Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017;358:j4008.
- Brouwers MC, Kho ME, Browman GP, et al. AGREE II: advancing guideline development, reporting, and evaluation in health care. *Prev Med* 2010;51:421–4.
- Mugambi MN, Musekiwa A, Lombard M, et al. Synbiotics, probiotics or prebiotics in infant formula for full term infants: a systematic review. *Nutr J* 2012;11:81.
- Skórka A, Pieścik-Lech M, Kotodziej M, et al. To add or not to add probiotics to infant formulae? An updated systematic review. *Benef Microbes* 2017;8:717–25.
- Posadzki P, Ernst E. Spinal manipulation: an update of a systematic review of systematic reviews. *N Z Med J* 2011;124:55–71.
- Salehi A, Hashemi N, Imanieh MH, et al. Chiropractic: is it efficient in treatment of diseases? Review of systematic reviews. *Int J Community Based Nurs Midwifery* 2015;3:244–54.
- Salvatore S, Barberi S, Borrelli O, et al. Pharmacological interventions on early functional gastrointestinal disorders. *Ital J Pediatr* 2016;42:68.
- Gutiérrez-Castrellón P, Indrio F, Bolio-Galvis A, et al. Efficacy of Lactobacillus reuteri DSM 17938 for infantile colic: systematic review with network meta-analysis. *Medicine* 2017;96:e9375.
- Harb T, Matsuyama M, David M, et al. Infant colic—what works: a systematic review of interventions for breast-fed infants. *J Pediatr Gastroenterol Nutr* 2016;62:668–86.
- Perry R, Hunt K, Ernst E. Nutritional supplements and other complementary medicines for infantile colic: a systematic review. *Pediatrics* 2011;127:720–33.
- Cruchet S, Furnes R, Maruy A, et al. The use of probiotics in pediatric gastroenterology: a review of the literature and recommendations by Latin-American experts. *Paediatr Drugs* 2015;17:199–216.
- Hall B, Chesters J, Robinson A. Infantile colic: a systematic review of medical and conventional therapies. *J Paediatr Child Health* 2012;48:128–37.
- Gieruszczak-Białek D, Konarska Z, Skórka A, et al. No effect of proton pump inhibitors on crying and irritability in infants: systematic review of randomized controlled trials. *J Pediatr* 2015;166:767–70.
- Gleberzon BJ, Arts J, Mei A, et al. The use of spinal manipulative therapy for pediatric health conditions: a systematic review of the literature. *J Can Chiropr Assoc* 2012;56:128–41.
- Posadzki P, Lee MS, Ernst E. Osteopathic manipulative treatment for pediatric conditions: a systematic review. *Pediatrics* 2013;132:140–52.
- Alcantara J, Alcantara JD, Alcantara J. The chiropractic care of infants with colic: a systematic review of the literature. *Explore* 2011;7:168–74.
- Clar C, Tsertsvadze A, Court R, et al. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: systematic review and update of UK evidence report. *Chiropr Man Therap* 2014;22:12.
- Parnell Prevost C, Gleberzon B, Carleo B, et al. Manual therapy for the pediatric population: a systematic review. *BMC Complement Altern Med* 2019;19:60.
- Ernst E. Chiropractic spinal manipulation for infant colic: a systematic review of randomised clinical trials. *Int J Clin Pract* 2009;63:1351–3.
- Driehuis F, Hoogeboom TJ, Nijhuis-van der Sanden MWG, et al. Spinal manual therapy in infants, children and adolescents: a systematic review and meta-analysis on treatment indication, technique and outcomes. *PLoS One* 2019;14:e0218940.

- 43 Carnes D, Plunkett A, Ellwood J, *et al.* Manual therapy for unsettled, distressed and excessively crying infants: a systematic review and meta-analyses. *BMJ Open* 2018;8:e019040.
- 44 Blank M-L, Parkin L. National study of off-label proton pump inhibitor use among New Zealand infants in the first year of life (2005-2012). *J Pediatr Gastroenterol Nutr* 2017;65:179-84.
- 45 Duncan DR, Mitchell PD, Larson K, *et al.* Association of proton pump inhibitors with hospitalization risk in children with oropharyngeal dysphagia. *JAMA Otolaryngol Head Neck Surg* 2018;144:1116-24.
- 46 Mitre E, Susi A, Kropp LE, *et al.* Association between use of Acid-Suppressive medications and antibiotics during infancy and allergic diseases in early childhood. *JAMA Pediatr* 2018;172:e180315.
- 47 Bell JC, Schneuer FJ, Harrison C, *et al.* Acid suppressants for managing gastro-oesophageal reflux and gastro-oesophageal reflux disease in infants: a national survey. *Arch Dis Child* 2018;103:archdischild-2017-314161.
- 48 Marchand V. Using probiotics in the paediatric population. *Paediatr Child Health* 2012;17:575-6.
- 49 Hinds R, Loveridge N, Lemberg DA, *et al.* Functional gastrointestinal disorders in infants: practice, knowledge and needs of Australian pharmacists. *J Paediatr Child Health* 2019. doi:10.1111/jpc.14536. [Epub ahead of print: 10 Jun 2019].
- 50 Rhoads JM, Collins J, Fatheree NY, *et al.* Infant colic represents gut inflammation and dysbiosis. *J Pediatr* 2018;203:55-61.
- 51 Pärtty A, Kalliomäki M, Salminen S, *et al.* Infantile colic is associated with low-grade systemic inflammation. *J Pediatr Gastroenterol Nutr* 2017;64:691-5.
- 52 Anabrees J, Indrio F, Paes B, *et al.* Probiotics for infantile colic: a systematic review. *BMC Pediatr* 2013;13:186.
- 53 Batchelor N, Kelly J, Choi H, *et al.* Towards evidence-based emergency medicine: best BETs from the Manchester Royal Infirmary. BET 2: probiotics and crying time in babies with infantile colic. *Emerg Med J* 2015;32:575-6.
- 54 Dryl R, Szajewska H. Probiotics for management of infantile colic: a systematic review of randomized controlled trials. *Arch Med Sci* 2018;14:1137-43.
- 55 Schreck Bird A, Gregory PJ, Jalloh MA, *et al.* Probiotics for the treatment of infantile colic: a systematic review. *J Pharm Pract* 2017;30:366-74.
- 56 Sung V, Collett S, de Gooyer T, *et al.* Probiotics to prevent or treat excessive infant crying: systematic review and meta-analysis. *JAMA Pediatr* 2013;167:1150-7.
- 57 Sung V, D'Amico F, Cabana MD, *et al.* *Lactobacillus reuteri* to treat infant colic: a meta-analysis. *Pediatrics* 2018;141:e20171811.
- 58 Xu M, Wang J, Wang N, *et al.* The efficacy and safety of the probiotic bacterium *Lactobacillus reuteri* DSM 17938 for infantile colic: a meta-analysis of randomized controlled trials. *PLoS One* 2015;10:e0141445.
- 59 Urbaiska M, Szajewska H. The efficacy of *Lactobacillus reuteri* DSM 17938 in infants and children: a review of the current evidence. *Eur J Pediatr* 2014;173:1327-37.