



ERRATUM

Frailty is independently associated with worse health-related quality of life in chronic kidney disease: a secondary analysis of the Frailty Assessment in Chronic Kidney Disease study

Andrew C. Nixon ^{1,2,3}, Theodoros M. Bampouras^{4,5}, Neil Pendleton⁶, Sandip Mitra^{7,8}, Mark E. Brady¹ and Ajay P. Dhaygude¹

¹Department of Renal Medicine, Lancashire Teaching Hospitals NHS Foundation Trust, Preston, UK, ²Centre for Health Research and Innovation, National Institute of Health Research Lancashire Clinical Research Facility, Lancashire Teaching Hospitals NHS Foundation Trust, Preston, UK, ³Division of Cardiovascular Sciences, University of Manchester, Manchester, UK, ⁴Active Ageing Research Group, University of Cumbria, Lancaster, UK, ⁵Lancaster Medical School, Lancaster University, Lancaster, UK, ⁶Division of Neuroscience and Experimental Psychology, University of Manchester, Manchester, UK, ⁷Manchester Academy of Health Sciences Centre, University of Manchester, Manchester, UK and ⁸Devices for Dignity, National Institute of Health Research MedTech & In-vitro Diagnostics Co-operative, UK

Correspondence and offprint requests to: Andrew C. Nixon; E-mail: andrew.nixon@lthtr.nhs.uk

In the original article, in the 'Methods' section of the abstract, there were some incorrect words added. It originally read 'Ninety participants with dialysis-dependent chronic kidney disease (CKD G4–5D) were recruited between December 2016 and December 2017'—the words 'dialysis-dependent' have been removed. The Publisher apologizes for the error.