















## Research review

## Enhancing crop yields through improvements in the efficiency of photosynthesis and respiration

Author for correspondence:  
Owen Atkin  
Email: owen.atkin@anu.edu.au

Received: 25 April 2022  
Accepted: 15 September 2022

Andres Garcia<sup>1,2</sup> , Oorbessy Gaju<sup>1,3</sup> , Andrew F. Bowerman<sup>1,2</sup> ,  
Sally A. Buck<sup>1,2</sup> , John R. Evans<sup>2,4</sup> , Robert T. Furbank<sup>2,4</sup> ,  
Matthew Gilliham<sup>5</sup> , A. Harvey Millar<sup>6</sup> , Barry J. Pogson<sup>1,2</sup> ,  
Matthew P. Reynolds<sup>7</sup> , Yong-Ling Ruan<sup>2</sup> , Nicolas L. Taylor<sup>6</sup> ,  
Stephen D. Tyerman<sup>5</sup>  and Owen K. Atkin<sup>1,2</sup> 

<sup>1</sup>ARC Centre of Excellence in Plant Energy Biology, Research School of Biology, The Australian National University, Canberra, ACT 2601, Australia; <sup>2</sup>Division of Plant Sciences, Research School of Biology, Australian National University, Canberra, ACT 2601, Australia; <sup>3</sup>College of Science, Lincoln Institute for Agri-Food Technology, University of Lincoln, Lincolnshire, LN2 2LG, UK; <sup>4</sup>ARC Centre of Excellence for Translational Photosynthesis, Research School of Biology, The Australian National University, Canberra, ACT 2601, Australia; <sup>5</sup>ARC Centre of Excellence in Plant Energy Biology, School of Agriculture, Food and Wine & Waite Research Institute, University of Adelaide, Glen Osmond, SA 5064, Australia; <sup>6</sup>ARC Centre of Excellence in Plant Energy Biology, School of Molecular Sciences & Institute of Agriculture, The University of Western Australia, Crawley, WA 6009, Australia; <sup>7</sup>International Maize and Wheat Improvement Center (CIMMYT), Km. 45, Carretera Mexico, El Batán, Texcoco, 56237, Mexico

## Summary

The rate with which crop yields per hectare increase each year is plateauing at the same time that human population growth and other factors increase food demand. Increasing yield potential ( $Y_p$ ) of crops is vital to address these challenges. In this review, we explore a component of  $Y_p$  that has yet to be optimised – that being improvements in the efficiency with which light energy is converted into biomass ( $\epsilon_c$ ) via modifications to  $\text{CO}_2$  fixed per unit quantum of light ( $\alpha$ ), efficiency of respiratory ATP production ( $\epsilon_{\text{prod}}$ ) and efficiency of ATP use ( $\epsilon_{\text{use}}$ ). For  $\alpha$ , targets include changes in photoprotective machinery, ribulose biphosphate carboxylase/oxygenase kinetics and photorespiratory pathways. There is also potential for  $\epsilon_{\text{prod}}$  to be increased via targeted changes to the expression of the alternative oxidase and mitochondrial uncoupling pathways. Similarly, there are possibilities to improve  $\epsilon_{\text{use}}$  via changes to the ATP costs of phloem loading, nutrient uptake, futile cycles and/or protein/membrane turnover. Recently developed high-throughput measurements of respiration can serve as a proxy for the cumulative energy cost of these processes. There are thus exciting opportunities to use our growing knowledge of factors influencing the efficiency of photosynthesis and respiration to create a step-change in yield potential of globally important crops.

New Phytologist (2023) 237: 60–77  
doi: 10.1111/nph.18545

**Key words:** alternative oxidase (AOX), carbon balance, photosynthesis, respiration, yield potential.

## Introduction

Anthropogenic carbon emissions since the Industrial Revolution have led to sustained global warming, with 2020 tying 2016 as the hottest year on record, at 1.25°C above preindustrial times (Voosen, 2021). How climate change affects our natural and managed systems is of major concern, particularly the effect of warming on major agricultural crop production. To keep pace with increasing crop demand for human and animal

consumption – which will potentially double between 2010 and 2050 (van Dijk *et al.*, 2021) – and to minimise the demand for arable land, grain yields per hectare need to increase by 2.4% each year. Yet, for the period 1989–2008, actual annual increases fell markedly short of this figure, being 1.6%, 1.0%, 0.9% and 1.3% for maize, rice, wheat and soybean, respectively (Ray *et al.*, 2013). Therefore, it is imperative that we improve major crop yields (Smith, 2013). This goal can be achieved through increasing both yield resilience in suboptimal

conditions (Munns & Gilliam, 2015; Rivero *et al.*, 2022) and yield potential, discussed herein.

To date, increases in crop yields have been driven by Green Revolution cultivars – that lent themselves to more intensive irrigation, fertilisation and mechanisation – and continued selective breeding (Tilman, 1998; Crespo-Herrera *et al.*, 2018). These strategies doubled wheat yield and tripled that of maize between 1961 and 2000 – outcomes that were achieved with only a 30% increase in cultivated land area (World Bank, 2007; Hazell, 2009). Global wheat production grew from 222 Mt in 1961 to 585 Mt in 2000 and then to 735 Mt as of 2018. However, in recent years, yield per hectare has plateaued and further investment in the current strategies may be ineffective at further boosting yield (Tester & Langridge, 2010; Móznér *et al.*, 2012; Iizumi *et al.*, 2017). To avoid farming more land and using more fertilisers, it is essential that new pathways are found to boost crop yields. Improving yield potential ( $Y_p$ ) is one such opportunity.

Breaking  $Y_p$  into components allows assessment of targeted multiplicative improvements.  $Y_p$  is the product of several underlying components:

$$Y_p = Q \times \epsilon_i \times \epsilon_c \times \epsilon_p \quad \text{Eqn 1}$$

where  $Q$  is the cumulative incident solar radiation ( $\text{MJ m}^{-2}$ ) throughout the growing season;  $\epsilon_p$  is the partitioning efficiency (commonly referred to as harvest index), the proportion of biomass allocated to harvested product;  $\epsilon_i$  is the efficiency with which the plant intercepts/absorbs radiation; and  $\epsilon_c$  is the efficiency with which the intercepted radiation is converted into biomass (Monteith, 1977; Long *et al.*, 2015) (see Box 1 for a mathematical expansion of  $\epsilon_c$ , and Fig. 1 for examples of factors that influence each underlying component of  $Y_p$ ). In theory, an increase in any of these factors would increase overall  $Y_p$ . However, not every term of the equation holds promise for a significant improvement in future years (Loomis & Amthor, 1999).

Both  $\epsilon_p$  and  $\epsilon_i$  have improved greatly since the Green Revolution through selective breeding, approaching 0.6 and 0.8–0.9, respectively (Zhu *et al.*, 2010). Increasing crop yields throughout the second half of the 20<sup>th</sup> century came with a concomitant increase in harvest index, attributable to biomass allocation patterns and dwarf genes that resulted in shorter stems that could support more grain (Sinclair, 1998). Meanwhile, because higher  $\epsilon_i$  increases the amount of incident solar radiation available for photosynthesis, improvements in canopy architecture and responses to light have been targets for selection. In many field-grown crops, plants with leaves perpendicular to incoming light will intercept most of the solar radiation at the uppermost layer of the canopy, shading lower levels and preventing them from achieving light-saturated levels of photosynthesis (Campbell & Ogren, 1990; Ort & Long, 2003). Structuring canopies such that leaves higher in the canopy arrange themselves at steeper angles allows for deeper penetration of solar radiation, potentially doubling  $\epsilon_i$  at peak irradiance or low latitudes. Through selective breeding during the Green Revolution,  $\epsilon_i$  was increased through reductions in plant stature and lodging.

By comparison, there are still many untapped opportunities for improvements in the underlying factors that influence the efficiency with which solar radiation is converted into biomass (i.e.  $\epsilon_c$ ). Currently, the maximum observed value of  $\epsilon_c$  lies at *c.* 0.024, yet the  $\epsilon_c$  for  $C_3$  crops has a theoretical maximum efficiency of 0.046–0.051 (Monteith, 1977; Long *et al.*, 2006; Zhu *et al.*, 2010). Given this gap,  $\epsilon_c$  values could be strategically targeted as they provide one of the best options for future improvement (Fig. 2). To enable this, opportunities to increase the efficiency of the biochemical processes underpinning  $\epsilon_c$  urgently need to be identified and prioritised.

## Steps to improve the efficiency of net photosynthetic $\text{CO}_2$ fixation

One way of improving  $\epsilon_c$  is to increase the efficiency of photosynthesis (Eqn B3). But how might this be achieved? Ultimately, improvements are needed in the ability of plants to convert absorbed light energy into products (ATP and NADPH) needed to drive  $\text{CO}_2$  fixation by ribulose biphosphate carboxylase/oxygenase (Rubisco) and in minimising carbon loss through processes such as photorespiration, while protecting the system from photoinhibition. Here, we discuss attributes that influence the efficiency of photosynthesis and strategies being employed to improve net photosynthesis.

### Making better use of intercepted light and improving carboxylation efficiency

The variable nature of light availability across space and time in the plant canopy requires consideration of biochemical responses to these fluctuations. Incoming solar radiation above saturating rates can cause photooxidative damage to light-harvesting complexes (Havaux & Niyogi, 1999; Baroli & Niyogi, 2000). Photoprotective mechanisms can decrease the quantum yield of photosystem II, accounting for *c.* 15% decrease in daily canopy carbon uptake, and even higher values under stress conditions (Long *et al.*, 1994; Zhu *et al.*, 2004). Photosynthetic induction, the ramp-up of photosynthetic activity after exposure to light, can delay a leaf from reaching optimal photosynthetic rates under given conditions on the scale of minutes (Retkute *et al.*, 2015; Soleh *et al.*, 2017). Modulating these responses could improve net photosynthetic performance in certain situations (Murchie & Niyogi, 2011). The issue of induction is related to canopy structure because lower canopy layers are more reliant on intermittent light interception for photosynthesis. For example, total canopy photosynthesis could increase up to 17% with increased light penetration to the lower canopy in rice (Burgess *et al.*, 2016). Additionally, the delay of photosynthetic induction ramping up to full capacity can cause a > 20% loss of potential carbon assimilation in wheat (Taylor & Long, 2017), which could potentially be ameliorated through overexpression or amino acid substitutions in Rubisco activase (Yamori *et al.*, 2012; Carmo-Silva & Salvucci, 2013). Because leaf N is not necessarily distributed to maximise photosynthesis, optimising nitrogen distribution throughout the canopy is another strategy for increased photosynthesis, particularly at key growth

**Box 1** Meeting agricultural demand for a growing population requires us to consider the components of yield

Improving the efficiency of these components will boost realised yield to keep pace with that demand. Yield potential  $\left(\frac{\text{g yield}}{\text{m}^2 \text{ ground area}}\right)$  can be thought of as a function of incident solar radiation  $\left(Q, \frac{\text{J light incoming}}{\text{m}^2 \text{ ground area}}\right)$ , efficiency of radiation interception and absorption  $\left(\epsilon_i, \frac{\text{J light absorbed}}{\text{J light incoming}}\right)$ , efficiency of converting intercepted radiation to biomass  $\left(\epsilon_c, \frac{\text{g plant biomass}}{\text{J light absorbed}}\right)$  and partitioning efficiency  $\left(\epsilon_p, \frac{\text{g yield}}{\text{g plant}}\right)$ . These factors are related by the following equation,

$$Y_p = Q \times \epsilon_i \times \epsilon_c \times \epsilon_p. \quad \text{Eqn B1}$$

Currently, the maximum observed value of  $\epsilon_c$  lies c. 0.024 (Monteith, 1977). Yet, the  $\epsilon_c$  for  $C_3$  crops has a theoretical efficiency of 0.046–0.051 (Long *et al.*, 2006; Zhu *et al.*, 2010). With current  $\epsilon_c$  values being only half of their theoretical maximum, there is clearly scope for improvement. Here, improvements in the efficiency of biochemical processes underpinning  $\epsilon_c$  are urgently needed. To understand ways of improving  $\epsilon_c$ , we can mathematically consider which factors contribute to variation in  $\epsilon_c$ . One way of achieving this is to define  $\epsilon_c$  according to:

$$\epsilon_c = \frac{\text{biomass}}{Q_{\text{absorbed}}} = \frac{\text{CO}_2 \text{ fixed}}{Q_{\text{absorbed}}} \times \frac{\text{g plant}}{\text{g CO}_2 \text{ fixed}}, \quad \text{Eqn B2}$$

where  $\epsilon_c$  (and thus the biomass produced per joule of light absorbed) is the product of how much  $\text{CO}_2$  is fixed by photosynthesis per joule of light, multiplied by the amount of plant mass per unit of  $\text{CO}_2$  fixed by photosynthesis. Eqn B2 shows that  $\epsilon_c$  could be improved through targeting aspects of photosynthetic biochemistry to improve the amount of  $\text{CO}_2$  fixed per unit quantum of light. Put another way:

$$\alpha = \frac{\text{CO}_2 \text{ fixed}}{Q_{\text{absorbed}}}. \quad \text{Eqn B3}$$

It also shows that  $\epsilon_c$  can be improved via changes in the efficiency with which plants accumulate biomass per unit of  $\text{CO}_2$  fixed:

$$\epsilon_{\text{biosynthesis}} = \frac{\text{g plant}}{\text{g CO}_2 \text{ fixed}}. \quad \text{Eqn B4}$$

Bringing Eqns B3 and B4 together, we see that  $\epsilon_c$  is related to  $\alpha$  and  $\epsilon_{\text{biosynthesis}}$  according to:

$$\epsilon_c = \alpha \times \epsilon_{\text{biosynthesis}}, \quad \text{Eqn B5}$$

where  $\epsilon_{\text{biosynthesis}}$  is dependent on the efficiency of ATP production (hereon termed  $\epsilon_{\text{prod}}$ ; i.e. how much substrate must be consumed for a certain ATP yield) and the efficiency of ATP use ( $\epsilon_{\text{use}}$ ), with the latter being the ATP yield necessary to drive processes of biosynthesis, transport and cellular maintenance. Bringing these factors together, we see that:

$$\epsilon_{\text{biosynthesis}} = \epsilon_{\text{prod}} \times \epsilon_{\text{use}}, \quad \text{Eqn B6}$$

and therefore:


$$\epsilon_c = \alpha \times \epsilon_{\text{prod}} \times \epsilon_{\text{use}}. \quad \text{Eqn B7}$$

These equations thus show that improvements in  $\epsilon_c$  could be achieved through  $\alpha$ ,  $\epsilon_{\text{prod}}$  and  $\epsilon_{\text{use}}$ . In subsequent sections, we discuss strategies available for improvement in these components of  $\epsilon_c$ .


stages (Johnson *et al.*, 2010; Moreau *et al.*, 2012). Potential strategies to improve induction of photosynthesis include speeding up the rate at which proteins are activated in the photosynthetic electron transport system, increasing carbon fixation capacity via increases in concentrations in Rubisco and Rubisco activase, and improving  $\text{CO}_2$  supply via increases in stomatal and mesophyll conductance (Yamori, 2021). Increasing nonphotochemical quenching (NPQ) relaxation under fluctuating light conditions could also be beneficial, in order to increase the efficiency of light energy use under light-limiting conditions (Yamori, 2021). A recent example is the study by De Souza *et al.* (2022) for soybean; they reported that the overexpression of genes linked to NPQ

relaxation (*AtVDE*, *AtPsbS* and *AtZEP*) improved photosynthetic efficiency of soybean under fluctuating light over 2 yr in field trials. Yield may also be improved, as shown by the 24% higher soybean yields of five independent transgenic lines at the Urbana site in 2020. While further work is needed across multiple sites and years, the work of De Souza *et al.* (2022) does highlight opportunities to improve yield potential via targeting photosynthetic properties such as NPQ relaxation.

Another approach to improve photosynthesis is via targeting the biochemical properties of Rubisco. As a central protein in the Calvin cycle, Rubisco's biochemical properties play a large role in defining photosynthetic performance. Variations in Rubisco's



**Fig. 1** Logic flow of improvements to the carbon economy of plants in order to increase yield potential. Minimising the ratio of respiration to either photosynthesis ( $R:P$ ) or growth ( $R:RGR$ ) increases the amount of photo assimilate available for yield (yellow). This goal can be achieved in two ways (green): by shrinking the numerator (decreased mitochondrial CO<sub>2</sub> loss) or by increasing the denominator (more CO<sub>2</sub> uptake). To date, increases in yield potential have been in part due to improvements in  $\epsilon_i$  (violet). Further improvements will require improving  $\epsilon_c$  through modifying the photorespiratory pathway ( $\alpha$ ; orange), increased  $\epsilon_{prod}$  (blue) or increased  $\epsilon_{use}$  (red). AOX, alternative oxidase; COX, cytochrome c oxidase; SWEET, Sugars Will Eventually be Exported Transporter; UQ, ubiquinone pool.



**Fig. 2** Theoretical efficiencies of yield potential factors. Yield potential can be improved by increasing any of its factors— $\epsilon_i$ ,  $\epsilon_c$  and  $\epsilon_p$ . However, only  $\epsilon_c$  shows appreciable difference between its observed and theoretical efficiency and therefore stands as the best target for improvement. Energy remaining after  $\epsilon_c$  processes is in the form of biomass, and a doubling would hypothetically flow downstream and double energy remaining during  $\epsilon_p$ , that is yield.  $Q$ , incoming solar radiation for growing season.

structure, catalytic activity and interactions with other proteins (such as Rubisco activase) present both opportunities and limitations for engineering improvements in photosynthesis (Whitney *et al.*, 2011; Conlan & Whitney, 2018). Collaborations such as the Realizing Increased Photosynthetic Efficiency project (<https://ripe.illinois.edu/>) highlight the role of strategies such as introducing novel Rubisco types (Prins *et al.*, 2016), selecting for shorter photosynthetic induction (Acevedo-Siaca *et al.*, 2020) and improving stomatal CO<sub>2</sub> response of mesophyll conductance of CO<sub>2</sub> (Bailey-Serres *et al.*, 2019). Hyperspectral reflectance has also shown promising correlations to related traits, such as the maximum catalytic activity of Rubisco ( $V_{\text{cmax}}$ ) and maximum rate

of photosynthetic electron transport ( $J_{\text{max}}$ ) (Silva-Perez *et al.*, 2018; Fu *et al.*, 2020). Focusing on such traits, and using emerging high-throughput, nondestructive methods of quantifying photosynthetic traits, provides opportunities to breed for increased  $\alpha$  and through it, improve  $\epsilon_c$ .

### Modifications to photorespiratory pathways to reduce CO<sub>2</sub> loss

Another route to improving the efficiency of net photosynthesis is via modifications to photorespiration – the pathway through which the oxygenation reaction by Rubisco consumes RuBP to form 3-



PGA and 2-phosphoglycolate (2PG), the latter being converted to glycolate. The 2PG inhibits triose-phosphate isomerase and disrupts the Calvin cycle, emphasising the need not only to minimise Rubisco oxygenation, but also to efficiently metabolise 2PG once produced. This is achieved, in part, via the export of glycolate to the peroxisome where it is converted to glyoxylate and then glycine (resulting in the release of  $\text{H}_2\text{O}_2$ ), with glycine being exported to mitochondria where it is decarboxylated by glycine decarboxylase (GDC) to yield serine,  $\text{NH}_3$ , NAD(H) and  $\text{CO}_2$ . Together, these reactions mean that photorespiration reduces the efficiency of net photosynthesis (and thus  $\alpha$  and  $\epsilon_c$ ). Because of this, minimising Rubisco oxygenation and associated rates of photorespiratory  $\text{CO}_2$  release are potential targets for improving yield potential. Importantly, any strategy to modify photorespiration needs to regulate the production and metabolism of toxic intermediates, the loss of nitrogen as volatile  $\text{NH}_3$ , and also consider the current role of photorespiration in wider amino acid biosynthesis and metabolism (Novitskaya *et al.*, 2002; Abadie & Tcherkez, 2019).

From a carbon metabolism perspective, photorespiration represents an energetically expensive process, either as an opportunity cost during Rubisco oxygenation (every  $\text{O}_2$  bound to Rubisco represents a potential  $\text{CO}_2$  that did not bind), or as  $\text{CO}_2$  release during glycine decarboxylation. Rubisco oxygenation will be influenced by the relative specificity of Rubisco ( $S_{c/o}$ ), leaf temperature and the intercellular and intracellular concentrations of  $\text{CO}_2$  (Parry *et al.*, 2013), with the subsequent rate of photorespiratory  $\text{CO}_2$  release varying depending on which metabolic intermediates are shuttled to other pathways (Peterhansel *et al.*, 2010; Busch, 2020). Reassimilation of photorespiratory  $\text{CO}_2$  by Rubisco will also influence the extent to which photorespiration reduces  $\alpha$  and  $\epsilon_c$  (Sage & Sage, 2009).

Because of its potential to reduce the efficiency of net photosynthesis, many researchers have investigated the effect of mutations to different points in the photorespiratory pathway over the past 30 yr. These studies have revealed a wide range of 'photorespiratory phenotypes' that vary depending on the site of disruption to photorespiratory pathway and the environmental conditions in which plants are grown (e.g. see table 2 in Timm & Bauwe (2013) for a list of mutants and associated phenotypes). In many cases, knockout of key steps in the photorespiratory pathway is deleterious and potentially lethal. For example, in *Arabidopsis thaliana*, the most severe phenotypes show conditional lethality in GDC double knockout mutants (*gldp1*  $\times$  *gldp2*) and serine hydroxymethyltransferase double knockout mutants (*shm1*  $\times$  *shm2*) (Engel *et al.*, 2007, 2011). Other deletions result in a milder photorespiratory phenotype because of functional redundancy with other members in the gene family (Timm & Bauwe, 2013) and/or because of the plasticity of phenotypic response in photorespiratory mutants (Peterhansel *et al.*, 2013; Walker *et al.*, 2016). In any case, the available mutant studies highlight that fully knocking out the entire photorespiratory pathway is not a viable strategy and that photorespiratory modifications (to reduce photorespiratory  $\text{CO}_2$  release) must consider reducing flux through the pathway, while regulating production and metabolism of toxic intermediates.


A range of studies provide examples of how modifications of photorespiration can increase biomass production (see Fig. 3), including: increasing  $\text{CO}_2$  concentration around Rubisco (to reduce the rate of oxygenation) based on the modelled introduction of cyanobacterial carboxysomes; changing associations between chloroplasts and mitochondria; and engineering photorespiratory bypasses (Sage & Sage, 2009; Price *et al.*, 2013; Xin *et al.*, 2015; Betti *et al.*, 2016; Batista-Silva *et al.*, 2020). One example is the expression of  $\text{CO}_2$  permeable aquaporins to improve membrane  $\text{CO}_2$  conductance (Groszmann *et al.*, 2017; Zhao *et al.*, 2017; Ermakova *et al.*, 2021) and reabsorb emitted  $\text{CO}_2$  in the chloroplast. Another example of the latter is recent work by South *et al.* (2019) who detail three alternative photorespiratory pathways (APs) in *Nicotiana tabacum*, two of which led to increased biomass and one which increased light-use efficiency. AP1 plants (with *Escherichia coli* glycolate oxidase genes targeted to the chloroplast) exhibited a 13% increase in plant dry mass. AP3 plants (with replaced malate synthase and glycolate oxidase) exhibited an 18% increase in the absence of an RNAi construct that reduced PLGG1 expression and 24% biomass increase with the RNAi construct – as well as increased glyoxylate and pyruvate, increased photosynthetic rate, increased chloroplastic  $\text{CO}_2$  concentration and decreased serine and glycerate. These findings highlight the potential and viability of modifications to the photorespiratory pathway to improve plant growth.

## Modifying respiratory metabolism to improve yield

Net photosynthesis determines the upper bounds of the plant carbon budget; however, understanding the fate of fixed carbon is just as important to ultimately improving  $Y_p$ . Photoassimilate from the Calvin cycle is stored or exported to sink tissues, where it can be respired to produce ATP, carbon skeletons and reductants. Between 30 and 60% of daily fixed carbon ends up being respired (Amthor *et al.*, 2019). The ultimate goal is to improve the efficiency of ATP synthesis and use, reducing the proportion of daily fixed carbon that is respired and improving yield through enhanced  $\epsilon_c$ . This could be accomplished through improvements in  $\epsilon_{\text{biosynthesis}}$  (Eqns B4 and B6). Knowing that respiration is a key component in determining available carbon for growth in a plant's carbon budget, we first explore the components of  $\epsilon_{\text{biosynthesis}}$  (i.e. processes that influence the efficiency of ATP production and consumption).

## Improving the efficiency of ATP production

To improve  $\epsilon_{\text{biosynthesis}}$ , one option is to consider  $\epsilon_{\text{prod}}$  – the efficiency with which ATP and other respiratory products are made (Eqn B7). During oxidative phosphorylation, high-energy electrons generated during glycolysis and from the tricarboxylic acid (TCA) cycle are transported through a series of complexes along the inner mitochondrial membrane (IMM) to the terminal electron acceptor, oxygen. The flow of electrons through Complexes I (NAD(H) dehydrogenase), III (cytochrome *b<sub>c</sub>1* complex) and IV (cytochrome *c* oxidase, COX) is coupled to the pumping of protons across the IMM to create the electrochemical gradient that drives generation of ATP by Complex V (ATP synthase). Importantly, the plant IMM also contains rotenone-resistant dehydrogenases



**Fig. 3** Photorespiratory modifications can minimise carbon loss. Ribulose 1,5-bisphosphate (RuBP) oxygenation by Rubisco results in the photorespiratory pathway (black) being engaged to process 2-phosphoglycolate (2PG) back into the Calvin cycle via the peroxisome and mitochondria. Changes to the pathway (coloured) represent improved efficiency of photosynthesis by modifying enzyme activity or abundance, colocalising CO<sub>2</sub> release to the site of Rubisco and/or modified flux through a branch of the pathway (thick, coloured lines). 3PG, 3-phosphoglycerate; Acetyl CoA, acetyl coenzyme A.


(alternative DHs) that reduce the ubiquinone pool without utilising Complex I (Rasmusson *et al.*, 2004) and the cyanide-resistant alternative oxidase (AOX) that diverts the flow of electrons from the ubiquinone pool directly to O<sub>2</sub> (McIntosh, 1994; Vanlerbergh & McIntosh, 1997; Berthold *et al.*, 2000). The alternative DHs and AOX pathway reduce the efficiency of ATP production per mol substrate because electrons bypass either Complex I, or both Complex III/IV, respectively, resulting in a decrease in the number of protons pumped across the membrane such that the amount of ATP per unit of CO<sub>2</sub> released and/or O<sub>2</sub> consumed decreases. Similarly, dissipation of the electrochemical gradient also occurs when protons are transported through the IMM-located uncoupling proteins (UCPs) or mechanosensitive ion channels and results in reduced ATP yield, with energy being dissipated as heat (Sweetlove *et al.*, 2006; Lee *et al.*, 2016). Because of this,  $\epsilon_{\text{prod}}$  has the potential to be inversely related to engagement of the AOX pathway and/or other nonphosphorylating/uncoupling pathways (Fig. 4). While we lack examples of where engagement of these pathways are linked to changes in  $\epsilon_{\text{prod}}$ , low activity of them could – in theory – contribute to increased  $\epsilon_{\text{prod}}$  and ultimately  $Y_p$ .

One challenge for improving  $\epsilon_{\text{prod}}$  through targeting the AOX and/or other nonphosphorylating/uncoupling pathways is the fact that nonphosphorylating and uncoupling pathways broadly increase stress resistance. For example, AOX transcripts increase under stressed conditions such as the presence of inhibitors of mitochondrial electron transport chain (Saisho *et al.*, 1997;

Polidoros *et al.*, 2005), chilling temperatures (Djajanegara *et al.*, 2002) and oxidative stress (Yukioka *et al.*, 1998). Loss of AOX makes plants susceptible to stress (Giraud *et al.*, 2008; Demircan *et al.*, 2020), and overexpression can improve plant stress tolerance (Dahal *et al.*, 2015; Gong *et al.*, 2020). Changes in UCP and NDH expression have similar effects (Taylor *et al.*, 2005; Arcuri *et al.*, 2021). Increases in the abundance of nonphosphorylating and uncoupling pathways enable plants to maintain flux through glycolysis and the TCA cycle under conditions of low ATP demand (i.e. enabling flexibility in the ATP yield and depending on demand). This flexibility comes with the trade-off of decreased efficiency in ATP production, and a decrease in  $\epsilon_{\text{prod}}$  means that more assimilated carbon must be consumed to produce ATP. However, suboptimal ATP yield is counteracted by increased stress tolerance and blunt excision of these pathways may do more harm than good, as the aforementioned mutant phenotypes show.

Assuming in specific scenarios that changes in  $\epsilon_{\text{prod}}$  could help increase  $\epsilon_{\text{biosynthesis}}$ , how might future studies incorporate screens for genotypic variability in  $\epsilon_{\text{prod}}$  into breeding strategies? One approach is to quantify the relative engagement of these pathways *in vivo*. In the case of terminal oxidases, COX and AOX exhibit differential isotope discrimination when consuming O<sub>2</sub> (Guy *et al.*, 1987; Cheah *et al.*, 2014). When placed in a closed system, the ratios of <sup>16</sup>O and <sup>18</sup>O change through time; when coupled to measurements of overall O<sub>2</sub> uptake, rates of COX and AOX can be calculated. While such an approach could be used to screen divergent genotypes of crops, the method is currently low

## Intermembrane space



**Fig. 4** Efficiency of ATP production decreases with increasing alternative oxidase (AOX) engagement. Electrons flow (yellow arrows) from Complex I and Complex II to ubiquinone pool (UQ). From UQ, electrons can be transported through Complex III to Complex IV as terminal oxidase. Complexes I, III and IV concurrently pump protons into the intermembrane space (dotted black lines) as they transfer electrons. Those protons drive ATP synthase and ATP production. However, electron flux through the alternative oxidase (dotted red lines) diverts electrons from Complexes III and IV and does not contribute to building proton gradient. Stoichiometrically, the AOX pathway is less efficient at ATP production than the COX pathway because significantly fewer electrons are pumped into the intermembrane space. Therefore, we hypothesise that to a first order,  $\epsilon_{\text{prod}}$  will increase as either: (1) a higher proportion of electron flux passes through the COX pathway; or (2) AOX protein abundance decreases. Additionally, the presence of nonphosphorylating alternative dehydrogenases and uncoupling proteins reduce the ubiquinone pool but produce heat instead of ATP, also lowering  $\epsilon_{\text{prod}}$ . ADP, adenosine diphosphate; Cyt c, cytochrome c;  $e^-$ , electrons;  $H^+$ , protons; NDH, nonphosphorylating alternative dehydrogenases;  $P_i$ , inorganic phosphate; UCP, uncoupling proteins.

throughput and requires sustained use of high-sensitivity mass spectrometers; thus, as currently available, the oxygen isotope discrimination method is unlikely to provide the rapid screening required by the crop breeding industry. Similarly, we lack methods to rapidly screen for variability in *in vivo* proton translocation via uncoupling pathways or the use of alternative NDHs. Thus, new approaches will need to be developed that enable *in vivo* rates through any or all of these alternative pathways in high-throughput measurements as part of screens of diverse germplasm. An alternative approach could be to employ a synthetic biology approach to modify their engagement. For example, Amthor *et al.* (2019) suggested engineering a light-specific AOX to replace constitutive expression, minimising AOX flux in the dark and improving  $\epsilon_{\text{prod}}$ . A dual-track approach – using screens of diversity panels and a synthetic biology approach – would allow us to take advantage of natural variation in engagement of alternative pathways and engineered variation. Finally, further efforts need to be made to adapt the genetically-encoded biosensors developed for live monitoring of photosynthetic redox and energy status in chloroplasts (Müller-Schüssele *et al.*, 2021) to the monitoring of respiratory energetics in mitochondria.

### Increasing the efficiency that energy is used

Improvements to  $\epsilon_{\text{biosynthesis}}$  can also be accomplished through changes to the efficiency by which respiratory energy is used (i.e.

$\epsilon_{\text{use}}$ ) for biosynthetic, maintenance and active transport. Cellular respiration not only provides the energy required for metabolic reactions, but also creates the carbon skeletons necessary for biosynthesis and regulates the cellular redox environment in which these reactions to take place (Plaxton & Podestá, 2006; Millar *et al.*, 2011; O'Leary *et al.*, 2019). Changes in the efficiency of processes that use products of respiration can thus influence  $\epsilon_{\text{use}}$ , with consequences for  $\epsilon_{\text{biosynthesis}}$ ,  $\epsilon_c$  and  $Y_p$ .

**Biosynthesis costs of plant tissues are generally invariant** Could  $\epsilon_{\text{use}}$  be improved via screening for variation in the energy costs of synthesising new tissues? The energy demand associated with biosynthesis depends on two factors: (1) the ATP costs of making individual chemical constituents; and (2) the relative abundance of individual chemicals in plant tissues (De Vries *et al.*, 1983; Lambers & Rychter, 1989). Beyond acquisition (described later), minerals have no construction cost, and organic acids, total nonstructural carbon and total structural carbon are relatively cheap to produce (0.91, 1.09 and 1.22 g glucose  $g^{-1}$  dry mass, respectively). Meanwhile, the costs of synthesis are higher for lignin, protein, soluble phenolics and lipids (2.12, 2.48, 2.60 and 3.03 g glucose  $g^{-1}$ , respectively), and key components of energy organelles. Theoretically, differences in the chemical makeup of different tissues could mean that some tissues are cheaper to produce than others. However, in studies that have quantified the construction costs of plant tissues in a range of contrasting plant species,



relatively minor differences have been observed. For example, Poorter & Villar (1994) found only small differences in the construction costs for different functional types: a 6% difference between herbaceous and woody plants (1.45 to 1.54 g glucose g<sup>-1</sup>, respectively) and a 4% difference between deciduous angiosperms and evergreens (1.52 and 1.58 g glucose g<sup>-1</sup>, respectively). Additionally, environmental-limiting conditions ((CO<sub>2</sub>), light and nutrients) have a minimal impact (≤10%) on construction costs. Underpinning the uniformity of construction costs is the fact that there are often trade-offs in the abundance of energetically expensive compounds. For example, fast-growing species are protein-rich, whereas slow-growing species contain high concentrations of lignin, with both compounds being energetically expensive to construct (Poorter & Villar, 1994), with there being a general trade-off between inexpensive and expensive constituents in both slow- and fast-growing species (Lambers *et al.*, 2008). Given these observations, targeting the efficiency of biosynthesis is unlikely to be the best strategy for improving  $\epsilon_{\text{use}}$ . Rather, improvements are likely to come through a focus on the efficiency with which respiratory energy is used by processes of cellular maintenance and/or ion transport.

**Improvements through targeting the energy costs of maintenance processes** Perhaps of greater significance in determining whether a particular plant has low or high  $\epsilon_{\text{use}}$  are the energy costs associated with *maintenance* of mature tissues (e.g. protein turnover and maintenance of solute gradients across membranes), which can consume 30–60% of daily fixed carbon (Penning de Vries, 1975; Amthor, 2000), with high costs being associated with slow growth rates (Lambers *et al.*, 2008). More recently, Amthor *et al.* (2019) outlined the opportunity to reduce energy costs of maintenance processes, including decreasing protein and membrane turnover, relocating and/or rescheduling metabolic activities, suppressing futile cycles in metabolism and reducing ion transport costs. As such, there is potential for  $\epsilon_{\text{use}}$  to be improved through screening for variation in the energy costs of these processes and/or genetic manipulation of the underlying components of each process (Jacoby *et al.*, 2016; Amthor *et al.*, 2019).

Plants employ a diverse assortment of proteins for daily growth, metabolism and function, as well as in response to dynamic stimuli. The plant proteome is responsive to abiotic and developmental factors as well as carbon and nitrogen economy (Kosová *et al.*, 2011). Total protein content in a plant is not constant and changes in response to environmental factors such as salinity (Doganlar *et al.*, 2010; Jouyban, 2012), drought (Akhzari & Pessaraki, 2016) and heat stress (Giri *et al.*, 2017) to name a few. Protein abundance depends on the rate of protein synthesis ( $k_s$ ) and degradation ( $k_d$ ) (Li *et al.*, 2017; Ross *et al.*, 2021). As Li *et al.* (2017) note, these two processes are often combined to loosely define protein turnover, the process by which degradation of existing proteins and replacement with newly synthesised proteins modify the proteome. For a given time period, Ross *et al.* (2021) define total protein produced simply as the integral of  $k_s$ , whereas total protein degraded is expressed as an exponential decay function, with  $k_d$  as the rate constant. By relating protein abundance to rates, the authors find that  $k_d$  controls turnover

under steady-state conditions, including after changes in  $k_s$ , as only  $k_d$  controls protein half-lives (see Ross *et al.*, 2021).

What do we know about the relationship between protein turnover and its associated ATP costs? Reflecting the factors mentioned earlier, it is clear that plants have evolved a complex system to regulate protein synthesis and degradation, allowing metabolism to dynamically adjust to changes in the type and abundance of proteins needed in different environments and growth stages (Vierstra, 2009; Marshall & Vierstra, 2018). Protein synthesis can be controlled at the transcriptional, post-transcriptional and translational levels, whereas degradation is dependent on post-translational regulation related to protein location and function, local environment and protein–protein interactions (Hinkson & Elias, 2011; Nelson & Millar, 2015; Huang *et al.*, 2020). The balance between synthesis and degradation dictates maintenance costs of protein turnover, either through differences in specific ATP cost of turnover or the absolute amount of turnover occurring.

While protein turnover can help plants adjust to new environments, it is energetically expensive. Each day, c. 6% of all proteins are replaced, with 6–10 mol ATP being consumed per mol peptide bonds undergoing turnover and 0.9–1.6 CO<sub>2</sub> per amino acid (Noguchi *et al.*, 2001; Nelson *et al.*, 2014; Ishihara *et al.*, 2015). Bouma *et al.* (1994) estimated that protein turnover in bean leaves accounted for 17–21% of darkened mature leaf respiration (c. 2.3 nmol CO<sub>2</sub> g<sup>-1</sup> s<sup>-1</sup>). In a review of previous research, Noguchi *et al.* (2001) reported that protein turnover consumes 3.5–60% of total ATP produced. In their own experiment, Noguchi *et al.* (2001) found protein turnover consumed 15.9–23.8% and 40.8–61.3% of total ATP produced in shaded and sunlit leaves, respectively. Quigg & Beardall (2003) reported that the proportion of maintenance respiration is attributable to protein turnover in higher plants, ranging from *Dactylis glomerata* roots (7%) to *Hordeum vulgare* full-grown leaves (30–60%). In wheat, protein turnover consumes approximately a third of total respiratory energy (Zagdańska, 1995). In *Arabidopsis*, 25–40% of ATP is used for protein turnover, depending on the age and growth rate of leaves, and this overall cost has been broken down to specific costs of maintaining different kinds of cellular organelles, metabolic pathways and enzymes (Li *et al.*, 2017). Specific proteins also play disproportionately large roles in turnover costs. For example, thiazole synthase (THI4) turnover can account for 2–10% of total maintenance respiration (Amthor *et al.*, 2019). Thus, while there is clearly considerable variability in the estimated ATP costs associated with protein turnover, what is clear is that reducing the costs of turnover could be a target for improving  $\epsilon_{\text{use}}$ , particularly when designing plants for future, warmer climates. This is because turnover costs are likely to increase under conditions of environmental stress (e.g. heat, salinity and hypoxia) that accelerate the turnover of proteins, thereby consuming more of the ATP budget (Hachiya *et al.*, 2007; Edwards *et al.*, 2012).

Bearing in mind the goal of maximising  $\epsilon_{\text{use}}$ , are there correlations between whole-plant growth and protein turnover? In *Arabidopsis*, larger (higher growth rate and rosette biomass) accessions reduced ribosome abundance at night, whereas smaller accessions maintained end-of-day levels of ribosomes (Ishihara



*et al.*, 2017). Smaller accessions also had rates of protein synthesis up to 30% higher than the estimated rate required for growth, and up to 25% of the observed difference in relative growth rate could be tied to increased ATP costs of protein turnover. Gibon *et al.* (2009) found a tight correlation between growth, starch metabolism and protein content in *Arabidopsis* grown under different photoperiods, although they observed protein *amount*, not turnover rate. There are currently few studies in this space, but it should be considered a priority for further study, considering the substantial proportion of ATP use devoted to protein turnover.

Other substantive energy demands are phospholipid membrane turnover (*c.* 0.7 nmol CO<sub>2</sub> g<sup>-1</sup> s<sup>-1</sup>) (Penning de Vries, 1975) and maintenance of ion gradients within cells; for the latter, Penning de Vries (1975) estimated that the total cost of maintaining intracellular ion gradients was *c.* 4.6 nmol CO<sub>2</sub> g<sup>-1</sup> s<sup>-1</sup> in some leaves, that is over 50% of respiratory rate. Lipids (a large part of membrane composition) are expensive to produce, costing nearly 3 g of glucose to produce 1 g of lipid (De Vries *et al.*, 1974). To our knowledge, little work exists on the specific construction costs or turnover rates of plant membranes. Prior studies show that whole-plant cell membranes can be fully turned over in 200 min (Warren & Glick, 1968; Steer, 1988). However, Hao & Maxfield (2000) found membrane turnover rates as low as 5–10 min in Chinese hamster ovarian cells. This large difference between very distinct cell types suggests more work remains to address gaps in how membrane turnover is supported by maintenance respiration. Similarly, considering protein turnover, improved measurement techniques will allow for fine-resolution, context-specific descriptions of synthesis and degradation rates,  $k_s$  and  $k_d$ , respectively (Nelson *et al.*, 2014; Ross *et al.*, 2021), across the multitude of subcellular and ambient environments experienced, allowing greater specificity of turnover characterisation. Variation in energy demands associated with maintenance can, therefore, have a substantive impact on  $\epsilon_{\text{use}}$  of leaves. From the context of increasing  $\epsilon_{\text{use}}$  and ultimately  $Y_p$ , minimising the energy costs associated with protein turnover, while maintaining adequate function, would present an opportunity to decrease the maintenance respiration costs incurred (Amthor *et al.*, 2019).

**Lowering the energy costs of phloem loading** A further factor that influences the demand for respiratory ATP is nocturnal carbohydrate export from mature leaves. Once synthesised in photosynthetic source leaves, sucrose (Suc), the main assimilate in most seed plants, needs to be loaded into the phloem for translocation towards nonphotosynthetic sink organs where Suc is unloaded. This source-to-sink allocation of assimilate plays critical roles in determining how biomass is translated into crop yield (Ruan, 2014; Fernie *et al.*, 2020). Opportunities exist to increase source-to-sink phloem translocation efficiency with no or little additional ATP cost as elaborated later.

In most species, phloem loading takes place either via the apoplast across plasma membranes or via the symplast through plasmodesmata (PD) at the interface of phloem parenchyma and the companion cell–sieve element (CC–SE) complex in developed leaves. The latter pathway also includes Suc moving from mesophyll and bundle sheath cells through PD into specialised

CCs called intermediary cells, where Suc is synthesised into a raffinose family oligosaccharide for loading into the SE, the so-called polymer trap model. This pathway has been observed in *Cucurbita* genus and some trees (Fu *et al.*, 2011; Zhang & Turgeon, 2018). In apoplastic phloem loading, the uptake of Suc into CCs is typically mediated by H<sup>+</sup>-Suc symporters energised by plasma membrane-bound ATPase, while efflux of Suc from phloem parenchyma is facilitated by the energy-independent clade III Sugars will Eventually be Exported transporter (SWEET). Knockout of clade III SWEETs significantly reduced phloem loading, leading to a severe growth phenotype in a range of species such as *Arabidopsis* (Chen *et al.*, 2012) and maize (Bezruczyk *et al.*, 2018). These findings indicate a major role of SWEETs in controlling Suc efflux for subsequent uptake into the CC–SE. It is therefore plausible that enhancing Suc efflux by increasing clade III SWEET expression or activity could improve phloem loading of Suc with no direct energy cost (and thus improve  $\epsilon_{\text{use}}$ ).

In parallel to apoplastic loading, Suc may also be loaded into CC–SE symplasmically via PD (Braun *et al.*, 2014; Bezruczyk *et al.*, 2018). In this case, phloem loading may be improved through increasing PD conductance by modifying sterol homeostasis or callose turnover or other PD-associated proteins (Zhang *et al.*, 2017; Fernie *et al.*, 2020). To this end,  $\beta$ -1,3 glucan synthase and  $\beta$ -1,3 glucanase are the two enzymes synthesising and degrading callose, respectively (Ruan, 2007; Zhang *et al.*, 2017). Thus, the knockout of  $\eta$ -1,3 glucan synthase could block callose synthesis at the PD neck region, while overexpressing  $\beta$ -1,3 glucanase could degrade callose at the PD neck region. Such changes would have the effect of ‘opening up’ or increasing PD conductance for symplasmic loading (potentially increasing  $\epsilon_{\text{use}}$ ).

The SWEETs also play an important role in unloading of Suc into sinks. Interestingly, clade III SWEETs could be bound and inactivated by SP6A, a homologue of FT and phloem-mobile tuberigen in potato tubers, which switches the apoplastic unloading route into symplasmic unloading, triggering tubulisation (Abelenda *et al.*, 2019). This shows potential to optimise phloem unloading for yield gain such as tuber formation via manipulating SWEET activity. On the contrary, PD appears to impose major restrictions on the delivery of assimilates to meristematic sinks such as shoot or root apex and ovules (Ruan, 2012). In this context, computational modelling has identified inefficient delivery of carbon as a cause of floral and seed abortion on the secondary and tertiary branches of rice panicles (Seki *et al.*, 2015), likely due to low PD conductance in those high-order branches (Fernie *et al.*, 2020). Collectively, the above analyses indicate potential scope and avenues for improving phloem loading and unloading for growth and yield gain in an energy-efficient manner, with positive implications for  $\epsilon_{\text{use}}$ .

**Reducing the energy costs associated with root nutrient uptake** Uptake of nutrients from the soil into roots is another major energy cost, with 50–60% of root respiratory ATP being used to support ion uptake (Poorter *et al.*, 1991). Importantly, the specific energy costs of ion uptake differ among species adapted to contrasting environments, with more respiratory ATP being needed to take up a given amount of nitrate in inherently slow-

growing species than their fast-growing counterparts, when plants are provided with high nutrient supply (Poorter *et al.*, 1991). The differences in the specific costs of ion uptake reflect the fact that: (1) there is an efflux of ions (e.g. nitrate) from roots, resulting in a reduction in the net rate of ion uptake; and (2) the extent of efflux is relatively greater in inherently slow-growing species than in fast-growing species (Scheurwater *et al.*, 1998, 1999). Such differences suggest that we may be able to increase  $\epsilon_{\text{use}}$  via understanding the mechanistic basis for these differences.

Roots collect water and nutrients from the surrounding soil matrix, not only through bulk flow passive transport but also through active transport facilitated by transporter proteins. Active transport impacts  $\epsilon_{\text{use}}$  by using ATP to transport nutrients across membranes, against a concentration gradient, to be assimilated by roots or transported to shoots (Xu *et al.*, 2012). In the case of nitrate, this movement is undertaken by nitrate transporter 1 (NRT1)/peptide transporter gene superfamily (NPF) that encodes transporters throughout the plant that allow its movement from root to shoot and ultimately nitrate reduction (Campbell, 1999; Plett *et al.*, 2010; Zhang *et al.*, 2020). However, as outlined earlier, nitrogen efflux from roots is common (Segonzac *et al.*, 2007; Xu *et al.*, 2012) and can contribute to high rates of respiration in slower growing plants as the ratio of nitrate influx to net nitrate uptake increases (Scheurwater *et al.*, 1998). Ideally,  $\epsilon_{\text{use}}$  would be maximised through maintaining high rates of net nitrate uptake and assimilation per ATP used. This could be achieved through increased influx, decreased efflux or more rapid translocation to the shoot. Strategies including suppressed expression of nitrate efflux transporters such as NAXT1 (Segonzac *et al.*, 2007) or other anion channels (Teakle & Tyerman, 2010; Wege *et al.*, 2017), targeted overexpression of influx transporters with high  $\text{NO}_3^-$  selectivity (Fan *et al.*, 2016), increasing structural barriers to impede leakage in roots such as suberised hypodermal or endodermal layers (Barberon, 2017) and relocation of nitrate assimilation from roots to shoots have been proposed to increase biomass gains from lower respiration rates (Amthor *et al.*, 2019). Microbiome interactions, for example with arbuscular mycorrhizal fungi, also aid with nutrient uptake and growth promotion (Pascale *et al.*, 2020). Griffiths *et al.* (2021) studied hydroponically-grown *Zea mays* and found genotypic variation in specific root ion uptake rates, which were heritable and positively correlated with specific root respiration rates. Variation between two rice subspecies in their *NRT1.1B* transporter showed increased nitrate uptake and transport to shoots, as well as upregulation expression of root and shoot nitrate reductase genes (Hu *et al.*, 2015). As Griffiths & York (2020) point out, the extent of genetic variation in transporter traits such as energy costs of active transport is not yet well-characterised. Thus, variations in uptake kinetics exist and present another target for improved  $\epsilon_{\text{use}}$ .

### Linking respiration rates to photosynthesis, growth and yield

The earlier sections have shown that there is potential for variability in both  $\epsilon_{\text{prod}}$  and  $\epsilon_{\text{use}}$  to occur in nature, with consequences for  $\epsilon_{\text{biosynthesis}}$  and  $\epsilon_{\text{c}}$ . Amthor *et al.* (2019) estimated that a 10% reduction in the energy costs of maintenance could increase

biomass accumulation by 6%, and a 20% maintenance respiration reduction could lead to gains up to 14%. This provides a rationale for screening genotypic variability in  $\epsilon_{\text{prod}}$  and/or  $\epsilon_{\text{use}}$  and therefore possibly identifying genotypes with improved yield. However, to be useful, such an approach needs an array of high-throughput methods capable of large-scale phenotyping of  $\epsilon_{\text{prod}}$  (e.g. rapid assays of nonphosphorylating or uncoupling pathway engagement) and/or  $\epsilon_{\text{use}}$  (e.g. measurements of the ratio of net to gross nitrate uptake and associated rates of root respiratory ATP production). However, such high-throughput tools are not available.

Consequently, we need to consider new ways of assessing variation in  $\epsilon_{\text{biosynthesis}}$ . Here, one option is to screen for variation in the relationship between plant respiration and photosynthesis (i.e. what fraction of daily fixed carbon is respired rather than conserved in plant biomass). Another is to look for variability in the relationship between respiration and relative growth rate (i.e. screen for genotypes that achieve high growth rates while also having low rates of respiration).

What evidence is there that rates of plant respiration vary among genotypes that differ in productivity? Here, insights can be drawn from the eco-physiology literature which has shown that inherently fast-growing species respire a lower fraction of daily fixed carbon than their slow-growing counterparts (Poorter *et al.*, 1990; Lambers & Poorter, 1992; Loveys *et al.*, 2002). Rates of whole-plant respiration are also *relatively* similar among species that differ markedly in relative growth rates, with high growth rates of some species being achieved without concomitant proportional increases in respiration rates. Such studies point to variation in  $\epsilon_{\text{biosynthesis}}$  being an outcome of evolution, with  $\epsilon_{\text{biosynthesis}}$  being highest in fast-growing species adapted to resource-rich, favourable habitats, and lowest in species adapted to resource-poor, unfavourable habitats.

Further evidence of the possible link between  $\epsilon_{\text{biosynthesis}}$ , biomass accumulation and yield comes from the work done on *Lolium perenne* (perennial ryegrass) in the 1980s. In a series of papers by Wilson, Jones and Robson, up to 13% increases in biomass accumulation and up to 40% increases in annual yield (in repeated years) were observed in plants grown under high-density conditions and selected for lower rates of mature leaf respiration (Wilson, 1975, 1982; Robson, 1982a,b; Wilson & Jones, 1982). In Wilson (1982), slow-respiring and fast-respiring progeny were produced by pair-crossing parents within either a slow- or fast-respiring cultivar. Additionally, progeny with low rates of mature leaf respiration were found to have higher net assimilation rate, relative growth rate, dry weight at harvest and leaf area index. Another study comparing selected fast- and slow-respiring ryegrass cultivars found the slow-respiring cultivar to have higher shoot and root mass at each harvest point (Day *et al.*, 1985). More recently, selection across multiple generations of lines with low respiration rates has been shown to be associated with higher yields in canola (Hauben *et al.*, 2009). Thus, there is some evidence that screening for variability in plant respiration rates may enable genotypes with higher  $\epsilon_{\text{c}}$  to be identified, with future work needing to use screens to ultimately identify what genes are responsible for variability in  $\epsilon_{\text{biosynthesis}}$ .

The link between respiration, growth and yield is not, however, always consistent. For example, in perennial ryegrass, the relationship between low rates of mature leaf respiration and improved biomass accumulation disappeared when plants were grown in low-density swards (Kraus *et al.*, 1993), a result that was supported by a later study (Kraus & Lambers, 2001). Similarly, there is evidence that the growth of pea plants is linked to the efficiency of the respiratory system in some studies (Musgrave *et al.*, 1986) but not others (Obenland *et al.*, 1988). Thus, future efforts to explore the link between respiration, growth and yield will need to ensure that this relationship holds over a wide range of industry-relevant planting regimes.

One of the challenges facing work designed to link respiratory rates to photosynthesis, biomass accumulation and yield is the need to screen large numbers of plants for variability in traits of interest and to establish the genetic basis of variability in  $\epsilon_{\text{biosynthesis}}$ . In the past, assays of photosynthesis and respiration were painstakingly slow, limiting our ability to screen large numbers of plants. Similarly, growth analysis required repeated destructive harvesting of plants. However, in recent years, methods have been developed that enable each of these parameters to be quantified in high throughput. Growth analyses are now possible using light detection and ranging technology to quantify aboveground biomass in field experiments (Jimenez-Berni *et al.*, 2018; Furbank *et al.*, 2019; Walter *et al.*, 2019; Jin *et al.*, 2020). For respiration, a breakthrough was the development of a high-throughput, robotic fluorophore method for measuring respiratory  $\text{O}_2$  uptake (Scafaro *et al.*, 2017). This method has provided the opportunity to screen hundreds of plant samples per day and has revealed that there is substantial variation in leaf respiration rates among genotypes of individual species, both in wheat (Scafaro *et al.*, 2017) and Arabidopsis (O'Leary *et al.*, 2017). Subsequent work has also shown that we can use hyperspectral reflectance signatures to predict variability in leaf respiration rates of laboratory and field-grown wheat (Coast *et al.*, 2019), further increasing our ability to screen respiration rates in field trials; similarly, hyperspectral reflectance can be used to predict photosynthetic capacity in wheat (Silva-Perez *et al.*, 2018). There is now an opportunity to use these methods to screen diversity panels and recombinant inbred line populations for variations in leaf respiration, photosynthesis and growth, and to identify quantitative trait loci for traits associated with high  $\epsilon_{\text{biosynthesis}}$ .

To fully exploit the potential for screens of variability in respiration rates to be used in strategies to improve  $\epsilon_{\text{biosynthesis}}$ , consideration needs to be given, not just to screening variations in leaf respiration ( $R_{\text{leaf}}$ ), but also  $R_{\text{stem}}$  and  $R_{\text{root}}$ . An understanding of the fraction of whole-plant biomass allocated to leaves (LMF), stems (SMF) and roots (RMF), with rates of whole-plant respiration ( $R_{\text{total}}$ ) being calculated according to:

$$R_{\text{total}} = (R_{\text{leaf}} \times \text{LMF}) + (R_{\text{stem}} \times \text{SMF}) + (R_{\text{root}} \times \text{RMF})$$

Eqn 2

The challenge going forward will be developing high throughput, (ideally) nondestructive ways of estimating stem and root

respiration, and the patterns of biomass allocation above and belowground.

### How would a warmer world temperature affect $\epsilon_c$ ?

Given that the world is warming, it is important to understand how rising temperatures affect  $\epsilon_c$ . Noting that  $\epsilon_c = \alpha \times \epsilon_{\text{prod}} \times \epsilon_{\text{use}}$  (Eqn B7), any temperature-dependent change in the factors and processes previously discussed would likewise influence  $\epsilon_c$ . On a short-term timescale, increasing temperature past the photosynthetic optimum will decrease photosynthetic efficiency by decreased Rubisco carboxylation, damage thylakoid membranes and induce changes to vapour pressure deficit that result in stomatal closure (Mathur *et al.*, 2014; Dusenage *et al.*, 2019). At the same time, respiration, along with photorespiration and reactive oxygen species, is stimulated by increasing temperatures (Tjoelker *et al.*, 2001; Mundim *et al.*, 2020). Importantly, the initial imposition of heat likely results in an increasing proportion of respiratory ATP being used to support maintenance processes instead of growth (Gibon *et al.*, 2009; Scafaro *et al.*, 2021). Rapid increases in temperature result in membranes becoming more fluid and therefore leaky (Allakhverdiev *et al.*, 2008), decreasing  $\epsilon_{\text{use}}$  as a result of more ATP being needed to maintain ion gradients across membranes and decreasing  $\epsilon_{\text{prod}}$  as protons leak across the IMM (Divakaruni & Brand, 2011). Heat also increases the rate of protein turnover and associated demand for respiratory ATP, lowering  $\epsilon_{\text{use}}$ . More severe temperature increases can denature proteins, deactivating enzymes and elevating the cost of maintenance chaperone and heat shock proteins, further lowering  $\epsilon_{\text{use}}$  (Wang *et al.*, 2004). Such factors – combined with knowledge of the mechanisms via which plants adjust membrane fluidity (via alterations in the degree of saturation of fatty acids) and stabilise proteins (e.g. via synthesis of heat shock proteins), as well as thermally acclimate photosynthesis and respiration (Smith & Keenan, 2020) – will need to be considered in work targeting higher  $\epsilon_c$  in crops developed to cope with more frequent and severe heat waves.

Considering abiotic stress more broadly, the complex, interacting nature of many of the components affecting energy efficiency in plants requires a multivariate investigative approach. Screening for trait variation in  $\epsilon_{\text{prod}}$  and  $\epsilon_{\text{use}}$  and isolation of genetic candidates via GWAS or QTL – while being an approach that can enhance classical breeding approaches – is complicated by the high degree of environmental sensitivity of  $\epsilon_{\text{prod}}$ - and  $\epsilon_{\text{use}}$ -related traits (O'Leary *et al.*, 2019; Scafaro *et al.*, 2021). As such, the experimental design used to screen trait  $\epsilon_{\text{prod}}$  and  $\epsilon_{\text{use}}$  variation needs to be done in ways that enable the roles of genetics and adverse environments to be clearly identified.

A further option to improve  $\epsilon_{\text{prod}}$  and  $\epsilon_{\text{use}}$  is to combine the above trait screens with additional approaches that harness the potential of genetic engineering. For example, modern genetic engineering techniques will permit the introduction of low-cost alleles that maintain stress responsiveness by targeting modifications to specific tissues only under permissive conditions, or by editing regulatory elements of promoters to better tune expression under such

**Table 1** Summary of targets and opportunities for increasing yield potential.

$\epsilon_c$ component	Process	Target	Cost	Opportunity
$\alpha$	Light and dark reactions of photosynthesis	Photoprotection and photoinhibition	15–20% loss of potential CO <sub>2</sub> fixation	Optimise photoprotective systems at the molecular and canopy level to avoid drop in quantum yield due to photoinhibition (Murchie <i>et al.</i> , 2015) and increase the speed that NPQ relaxes following shift from sun to shade (de Souza <i>et al.</i> , 2022).
		Photosynthetic induction	21% loss of potential CO <sub>2</sub> capture in wheat when induction is slow	Screen for natural variation in induction rate (Taylor & Long, 2017) – for example in amino acid substitutions and overexpression of Rca – to speed induction of photosynthesis following illumination.
	Carboxylation	Rubisco properties	Competitive inhibition, slow turnover, inactivation resulting in lower carboxylation rates	Study structure and catalytic properties of Rubisco (as in RIPE) to ultimately improve enzyme CO <sub>2</sub> fixing efficiency.
	Photorespiration	Oxygenation	Low Rubisco specificity to CO <sub>2</sub> leading to reduced net CO <sub>2</sub> uptake and increased H <sub>2</sub> O <sub>2</sub> formation	Increase local CO <sub>2</sub> concentration – for example introducing cyanobacterial carboxysomes (Price <i>et al.</i> , 2013; Long <i>et al.</i> , 2021) which can lead to reduced photorespiration.  Target <i>Escherichia coli</i> glycolate oxidase to chloroplast (Kebeish <i>et al.</i> , 2007) to convert glycolate into glycerate and reduce photorespiratory carbon flux to peroxisome and mitochondria. Replace glycolate oxidase with <i>Chlamydo reinhardtii</i> glycolate dehydrogenase (Maier <i>et al.</i> , 2012) to eliminate photorespiratory CO <sub>2</sub> release.
$\epsilon_{\text{prod}}$	Mitochondrial electron transport and dissipation of the proton motive force	Alternative dehydrogenases and UCP	Reduce ubiquinone pool without engaging ATP synthase	Knockout of UCP (Sweetlove <i>et al.</i> , 2006) lead to photorespiratory defects and decreased expression of type A NAD(H) dehydrogenases and AtNDB1 (Wallström <i>et al.</i> , 2014) which resulted in delayed growth, knockdown of AtNDB4 resulted in better or unaltered growth under standard and stress growth conditions (Smith <i>et al.</i> , 2011), while knockout of AtNDB2 lead to susceptibility to environmental stress (Sweetman <i>et al.</i> , 2019). Therefore, a strategy for improvement will need to be tissue specific or timed to avoid these detriments.
$\epsilon_{\text{use}}$	Growth	Alternative oxidase pathway	Flux through AOX does not pump protons into IMM	Replace constitutive AOX isoform with a diel-scheduled isoform (such as with a light-specific promoter region) to increase $\epsilon_{\text{prod}}$ at night (Amthor <i>et al.</i> , 2019).
		Construction costs	Expensive constituents (lignin, protein, lipids) cost > 2 g glucose g <sup>-1</sup> to construct	Employ Fourier transform (near-) infrared spectroscopy for high-throughput screening for variation of chemical composition in tissues (Collins <i>et al.</i> , 2014).
		Membrane composition	Lipids in membranes are expensive to synthesise and leaky membranes can reduce efficiency of cellular processes	Engineer DNA nanostructures to mimic transmembrane proteins, regulate liposomes, modify membrane morphology and mediate membrane interface chemistry (Feng <i>et al.</i> , 2021).
	Maintenance	Protein turnover	Large proportion of maintenance respiration	Slowing protein turnover may translate to higher growth rate and biomass (Jacoby <i>et al.</i> , 2016), for example by modifying synthesis of THI4, a high-turnover protein (Hanson <i>et al.</i> , 2018).
		Metabolic futile cycles	Inefficient energy use with no net gain in useful product	Minimising enzymatic sucrose degradation and resynthesis (especially that paired with active transport), or F6P/F16BP cycling under specific developmental or stress conditions (Amthor <i>et al.</i> , 2019).
		Phloem loading	Active transport costs ATP	Upregulate SWEET/SUC/WAKL proteins to increase phloem loading, biomass and yield (Xu <i>et al.</i> , 2020).



Table 1 (Continued)

$\epsilon_c$ component	Process	Target	Cost	Opportunity
		Nutrient uptake and translocation	ATP cost of ion uptake	Replace hydrolase with phosphorylase to avoid ATP-dependent hexokinase phosphorylation (Marques <i>et al.</i> , 2018). Quantify unexplored genetic variation in transporter profile (Griffiths <i>et al.</i> , 2021) to select for optimised traits of interest.
			High efflux of ions necessitates reassimilation	Suppress expression of efflux transporters such as NAXT1 and overexpress selective influx transporters (Segonzac <i>et al.</i> , 2007; Fan <i>et al.</i> , 2016) to reduce futile cycling of nutrients.

AOX, alternative oxidase; F16BP, fructose 1,6-biphosphate; F6P, fructose 6-phosphate; IMM, mitochondrial intermembrane matrix; NPQ, nonphotochemical quenching; Rca, Rubisco activase; RIPE, Realizing Increased Photosynthetic Efficiency; SUC, sucrose transporter; SWEET, Sugars Will Eventually be Exported Transporter; UCP, uncoupling proteins; WAKL, wall-associated kinase-like gene.

conditions (Zhang *et al.*, 2018). The generation of a range of cis-regulatory element edits provides the opportunity to produce a wide range of expression levels of endogenous genes, producing a broader range of potential trait variation than can be seen in natural populations (Rodríguez-Leal *et al.*, 2017; Liu *et al.*, 2021). Because of this, trait stacking (i.e. incorporation of multiple genetic modifications into a single variety of a crop) via transgenesis has enormous potential to speed up the process of crop improvement compared with approaches that rely on traditional breeding approaches. Trait stacking has proven successful in improving crop resistance/tolerance to biotic stresses (Dormatey *et al.*, 2020) where a binary response – immune or susceptible – can be easily assessed; with improvements in high-throughput phenotyping methods, the same is likely to be true for assessing the outcomes of stacking of traits that contribute to improved  $\epsilon_c$ . While ‘trait stacking’ will not always deliver overall positive outcomes for plant metabolism and biomass accumulation – with balance between positive vs negative likely to depend on developmental stage, environment and management practices – there is clearly potential to create crop lines with marked improvements in  $\epsilon_c$  through exploring the effect of multiple changes in energy-related traits.

## Conclusions

As outlined earlier, annual increases in crop production are well behind that needed to meet future global demand for food, with new ways needed to improve yield potential ( $Y_p$ ). Our review has highlighted opportunities to increase  $Y_p$  using a range of approaches – summarised in Table 1 – that target photosynthesis and respiration to improve the efficiency with which light energy is converted into biomass ( $\epsilon_c$ ). For photosynthesis, this includes changes in the properties of Rubisco and photosynthetic induction rates to increase carbon fixation (Table 1). Minimising photorespiratory carbon loss through metabolic engineering to increase carboxylation rates while still allowing oxygenation at a reduced but viable rate also shows promise (Table 1). Similarly, reducing the percentage of daily fixed carbon that is released through respiration could improve  $\epsilon_{\text{biosynthesis}}$  – with there being potential to improve the efficiency of ATP synthesis (i.e.  $\epsilon_{\text{prod}}$ ) and/or ATP use (i.e.  $\epsilon_{\text{use}}$ ) to improve  $\epsilon_c$  (Table 1). When such changes are brought

together through a ‘trait stacking’ approach, there is potential to create additive or even multiplicative improvements to  $\epsilon_c$ , and through it, a step change in  $Y_p$ .

Key to exploiting the potential to improve yields through increases in efficiency of photosynthesis and respiration will be the development of high-throughput assays that quantify components of  $\epsilon_c$ . We suggest prioritising high-throughput and digital phenotyping of these components through hyperspectral and automated fluorophore systems and associated growth. The ability to phenotype variation in the balance between respiration, photosynthesis and growth across many plants would provide opportunities to exploit genetic variation in  $\alpha$  and  $\epsilon_{\text{biosynthesis}}$  – and through them  $\epsilon_c$  and  $Y_p$ . Collectively, a focus on these strategies will inform breeding programmes targeting the increases in annual yield necessary to keep pace with increasing demand for food, reduce the amount of land used for agriculture, limit the demand for inorganic fertilisers and restrict greenhouse gas emissions from the agriculture sector.

## Acknowledgements

This work was funded by grants from the Australian Research Council and was supported by the ARC Centre of Excellence in Plant Energy Biology (CE140100008). AG was supported by a Grains Research Development Corporation PhD stipend, ANU International PhD Scholarship and HDR Fee Remission Merit Scholarship. RTF acknowledges the ARC Centre of Excellence for Translational Photosynthesis (CE140100015). MPR acknowledges support from the Foundation for Food and Agricultural Research. Y-LR was supported by the Australian Research Council (DP180103834). Open access publishing facilitated by Australian National University, as part of the Wiley - Australian National University agreement via the Council of Australian University Librarians.














## Author contributions

AG, OG and OKA initiated the review. AG and OKA wrote the manuscript. AFB, SAB, JRE, RTF, MG, AHM, BJP, MPR, Y-LR, NLT and SDT provided substantial contributions to the editing and revision of the manuscript.

## Competing interests

None declared.

## ORCID

Owen K. Atkin  <https://orcid.org/0000-0003-1041-5202>  
 Andrew F. Bowerman  <https://orcid.org/0000-0003-1729-7843>  
 Sally A. Buck  <https://orcid.org/0000-0001-9569-0160>  
 John R. Evans  <https://orcid.org/0000-0003-1379-3532>  
 Robert T. Furbank  <https://orcid.org/0000-0001-8700-6613>  
 Oorbessy Gaju  <https://orcid.org/0000-0003-0493-7354>  
 Andres Garcia  <https://orcid.org/0000-0002-1554-6511>  
 Matthew Gilliham  <https://orcid.org/0000-0003-0666-3078>  
 A. Harvey Millar  <https://orcid.org/0000-0001-9679-1473>  
 Barry J. Pogson  <https://orcid.org/0000-0003-1869-2423>  
 Matthew P. Reynolds  <https://orcid.org/0000-0002-4291-4316>  
 Yong-Ling Ruan  <https://orcid.org/0000-0002-8394-4474>  
 Nicolas L. Taylor  <https://orcid.org/0000-0003-2004-5217>  
 Stephen D. Tyerman  <https://orcid.org/0000-0003-2455-1643>

## References

- Abadie C, Tcherkez G. 2019. Plant sulphur metabolism is stimulated by photorespiration. *Communications Biology* 2: 1–7.
- Abelenda JA, Bergonzi S, Oortwijn M, Sonnewald S, Du M, Visser RG, Sonnewald U, Bachem CW. 2019. Source-sink regulation is mediated by interaction of an FT homolog with a SWEET protein in potato. *Current Biology* 29: 1178–1186.
- Acevedo-Siaca LG, Coe R, Wang Y, Kromdijk J, Quick WP, Long SP. 2020. Variation in photosynthetic induction between rice accessions and its potential for improving productivity. *New Phytologist* 227: 1097–1108.
- Akhzari D, Pessarakli M. 2016. Effect of drought stress on total protein, essential oil content, and physiological traits of *Levisticum officinale* Koch. *Journal of Plant Nutrition* 39: 1365–1371.
- Allakhverdiev SI, Kreslavski VD, Klimov VV, Los DA, Carpentier R, Mohanty P. 2008. Heat stress: an overview of molecular responses in photosynthesis. *Photosynthesis Research* 98: 541–550.
- Amthor JS. 2000. The McCree-de Wit-Penning de Vries-Thornley respiration paradigms: 30 years later. *Annals of Botany* 86: 1–20.
- Amthor JS, Bar-Even A, Hanson AD, Millar AH, Stitt M, Sweetlove LJ, Tyerman SD. 2019. Engineering strategies to boost crop productivity by cutting respiratory carbon loss. *Plant Cell* 31: 297–314.
- Arcuri MLC, Nunes-Laitz AV, Lima RPM, Barreto P, Marinho AN, Arruda P, Maia IG. 2021. The knockdown of mitochondrial uncoupling proteins 1 and 2 (AtUCP1 and 2) in *Arabidopsis thaliana* impacts vegetative development and fertility. *Plant and Cell Physiology* 62: 1630–1644.
- Bailey-Serres J, Parker JE, Ainsworth EA, Oldroyd GE, Schroeder JI. 2019. Genetic strategies for improving crop yields. *Nature* 575: 109–118.
- Barberon M. 2017. The endodermis as a checkpoint for nutrients. *New Phytologist* 213: 1604–1610.
- Baroli I, Niyogi KK. 2000. Molecular genetics of xanthophyll-dependent photoprotection in green algae and plants. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 355: 1385–1394.
- Batista-Silva W, da Fonseca-Pereira P, Martins AO, Zsögön A, Nunes-Nesi A, Araújo WL. 2020. Engineering improved photosynthesis in the era of synthetic biology. *Plant Communications* 1: 100032.
- Berthold DA, Andersson ME, Nordlund P. 2000. New insight into the structure and function of the alternative oxidase. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* 1460: 241–254.
- Betti M, Bauwe H, Busch FA, Fernie AR, Keech O, Levey M, Ort DR, Parry MA, Sage R, Timm S. 2016. Manipulating photorespiration to increase plant productivity: recent advances and perspectives for crop improvement. *Journal of Experimental Botany* 67: 2977–2988.
- Bezruczyk M, Hartwig T, Horschman M, Char SN, Yang J, Yang B, Frommer WB, Sosso D. 2018. Impaired phloem loading in *zmsweet13a, b, c* sucrose transporter triple knock-out mutants in *Zea mays*. *New Phytologist* 218: 594–603.
- Bouma T, De Visser R, Janssen J, De Kock M, Van Leeuwen P, Lambers H. 1994. Respiratory energy requirements and rate of protein turnover *in vivo* determined by the use of an inhibitor of protein synthesis and a probe to assess its effect. *Physiologia Plantarum* 92: 585–594.
- Braun DM, Wang L, Ruan Y-L. 2014. Understanding and manipulating sucrose phloem loading, unloading, metabolism, and signalling to enhance crop yield and food security. *Journal of Experimental Botany* 65: 1713–1735.
- Burgess AJ, Retkute R, Preston SP, Jensen OE, Pound MP, Pridmore TP, Murchie EH. 2016. The 4-dimensional plant: effects of wind-induced canopy movement on light fluctuations and photosynthesis. *Frontiers in plant science* 7: 1392.
- Busch FA. 2020. Photorespiration in the context of Rubisco biochemistry, CO<sub>2</sub> diffusion and metabolism. *The Plant Journal* 101: 919–939.
- Campbell WH. 1999. Nitrate reductase structure, function and regulation: bridging the gap between biochemistry and physiology. *Annual Review of Plant Biology* 50: 277–303.
- Campbell WJ, Ogren WL. 1990. A novel role for light in the activation of ribulosebiphosphate carboxylase/oxygenase. *Plant Physiology* 92: 110–115.
- Carmo-Silva AE, Salvucci ME. 2013. The regulatory properties of Rubisco activase differ among species and affect photosynthetic induction during light transitions. *Plant Physiology* 161: 1645–1655.
- Cheah MH, Millar AH, Myers RC, Day DA, Roth J, Hillier W, Badger MR. 2014. Online oxygen kinetic isotope effects using membrane inlet mass spectrometry can differentiate between oxidases for mechanistic studies and calculation of their contributions to oxygen consumption in whole tissues. *Analytical Chemistry* 86: 5171–5178.
- Chen L-Q, Qu X-Q, Hou B-H, Sosso D, Osorio S, Fernie AR, Frommer WB. 2012. Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. *Science* 335: 207–211.
- Coast O, Shah S, Ivakov A, Gaju O, Wilson PB, Posch BC, Bryant CJ, Negrini ACA, Evans JR, Condon AG. 2019. Predicting dark respiration rates of wheat leaves from hyperspectral reflectance. *Plant, Cell & Environment* 42: 2133–2150.
- Collins SR, Wellner N, Martinez Bordonado I, Harper AL, Miller CN, Bancroft I, Waldron KW. 2014. Variation in the chemical composition of wheat straw: the role of tissue ratio and composition. *Biotechnology for Biofuels* 7: 1–14.
- Conlan B, Whitney S. 2018. Preparing Rubisco for a tune up. *Nature Plants* 4: 12–13.
- Crespo-Herrera L, Crossa J, Huerta-Espino J, Vargas M, Mondal S, Velu G, Payne T, Braun H, Singh R. 2018. Genetic gains for grain yield in CIMMYT's semi-arid wheat yield trials grown in suboptimal environments. *Crop Science* 58: 1890.
- Dahal K, Martyn GD, Vanlerberghe GC. 2015. Improved photosynthetic performance during severe drought in *Nicotiana tabacum* overexpressing a nonenergy conserving respiratory electron sink. *New Phytologist* 208: 382–395.
- Day DA, De Vos OC, Wilson D, Lambers H. 1985. Regulation of respiration in the leaves and roots of two *Lolium perenne* populations with contrasting mature leaf respiration rates and crop yields. *Plant Physiology* 78: 678–683.
- De Souza AP, Burgess SJ, Doran L, Hansen J, Manukyan L, Maryn N, Gotarkar D, Leonelli N, Niyogi KK, Long SP. 2022. Soybean photosynthesis and crop yield are improved by accelerating recovery from photoprotection. *Science* 377: 851–854.
- De Vries FP, Brunsting A, Van Laar H. 1974. Products, requirements and efficiency of biosynthesis a quantitative approach. *Journal of Theoretical Biology* 45: 339–377.
- De Vries FP, Van Laar H, Chardon M. 1983. Bioenergetics of growth of seeds, fruits and storage organs. In: Smith WH, ed. *Potential productivity of field crops under different environments*. Los Baños, Philippines: International Rice Research Institute, 37–59.
- Demircan N, Cucun G, Uzilday B. 2020. Mitochondrial alternative oxidase (AOX1a) is required for the mitigation of arsenic-induced oxidative stress in *Arabidopsis thaliana*. *Plant Biotechnology Reports* 14: 235–245.
- van Dijk M, Morley T, Rau ML, Saghai Y. 2021. A meta-analysis of projected global food demand and population at risk of hunger for the period 2010–2050. *Nature Food* 2: 494–501.

- Divakaruni AS, Brand MD. 2011. The regulation and physiology of mitochondrial proton leak. *Physiology* 26: 192–205.
- Djajaneegara I, Finnegan PM, Mathieu C, McCabe T, Whelan J, Day DA. 2002. Regulation of alternative oxidase gene expression in soybean. *Plant Molecular Biology* 50: 735–742.
- Doganlar ZB, Demir K, Basak H, Gul I. 2010. Effects of salt stress on pigment and total soluble protein contents of three different tomato cultivars. *African Journal of Agricultural Research* 5: 2056–2065.
- Dormatey R, Sun C, Ali K, Coulter JA, Bi Z, Bai J. 2020. Gene pyramiding for sustainable crop improvement against biotic and abiotic stresses. *Agronomy* 10: 1255.
- Dusenge ME, Duarte AG, Way DA. 2019. Plant carbon metabolism and climate change: elevated CO<sub>2</sub> and temperature impacts on photosynthesis, photorespiration and respiration. *New Phytologist* 221: 32–49.
- Edwards JM, Roberts TH, Atwell BJ. 2012. Quantifying ATP turnover in anoxic coleoptiles of rice (*Oryza sativa*) demonstrates preferential allocation of energy to protein synthesis. *Journal of Experimental Botany* 63: 4389–4402.
- Engel N, Ewald R, Gupta KJ, Zrenner R, Hagemann M, Bauwe H. 2011. The presence of Arabidopsis serine hydroxymethyltransferase SHM2 selectively prevents import into mesophyll mitochondria. *Plant Physiology* 157: 1711–1720.
- Engel N, van den Daele K, Kolukisaoglu Ü, Morgenthal K, Weckwerth W, Pärnik T, Keerbergh O, Bauwe H. 2007. Deletion of glycine decarboxylase in Arabidopsis is lethal under nonphotorespiratory conditions. *Plant Physiology* 144: 1328–1335.
- Ermakova M, Osborn H, Groszmann M, Bala S, Bowerman A, McGaughey S, Byrt C, Alonso-Cantabrana H, Tyerman S, Furbank RT. 2021. Expression of a CO<sub>2</sub>-permeable aquaporin enhances mesophyll conductance in the C<sub>4</sub> species *Setaria viridis*. *eLife* 10: e70095.
- Fan X, Tang Z, Tan Y, Zhang Y, Luo B, Yang M, Lian X, Shen Q, Miller AJ, Xu G. 2016. Overexpression of a pH-sensitive nitrate transporter in rice increases crop yields. *Proceedings of the National Academy of Sciences, USA* 113: 7118–7123.
- Feng L, Li J, Sun J, Wang L, Fan C, Shen J. 2021. Recent advances of DNA nanostructure-based cell membrane engineering. *Advanced Healthcare Materials* 10: 2001718.
- Fernie AR, Bachem CW, Helariutta Y, Neuhaus HE, Prat S, Ruan Y-L, Stitt M, Sweetlove LJ, Tegeder M, Wahl V. 2020. Synchronization of developmental, molecular and metabolic aspects of source–sink interactions. *Nature Plants* 6: 55–66.
- Fu P, Meacham-Hensold K, Guan K, Wu J, Bernacchi C. 2020. Estimating photosynthetic traits from reflectance spectra: a synthesis of spectral indices, numerical inversion, and partial least square regression. *Plant, Cell & Environment* 43: 1241–1258.
- Fu Q, Cheng L, Guo Y, Turgeon R. 2011. Phloem loading strategies and water relations in trees and herbaceous plants. *Plant Physiology* 157: 1518–1527.
- Furbank RT, Jimenez-Berni JA, George-Jaeggli B, Potgieter AB, Deery DM. 2019. Field crop phenomics: enabling breeding for radiation use efficiency and biomass in cereal crops. *New Phytologist* 223: 1714–1727.
- Gibon Y, Pyl ET, Sulpice R, Lunn JE, Hoehne M, Günther M, Stitt M. 2009. Adjustment of growth, starch turnover, protein content and central metabolism to a decrease of the carbon supply when Arabidopsis is grown in very short photoperiods. *Plant, Cell & Environment* 32: 859–874.
- Giraud E, Ho LH, Clifton R, Carroll A, Estavillo G, Tan Y-F, Howell KA, Ivanova A, Pogson BJ, Millar AH. 2008. The absence of ALTERNATIVE OXIDASE1a in Arabidopsis results in acute sensitivity to combined light and drought stress. *Plant Physiology* 147: 595–610.
- Giri A, Heckathorn S, Mishra S, Krause C. 2017. Heat stress decreases levels of nutrient-uptake and-assimilation proteins in tomato roots. *Plants* 6: 6.
- Gong Q, Li S, Zheng Y, Duan H, Xiao F, Zhuang Y, He J, Wu G, Zhao S, Zhou H. 2020. SUMOylation of MYB30 enhances salt tolerance by elevating alternative respiration via transcriptionally upregulating AOX1a in Arabidopsis. *The Plant Journal* 102: 1157–1171.
- Griffiths M, Roy S, Guo H, Seethepalli A, Huhman D, Ge Y, Sharp RE, Fritschi FB, York LM. 2021. A multiple ion-uptake phenotyping platform reveals shared mechanisms affecting nutrient uptake by roots. *Plant Physiology* 185: 781–795.
- Griffiths M, York LM. 2020. Targeting root ion uptake kinetics to increase plant productivity and nutrient use efficiency. *Plant Physiology* 182: 1854–1868.
- Groszmann M, Osborn HL, Evans JR. 2017. Carbon dioxide and water transport through plant aquaporins. *Plant, Cell & Environment* 40: 938–961.
- Guy RD, Fogel MF, Berry JA, Hoering TC. 1987. Isotope fractionation during oxygen production and consumption by plants. In: Biggins J, ed. *Progress in photosynthesis research*. Providence, RI, USA: Springer, 597–600.
- Hachiya T, Terashima I, Noguchi K. 2007. Increase in respiratory cost at high growth temperature is attributed to high protein turnover cost in *Petunia × hybrida* petals. *Plant, Cell & Environment* 30: 1269–1283.
- Hanson AD, Amthor JS, Sun J, Niehaus TD, Gregory JF III, Bruner SD, Ding Y. 2018. Redesigning thiamin synthesis: prospects and potential payoffs. *Plant Science* 273: 92–99.
- Hao M, Maxfield FR. 2000. Characterization of rapid membrane internalization and recycling. *Journal of Biological Chemistry* 275: 15279–15286.
- Hauben M, Haesendonckx B, Staeldert E, Van Der Kelen K, Azmi A, Akpo H, Van Breusegem F, Guisez Y, Bots M, Lambert B. 2009. Energy use efficiency is characterized by an epigenetic component that can be directed through artificial selection to increase yield. *Proceedings of the National Academy of Sciences, USA* 106: 20109–20114.
- Havaux M, Niyogi KK. 1999. The violaxanthin cycle protects plants from photooxidative damage by more than one mechanism. *Proceedings of the National Academy of Sciences, USA* 96: 8762–8767.
- Hazell PB. 2009. *The Asian green revolution*. IFPRI Discussion Paper: International Food Policy Research Institute. [WWW document] URL <https://www.ifpri.org/publication/asian-green-revolution> [accessed 16 March 2019].
- Hinkson IV, Elias JE. 2011. The dynamic state of protein turnover: it's about time. *Trends in Cell Biology* 21: 293–303.
- Hu B, Wang W, Ou S, Tang J, Li H, Che R, Zhang Z, Chai X, Wang H, Wang Y. 2015. Variation in NRT1. 1B contributes to nitrate-use divergence between rice subspecies. *Nature Genetics* 47: 834.
- Huang S, Li L, Peterleit J, Millar AH. 2020. Protein turnover rates in plant mitochondria. *Mitochondrion* 53: 57–65.
- Iizumi T, Furuya J, Shen Z, Kim W, Okada M, Fujimori S, Hasegawa T, Nishimori M. 2017. Responses of crop yield growth to global temperature and socioeconomic changes. *Scientific Reports* 7: 1–10.
- Ishihara H, Moraes TA, Pyl ET, Schulze WX, Obata T, Scheffel A, Fernie AR, Sulpice R, Stitt M. 2017. Growth rate correlates negatively with protein turnover in Arabidopsis accessions. *The Plant Journal* 91: 416–429.
- Ishihara H, Obata T, Sulpice R, Fernie AR, Stitt M. 2015. Quantifying protein synthesis and degradation in Arabidopsis by dynamic <sup>13</sup>CO<sub>2</sub> labeling and analysis of enrichment in individual amino acids in their free pools and in protein. *Plant Physiology* 168: 74–93.
- Jacoby RP, Millar AH, Taylor NL. 2016. Opportunities for wheat proteomics to discover the biomarkers for respiration-dependent biomass production, stress tolerance and cytoplasmic male sterility. *Journal of Proteomics* 143: 36–44.
- Jimenez-Berni JA, Deery DM, Rozas-Larraondo P, Condon ATG, Rebetzke GJ, James RA, Bovill WD, Furbank RT, Sirault XR. 2018. High throughput determination of plant height, ground cover, and above-ground biomass in wheat with LiDAR. *Frontiers in Plant Science* 9: 237.
- Jin S, Su Y, Song S, Xu K, Hu T, Yang Q, Wu F, Xu G, Ma Q, Guan H. 2020. Non-destructive estimation of field maize biomass using terrestrial lidar: an evaluation from plot level to individual leaf level. *Plant Methods* 16: 1–19.
- Johnson IR, Thornley JH, Frantz JM, Bugbee B. 2010. A model of canopy photosynthesis incorporating protein distribution through the canopy and its acclimation to light, temperature and CO<sub>2</sub>. *Annals of Botany* 106: 735–749.
- Jouyban Z. 2012. The effects of salt stress on plant growth. *Technical Journal of Engineering and Applied Sciences* 2: 7–10.
- Kebeish R, Niessen M, Thiruveedhi K, Bari R, Hirsch H-J, Rosenkranz R, Stäbler N, Schönfeld B, Kreuzaler F, Peterhänsel C. 2007. Chloroplastic photorespiratory bypass increases photosynthesis and biomass production in *Arabidopsis thaliana*. *Nature Biotechnology* 25: 593–599.
- Kosová K, Vítámvás P, Prášil IT, Renaut J. 2011. Plant proteome changes under abiotic stress – contribution of proteomics studies to understanding plant stress response. *Journal of Proteomics* 74: 1301–1322.
- Kraus E, Lambers H. 2001. Leaf and root respiration of *Lolium perenne* populations selected for contrasting leaf respiration rates are affected by intra- and interpopulation interactions. *Plant and Soil* 231: 267–274.
- Kraus E, Lambers H, Kollöffel C. 1993. The effect of handling on the yield of two populations of *Lolium perenne* selected for differences in mature leaf respiration rate. *Physiologia Plantarum* 89: 341–346.



- Lambers H, Chapin FS, Pons TL. 2008. *Plant physiological ecology*. New York, NY, USA: Springer.
- Lambers H, Poorter H. 1992. Inherent variation in growth rate between higher plants: a search for physiological causes and ecological consequences. In: Begon M, ed. *Advances in ecological research*. London, UK: Academic Press, 187–261.
- Lambers H, Rychter A. 1989. The biochemical background of variation in respiration rate: respiratory pathways and chemical composition. In: Lambers H, ed. *Causes and consequences of variation in growth rate and productivity of higher plants*. The Hague, the Netherlands: SPB Academic, 199–225.
- Lee CP, Makshev G, Jensen GS, Murcha MW, Wilson ME, Fricker M, Hell R, Haswell ES, Millar AH, Sweetlove LJ. 2016. MSL1 is a mechanosensitive ion channel that dissipates mitochondrial membrane potential and maintains redox homeostasis in mitochondria during abiotic stress. *The Plant Journal* 88: 809–825.
- Li L, Nelson CJ, Trösch J, Castleden I, Huang S, Millar AH. 2017. Protein degradation rate in *Arabidopsis thaliana* leaf growth and development. *Plant Cell* 29: 207–228.
- Liu L, Gallagher J, Arevalo ED, Chen R, Skopelitis T, Wu QY, Bartlett M, Jackson D. 2021. Enhancing grain-yield-related traits by CRISPR-Cas9 promoter editing of maize CLE genes. *Nature Plants* 7: 287–294.
- Long BM, Förster B, Pulsford SB, Price GD, Badger MR. 2021. Rubisco proton production can drive the elevation of CO<sub>2</sub> within condensates and carboxysomes. *Proceedings of the National Academy of Sciences, USA* 118: 1–12.
- Long S, Humphries S, Falkowski PG. 1994. Photoinhibition of photosynthesis in nature. *Annual Review of Plant Biology* 45: 633–662.
- Long SP, Marshall-Colon A, Zhu X-G. 2015. Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. *Cell* 161: 56–66.
- Long SP, Zhu XG, Naidu SL, Ort DR. 2006. Can improvement in photosynthesis increase crop yields? *Plant, Cell & Environment* 29: 315–330.
- Loomis R, Amthor J. 1999. Yield potential, plant assimilatory capacity, and metabolic efficiencies. *Crop Science* 39: 1584–1596.
- Loveys B, Scheurwater I, Pons T, Fitter A, Atkin O. 2002. Growth temperature influences the underlying components of relative growth rate: an investigation using inherently fast- and slow-growing plant species. *Plant, Cell & Environment* 25: 975–988.
- Maier A, Fahnenstich H, Von Caemmerer S, Engqvist MK, Weber AP, Flügge U-I, Maurino VG. 2012. Transgenic introduction of a glycolate oxidative cycle into *A. thaliana* chloroplasts leads to growth improvement. *Frontiers in Plant Science* 3: 38.
- Marques WL, Mans R, Henderson RK, Marella ER, Ter Horst J, de Hulster E, Poolman B, Daran J-M, Pronk JT, Gombert AK. 2018. Combined engineering of disaccharide transport and phosphorylation for enhanced ATP yield from sucrose fermentation in *Saccharomyces cerevisiae*. *Metabolic Engineering* 45: 121–133.
- Marshall RS, Vierstra RD. 2018. Autophagy: the master of bulk and selective recycling. *Annual Review of Plant Biology* 69: 173–208.
- Mathur S, Agrawal D, Jajoo A. 2014. Photosynthesis: response to high temperature stress. *Journal of Photochemistry and Photobiology B: Biology* 137: 116–126.
- McIntosh L. 1994. Molecular biology of the alternative oxidase. *Plant Physiology* 105: 781.
- Millar AH, Whelan J, Soole KL, Day DA. 2011. Organization and regulation of mitochondrial respiration in plants. *Annual Review of Plant Biology* 62: 79–104.
- Monteith JL. 1977. Climate and the efficiency of crop production in Britain. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 281: 277–294.
- Moreau D, Allard V, Gaju O, Le Gouis J, Foulkes MJ, Martre P. 2012. Acclimation of leaf nitrogen to vertical light gradient at anthesis in wheat is a whole-plant process that scales with the size of the canopy. *Plant Physiology* 160: 1479–1490.
- Móznér Z, Tabi A, Csutóra M. 2012. Modifying the yield factor based on more efficient use of fertilizer – The environmental impacts of intensive and extensive agricultural practices. *Ecological Indicators* 16: 58–66.
- Müller-Schüssele SJ, Schwarzländer M, Meyer AJ. 2021. Live monitoring of plant redox and energy physiology with genetically encoded biosensors. *Plant Physiology* 186: 93.
- Mundim KC, Baraldi S, Machado HG, Vieira FM. 2020. Temperature coefficient (Q<sub>10</sub>) and its applications in biological systems: beyond the Arrhenius theory. *Ecological Modelling* 431: 109127.
- Munns R, Gilliam M. 2015. Salinity tolerance of crops—what is the cost? *New Phytologist* 208: 668–673.
- Murchie EH, Ali A, Herman T. 2015. Photoprotection as a trait for rice yield improvement: status and prospects. *Rice* 8: 1–9.
- Murchie EH, Niyogi KK. 2011. Manipulation of photoprotection to improve plant photosynthesis. *Plant Physiology* 155: 86–92.
- Musgrave ME, Strain BR, Siedow JN. 1986. Response of two pea hybrids to CO<sub>2</sub> enrichment: a test of the energy overflow hypothesis for alternative respiration. *Proceedings of the National Academy of Sciences, USA* 83: 8157–8161.
- Nelson CJ, Li L, Millar AH. 2014. Quantitative analysis of protein turnover in plants. *Proteomics* 14: 579–592.
- Nelson CJ, Millar AH. 2015. Protein turnover in plant biology. *Nature Plants* 1: 15017.
- Noguchi K, Go C-S, Miyazawa S-I, Terashima I, Ueda S, Yoshinari T. 2001. Costs of protein turnover and carbohydrate export in leaves of sun and shade species. *Functional Plant Biology* 28: 37–47.
- Novitskaya L, Trevanion S, Driscoll S, Foyer C, Noctor G. 2002. How does photorespiration modulate leaf amino acid contents? A dual approach through modelling and metabolite analysis. *Plant, Cell & Environment* 25: 821–835.
- O’Leary BM, Lee CP, Atkin OK, Cheng R, Brown TB, Millar AH. 2017. Variation in leaf respiration rates at night correlates with carbohydrate and amino acid supply. *Plant Physiology* 174: 2261–2273.
- Obenland D, Hiser C, McIntosh L, Shibles R, Stewart CR. 1988. Occurrence of alternative respiratory capacity in soybean and pea. *Plant Physiology* 88: 528–531.
- O’Leary BM, Asao S, Millar AH, Atkin OK. 2019. Core principles which explain variation in respiration across biological scales. *New Phytologist* 222: 670–686.
- Ort D, Long S. 2003. Converting solar energy into crop production. In: Chrispeels MJ, ed. *Plants, genes, and crop biotechnology*. Sudbury, MA, USA: Jones and Bartlett Publisher International, 240–269.
- Parry MA, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney SM. 2013. Rubisco activity and regulation as targets for crop improvement. *Journal of Experimental Botany* 64: 717–730.
- Pascale A, Proietti S, Pantelides IS, Stringlis IA. 2020. Modulation of the root microbiome by plant molecules: the basis for targeted disease suppression and plant growth promotion. *Frontiers in Plant Science* 10: 1741.
- Penning de Vries FWT. 1975. The cost of maintenance processes in plant cells. *Annals of Botany* 39: 77–92.
- Peterhansel C, Horst I, Niessen M, Blume C, Kebeish R, Kürkcüoglu S, Kreuzaler F. 2010. Photorespiration. In: *The Arabidopsis book*, vol. 8. Rockville, MD, USA: American Society of Plant Biologists, 1–29.
- Peterhansel C, Krause K, Braun HP, Espie G, Fernie A, Hanson D, Keech O, Maurino V, Mielewicz M, Sage R. 2013. Engineering photorespiration: current state and future possibilities. *Plant Biology* 15: 754–758.
- Plaxton WC, Podestá FE. 2006. The functional organization and control of plant respiration. *Critical Reviews in Plant Sciences* 25: 159–198.
- Plett D, Toubia J, Garnett T, Tester M, Kaiser BN, Baumann U. 2010. Dichotomy in the NRT gene families of dicots and grass species. *PLoS ONE* 5: e15289.
- Polidoros AN, Mylona PV, Pasentsis K, Scandalios JG, Tsafaris AS. 2005. The maize alternative oxidase 1a (Aox1a) gene is regulated by signals related to oxidative stress. *Redox Report* 10: 71–78.
- Poorter H, Remkes C, Lambers H. 1990. Carbon and nitrogen economy of 24 wild species differing in relative growth rate. *Plant Physiology* 94: 621–627.
- Poorter H, Van der Werf A, Atkin OK, Lambers H. 1991. Respiratory energy requirements of roots vary with the potential growth rate of a plant species. *Physiologia Plantarum* 83: 469–475.
- Poorter H, Villar R. 1994. The fate of acquired carbon in plants: chemical composition and construction costs. In: Fakhri A, Bazzaz JG, eds. *Plant resource allocation*. San Diego, CA, USA: Academic Press, 39–72.
- Price GD, Pengelly JJ, Forster B, Du J, Whitney SM, von Caemmerer S, Badger MR, Howitt SM, Evans JR. 2013. The cyanobacterial CCM as a source of genes for improving photosynthetic CO<sub>2</sub> fixation in crop species. *Journal of Experimental Botany* 64: 753–768.
- Prins A, Orr DJ, Andralojc PJ, Reynolds MP, Carmo-Silva E, Parry MA. 2016. Rubisco catalytic properties of wild and domesticated relatives provide scope for improving wheat photosynthesis. *Journal of Experimental Botany* 67: 1827–1838.



- Quigg A, Beardall J. 2003. Protein turnover in relation to maintenance metabolism at low photon flux in two marine microalgae. *Plant, Cell & Environment* **26**: 693–703.
- Rasmusson AG, Soole KL, Elthon TE. 2004. Alternative NAD (P) H dehydrogenases of plant mitochondria. *Annual Review of Plant Biology* **55**: 23–39.
- Ray DK, Mueller ND, West PC, Foley JA. 2013. Yield trends are insufficient to double global crop production by 2050. *PLoS ONE* **8**: e66428.
- Retkute R, Smith-Unna SE, Smith RW, Burgess AJ, Jensen OE, Johnson GN, Preston SP, Murchie EH. 2015. Exploiting heterogeneous environments: does photosynthetic acclimation optimize carbon gain in fluctuating light? *Journal of Experimental Botany* **66**: 2437–2447.
- Rivero RM, Mittler R, Blumwald E, Zandalinas SI. 2022. Developing climate-resilient crops: improving plant tolerance to stress combination. *The Plant Journal* **109**: 373–389.
- Robson M. 1982a. The growth and carbon economy of selection lines of *Lolium perenne* cv. S23 with differing rates of dark respiration: 1. Grown as simulated swards during a regrowth period. *Annals of Botany* **49**: 321–329.
- Robson M. 1982b. The growth and carbon economy of selection lines of *Lolium perenne* cv. S23 with differing rates of dark respiration: 2. Grown as young plants from seed. *Annals of Botany* **49**: 331–339.
- Rodriguez-Leal D, Lemmon ZH, Man J, Bartlett ME, Lippman ZB. 2017. Engineering quantitative trait variation for crop improvement by genome editing. *Cell* **171**: 470–480.
- Ross AB, Langer JD, Jovanovic M. 2021. Proteome turnover in the spotlight: approaches, applications, and perspectives. *Molecular & Cellular Proteomics* **20**: 1–23.
- Ruan Y-L. 2007. Rapid cell expansion and cellulose synthesis regulated by plasmodesmata and sugar: insights from the single-celled cotton fibre. *Functional Plant Biology* **34**: 1–10.
- Ruan Y-L. 2012. Signaling role of sucrose metabolism in development. *Molecular Plant* **5**: 763–765.
- Ruan Y-L. 2014. Sucrose metabolism: gateway to diverse carbon use and sugar signaling. *Annual Review of Plant Biology* **65**: 33–67.
- Sage TL, Sage RF. 2009. The functional anatomy of rice leaves: implications for refixation of photorespiratory CO<sub>2</sub> and efforts to engineer C<sub>4</sub> photosynthesis into rice. *Plant and Cell Physiology* **50**: 756–772.
- Saisho D, Nambara E, Naito S, Tsutsumi N, Hirai A, Nakazono M. 1997. Characterization of the gene family for alternative oxidase from *Arabidopsis thaliana*. *Plant Molecular Biology* **35**: 585–596.
- Scafaro AP, Fan Y, Posch BC, Garcia A, Coast O, Atkin OK. 2021. Responses of leaf respiration to heatwaves. *Plant, Cell & Environment* **44**: 2090–2101.
- Scafaro AP, Negrini ACA, O'Leary B, Rashid FAA, Hayes L, Fan Y, Zhang Y, Chochois V, Badger MR, Millar AH. 2017. The combination of gas-phase fluorophore technology and automation to enable high-throughput analysis of plant respiration. *Plant Methods* **13**: 16.
- Scheurwater I, Clarkson DT, Purves JV, Van Rijt G, Saker LR, Welschen R, Lambers H. 1999. Relatively large nitrate efflux can account for the high specific respiratory costs for nitrate transport in slow-growing grass species. *Plant and Soil* **215**: 123–134.
- Scheurwater I, Cornelissen C, Dictus F, Welschen R, Lambers H. 1998. Why do fast- and slow-growing grass species differ so little in their rate of root respiration, considering the large differences in rate of growth and ion uptake? *Plant, Cell & Environment* **21**: 995–1005.
- Segonzac C, Boyer J-C, Ipotesi E, Szponarski W, Tillard P, Touraine B, Sommerer N, Rossignol M, Gibrat R. 2007. Nitrate efflux at the root plasma membrane: identification of an *Arabidopsis* excretion transporter. *Plant Cell* **19**: 3760–3777.
- Seki M, Feugier FG, Song X-J, Ashikari M, Nakamura H, Ishiyama K, Yamaya T, Inari-Ikeda M, Kitano H, Satake A. 2015. A mathematical model of phloem sucrose transport as a new tool for designing rice panicle structure for high grain yield. *Plant and Cell Physiology* **56**: 605–619.
- Silva-Perez V, Molero G, Serbin SP, Condon AG, Reynolds MP, Furbank RT, Evans JR. 2018. Hyperspectral reflectance as a tool to measure biochemical and physiological traits in wheat. *Journal of Experimental Botany* **69**: 483–496.
- Sinclair TR. 1998. Historical changes in harvest index and crop nitrogen accumulation. *Crop Science* **38**: 638–643.
- Smith C, Barthet M, Melino V, Smith P, Day D, Soole K. 2011. Alterations in the mitochondrial alternative NAD(P)H dehydrogenase NDB4 lead to changes in mitochondrial electron transport chain composition, plant growth and response to oxidative stress. *Plant and Cell Physiology* **52**: 1222–1237.
- Smith NG, Keenan TF. 2020. Mechanisms underlying leaf photosynthetic acclimation to warming and elevated CO<sub>2</sub> as inferred from least-cost optimality theory. *Global Change Biology* **26**: 5202–5216.
- Smith P. 2013. Delivering food security without increasing pressure on land. *Global Food Security* **2**: 18–23.
- Soleh M, Tanaka Y, Kim SY, Huber SC, Sakoda K, Shiraiwa T. 2017. Identification of large variation in the photosynthetic induction response among 37 soybean [*Glycine max* (L.) Merr.] genotypes that is not correlated with steady-state photosynthetic capacity. *Photosynthesis Research* **131**: 305–315.
- South PF, Cavanagh AP, Liu HW, Ort DR. 2019. Synthetic glycolate metabolism pathways stimulate crop growth and productivity in the field. *Science* **363**: 1–9.
- Steer MW. 1988. Plasma membrane turnover in plant cells. *Journal of Experimental Botany* **39**: 987–996.
- Sweetlove LJ, Lytovchenko A, Morgan M, Nunes-Nesi A, Taylor NL, Baxter CJ, Eickmeier I, Fernie AR. 2006. Mitochondrial uncoupling protein is required for efficient photosynthesis. *Proceedings of the National Academy of Sciences, USA* **103**: 19587–19592.
- Sweetman C, Waterman CD, Rainbird BM, Smith PMC, Jenkins CD, Day DA, Soole KL. 2019. AtNDB2 is the main external NADH dehydrogenase in mitochondria and is important for tolerance to environmental stress. *Plant Physiology* **181**: 774–788.
- Taylor NL, Heazlewood JL, Day DA, Millar AH. 2005. Differential impact of environmental stresses on the pea mitochondrial proteome. *Molecular & Cellular Proteomics* **4**: 1122–1133.
- Taylor SH, Long SP. 2017. Slow induction of photosynthesis on shade to sun transitions in wheat may cost at least 21% of productivity. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **372**: 20160543.
- Teakle NL, Tyerman SD. 2010. Mechanisms of Cl<sup>-</sup> transport contributing to salt tolerance. *Plant, Cell & Environment* **33**: 566–589.
- Tester M, Langridge P. 2010. Breeding technologies to increase crop production in a changing world. *Science* **327**: 818–822.
- Tilman D. 1998. The greening of the green revolution. *Nature* **396**: 211.
- Timm S, Bauwe H. 2013. The variety of photorespiratory phenotypes—employing the current status for future research directions on photorespiration. *Plant Biology* **15**: 737–747.
- Tjoelker MG, Oleksyn J, Reich PB. 2001. Modelling respiration of vegetation: evidence for a general temperature-dependent Q<sub>10</sub>. *Global Change Biology* **7**: 223–230.
- Vanlerberghe GC, McIntosh L. 1997. Alternative oxidase: from gene to function. *Annual Review of Plant Biology* **48**: 703–734.
- Vierstra RD. 2009. The ubiquitin–26S proteasome system at the nexus of plant biology. *Nature Reviews Molecular Cell Biology* **10**: 385–397.
- Voosen P. 2021. *Global temperatures in 2020 tied record highs*. Science: American Association for the Advancement of Science. [WWW document] URL <https://www.science.org/content/article/global-temperatures-2020-tied-record-highs> [accessed 23 February 2021].
- Walker BJ, South PF, Ort DR. 2016. Physiological evidence for plasticity in glycolate/glycerate transport during photorespiration. *Photosynthesis Research* **129**: 93–103.
- Wallström SV, Florez-Sarasa I, Araújo WL, Aidemark M, Fernández-Fernández M, Fernie AR, Ribas-Carbó M, Rasmusson AG. 2014. Suppression of the external mitochondrial NADPH dehydrogenase, NDB1, in *Arabidopsis thaliana* affects central metabolism and vegetative growth. *Molecular Plant* **7**: 356–368.
- Walter JD, Edwards J, McDonald G, Kuchel H. 2019. Estimating biomass and canopy height with LiDAR for field crop breeding. *Frontiers in Plant Science* **10**: 1145.
- Wang W, Vinocur B, Shoseyov O, Altman A. 2004. Role of plant heat-shock proteins and molecular chaperones in the abiotic stress response. *Trends in Plant Science* **9**: 244–252.
- Warren L, Glick M. 1968. Membranes of animal cells: II. The metabolism and turnover of the surface membrane. *The Journal of Cell Biology* **37**: 729–746.
- Wege S, Gilliam M, Henderson SW. 2017. Chloride: not simply a 'cheap osmoticum', but a beneficial plant macronutrient. *Journal of Experimental Botany* **68**: 3057–3069.

- Whitney SM, Houtz RL, Alonso H. 2011. Advancing our understanding and capacity to engineer nature's CO<sub>2</sub>-sequestering enzyme, Rubisco. *Plant Physiology* 155: 27–35.
- Wilson D. 1975. Variation in leaf respiration in relation to growth and photosynthesis of *Lolium*. *Annals of Applied Biology* 80: 323–338.
- Wilson D. 1982. Response to selection for dark respiration rate of mature leaves in *Lolium perenne* and its effects on growth of young plants and simulated swards. *Annals of Botany* 49: 303–312.
- Wilson D, Jones J. 1982. Effect of selection for dark respiration rate of mature leaves on crop yields of *Lolium perenne* cv. S23. *Annals of Botany* 49: 313–320.
- World Bank. 2007. *World development report 2008: agriculture for development*. Washington, DC, USA: The World Bank.
- Xin C-P, Tholen D, Devloo V, Zhu X-G. 2015. The benefits of photorespiratory bypasses: how can they work? *Plant Physiology* 167: 574–585.
- Xu G, Fan X, Miller AJ. 2012. Plant nitrogen assimilation and use efficiency. *Annual Review of Plant Biology* 63: 153–182.
- Xu Q, Yin S, Ma Y, Song M, Song Y, Mu S, Li Y, Liu X, Ren Y, Gao C *et al.* 2020. Carbon export from leaves is controlled via ubiquitination and phosphorylation of sucrose transporter SUC2. *Proceedings of the National Academy of Sciences, USA* 117: 6223–6230.
- Yamori W. 2021. Strategies for engineering photosynthesis for enhanced plant biomass production. In: Jauhar Ali SHW, ed. *Rice improvement: physiological, molecular breeding and genetic perspectives*. Cham, Switzerland: Springer, 31–58.
- Yamori W, Masumoto C, Fukayama H, Makino A. 2012. Rubisco activase is a key regulator of non-steady-state photosynthesis at any leaf temperature and, to a lesser extent, of steady-state photosynthesis at high temperature. *The Plant Journal* 71: 871–880.
- Yukioka H, Inagaki S, Tanaka R, Katoh K, Miki N, Mizutani A, Masuko M. 1998. Transcriptional activation of the alternative oxidase gene of the fungus *Magnaporthe grisea* by a respiratory-inhibiting fungicide and hydrogen peroxide. *Biochimica et Biophysica Acta (BBA)-Gene Structure and Expression* 1442: 161–169.
- Zagdańska B. 1995. Respiratory energy demand for protein turnover and ion transport in wheat leaves upon water deficit. *Physiologia Plantarum* 95: 428–436.
- Zhang C, Turgeon R. 2018. Mechanisms of phloem loading. *Current Opinion in Plant Biology* 43: 71–75.
- Zhang H, Li S, Shi M, Wang S, Shi L, Xu F, Ding G. 2020. Genome-wide systematic characterization of the NPF family genes and their transcriptional responses to multiple nutrient stresses in allotetraploid rapeseed. *International Journal of Molecular Sciences* 21: 5947.
- Zhang HW, Si XM, Ji X, Fan R, Liu JX, Chen KL, Wang DW, Gao CX. 2018. Genome editing of upstream open reading frames enables translational control in plants. *Nature Biotechnology* 36: 894–898.
- Zhang Z, Ruan Y-L, Zhou N, Wang F, Guan X, Fang L, Shang X, Guo W, Zhu S, Zhang T. 2017. Suppressing a putative sterol carrier gene reduces plasmodesmal permeability and activates sucrose transporter genes during cotton fiber elongation. *Plant Cell* 29: 2027–2046.
- Zhao M, Tan HT, Scharwies J, Levin K, Evans JR, Tyerman SD. 2017. Association between water and carbon dioxide transport in leaf plasma membranes: assessing the role of aquaporins. *Plant, Cell & Environment* 40: 789–801.
- Zhu X-G, Long SP, Ort DR. 2010. Improving photosynthetic efficiency for greater yield. *Annual Review of Plant Biology* 61: 235–261.
- Zhu XG, Ort DR, Whitmarsh J, Long SP. 2004. The slow reversibility of photosystem II thermal energy dissipation on transfer from high to low light may cause large losses in carbon gain by crop canopies: a theoretical analysis. *Journal of Experimental Botany* 55: 1167–1175.