

CORRECTION

Open Access



Correction: Antioxidant effect of nicotinamide mononucleotide in tendinopathy

Kohei Yamaura, Yutaka Mifune^{*}, Atsuyuki Inui, Hanako Nishimoto, Takashi Kurosawa, Shintaro Mukohara, Yuichi Hoshino, Takahiro Niikura and Ryosuke Kuroda

Correction: BMC Musculoskelet Disord 23, 249 (2022)
<https://doi.org/10.1186/s12891-022-05205-z>

Following the publication of the original article [1] the author noticed that the published Fig. 8 is incorrect.

The original article [1] has been updated.

Below is the correct Fig. 8.

Published online: 06 June 2022

Reference

1. Yamaura K, Mifune Y, Inui A, et al. Antioxidant effect of nicotinamide mononucleotide in tendinopathy. *BMC Musculoskelet Disord*. 2022;23:249. <https://doi.org/10.1186/s12891-022-05205-z>.

The original article can be found online at <https://doi.org/10.1186/s12891-022-05205-z>.

*Correspondence: m-ship@kf7.so-net.ne.jp

Department of Orthopaedic Surgery, Kobe University Graduate School of Medicine, 7-5-1, Kusunoki-cho, Chuo-ku, Kobe 650-0017, Japan



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 8 Quantitative expression of SIRT1, SIRT6, and IL6 mRNA in vivo. Quantitative RT-PCR results showing the mRNA expression of SIRT1, SIRT6, and IL6 in an in vivo collagenase-induced tendinopathy model. Expression is shown for the NMN group and the control group. The expression of IL6 was significantly suppressed in the NMN group compared to the control group, whereas the expression of SIRT1 and SIRT6 was significantly increased in the NMN group compared to the control group. * $p < 0.05$; ** $p < 0.0001$