

RESEARCH

Open Access



The association between residential eviction and syringe sharing among a prospective cohort of street-involved youth

Andreas Pilarinos^{1,2}, Mary Clare Kennedy^{1,3}, Ryan McNeil^{1,4}, Huiru Dong¹, Thomas Kerr^{1,4} and Kora DeBeck^{1,5*}

Abstract

Background: Syringe sharing is a high-risk practice associated with the transmission of infectious diseases, such as HIV and HCV. While youth who contend with housing instability are known to be more likely to engage in high-risk substance use, the potential relationship between being evicted from housing and syringe sharing has not been examined. This study assessed whether residential eviction was associated with syringe sharing among street-involved youth in Vancouver, Canada.

Methods: Data were derived from the At-Risk Youth Study (ARYS), a prospective cohort of street-involved youth who use drugs age 14–26 in Vancouver, Canada. The study period was June 2007 to May 2014, and the potential relationship between residential eviction and syringe sharing was analyzed using multivariable generalized estimating equations (GEE) logistic regression.

Results: Among 405 street-involved youth who injected drugs, 149 (36.8%) reported syringe sharing, defined as borrowing or lending a syringe, at some point during the study period. In a multivariable GEE analysis, recent residential eviction remained independently associated with syringe sharing (adjusted odds ratio (AOR) = 1.72, 95% confidence interval (CI): 1.16–2.57), after adjusting for potential confounders.

Conclusions: Syringe sharing was significantly elevated among youth who had recently been evicted from housing. These findings indicate that policy and programmatic interventions that increase housing stability may help mitigate high-risk substance use practices among vulnerable youth.

Keywords: Eviction, Street-involved youth, Syringe sharing, Housing policy

Background

Injection drug use constitutes a significant public health concern due to its association with the transmission of blood-borne pathogens such as HIV and HCV as well as other severe health-related harms [1–4]. These negative health consequences are particularly pronounced among people who inject drugs (PWID) and share syringes [5]. Previous research has identified risk factors associated with syringe sharing that include difficulty accessing needles, binge drug use, injection cocaine use, and homelessness [6, 7]. While evidence suggests that there have been

significant reductions in syringe sharing among adult populations in Vancouver, Canada, [8–10], research among street-involved youth indicates that rates of syringe sharing have persisted at concerning levels [11, 12].

Previous studies of street-involved populations have documented numerous health-related risks and harms associated with housing insecurity. To date, this work has predominantly focused on the adverse health impacts of homelessness, including increased likelihood of drug-related risk behaviors such as intensified substance use [13, 14], initiation of injection drug use [15, 16], public injecting [2], and syringe sharing [17]. More recently, there has been increased research interest in the health consequences of residential eviction (i.e., forced displacement of a tenant from a leased residence through legal or extra-legal mechanisms). For instance, eviction

* Correspondence: uhri-kd@cfenet.ubc.ca

¹BC Centre on Substance Use, British Columbia Centre for Excellence in HIV/AIDS, 608-1081 Burrard Street, Vancouver, BC V6Z 1Y6, Canada

⁵School of Public Policy, Simon Fraser University, 3277-515 Hastings W Street, Vancouver, BC V6B 5K3, Canada

Full list of author information is available at the end of the article



has been associated with an increased likelihood of experiencing violence [18] and exhibiting a detectable HIV-1 RNA viral load among adult PWID in Vancouver, Canada [19]. In addition, studies of other marginalized populations have demonstrated associations between housing displacement and other mental and physical health concerns including depression, anxiety, suicide, high blood pressure, and poorer self-rated health status [20]. At present, however, we know of no existing studies that have investigated the relationship between residential eviction and syringe sharing. Therefore, we sought to examine whether residential eviction was associated with syringe sharing among a prospective cohort of street-involved youth in Vancouver, Canada.

Methods

This study is based on data from the At-Risk Youth Study (ARYS), an open prospective cohort of street-involved youth in Vancouver, Canada, that began in 2005. The study design of this prospective cohort has been previously described in detail [21]. In brief, recruitment consists of street-based outreach and snowball sampling. Eligibility is restricted to street-involved youth who are age 14–26 years at baseline, have used illicit drugs other than or in addition to marijuana in the past 30 days, and who provide written and informed consent. The term “street-involved” refers to youth who are currently or recently homeless or who recently accessed services for homeless youth. Upon enrolment, and bi-annually thereafter, an interview-administered questionnaire is conducted with participants and includes demographic information, drug use patterns, and practices, as well as engagement in health and social services. A stipend (\$30 CAD) is provided to participants at each study visit. This study has been approved by the University of British Columbia and the Providence Health Care Research Ethics Board.

The study period for the present analysis was restricted to between June 2007 and May 2014, as the measures for residential eviction were only available during this period. To examine the potential relationship between residential eviction and syringe sharing, all analyses were restricted to study observations in which participants reported injection drug use in the last 6 months. The primary outcome of interest for this analysis was syringe sharing, defined as responding affirmatively to the question: “In the last 6 months, have you fixed with a rig that had already been used by someone else?” or “In the last 6 months have you lent your used rig to someone else?” (yes vs. no). The primary explanatory variable of interest was recent residential eviction, defined as responding affirmatively to the question: “Have you been evicted in the last 6 months” (yes vs. no).

To determine whether there was a significant relationship between our outcome of interest and our primary

explanatory variable, we *a priori* selected other secondary explanatory factors we hypothesized might be associated with both residential eviction and syringe sharing. Secondary explanatory factors included: age (per year older), gender (female vs. male), ethnicity (Aboriginal Ancestry vs. other), binge drug use (yes vs. no), daily heroin use (yes vs. no), daily cocaine use (yes vs. no), daily crystal methamphetamine use (yes vs. no), difficulty accessing syringes (yes vs. no), public injecting (yes vs. no), incarceration (yes vs. no), and accessing drug treatment (e.g., pharmacotherapy, residential treatment, and counseling) (yes vs. no). All variables, excluding age, gender, and Aboriginal Ancestry, refer to circumstances and behaviors over the previous 6 months and were treated as time-updated covariates on the basis of semiannual follow-up data.

Initially, we examined the descriptive characteristics, stratified by reports of syringe sharing at the first study visit. Comparisons were made using the Pearson's χ^2 test for binary variables (Fisher's exact test when cell counts were less than or equal to 5) and the Wilcoxon rank-sum test for continuous variables.

Next, we used generalized estimating equation (GEE) analyses with logit link function to assess the independent association between residential eviction and syringe sharing. These methods provide standard errors adjusted by multiple observations per person using an exchangeable working correlation structure [22, 23]. Therefore, data from every participant follow-up visit was considered.

To examine the associations between syringe sharing and each explanatory variable, we first conducted bivariate GEE analyses. To fit the multivariable model, we employed a conservative variable selection approach [24]. Specifically, we initially included the primary explanatory variable and all secondary variables where $p < 0.10$ in bivariate analyses in a multivariable model. We then used a stepwise approach to fit a series of reduced models. After comparing the value of the coefficient associated with the main independent variable of interest (residential eviction) in the full model to the value of the coefficient in each of the reduced models, we dropped the secondary variable associated with the smallest relative change. We continued this iterative process until the minimum change exceeded 5%. Remaining variables were considered confounders in the final multivariable model. P values are all two-sided. Statistical analyses were conducted using SAS software version 9.4 (SAS, Cary, NC).

Results

Among 938 street-involved, 405 (43.2%) reported injection drug use over the study period. Of these, 142 (35.1%) were female and 91 (22.5%) reported being of Aboriginal Ancestry. The median age of the sample was 22.7 years (interquartile range (IQR) = 20.9–24.4). This sample contributed 1131 observations, and the median number of study visits

per participant during the study period that included a report of active injection drug use was 2 (IQR = 1–3). The median follow-up time per participant was 19.2 months (IQR = 6.6–48.3).

In total, 149 (36.8%) youth reported syringe sharing at some point during the study period and a total of 208 (18.4%) observations included a report of syringe sharing. Additionally, 114 (28.1%) unique participants in our sample reported being evicted at least once over the study period. Of these, 86 (75.4%) reported one eviction event, 19 (16.7%) reported two evictions, 7 (6.1%) reported three evictions, and 2 (1.8%) reported four evictions during follow-up. Among the 533 (56.8%) participants excluded from the study, the proportion who experienced residential eviction was not significantly different from those who were included in the study (12.2 vs. 13.5%, $P = 0.304$).

The baseline characteristics, stratified by syringe sharing in the last 6 months, are presented in Table 1. The results of the bivariate and multivariable GEE analyses are presented in Table 2. In bivariate analyses, recent residential eviction (odds ratio (OR) = 1.78; 95% confidence interval (CI): 1.20–2.65) was positively associated with syringe sharing. In the multivariable analyses, recent residential eviction (adjusted odds ratio (AOR) = 1.72, 95% confidence interval (CI): 1.16–2.57) remained significantly and positively associated with syringe sharing after adjusting for Aboriginal Ancestry, which was the only identified confounder.

Discussion

Among this community-recruited cohort of 405 drug-using street-involved youth in Vancouver, Canada, syringe sharing was common with over one-third of study participants

reporting borrowing or lending used syringes during the study period. In multivariable analyses, residential eviction was independently associated with syringe sharing.

Whereas existing literature has demonstrated that eviction may have harmful health-related consequences for drug-using populations [18–20] and that homeless and unstably housed street-involved youth experience higher rates of risky substance use [13, 25], the current study expands on this area of research by demonstrating that experiencing residential eviction is linked with an increased likelihood of syringe sharing. One potential explanation for this finding is that evicted street-involved youth may be displaced to other neighborhoods [26], reducing spatial access to services that provide sterile injecting equipment. Additionally, youth who are displaced due to residential eviction may subsequently experience homelessness, which has been positively associated with syringe sharing [7]. For example, homeless individuals may be more likely to come into contact with police or experience violence, which may result in the loss of sterile injecting equipment and an increased likelihood of sharing syringes [27–29].

Another possible explanation is that evicted street-involved youth may subsequently become homeless or unstably housed [30–32] and, as a result, may be more likely to use drugs in public settings, which has previously been associated with syringe sharing [33]. However, given that the analyses employed herein limit interpretation of temporal relationships, it could also be that street-involved youth who share syringes are more heavily involved in the street-based drug scene or are more likely to experience financial instability, which might increase their vulnerability to residential eviction [32]. Further research is needed to

Table 1 Baseline characteristics of street-involved youth in Vancouver, Canada, stratified by syringe sharing ($n = 405$)

Characteristic	Total (%) ($n = 405$)	Syringe sharing ^a		Odds Ratio (95% CI)	p value
		Yes (%) ($n = 95$)	No (%) ($n = 310$)		
Age (median, IQR)	22.7 (20.9–24.4)	22.2 (20.4–23.7)	22.9 (21.1–24.5)	–	0.055 ^a
Female gender	142 (35.1)	37 (38.9)	105 (33.9)	1.25 (0.77, 2.00)	0.364
Aboriginal Ancestry	91 (22.5)	12 (12.6)	79 (25.5)	0.43 (0.22, 0.83)	0.010
Binge drug use ^b	224 (55.3)	63 (66.3)	161 (51.9)	1.81 (1.12, 2.93)	0.015
Daily heroin use ^b	91 (22.5)	27 (28.4)	64 (20.6)	1.52 (0.90, 2.57)	0.116
Daily cocaine use ^b	9 (2.2)	2 (2.1)	7 (2.3)	0.93 (0.19, 4.54)	1.000 [†]
Daily crystal methamphetamine use ^b	79 (19.5)	19 (20.0)	60 (19.4)	1.04 (0.59, 1.85)	0.890
Residential eviction ^b	67 (16.5)	21 (22.1)	46 (14.8)	1.62 (0.91, 2.90)	0.099
Difficulty accessing syringes ^b	101 (24.9)	28 (29.5)	73 (23.5)	1.37 (0.82, 2.30)	0.225
Public injecting ^b	274 (67.7)	77 (81.1)	197 (63.6)	2.55 (1.44, 4.53)	0.001
Incarceration ^b	93 (23.0)	20 (21.1)	73 (23.6)	0.87 (0.50, 1.51)	0.613
Accessing drug treatment ^{b, c}	171 (42.2)	50 (52.6)	121 (39.0)	1.75 (1.10, 2.78)	0.018

[†] p value is generated from Fisher's Exact Test because of small cell count

^aRefers to continuous variable, p value is generated from Wilcoxon rank-sum test

^bRefers to activities in the last 6 months

^cExcludes detoxification services

Table 2 Bivariate and multivariable generalized estimating equation (GEE) analyses of factors associated with syringe sharing among street-involved youth in Vancouver, Canada, ($n = 405$)

Characteristic	Unadjusted		Adjusted	
	Odds ratio (95% CI)	<i>p</i> value	Odds ratio (95% CI)	<i>p</i> value
Residential eviction ^{a,b}	1.78 (1.20, 2.65)	0.004	1.72 (1.16, 2.57)	0.007
Age (per year older)	0.90 (0.84, 0.96)	0.001		
Female gender ^b	1.09 (0.75, 1.58)	0.656		
Aboriginal Ancestry ^b	0.51 (0.33, 0.80)	0.003	0.50 (0.32, 0.80)	0.003
Binge drug use ^{a,b}	1.58 (1.21, 2.07)	<0.001		
Daily heroin use ^{a,b}	0.93 (0.68, 1.25)	0.619		
Daily cocaine use ^{a,b}	0.89 (0.38, 2.10)	0.793		
Daily crystal methamphetamine use ^{a,b}	0.86 (0.57, 1.30)	0.471		
Difficulty accessing syringes ^{a,b}	2.11 (1.51, 2.94)	<0.001		
Public injecting ^{a,b}	2.88 (1.95, 4.25)	<0.001		
Incarceration ^{a,b}	1.26 (0.88, 1.80)	0.214		
Accessing drug treatment ^{a,b,c}	1.09 (0.80, 1.48)	0.589		

^aRefers to activities in the last 6 months^bComparison is yes vs. no^cExcludes detoxification services

better understand the mechanisms underlying the observed association between residential eviction and syringe sharing.

While accessing housing remains imperative to reducing risky substance use practices among youth, the findings of the present study suggest that maintaining housing stability may also play an important role in reducing such practices. Previous research among street-involved youth in this setting indicate that financial instability is among the most commonly reported reasons for eviction [32]. A recent study also found that 52% of street-involved youth in Vancouver reported being unable to access housing and housing referral services [34]. The current study findings extend this work and underscore the importance of ensuring that youth receive necessary social and financial supports to access and maintain housing in order to prevent eviction and reduce risky substance use behaviors.

Given that over one-third of study participants reported syringe sharing at some point during the study period, it is evident that interventions that reduce syringe sharing among street-involved youth are needed, particularly among those experiencing residential eviction. Among adult populations with concurrent disorders, existing literature has pointed to the effectiveness of providing housing supports. Examples of housing supports include access to case management supports, home support services, crisis intervention services, and health care services [35, 36]. The provision of housing supports have been associated with significant reductions in substance use and risky practices [34, 37–39], as well as improved housing stability among street-involved populations [40–42].

While providing housing supports has realized significant successes among adult populations and has been integrated into youth homelessness strategies in the USA, Australia, and the UK, Canada has been slow to implement such a model for street-involved youth [43]. Therefore, expanding housing options and supports for youth may have the potential to reduce experiences of eviction and related risky substance use practices among this population. Particular emphasis should be placed on ensuring that housing options espouse a continuum of care that ranges from low-threshold to abstinence-based housing in order to meet diverse range of housing needs facing marginalized populations of youth, particularly those who inject drugs.

This study has a number of limitations. First, because self-reported responses were solicited for this survey, results are subject to response biases. Existing literature, however, suggests that self-reported responses among street-involved populations are generally reflective of actual behaviors [44]. Nonetheless, we suspect that socially desirable responses would likely lead to an underreporting of high-risk practices thereby potentially leading to conservative estimates of syringe sharing. Second, the ARYS cohort is a non-randomized, community-recruited sample of street-involved youth in Vancouver and therefore may not be generalizable to other populations. Third, given the observational nature of the study, there may be unmeasured confounding. In addition, the methods employed herein did not allow us to determine the temporality of the association between residential eviction and syringe sharing.

Conclusions

In sum, this study found that youth who recently experienced residential eviction had a significantly higher likelihood of sharing syringes. These findings suggest that policy approaches aiming to provide youth with access to a continuum of housing supports may have significant potential to mitigate residential eviction and high-risk substance use among street-involved youth.

Abbreviations

ARYS: At-Risk Youth Study; PWID: People who inject drugs

Acknowledgements

The authors thank the study participants for their participation in this study, as well as both current and past study staff and researchers. We extend our appreciation to Cody Callon, Ana Prado, Deborah Graham, Peter Vann, Steve Kain, and Tricia Collingham for their administrative and research assistance.

Funding

The authors thank the study participants for their contribution to the research, as well as current and past researchers and staff. The study was supported by the US National Institutes of Health (U01DA038886). Dr. Kora DeBeck is supported by a MSFHR/St. Paul's Hospital Foundation—Providence Health Care Career Scholar Award and a Canadian Institutes of Health Research New Investigator Award. Mary Clare Kennedy is supported by a Mitacs Accelerate Award from Mitacs Canada.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

AP, KD, and TK conceptualized the study design. HD performed the statistical analyses, and AP and KD interpreted the results. AP drafted the initial manuscript. KD, TK, RM, and MCK provided substantial revisions of the manuscript and guidance throughout the writing. All authors have read and approved the final manuscript.

Competing interests

All authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The At-Risk Youth Study has been approved by the Providence Health Care/University of British Columbia's Ethics Board. All participants provided informed consent prior to participating in this study.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹BC Centre on Substance Use, British Columbia Centre for Excellence in HIV/AIDS, 608-1081 Burrard Street, Vancouver, BC V6Z 1Y6, Canada.

²Interdisciplinary Studies Graduate Program, University of British Columbia, 270-2357 Main Mall, Vancouver, BC V6T 1Z4, Canada. ³School of Population and Public Health, University of British Columbia, 2206 East Mall, Vancouver, BC V6T 1Z3, Canada. ⁴Division of AIDS, Department of Medicine, University of British Columbia, 667-1081 Burrard Street, Vancouver, BC V6Z 1Y6, Canada.

⁵School of Public Policy, Simon Fraser University, 3277-515 Hastings W Street, Vancouver, BC V6B 5K3, Canada.

Received: 16 February 2017 Accepted: 1 May 2017

Published online: 12 May 2017

References

- Cunningham EB, Jacka B, DeBeck K, Applegate TL, Harrigan PR, Krajden M, Marshall BDL, Montaner J, Lima VD, Olmstead AD, et al. Methamphetamine injecting is associated with phylogenetic clustering of hepatitis C virus infection among street-involved youth in Vancouver, Canada. *Drug Alcohol Depend.* 2015;152:272.
- Marshall BDL, Kerr T, Qi J, Montaner JSG, Wood E. Public injecting and HIV risk behaviour among street-involved youth. *Drug Alcohol Depend.* 2010;110:254–8.
- Spijkerman IJ, van Ameijden EJ, Mientjes GH, Coutinho RA, van den Hoek A. Human immunodeficiency virus infection and other risk factors for skin abscesses and endocarditis among injection drug users. *J Clin Epidemiol.* 1996;49:1149–54.
- Mahanta J, Borkakoty B, Das HK, Chelleng PK. The risk of HIV and HCV infections among injection drug users in northeast India. *AIDS Care.* 2009; 21:1420–4.
- UNAIDS. Report on the global AIDS epidemic. Geneva: UNAIDS; 2006. 2006.
- Wood E, Tyndall MW, Spittal PM, Li K, Hogg RS, Montaner JS, O'shaughnessy MV, Schechter MT. Factors associated with persistent high-risk syringe sharing in the presence of an established needle exchange programme. *Aids.* 2002;16:941–3.
- Bozinoff N, Wood E, Dong H, Richardson L, Kerr T, DeBeck K. Syringe sharing among a prospective cohort of street-involved youth: implications for needle distribution programs. *AIDS Behav.* 2017;1–9.
- Stoltz J-A, Wood E, Small W, Li K, Tyndall M, Montaner J, Kerr T. Changes in injecting practices associated with the use of a medically supervised safer injection facility. *J Public Health.* 2007;29:35–9.
- Kerr T, Small W, Buchner C, Zhang R, Li K, Montaner J, Wood E. Syringe sharing and HIV incidence among injection drug users and increased access to sterile syringes. *Am J Public Health.* 2010;100:1449–53.
- Kerr T, Tyndall M, Li K, Montaner J, Wood E. Safer injection facility use and syringe sharing in injection drug users. *Lancet.* 2005;366:316–8.
- Lloyd-Smith E, Kerr T, Zhang R, Montaner JS, Wood E. High prevalence of syringe sharing among street involved youth. *Addict Res Theory.* 2008;16:353–8.
- Marshall BDL, Shoveller JA, Wood E, Patterson TL, Kerr T. Difficulty accessing syringes mediates the relationship between methamphetamine use and syringe sharing among young injection drug users. *AIDS Behav.* 2011;15: 1546–53.
- Cheng T, Wood E, Nguyen P, Kerr T, DeBeck K. Increases and decreases in drug use attributed to housing status among street-involved youth in a Canadian setting. *Harm Reduct J.* 2014;11:12.
- Johnson G, Chamberlain C. Homelessness and substance abuse: which comes first? *Aust Soc Work.* 2008;61:342–56.
- Feng C, DeBeck K, Kerr T, Mathias S, Montaner J, Wood E. Homelessness independently predicts injection drug use initiation among street-involved youth in a Canadian setting. *J Adolesc Health.* 2013;52:499–501.
- Roy É, Haley N, Leclerc MP, Cédras ML, Blais L, Boivin J-F. Drug injection among street youths in Montreal: predictors of initiation. *J urban health.* 2003;80:92–105.
- Des Jarlais DC, Braine N, Friedmann P. Unstable housing as a factor for increased injection risk behavior at US syringe exchange programs. *AIDS Behav.* 2007;11:78–84.
- Kennedy M, McNeil R, Milloy M, Dong H, Kerr T, Hayashi K. Residential eviction and exposure to violence among people who inject drugs in Vancouver, Canada. *Int J Drug Policy.* 2017;41:59.
- Kennedy M, Kerr T, McNeil R, Parashar S, Montaner J, Wood E, Milloy M. Residential eviction and risk of detectable plasma HIV-1 RNA viral load among HIV-positive people who use drugs. *AIDS Behav.* 2016;1–10.
- Vásquez-Vera H, Palència L, Magna I, Mena C, Neira J, Borrell C. The threat of home eviction and its effects on health through the equity lens: a systematic review. *Soc Sci Med.* 2017;175:199–208.
- Wood E, Stoltz J-A, Montaner JS, Kerr T. Evaluating methamphetamine use and risks of injection initiation among street youth: the ARYS study. *Harm Reduct J.* 2006;3:1.
- Ballinger GA. Using generalized estimating equations for longitudinal data analysis. *Organ Res Methods.* 2004;7:127–50.
- Hanley JA, Negassa A, Forrester JE. Statistical analysis of correlated data using generalized estimating equations: an orientation. *Am J Epidemiol.* 2003;157:364–75.

24. Maldonado G, Greenland S. Simulation study of confounder-selection strategies. *Am J Epidemiol*. 1993;138:923–36.
25. Marshall BDL, Kerr T, Shoveller JA, Patterson TL, Buxton JA, Wood E. Homelessness and unstable housing associated with an increased risk of HIV and STI transmission among street-involved youth. *Health Place*. 2009;15:753–60.
26. Desmond M, Shollenberger T. Forced displacement from rental housing: prevalence and neighborhood consequences. *Demography*. 2015;52:1751–72.
27. Werb D, Wood E, Small W, Strathdee S, Li K, Montaner J, Kerr T. Effects of police confiscation of illicit drugs and syringes among injection drug users in Vancouver. *Int J Drug Policy*. 2008;19:332–8.
28. Aitken C, Moore D, Higgs P, Kelsall J, Kerger M. The impact of a police crackdown on a street drug scene: evidence from the street. *Int J Drug Policy*. 2002;13:193–202.
29. Small W, Kerr T, Charette J, Schechter MT, Spittal PM. Impacts of intensified police activity on injection drug users: evidence from an ethnographic investigation. *Int J Drug Policy*. 2006;17:85–95.
30. Kleysteuber R. Tenant screening thirty years later: a statutory proposal to protect public records. *Yale Law J*. 2007;116:1344–88.
31. Desmond M, Gershenson C, Kiviat B. Forced relocation and residential instability among urban renters. *Soc Serv Rev*. 2015;89:227–62.
32. Zivanovic R, Omura J, Wood E, Nguyen P, Kerr T, DeBeck K. Eviction and loss of income assistance among street-involved youth in Canada. *J Public Health Policy*. 2016;37:244–59.
33. Montaner JS, Kerr T. Factors associated with syringe sharing among users of a medically supervised safer injecting facility. 2005.
34. Barker B, Kerr T, Nguyen P, Wood E, DeBeck K. Barriers to health and social services for street-involved youth in a Canadian setting. *J Public Health Policy*. 2015;36:350–63.
35. Kidd SA, Frederick T, Karabanow J, Hughes J, Naylor T, Barbic S. A mixed methods study of recently homeless youth efforts to sustain housing and stability. *Child Adolesc Soc Work J*. 2016;33:207–18.
36. Tabol C, Drebing C, Rosenheck R. Studies of “supported” and “supportive” housing: a comprehensive review of model descriptions and measurement. *Eval Program Plann*. 2010;33:446–56.
37. Atherton I, Nicholls CM. ‘Housing First’ as a means of addressing multiple needs and homelessness. 2008.
38. Goering P, Veldhuizen S, Watson A, Adair C, Kopp B, Latimer E, Aubry T. National at home/Chez Soi final report. Calgary: Mental Health Commission of Canada; 2014. p. 5.
39. Padgett DK, Stanhope V, Henwood BF, Stefancic A. Substance use outcomes among homeless clients with serious mental illness: comparing housing first with treatment first programs. *Community Ment Health J*. 2011;47:227–32.
40. Stefancic A, Tsemberis S. Housing First for long-term shelter dwellers with psychiatric disabilities in a suburban county: a four-year study of housing access and retention. *J Prim Prev*. 2007;28:265–79.
41. Pearson C, Montgomery AE, Locke G. Housing stability among homeless individuals with serious mental illness participating in housing first programs. *J Community Psychol*. 2009;37:404–17.
42. Kertesz SG, Crouch K, Milby JB, Cusimano RE, Schumacher JE. Housing first for homeless persons with active addiction: are we overreaching? *Milbank Q*. 2009;87:495–534.
43. Gaetz S. Coming of age: reimagining the response to youth homelessness in Canada; Toronto: Canadian Homelessness Research Network. 2014.
44. Brener ND, Billy JO, Grady WR. Assessment of factors affecting the validity of self-reported health-risk behavior among adolescents: evidence from the scientific literature. *J Adolesc Health*. 2003;33:436–57.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

