

# A comparative study of the effect of some nutritional medicinal plants effect on lead accumulation in the liver following different modes of administration

Chukwuemeka Nwokocha, Novie Younger-Coleman<sup>1</sup>, Magdalene Nwokocha<sup>2</sup>, Daniel Owu<sup>3</sup>, Moses Iwuala<sup>4</sup>

Department of Basic Medical Sciences, <sup>1</sup>Tropical Medicine Research Institute, <sup>2</sup>Department of Haematology, The University of The West Indies Mona Campus, Kingston Jamaica, <sup>3</sup>Department of Physiology, University of Calabar, Calabar, <sup>4</sup>Federal University of Technology, Owerri, Nigeria

Submitted: 06-01-2014

Revised: 11-03-2014

Published: 06-08-2014

## ABSTRACT

**Context and Objectives:** Lead (Pb) toxicity leads to cell damage in many organs of the body. Using different treatment interventions and modes of administration we comparatively examined the protective ability of some medicinal plants on liver Pb accumulation. **Materials and Methods:** Rats were fed on either 7% w/w *Zingiber officinale*, 7% w/w *Allium sativum*, 10% w/w *Lycopersicon esculentum*, 5%, w/w *Garcinia kola* (all in rat chow), while Pb (100 ppm) was given in drinking water. The additives were administered together with (mode 1), a week after exposure to (mode 2) or a week before metal exposure to (mode 3) the metal for a period of 6 weeks. The metal accumulations in the liver were determined using atomic absorption spectrometry and compared using analysis of variance. **Results:** Some additives significantly ( $P < 0.05$ ) reduced, while others enhanced Pb accumulation. Mode 2 yielded the highest mean % protection and mode 3 the lowest, no significant interaction between modes of administration and time of measurement in their relationships to percentage protection, but there was statistically significant ( $P < 0.05$ ) interaction between modes of administration and additive used in their relationships to percentage protection. **Conclusion:** Protective effects of medicinal plants are varied and depend on the nature of lead exposure.

**Key words:** Accumulation pattern, *Allium sativum*, *Garcinia kola*, lead, liver, *Lycopersicon esculentum*, mode of administration, *Zingiber officinale*

## INTRODUCTION

Lead (Pb) has been shown to have a multisystem effect seriously affecting the nervous, circulatory, skeletal, renal, hematopoietic, and endocrine systems.<sup>[1]</sup> Long-term exposure to Pb may also result in a nephropathy or renal adenocarcinoma, neuromuscular weakness,<sup>[2,3]</sup> behavioral, and neurochemical alterations.<sup>[3-5]</sup> These can occur through a derangement of the antioxidant enzymes and neurotransmitter balance.<sup>[4,5]</sup> Pb toxicity is reported to occur through its affinity for proteins and its capacity to simulate calcium and iron (Fe) channels.<sup>[6]</sup> The increase in intracellular calcium together with the increase in reactive oxygen species, which is reported to occur with increase in lead exposure,

can trigger apoptosis through the cytochrome C release and a fall in the mitochondrial potential.<sup>[6,7]</sup>

The use of chelators are usually the primary cause of management with metal poisoning and these chelators can shield the metal from the biological targets by way of mobilization and excretion,<sup>[8]</sup> but they are not always useful in reversing the damage done by exposures to these toxicants.<sup>[6]</sup> Reports on the increasing role and usefulness of medicinal plant products in maintenance and optimization of health have led to research on the efficacy of these medicinal plants/nutrient products' effects on the amelioration of the metal-induced toxicity.<sup>[9-13]</sup> These plant products are reported to affect the bioavailability, transport, and toxicodynamics of this metal in the tissues.

*Allium sativum*'s (garlic) metal chelating properties,<sup>[9,14]</sup> *Zingiber officinale*'s (ginger) radical scavenging properties,<sup>[11,15]</sup> *Garcinia kola* characteristics to act as chelators of divalent

### Access this article online

#### Website:

www.phcogres.com

DOI: 10.4103/0974-8490.138278

#### Quick Response Code:



### Address for correspondence:

Dr. Chukwuemeka R. Nwokocha, Department of Basic Medical Sciences, University of the West Indies, Mona Campus, Mona, Kingston 7, Jamaica.  
E-mail: chukwuemeka.nwokocha@uwimona.edu.jm

metal,<sup>[9,16,17]</sup> and Tomato's, (*Lycopersicon esculentum*) abilities to synthesize metal chelating proteins, metallothioneins, peptides, phytochelatin<sup>[12,18,19]</sup> have all been reported and documented. These medicinal plant products are also reported for their prophylactic and curative uses, which is related to their antioxidant properties.<sup>[9,20,21]</sup>

We have recently shown that *G. kola*,<sup>[9]</sup> *A. sativum*,<sup>[10]</sup> *Z. officinale*<sup>[11]</sup> and *L. esculentum*<sup>[12]</sup> reduced hepatotoxic effect of lead through a hepato-protective role. This study is to comparatively study the effects of these nutrients and medicinal plants on lead accumulation in the liver of rats following different treatment interventions. The study also aimed to determine whether the modes of administration yielded different results and whether these differences varied with the additive used or with the time of measurement.

## MATERIALS AND METHODS

### Animals and experimental design

Male Wistar rats of about 7 weeks old weighing 150-180 g were obtained from the animal house of the Faculty and used for the study. The animals were kept at constant room temperature with 12-h light/dark cycles. All animals were fed with normal rat chow and had access to tap water *ad libitum* during the period of acclimatization. We sought and received ethical approval from the Federal University of Technology Faculty Ethical Committee for this study; and "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable.

### Preparation of nutrient substance and heavy metals

Fresh ginger rhizome, Garlic, *G. kola* and Tomato were purchased from the market at Okija, Nigeria, from January to April (mainly during the dry season). Professor C. Ufearo of the Nnamdi Azikiwe University Nigeria did the authentication. Materials were ground with a kitchen blender and sieved using a very fine sieve (particulate size of 250 µm). Each additive –7% w/w of *Z. officinale*, 7% w/w of *A. sativum*, 10% w/w of *L. esculentum*, 5% w/w of *G. kola* – was mixed with rat chow and fed to different groups of animals. Each group received directly from the drinking water bottle tap water that contained 100 ppm lead acetate. The lead was prepared from Lead acetate. Ten millilitre of 1000 ppm lead was further diluted with 90 ml of distilled water. Lead estimation was calculated from the calibration curves. These concentrations were arrived at following our earlier reported studies<sup>[9-12]</sup> with a calibration curve, which was prepared for the estimation of the sample and control materials.

### Experimental protocol

Group 1 was fed with normal rat chow and lead (Pb = 100 ppm,) only. Group 2 was fed with rat chow and one of the nutritional medicinal plants (7% w/w of *Z. officinale*, 7% w/w of Garlic, 10% w/w of *L. esculentum*, 5%, w/w of *G. kola*) mixed with rat chow along with the lead acetate water, the exposure to lead and feeding starting same time. Group 3 was fed with normal rat chow and water mixed with Pb (100 ppm), for the 1<sup>st</sup> week and then with rat chow mixed with one of the additives and tap water without lead from the 2<sup>nd</sup> to the 6<sup>th</sup> week. Group 4 was fed with rat chow mixed with one of the additives for 1 week, and after that they were fed with normal rat chow and Pb (100 ppm) in drinking water for the remaining 5 weeks. The grouping and feeding patterns are summarized in Table 1. All administrations were through the oral route.

### Tissue preparation

At the end of the experimental period, the rats were sacrificed under chloroform anesthesia. Liver (1 g) was excised and transferred in polypropylene vials for analysis. Before acid digestion, a porcelain mortar was employed to grind and homogenize the dry tissue samples in 5 ml of normal saline. After digestion of all samples the concentrations of Pb was analyzed using flame atomic absorption spectrophotometer (AAS) (Perkin Elmer A.A. 3030 Waltham, MA USA) with D2 background correction device. Cold vapor technique was used for the analysis of Pb (Kingston and Jessie, 1988; Medham, 2000). Lead was estimated using the AAS at 283 nm wavelength.

### Statistical analysis

The percentage protection provided by the presence of each additive was obtained using the formula:

$$\frac{Pb_{1j} - Pb_{ij}}{Pb_{1j}} \times 100 \quad (1)$$

Where,

$Pb_{1j}$  is the concentration of accumulated lead in the liver sample when exposed to lead alone (treatment Group 1)

**Table 1: Summary of specimen grouping and 6 weeks feeding pattern**

Week	Group 1	Group 2	Group 3	Group 4
1	F+WPb	FGa+WPb	F+WPb	FGa+W
2	F+WPb	FGa+WPb	FGa+W	F+WPb
3	F+WPb	FGa+WPb	FGa+W	F+WPb
4	F+WPb	FGa+WPb	FGa+W	F+WPb
5	F+WPb	FGa+WPb	FGa+W	F+WPb
6	F+WPb	FGa+WPb	FGa+W	F+WPb

F=Feed (rat chow); W=Water; FGa=Feed-nutrient concentrate; WPb=Lead in water (Pb=100 ppm)

at time  $j$ , ( $j = 2, 4, 6$ , weeks).  $Pb_{ij}$  is the mean value for accumulated lead concentration in the liver obtained from the sample of five rats sacrificed at each time point.


$Pb_j$  is the concentration of accumulated lead in the liver sample when exposed to lead in the treatment group  $i$ , ( $i = 2, 3, 4$ ) at time  $j$ , ( $j = 2, 4, 6$ , weeks).  $Pb_{ij}$  is the mean value for accumulated lead concentration in the liver obtained from the sample of five rats fed a given additive using a particular mode of administration at time points 2, 4 and 6, respectively.

Means and standard errors for the percentage protection were obtained for the various additives, modes of treatment and times of administration. Analysis of variance (ANOVA) with Bonferonni's posttest analysis was performed to determine whether the means differed with respect to each of the aforementioned sources of variation - additives, modes of treatment and times of administration. In addition the ANOVA was used to examine whether the relationship of each of additive and time of measurement with percentage protection differed with respect to mode of treatment. The non-parametric test for trend was also used to examine whether the percentage protection increased or decreased with time of administration. Stata version 12.0 was used to carry out data analysis and Origin™ 5.0 (Microcal Software Inc., Northampton, USA) created graphical displays of the data.  $P < 0.05$  was considered as statistically significant.


## RESULTS

### Effect of various additives and lead co-administration on accumulation in the liver


Figures 1-3, each show that the concentration of lead accumulated in the liver tissue over time for each of the additives and treatment groups (modes of administration). Excepting for tomato given using mode 2, the concentration of lead generally fell over time after the 2<sup>nd</sup> week. The additives used all significantly ( $P < 0.05$ ) reduced the accumulation of lead in the first 2 weeks of the study for all modes of administration except for garlic in mode 3. This pattern was observed at weeks 4 and 6, but not at week 2. In addition, confidence intervals (CIs) for mode 3 at weeks 4 and 6 and mode 1 at week 4 indicated that the means for these modes at these times did not differ significantly from 0, due to the fact that, in the presence of lead some of the additives gave little or no protection. As one of the study objectives was to determine whether there was interaction between mode of administration and time, ANOVA was also used to examine whether the nature of the variation in means for mode of administration was the same at each of the three measurement times -2, 4 and 6 weeks.



**Figure 1:** Effects of the various nutrient substances on lead accumulation in the liver of rats when animals were exposed to lead and nutrients at same time



**Figure 2:** Effects of the various nutrient substances on lead accumulation in the liver of rats when nutrients were used after a week's exposure to lead



**Figure 3:** Effects of the various nutrient substances on lead accumulation in the liver of rats when animals were exposed to lead after a week's exposure to the nutrients

### Comparative analysis of percentage protection to lead

Table 2a gives the mean percentage accumulation for the samples of rats sacrificed at the different time periods within groups defined by additive and mode of administration. For ginger, the level of protection tended to fall to zero as time passed and for tomato, an actual increase

in the lead accumulation, leading to negative percentage protection was observed.

Analysis of variance was used to determine whether the nature of the variation in means for mode of administration was the same for each additive. There was statistically significant interaction between modes of administration and additive used, in their relationships to percentage protection. This means that the differences between the modes were not the same for all additives. The adjusted mean values, produced by the ANOVA, for the percentage protection for groups defined by additive and mode combinations revealed that the percentage protection offered by mode 3 in the presence of garlic was actually negative – indicative of increasing lead levels relative to the amounts when lead is given alone - and significantly different from the percentage protection obtained for the other modes in the presence of garlic [Tables 1 and 2]. Results in Table 2b further showed

that the differences between the modes for the other additives were not significantly different.

## DISCUSSION

Increasing drug toxicity due to metabolic activation and other unwanted side-effects of synthetic drugs has led to the increasing global demands and use of herbal remedies in the management and treatment of various ailments and to promote health.<sup>[9-12]</sup> Our results showed that the additives used affected the accumulation levels of lead in the liver. Some of them significantly reduced the accumulation, while others had no effect or, interestingly, increased its accumulation. The additives used all significantly reduced the accumulation of lead in the first 2 weeks of the study for all modes of administration except for garlic in mode 3. Though nutrients and nutritional status<sup>[9,22]</sup> affect the rate of accumulation and or excretion of metals, the observed decrease in the lead concentrations seen across all modes is not due to any analytical error. Jezierska and Witeska,<sup>[23]</sup> had reported that accumulation is a function of uptake and elimination, the unexpected patterns of metal accumulation seen here has been reported by others. We also observed Pb accumulation increased with initial exposure but decreased with further exposure in our studies, this is not due to any experimental or analytical error as such patterns have been reported by others.<sup>[24,25,29]</sup>

Many reports exist in literature about the efficacy of medicinal plant products in ameliorating or reducing the toxicity and accumulation of lead in tissues<sup>[9,14]</sup> due to their metal chelating, antioxidant and scavenging properties,<sup>[18,19]</sup> and have also highlighted that these properties are not

**Table 2a: Summary statistics (sample means) of percentage protection by the nutritional medicinal plants, time and mode of administration**

	Garlic	Ginger	Tomato	Garcina kola
Group 2				
Week 2	54.1625	56.25	97.5	33.75
Week 4	0	0	-33.3	44
Week 6	44.1	0	48.9	47.9
Group 3				
Week 2	12.4938	65.625	29.1875	87.5
Week 4	33.3	0	20	33.3
Week 6	80	0	20	72
Group 4				
Week 2	0	96.875	97.9188	56.25
Week 4	80	50	43.2	0
Week 6	33.9	0	66.1	0

**Table 2b: Means and 95% CIs (in brackets) for percentage protection for the different modes, additives and times**

Factor	Mode 1		Mode 2		Mode 3		Total	
	n	Mean (95% CI)	n	Mean (95% CI)	n	Mean (95% CI)	n	Mean (95% CI)
Additive								
Garlic								
Unadjusted	3	32.8 (-1.0-66.5)	3	41.9 (1.4-82.5)	3	-55.4 (-115.0-4.2)	9	6.4 (-32.5-45.4)
Adjusted		32.8 (3.3-62.2)		41.9 (12.5-71.4)		-55.4 (-84.8-25.9)**a		
Ginger								
Unadjusted	3	18.8 (-19.3-56.8)	3	21.9 (-22.5-66.3)	3	49.0 (-7.8-105.7)	9	29.9 (4.4-55.3)
Adjusted		18.8 (-10.7-48.2)		21.9 (-7.6-51.3)		49.0 (19.5-78.4)		
Garcinia Kola								
Unadjusted	3	41.9 (33.3-50.4)	3	64.3 (31.5-97.0)	3	18.8 (-19.3-56.9)	9	41.6 (21.8-61.4)
Adjusted		41.9 (12.4-71.4)		64.3 (34.8-93.7)		18.7 (-10.7-48.2)		
Tomato								
Unadjusted	3	37.7 (-39.8-115.2)	3	23.1 (16.8-29.3)	3	69.1 (36.9-101.3)	9	43.3 (15.4-71.2)
Adjusted		37.7 (8.2-67.2)		23.1 (-6.4-52.5)		69.1 (39.6-98.5)		
Time								
2 weeks	4	60.4 (33.3-87.5)	4	48.7 (14.1-83.3)	4	62.8 (16.0-109.6)	12	57.3 (37.6-77.0)
4 weeks	4	2.7 (-29.5-34.9)	4	21.7 (5.7-37.6)	4	-1.7 (-71.9-68.5)	12	7.5 (-17.0-32.1)*b
6 weeks	4	35.2 (11.3-59.2)	4	43.0 (3.3-82.7)	4	0.0 (-54.8-54.8)	12	26.1 (1.6-50.6)
Total	12	32.8 (12.2-53.3)	12	37.8 (19.7-55.8)	12	20.4 (-15.1-55.8)		

\*P<0.05, \*\*P<0.01, \*\*\*P<0.001, <sup>a</sup>Significant difference between mode 3 and mode 1 for garlic. <sup>b</sup>Significant difference between week 4 and week 2, means. CI=Confidence interval

same for all nutrient substances.<sup>[26,27]</sup> Enhancement of the antioxidant capacity of the liver, reduction of hepatocyte injury and lipid peroxidation, improvement of barrier functions and antioxidant activity, decrease oxidative DNA damage in the liver and increased hepatic detoxification and bile production are some of the ways by which nutrient substances can offer a hepatoprotective ability. Our result shows that these products gave protection in varying degrees. It was observed that ginger gave less protection as the weeks progressed in all modes of treatment, while tomato was associated with an increase in accumulation at week 4 using mode 2 when compared with the control values.

The mode of administration of the additive substances and metals was designed to give an understanding as to how these plant products/additives affect the bioavailability of lead.<sup>[30]</sup> Administration of the lead and the additive at the same time, as per mode 1, would provide indication of whether the additives prevented the absorption of lead. Administration of lead during the 1<sup>st</sup> week only, as per mode 2, followed by administration of the additive starting from the 2<sup>nd</sup> week until the end of the study would indicate whether the additive enhanced excretion of the accumulated metal, while administration of the additive in the 1<sup>st</sup> week only, as per mode 3, followed by administration of lead from week 2 onwards would indicate whether the additives prevented the absorption and accumulation of the metal. The summary statistics shown indicated that mode 2 yielded the highest mean percentage protection and mode 3 the lowest; this showed that the additives used in the study were effective mainly through reducing the absorption of Pb in the liver. Results revealed there was no statistically significant interaction between modes of administration and time of measurement in their relationships to percentage protection. The frequency and mode of administration is reported to affect efficacy of treatment.<sup>[28]</sup>

The trend of increased mean percentage protection for modes 1 and 2 relative to mode 3 was observed in the presence of garlic and *G. kola* only. In the presence of lead some of the combinations of modes and additives can be expected to give little or no protection as the CIs produced from these data were relatively wide and included zero for some of these estimates. Our study showed there was statistically significant interaction between modes of administration and additive used with percentage protection being negative for garlic used according to mode 3 and significantly different from the percentage protection obtained when garlic is used based on mode 1. This difference between modes was not observed for the other additives.

Medicinal plant products affect the toxic dynamics of metals in tissues at different degrees, some by decreasing

accumulation and uptake of these metals and also enhancing their elimination from the tissues. Medicinal plant products are not always useful in the reduction of metal accumulation and toxicity; their use can also be harmful, and the identification of the roles played by these substances becomes more important if they are to be used in this regard. More studies are needed to understand these interactions and efficacy of these substances together with the best modes of administration that will significantly impact on their usefulness in the management of metal exposures and toxicity.

## REFERENCES

1. Flora SJ, Flora G, Saxena G. Environmental occurrence, health effects and management of lead poisoning. In: Cascas SB, Sordo J, editors. Lead Chemistry, Analytical Aspects, Environmental Impacts and Health Effects. Netherlands: Elsevier Publication; 2006. p. 158-228.
2. Damek-Poprawa M, Sawicka-Kapusta K. Histopathological changes in the liver, kidneys, and testes of bank voles environmentally exposed to heavy metal emissions from the steelworks and zinc smelter in Poland. Environ Res 2004;96:72-8.
3. Flora SJ, Saxena G, Gautam P, Kaur P, Gill KD. Response of lead-induced oxidative stress and alterations in biogenic amines in different rat brain regions to combined administration of DMSA and MiADMSA. Chem Biol Interact 2007;170:209-20.
4. Brennan PA, Kendrick KM, Keever EB. Neurotransmitter release in the accessory olfactory bulb during and after the formation of an olfactory memory in mice. Neuroscience 1995;69:1075-86.
5. Moreira EG, Vassilief I, Vassilief VS. Developmental lead exposure: Behavioral alterations in the short and long term. Neurotoxicol Teratol 2001;23:489-95.
6. Flora SJ, Saxena G, Mehta A. Reversal of lead-induced neuronal apoptosis by chelation treatment in rats: role of reactive oxygen species and intracellular Ca (2+). J Pharmacol Exp Ther 2007;322:108-16.
7. Xu J, Ji LD, Xu LH. Lead-induced apoptosis in PC 12 cells: involvement of p53, Bcl-2 family and caspase-3. Toxicol Lett 2006;166:160-7.
8. Andersen O. Principles and recent developments in chelation treatment of metal intoxication. Chem Rev 1999;99:2683-710.
9. Nwokocha CR, Owu DU, Ufearo CS, Iwuala MO. Comparative study on the efficacy of *Garcinia kola* in reducing some heavy metal accumulation in liver of Wistar rats. J Ethnopharmacol 2011;135:488-91.
10. Nwokocha CR, Owu DU, Nwokocha MI, Ufearo CS, Iwuala MO. Comparative study on the efficacy of *Allium sativum* (garlic) in reducing some heavy metal accumulation in liver of wistar rats. Food Chem Toxicol 2012;50:222-6.
11. Nwokocha CR, Owu DU, Nwokocha MI, Ufearo CS, Iwuala MO. Comparative study on the hepatoprotection to heavy metals of *Zingiber officinale*. Pharmacognosy Res 2012;4:208-13.
12. Nwokocha CR, Nwokocha MI, Aneto I, Obi J, Udekweleze DC, Olatunde B, *et al.* Comparative analysis on the effect of *Lycopersicon esculentum* (tomato) in reducing cadmium, mercury and lead accumulation in liver. Food Chem Toxicol 2012;50:2070-3.
13. Nwokocha CR, Nwokocha MI, Owu DU, Obi J, Olatunde B, Ebe C, *et al.* Comparative analysis on the effect of palm oil (*Elaeis*

- guineensis*) in reducing cadmium and lead accumulation in liver of Wistar rats. *Pharmacognosy Res* 2012;4:214-8.
14. Obioha UE, Suru SM, Ola-Mudathir KF, Faremi TY. Hepatoprotective potentials of onion and garlic extracts on cadmium-induced oxidative damage in rats. *Biol Trace Elem Res* 2009;129:143-56.
  15. Sultana S, Ripa FA, Hamid K. Comparative antioxidant activity study of some commonly used spices in Bangladesh. *Pak J Biol Sci* 2010;13:340-3.
  16. Monago CC, Akhidue V. Estimation of tannin, saponin, oxalate, cyanogenic and cardiac glycosides in *Garcinia kola*. *J Appl Sci Environ Manage* 2002;6:22-5.
  17. Nwokocha C, Ejebe DE, Nwangwa EK, Ekene N, Akonoghre R, Ukwu J. The effects of bitter kola supplemented diet on hepatotoxicity of mercury in Wistar rats. *J Appl Sci Environ Manage* 2010;14:89-95.
  18. Scheller HV, Huang B, Hatch E, Goldsbrough PB. Phytochelatin synthesis and glutathione levels in response to heavy metals in tomato cells. *Plant Physiol* 1987;85:1031-5.
  19. Tito A, Carola A, Bimonte M, Barbulova A, Arciello S, de Laurentis F, *et al.* A tomato stem cell extract, containing antioxidant compounds and metal chelating factors, protects skin cells from heavy metal-induced damages. *Int J Cosmet Sci* 2011;33:543-52.
  20. Shi J, Le Maguer M. Lycopene in tomatoes: Chemical and physical properties affected by food processing. *Crit Rev Food Sci Nutr* 2000;40:1-42.
  21. Nwokocha CR, Ozolua RI, Owu DU, Nwokocha MI, Ugwu AC. Antihypertensive properties of *Allium sativum* (garlic) on normotensive and two kidney one clip hypertensive rats. *Niger J Physiol Sci* 2011;26:213-8.
  22. Zalups RK, Ahmad S. Molecular handling of cadmium in transporting epithelia. *Toxicol Appl Pharmacol* 2003;186:163-88.
  23. Jezierska B, Witeska M. The metal uptake and accumulation in fish living in polluted waters. In: Twardowska I, Allen HE, Häggblom MM, Stefaniak S, editors. *Soil and Water Pollution Monitoring, Protection and Remediation*. Netherlands: Springer; 2006. p. 107-14.
  24. Spurgeon DJ, Hopkin SP. Comparisons of metal accumulation and excretion kinetics in earthworms (*Eisenia fetida*) exposed to contaminated field and laboratory soils. *Appl Soil Ecol* 1999;11:227-43.
  25. Lagisz M, Kramarz P, Niklinska M. Metal kinetics and respiration rates in F1 generation of carabid beetles (*Pterostichus oblongopunctatus* F.) originating from metal-contaminated and reference areas. *Arch Environ Contam Toxicol* 2005;48:484-9.
  26. Højbjerg S, Nielsen JB, Andersen O. Effects of dietary lipids on whole-body retention and organ distribution of organic and inorganic mercury in mice. *Food Chem Toxicol* 1992;30:703-8.
  27. Turner CJ, Bhatnagar MK, Speisky H. Effect of subchronic administration of ethanol and methylmercury in combination on the tissue distribution of mercury in rats. *Can J Physiol Pharmacol* 1990;68:1558-62.
  28. Allen PB, Lindsay H, Tham TC. How do patients with inflammatory bowel disease want their biological therapy administered? *BMC Gastroenterol* 2010;10:1.
  29. Nwokocha CR, Ufearo CS, Owu DU, Idemudo NC, Ojukwu LC. *In vivo* distribution of lead in male and female rats after intraperitoneal and oral administration. *Toxicol Ind Health* 2012;28:131-5.
  30. Nwokocha CR, Younger-Coleman N, Owu DU, Asemota HN, Iwuala MO, Lay-Ekuakille A. Analysis of time of measurement and modes of administration of some medicinal plants additives on mercury accumulation in the liver. *Int J Meas Technol Instrum Eng* 2013;3:60-70.

**Cite this article as:** Nwokocha C, Younger-Coleman N, Nwokocha M, Owu D, Iwuala M. A comparative study of the effect of some nutritional medicinal plants effect on lead accumulation in the liver following different modes of administration. *Phcog Res* 2014;6:306-11.

**Source of Support:** Nil, **Conflict of Interest:** None declared.