



REVIEW

REVISED Variation in RAD51 details a hub of functions: opportunities to advance cancer diagnosis and therapy [version 2; referees: 4 approved]

Nick LL van der Zon¹, Roland Kanaar ^{1,2}, Claire Wyman^{1,3}

¹Department of Molecular Genetics, Erasmus MC, Rotterdam, PO Box 2040, 3000 CA, The Netherlands

²Oncode Institute, Erasmus MC, Rotterdam, PO Box 2040, 3000 CA, The Netherlands

³Department of Radiation Oncology, Erasmus MC, Rotterdam, PO Box 2040, 3000 CA, The Netherlands

v2 **First published:** 12 Sep 2018, 7(F1000 Faculty Rev):1453 (<https://doi.org/10.12688/f1000research.15650.1>)
Latest published: 03 Dec 2018, 7(F1000 Faculty Rev):1453 (<https://doi.org/10.12688/f1000research.15650.2>)

Abstract

Loss of genome stability is one of the hallmarks of the enabling characteristics of cancer development. Homologous recombination is a DNA repair process that often breaks down as a prelude to developing cancer. Conversely, homologous recombination can be the Achilles' heel in common anti-cancer therapies, which are effective by inducing irreparable DNA damage. Here, we review recent structural and functional studies of RAD51, the protein that catalyzes the defining step of homologous recombination: homology recognition and DNA strand exchange. Specific mutations can be linked to structural changes and known essential functions. Additional RAD51 interactions and functions may be revealed. The identification of viable mutations in this essential protein may help define the range of activity and interactions needed. All of this information provides opportunities to fine-tune existing therapies based on homologous recombination status, guide diagnosis, and hopefully develop new clinical tools.

Keywords

Homologous recombination, BRCA2, variants of unknown significance, nucleoprotein complexes, structural biology, cancer

Open Peer Review

Referee Status:

	Invited Referees			
	1	2	3	4
REVISED				
version 2 published 03 Dec 2018				
version 1 published 12 Sep 2018				

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty. In order to make these reviews as comprehensive and accessible as possible, peer review takes place before publication; the referees are listed below, but their reports are not formally published.

- 1 **Scott W Morrical**, University of Vermont College of Medicine, USA
- 2 **Maria Spies**, University of Iowa, USA
- 3 **Mauro Modesti**, Cancer Research Center of Marseille, France
- 4 **Michael G Sehorn**, Clemson University, USA

Discuss this article

Comments (0)

Corresponding author: Roland Kanaar (r.kanaar@erasmusmc.nl)

Author roles: **van der Zon NL:** Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Kanaar R:** Conceptualization, Funding Acquisition, Resources, Supervision, Writing – Review & Editing; **Wyman C:** Conceptualization, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Work in the RK laboratory is supported by the gravitation program CancerGenomiCs.nl from the Netherlands Organisation for Scientific Research (NWO) and is part of the OncoCode Institute, which is partly financed by the Dutch Cancer Society. Work in the CW laboratory is supported by NWO Exact and Natural Sciences Project grant 680-91-102.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 van der Zon NL *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: van der Zon NL, Kanaar R and Wyman C. **Variation in RAD51 details a hub of functions: opportunities to advance cancer diagnosis and therapy [version 2; referees: 4 approved]** *F1000Research* 2018, 7(F1000 Faculty Rev):1453 (<https://doi.org/10.12688/f1000research.15650.2>)

First published: 12 Sep 2018, 7(F1000 Faculty Rev):1453 (<https://doi.org/10.12688/f1000research.15650.1>)

REVISED Amendments from Version 1

In the new version we corrected a misstatement regarding the interpretation of the data presented by [Brouwer *et al.*, 2018](#). The sentence "Slight variation at this interface was observed in two different crystal forms determined by Brouwer *et al.*, but the forms did not alternate in the same filament¹⁵." has been deleted. As a matter of fact, RAD51-ATP filament crystallized in one crystal form, not two. This one crystal form contained both interfaces referred to in our original sentence. In the crystal, the two interface types alternate between successive protomer-protomer interfaces. Figure 5C in the publication by [Brouwer *et al.*, 2018](#) represents two models of the RAD51-ATP filament, obtained by building a filament using only one or the other type of interface seen in the crystal. In our publication we do describe the two conformations described by the [Brouwer *et al.*, 2018](#) from the structure and single-molecule analyses in the subsequent paragraph.

See referee reports

Introduction

Homologous recombination (HR) is essential for the maintenance of genome stability. By taking advantage of DNA sequence complementarity, HR is uniquely suited to repair a variety of DNA lesions—such as DNA double-strand breaks (DSBs) and stalled or collapsed DNA replication forks—that affect both strands of the double helix¹. A key intermediate in the complex choreography of the DNA strands underlying HR is single-stranded DNA (ssDNA), generated near the lesion, bound by RAD51 protein. Initial nucleation of RAD51 on the ssDNA is followed by binding of additional protomers, resulting in a dynamic nucleoprotein filament with homology search and DNA strand exchange capabilities². The product of the breast cancer-associated gene 2 (*BRCA2*) acts as a molecular scaffold (mediator) in this process by chaperoning RAD51 onto replication protein A (RPA)-coated ssDNA³. Although HR is required for normal cell function, genome sequencing approaches⁴ and functional HR assays performed on viable tumor material⁵ reveal that HR deficiency is more widespread in tumors than originally anticipated. This insight is important because patients with HR-deficient tumors can benefit greatly from treatment with poly ADP-ribose polymerase (PARP) inhibitors, which selectively kill HR-deficient cells⁶. Because worldwide sequencing efforts of germline and tumor DNA identify more and more variants of RAD51, assessing the functional consequences of these amino acid variations is of utmost importance for prognosis, diagnosis, and precision therapies for patients with cancer. This review will provide the latest advances and insights gained on the functional implications of RAD51 mutations.

The biochemical and biological functions of RAD51 can be related to the structure of individual protomers, the filaments they form, and their dynamic filament properties. It has long been appreciated that the particular structural features of filaments formed by recombinase proteins like RAD51 reflect its function. Recombinase nucleoprotein filaments are arranged with proteins in a right-handed helix around DNA. Functionally important parameters, such as the number of monomers per turn of the helix, the helical pitch, and rise per monomer, define these nucleoprotein filaments⁷. More detailed structural information is

available on the interface between monomers, which comprises the ATPase active site of DNA-bound recombinases⁸. Additional dynamic properties such as kinetic parameters for filament assembly and disassembly as well as continuity and flexibility have more recently been defined on the basis of single-molecule methods⁹. These all proved to be rich quantitative characteristics of wild-type RAD51 for comparison with any new variants and predicting the effect of amino acid changes.

RAD51 nucleoprotein filaments: variety in form and function

Bacterial RecA is the paradigm recombinase first described in detail to define the filament structure. Nucleoprotein filament structures of RecA and RAD51 have been reconstructed from negative stain electron microscope images. This revealed, among other features, that both proteins form filaments with a variety of conformations⁷. The first atomic resolution crystal structure of a eukaryotic nucleoprotein filament, yeast Rad51, revealed that the protomer-protomer interface created an ATPase active site⁸. This nucleotide-binding protomer interface was in two different conformations. Specifically, the movement of two amino acid side chains, Phe187 and His352, resulted in a rigid body movement of one protomer relative to the other, creating the two conformations⁸. That structure was solved for a version of Rad51 missing 79 N-terminal amino acids. Subsequent crystal structures of full-length Rad51, including these N-terminal amino acids, showed only one conformation for the ATPase interface¹⁰. Although there were other differences in the proteins from these two studies, it is possible that the N-terminus constrains the filaments to a single conformation at the ATPase interface^{8,10}. More recently, high-resolution structural models of human RAD51 filaments have been obtained from cryo-electron microscopy reconstructions^{11,12}. Here, filaments formed with human RAD51 had only one conformation at the nucleotide-binding site. However, there is evidence that this region influences conformation at the dimer interface and filament dynamics. It has been shown that interaction of the N-terminal domain of RAD51 with the BRC4 peptide of BRCA2 at the protomer-protomer interface restricts RAD51 dynamics, locking it into a single conformation in the filament¹³. Together, the structural studies of yeast Rad51 with and without the N-terminal 79 amino acids and the effect of BRC4 interaction with the N-terminus of human RAD51 show a role for these regions in filament arrangement and dynamics. Control and tuning of dynamic variation at the protomer contacts in the filament are certainly important aspects of RAD51 catalyzed homology recognition and DNA strand exchange functions.

More globally, RAD51 nucleoprotein filaments can be described by helical parameters and dynamic rearrangements. Specifically, active filaments formed with ATP or an analog bound at the interface have a longer helical pitch and an "open" conformation^{12,14}. Recently, RAD51 nucleoprotein filaments with distinct mechanical properties have been defined for the wild-type protein (for example, persistence length, helical pitch, and rise per monomer) by a combination of single-molecule force spectroscopy and crystallography¹⁵. One form is described as the "open" or "extended" conformation, which displays a helical pitch of 134 Å and a rise per monomer of 18.4 Å. This

conformation is presumed to be the active state required for strand exchange. The second—considered to be the “closed”, “condensed”, or “collapsed” conformation—displays a helical pitch and rise per monomer of 112 Å and 16.9 Å, respectively. This difference results in a change of the monomers per helical turn of 7.3 versus 6.6 for the “open” versus “closed” conformation, respectively¹⁵. These two forms differ in position of a hinge at the protomer interface, resulting in tilting of one protomer relative to the other in the “closed” conformation with respect to the “open” conformation. Because the measured energy difference between these two forms was small, they could readily switch between the two conformations by thermal excitation, independent of other energy input or ATP hydrolysis¹⁵. These data lead to the conclusion that RAD51 nucleoprotein filaments can interconvert between two different conformational states. The functional role of this intriguing dynamic interchange has yet to be defined but is consistent with larger filament dynamics associated with ATP hydrolysis, which has already been linked to function^{16,17}. Flexible and dynamic arrangements of filament structure are likely to be important, even essential, to the RAD51 catalyzed function: homology recognition and DNA strand exchange. Recent work has modified the ATPase active site to see what effect this has on filament form and function. Specifically, filaments formed with RAD51 mutants, K113A and K113R, which are located in the ATPase domain, have been characterized. RAD51 K131A does not bind ATP, and RAD51 K113R can bind but not hydrolyze ATP. Both mutants form filaments with a “closed” conformation, as does the wild-type protein with the non-hydrolyzable ATP analog AMP-PNP¹². It is suggested that these forms, with different helical pitch and rise, could represent intermediate steps in dynamic filament assembly and rearrangement important for function.

Like many other proteins, RAD51 is modified by phosphorylation in response to genotoxic stress. The importance and biochemical effect of the several phosphorylation sites have been difficult to sort out (as discussed in Subramanyam *et al.*¹⁸). Clear effects of one RAD51 phosphorylation site, Y54, at the protomer interface were obtained by mimicking phosphorylation with a non-natural amino acid (p-carboxymethyl-L-phenylalanine, or

pCMF). RAD51 with Y54 replaced by pCMF has enhanced DNA strand exchange activity. The enhanced activity is coupled to weaker DNA binding and formation of less stable/more dynamic filaments, emphasizing the essential role of dynamic interactions among RAD51 itself and DNA to perform its core biochemical function¹⁸. Another example illustrating the importance of the RAD51–RAD51 interface is demonstrated by using small-molecule chemical inhibitor RI-1¹⁹. This inhibitor covalently targets C319. C319 is located at the protomer–protomer interface, where it serves as an ATP cap, overlaying the nucleotide-binding site. The inhibitor thereby disrupts the protomer–protomer interface¹⁹. RAD51 filaments apparently have a “plastic” structure, and interconvertible conformations are influenced by the state of nucleotide bound at their interface, specific amino acid interactions, post-translational modifications, and chemical compounds. Mutations that potentially affect RAD51 filament dynamics will be discussed in the next section.

RAD51 mutations associated with human disease

Because RAD51 nucleoprotein filament formation and dynamics are essential elements of its DNA strand exchange function, small changes to the protomers could cause dramatic effects. Mutations affecting DNA binding, filament formation or structure, ATP hydrolysis, or strand exchange would all influence function in HR. Here, we will specifically discuss recently identified mutations and what is known about their effects on RAD51 function (Table 1 and Figure 1). New RAD51 mutants have been discovered from functional HR assays on tumor material, genome sequencing of patients with phenotypes like those of Fanconi anemia (FA) (a genetic disease including HR defects), and large-scale cancer genome sequencing. The mutations causing FA-like phenotypes are perhaps the easiest to understand functionally. They are T131P and A293T, located at the ATPase active site interface, close to the amino acids F129 and H294, equivalent to F187 and H352 in yeast Rad51. These amino acids are expected to influence the conformation at this interface and its alternating conformations, which they directly contributed to in yeast Rad51 filaments^{8,12}. Mutations creating different chemical environments in close proximity of this

Table 1. Recently identified RAD51 mutations.

Mutation	Sort of mutation	DNA-binding activity	ATPase activity	Strand exchange activity	Reference(s)
F86L	Somatic	Impaired	Normal	Impaired	20
T131P	Germline	Normal	DNA-independent	Impaired	21
D149N	Somatic	Impaired	Normal	Normal	22
R150Q	Germline	Impaired	Impaired	Normal	22
G151D	Somatic	Impaired	Impaired	Enhanced	22,23
E258A	Germline	Impaired	DNA-independent	Impaired	20
Q268P	Somatic	Impaired	Normal	Impaired	24
Q272L	Somatic	Impaired	Normal	Impaired	24
A293T	Germline	Impaired	Impaired	Impaired	25




Figure 1. Overview of RAD51 structures and interactions. Overview of RAD51 mutations mapped onto structures based on PDB ID 5NWL¹⁵ and 1NOW²⁶. **(a)** Model based on the crystal structure of a heptameric right-handed RAD51 nucleoprotein filament. Two adjacent protomers are colored green and light gray; the other five protomers are shown in darker gray. ATP is highlighted by the blue areas at the interfaces of distinct protomers, and sites of mutation are highlighted in red. The red patch on the top of the white-colored protomer is the patch of amino acids 149–151 that is assumed to affect interaction of RAD51 with other binding partners. Note that Q272 is not highlighted in the figure, as that residue was not resolved in the crystal structure. **(b)** Focus on the interface between two protomers, where ATP is shown in blue and the discussed sites of mutation are highlighted with red spheres. Most of these mutations can be found at the direct interface between the two protomers and thus close to the nucleotide-binding pocket. **(c)** Interaction of RAD51 (light gray) with a BRCA2 BRC repeat (cyan). The locations of mutation discussed are highlighted in red. Here, one mutation is in close proximity of the binding region with the BRC repeat (E258) and therefore might impair RAD51 binding by BRCA2.

interaction might affect conformational dynamics of human RAD51 nucleoprotein filaments. For example, A293T at the protomer–protomer interface is directly adjacent to a conserved loop involved in DNA binding and might explain the impaired filament assembly and stability²⁵. These mutations have a dominant negative effect in some biochemical assays^{21,25}.

Genome analysis of patients with cancer has revealed RAD51 somatic mutations with interesting functional implications. Changes in three adjacent residues (D149, R150, and G151) located in the so-called Schellman loop were identified in breast cancer²². This region is reported to include a p53-binding site^{27–29}. These three adjacent amino acids map onto RAD51 at a position distinct from the DNA-binding or ATPase interface sites²². The last of these three mutations, G151D, does have altered DNA-binding and ATPase activity, indicating that filament architecture or dynamics or both are affected²³. Mutations found in kidney and lung tumors, Q268P and Q272L, are located in a DNA-binding loop important for allosteric activation of ATP hydrolysis and DNA strand exchange²⁴. Additional breast cancer-associated mutations F86L and E258A are located in the multimerization interface of RAD51 nucleoprotein filament. The DNA-binding loop and interface mutations, like the mutants described above, have a dominant negative effect in biochemical assays for RAD51 function^{20,24}. Because RAD51 is an essential protein, these viable mutants may identify a functional range of activity. This aspect will be important to better define if such changes are to be targeted for eventual therapeutic intervention. Interestingly, F86L and E258A change the RAD51 protomer interface as well as the interaction interface with BRCA2²⁰. This example points out that RAD51 mutations not only impair

nucleoprotein filament structure or stability but also can affect interactions with other partner proteins such as BRCA2 or possibly p53. BRCA2 is also an essential protein and its presumed essential role is as a RAD51 chaperone^{3,4}. Our own work indicates that all diffusing nuclear RAD51 is bound to BRCA2, indicating that even minor changes to structure at the interface may have large consequences³⁰.

The possible effects of RAD51 mutations on the interaction with BRCA2

The recent analysis of RAD51 we have reviewed focused mainly on the effects on filament formation and function. Nucleoprotein filament dynamics and activities, essentially RAD51–RAD51 and RAD51–DNA interactions, are not the only interactions needed in normal HR function. Interaction partners, such as BRCA2, are obviously also needed for proper HR. BRCA2 is a large protein (3,418 amino acids) that includes a central region with eight BRC repeats that bind RAD51 and at least one additional RAD51-binding motif at the C-terminus. BRCA2–RAD51 interaction is needed to transfer RAD51 to replace RPA on ssDNA originating from the DSBs^{31,32}. Partial structures of a complex including one BRC repeat and RAD51 are available^{26,33}. The interface between RAD51 and BRCA2–BRC repeats overlaps with the RAD51–RAD51 interaction. Thus, changes in RAD51–RAD51 interface may also affect its interaction with BRCA2 (Figure 1C). Additional binding sites of RAD51 are present in BRCA2, such as the C-terminal domain of BRCA2, which is suggested to bind across a RAD51–RAD51 interface in the nucleoprotein filament³⁴. The importance of interaction between RAD51 and BRCA2 is demonstrated by observations that mutated BRC repeats conferred DNA-damage

sensitivity *in vivo*³⁵ and altered RAD51 nucleoprotein filament formation *in vitro*^{36–38}. The importance of RAD51 mutations on its interaction with additional partners and mediators, such as the RAD51 paralogs PALB2, MMS22L-TONSL, RADX, RAD54, RAD51AP1, and others^{39,40}, awaits structural information on their relevant interfaces. The RAD51 mutations we have reviewed may indicate important additional interactions or identify the dynamic range of interaction strengths tolerated for RAD51 function in cell viability. Altered RAD51 is linked to cancer and other diseases affected by impaired genome stability. A detailed definition of RAD51 mutations linked to functions, emerging from the work we review here, is needed to guide personalized anti-cancer therapies and guide diagnosis based on variants of currently unknown significance.

Abbreviations

BRCA2, breast cancer-associated gene 2; DSB, double-strand break; FA, Fanconi anemia; HR, homologous recombination;

pCMF, p-carboxymethyl-L-phenylalanine; RPA, replication protein A; ssDNA, single-stranded DNA.

Grant information

Work in the RK laboratory is supported by the gravitation program CancerGenomiCs.nl from the Netherlands Organisation for Scientific Research (NWO) and is part of the Oncode Institute, which is partly financed by the Dutch Cancer Society. Work in the CW laboratory is supported by NWO Exact and Natural Sciences Project grant 680-91-102.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank the reviewers of our manuscript for their valuable insights and comments.

References



1. Piazza A, Heyer WD: **Multi-Invasion-Induced Rearrangements as a Pathway for Physiological and Pathological Recombination.** *BioEssays.* 2018; 40(5): e1700249.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
2. Ranjha L, Howard SM, Cejka P: **Main steps in DNA double-strand break repair: an introduction to homologous recombination and related processes.** *Chromosoma.* 2018; 127(2): 187–214.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
3. Bell JC, Kowalczykowski SC: **Mechanics and Single-Molecule Interrogation of DNA Recombination.** *Annu Rev Biochem.* 2016; 85: 193–226.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
4. Davies H, Glodzik D, Morganello S, *et al.*: **HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on mutational signatures.** *Nat Med.* 2017; 23(4): 517–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
5. Naipal KA, Verkaik NS, Ameziane N, *et al.*: **Functional ex vivo assay to select homologous recombination-deficient breast tumors for PARP inhibitor treatment.** *Clin Cancer Res.* 2014; 20(18): 4816–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. O'Connor MJ: **Targeting the DNA Damage Response in Cancer.** *Mol Cell.* 2015; 60(4): 547–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
7. Yu X, Jacobs SA, West SC, *et al.*: **Domain structure and dynamics in the helical filaments formed by RecA and Rad51 on DNA.** *Proc Natl Acad Sci U S A.* 2001; 98(15): 8419–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
8. Conway AB, Lynch TW, Zhang Y, *et al.*: **Crystal structure of a Rad51 filament.** *Nat Struct Mol Biol.* 2004; 11(8): 791–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
9. Candelli A, Modesti M, Peterman EJ, *et al.*: **Single-molecule views on homologous recombination.** *Q Rev Biophys.* 2013; 46(4): 323–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
10. Chen J, Villanueva N, Rould MA, *et al.*: **Insights into the mechanism of Rad51 recombinase from the structure and properties of a filament interface mutant.** *Nucleic Acids Res.* 2010; 38(14): 4889–906.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Short JM, Liu Y, Chen S, *et al.*: **High-resolution structure of the presynaptic RAD51 filament on single-stranded DNA by electron cryo-microscopy.** *Nucleic Acids Res.* 2016; 44(19): 9017–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
12. Špirek M, Mlcoušková J, Belán O, *et al.*: **Human RAD51 rapidly forms intrinsically dynamic nucleoprotein filaments modulated by nucleotide binding state.** *Nucleic Acids Res.* 2018; 46(8): 3967–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
13. Subramanyam S, Jones WT, Spies M, *et al.*: **Contributions of the RAD51 N-terminal domain to BRCA2-RAD51 interaction.** *Nucleic Acids Res.* 2013; 41(19): 9020–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
14. Yu X, Egelman EH: **Structural data suggest that the active and inactive forms of the RecA filament are not simply interconvertible.** *J Mol Biol.* 1992; 227(1): 334–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Brouwer I, Moschetti T, Candelli A, *et al.*: **Two distinct conformational states define the interaction of human RAD51-ATP with single-stranded DNA.** *EMBO J.* 2018; 37(7): pii: e98162.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
16. Ristic D, Modesti M, van der Heijden T, *et al.*: **Human Rad51 filaments on double- and single-stranded DNA: correlating regular and irregular forms with recombination function.** *Nucleic Acids Res.* 2005; 33(10): 3292–302.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. van der Heijden T, Seidel R, Modesti M, *et al.*: **Real-time assembly and disassembly of human RAD51 filaments on individual DNA molecules.** *Nucleic Acids Res.* 2007; 35(17): 5646–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Subramanyam S, Ismail M, Bhattacharya I, *et al.*: **Tyrosine phosphorylation stimulates activity of human RAD51 recombinase through altered nucleoprotein filament dynamics.** *Proc Natl Acad Sci U S A.* 2016; 113(41): E6045–E6054.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
19. Budke B, Logan HL, Kallin JH, *et al.*: **RI-1: a chemical inhibitor of RAD51 that disrupts homologous recombination in human cells.** *Nucleic Acids Res.* 2012; 40(15): 7347–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Silva MC, Bryan KE, Morrical MD, *et al.*: **Defects in recombination activity caused by somatic and germline mutations in the multimerization/BRCA2 binding region of human RAD51 protein.** *DNA Repair (Amst).* 2017; 60: 64–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
21. Wang AT, Kim T, Wagner JE, *et al.*: **A Dominant Mutation in Human RAD51 Reveals Its Function in DNA Interstrand Crosslink Repair Independent of Homologous Recombination.** *Mol Cell.* 2015; 59(3): 478–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
22. Chen J, Morrical MD, Donigan KA, *et al.*: **Tumor-associated mutations in a conserved structural motif alter physical and biochemical properties of human RAD51 recombinase.** *Nucleic Acids Res.* 2015; 43(2): 1098–111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
23. Marsden CG, Jensen RB, Zagebaum J, *et al.*: **The Tumor-Associated Variant RAD51 G151D Induces a Hyper-Recombination Phenotype.** *PLoS Genet.* 2016; 12(8): e1006208.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

24. **F** Silva MC, Morrical MD, Bryan KE, *et al.*: **RAD51 variant proteins from human lung and kidney tumors exhibit DNA strand exchange defects.** *DNA Repair (Amst)*. 2016; **42**: 44–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
25. Ameziane N, May P, Haitjema A, *et al.*: **A novel Fanconi anaemia subtype associated with a dominant-negative mutation in RAD51.** *Nat Commun*. 2015; **6**: 8829.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. **F** Pellegrini L, Yu DS, Lo T, *et al.*: **Insights into DNA recombination from the structure of a RAD51-BRCA2 complex.** *Nature*. 2002; **420**(6913): 287–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
27. **F** Friedler A, Veprintsev DB, Rutherford T, *et al.*: **Binding of Rad51 and other peptide sequences to a promiscuous, highly electrostatic binding site in p53.** *J Biol Chem*. 2005; **280**(9): 8051–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
28. Yoon D, Wang Y, Stapleford K, *et al.*: **P53 inhibits strand exchange and replication fork regression promoted by human Rad51.** *J Mol Biol*. 2004; **336**(3): 639–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Buchhop S, Gibson MK, Wang XW, *et al.*: **Interaction of p53 with the human Rad51 protein.** *Nucleic Acids Res*. 1997; **25**(19): 3868–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Reuter M, Zelensky A, Smal I, *et al.*: **BRCA2 diffuses as oligomeric clusters with RAD51 and changes mobility after DNA damage in live cells.** *J Cell Biol*. 2014; **207**(5): 599–613.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Shahid T, Soroka J, Kong E, *et al.*: **Structure and mechanism of action of the BRCA2 breast cancer tumor suppressor.** *Nat Struct Mol Biol*. 2014; **21**(11): 962–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Sánchez H, Paul MW, Grosbart M, *et al.*: **Architectural plasticity of human BRCA2-RAD51 complexes in DNA break repair.** *Nucleic Acids Res*. 2017; **45**(8): 4507–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Galkin VE, Esashi F, Yu X, *et al.*: **BRCA2 BRC motifs bind RAD51-DNA filaments.** *Proc Natl Acad Sci U S A*. 2005; **102**(24): 8537–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. **F** Esashi F, Galkin VE, Yu X, *et al.*: **Stabilization of RAD51 nucleoprotein filaments by the C-terminal region of BRCA2.** *Nat Struct Mol Biol*. 2007; **14**(6): 468–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
35. Tal A, Arbel-Goren R, Stavans J: **Cancer-associated mutations in BRC domains of BRCA2 affect homologous recombination induced by Rad51.** *J Mol Biol*. 2009; **393**(5): 1007–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Thorslund T, Mcllwraith MJ, Compton SA, *et al.*: **The breast cancer tumor suppressor BRCA2 promotes the specific targeting of RAD51 to single-stranded DNA.** *Nat Struct Mol Biol*. 2010; **17**(10): 1263–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Jensen RB, Carreira A, Kowalczykowski SC: **Purified human BRCA2 stimulates RAD51-mediated recombination.** *Nature*. 2010; **467**(7316): 678–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Liu J, Doty T, Gibson B, *et al.*: **Human BRCA2 protein promotes RAD51 filament formation on RPA-covered single-stranded DNA.** *Nat Struct Mol Biol*. 2010; **17**(10): 1260–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Dray E, Etchin J, Wiese C, *et al.*: **Enhancement of RAD51 recombinase activity by the tumor suppressor PALB2.** *Nat Struct Mol Biol*. 2010; **17**(10): 1255–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. **F** Godin SK, Sullivan MR, Bernstein KA: **Novel insights into RAD51 activity and regulation during homologous recombination and DNA replication.** *Biochem Cell Biol*. 2016; **94**(5): 407–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Referee Status:



Editorial Note on the Review Process

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty and are edited as a service to readers. In order to make these reviews as comprehensive and accessible as possible, the referees provide input before publication and only the final, revised version is published. The referees who approved the final version are listed with their names and affiliations but without their reports on earlier versions (any comments will already have been addressed in the published version).

The referees who approved this article are:

Version 1

- 1 **Michael G Sehorn** Department of Genetics and Biochemistry, Clemson University, Clemson, USA
Competing Interests: No competing interests were disclosed.
- 2 **Mauro Modesti** Cancer Research Center of Marseille, Marseille, France
Competing Interests: No competing interests were disclosed.
- 3 **Maria Spies** Department of Biochemistry, University of Iowa, Iowa City, USA
Competing Interests: No competing interests were disclosed.
- 4 **Scott W Morrical** Department of Biochemistry, University of Vermont College of Medicine, Vermont, USA
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research