

# Mycoflora and Enzymatic Characterization of Fungal Isolates in Commercial Meju, Starter for a Korean Traditional Fermented Soybean Product

Jin-Ho Baek<sup>1</sup>, Kum-Kang So<sup>1</sup>, Yo-Han Ko<sup>1</sup>, Jung-Mi Kim<sup>2</sup> and Dae-Hyuk Kim<sup>1,\*</sup>

<sup>1</sup>Department of Molecular Biology, Department of Bioactive Material Sciences, Center for Fungal Pathogenesis, Chonbuk National University, Jeonju 561-756, Korea

<sup>2</sup>Department of Bio-Environmental Chemistry, Wonkwang University, Iksan 570-749, Korea

**Abstract** Mycoflora was assessed in the commercial meju from four well-separated geographic origins. A total of 112 fungal isolates were identified by phenotypic characteristics and molecular taxonomy using sequencing the internal transcribed spacer of the rDNA and revealed 19 species from 13 genera. Enzymatic characteristics of protease and amylase, and mycotoxin production were analyzed.

**Keywords** Amylase, Meju, Mycoflora, Mycotoxin, Protease

Korean traditional meju, a brick of dried fermented soybeans, serves as an important starting material of several Korean condiments such as soybean paste (doenjang), soy sauce (ganjang), and hot pepper paste (gochujang) [1]. Meju is made by soaking, steaming, and mashing soybeans, and then fermenting them with various microorganisms such as bacteria, yeast, and fungi for a prolonged term, usually 2–3 mon, until it used for further process. Fungi particularly perform a crucial role in decomposition organic matter on meju [1, 2].

Although several factors determine differences in the taste and aroma of traditional fermented soybean products such as the raw materials and process techniques, the most significant factor is the extracellular enzymatic activities of the various microorganisms involved in the fermentation process with starter material, meju [3]. The fungi of meju

or soybean products are commonly evaluated as being safe. However, it is possible to be contaminated with mycotoxin-producing fungi under the natural fermentation [4].

It has been known that fungi perform an important role in fermentation and decomposition of organic matters of the soybeans into small nutrient functional molecules as amino acids, sugars, organic acids, alcohol, and esters, which underline the typical flavor of meju [1, 3]. Therefore, the mycoflora study of fermented soybean products have been reported that *Aspergillus* spp., *Cladosporium* spp., *Eurotium* spp., *Fusarium* spp., *Lichtheimia* spp., *Mucor* spp., *Penicillium* spp., *Rhizopus* spp., and *Scopulariopsis* spp. participated in general meju fermentation [5–9].

In order to better comprehend the complex progresses that appear during the slow fermentation of commercial meju, analysis of the fungal diversity in each product will have taken a great interest. Therefore, the aims of this study were to assess the mycoflora of meju fermentation on traditionally manufactured commercially available meju samples and to compare these fungal species with those analyzed in meju. In addition, extracellular enzymatic characteristics of the fungal isolates related to proteins and carbohydrates degradation were carried out. Finally, mycotoxin producing capacity was evaluated using thin layer chromatography (TLC).

Fungal strains were isolated from commercial manufactured meju samples from four different regions in Korea (Table 1). Ten loaves of each molded meju sample, including the surface and the inside of the sample, were collected. Each loaf was smashed to paste and 2 g from smashed each loaf was mixed for fungal isolation. Twenty grams of the mixed preparation for each sample was added to 180 mL of sterile

Mycobiology 2014 September, 42(3): 291-295  
<http://dx.doi.org/10.5941/MYCO.2014.42.3.291>  
pISSN 1229-8093 • eISSN 2092-9323  
© The Korean Society of Mycology

\*Corresponding author

E-mail: dhkim@jbnu.ac.kr

Received May 27, 2014

Revised June 23, 2014

Accepted July 23, 2014

©This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Table 1.** Representative fungal strains isolated from meju used in this study

Sample designation	Geographic origin	Morphological group	No. of isolates	Scientific nomenclature	ITS identification				Enzyme activity <sup>a</sup>	
					GenBank accession No.	Sequence length (bp)	Coverage (%)	Identity (%)	Protease activity	Amylase activity
A	Gyeonggi, Paju	A - a	4	<i>Rhizopus stolonifer</i>	FN401529.1	882	100	100	–	–
		A - b	2	<i>Mucor circinelloides</i>	FN598919.1	639	92	99	+	–
		A - c	14	<i>Aspergillus flavus</i>	JQ269826.1	597	84	99	++	+++
		A - d	3	<i>Mucor racemosus</i>	AY213659.1	638	98	100	+	+
		A - e	12	<i>Mucor racemosus</i>	HM641690.1	638	99	99	+	+
		A - f	1	<i>Penicillium chrysogenum</i>	HM371375.1	540	79	100	+	+
		A - g	1	<i>Mucor racemosus</i>	HM641690.1	638	98	99	+	+
		A - h	1	<i>Penicillium polonicum</i>	AF033475.1	531	91	100	+	+
		A - i	1	<i>Penicillium polonicum</i>	JQ082508.1	531	100	99	+	+
		A - j	2	<i>Penicillium crustosum</i>	JN252103.1	532	100	100	++	++
B	Gyeongbuk, Yecheon	B - a	7	<i>Mucor circinelloides</i>	AY213658.1	639	95	99	+	–
		B - b	1	<i>Rhizomucor variabilis</i>	DQ119007.1	640	95	99	+	–
		B - c	4	<i>Lichtheimia</i> sp.	FJ719383.1	743	90	100	+	–
		B - d	7	<i>Aspergillus flavus</i>	JQ269826.1	596	82	100	++	+++
		B - e	3	<i>Aspergillus oryzae</i>	GU385811.1	638	83	99	++	+++
		B - f	4	<i>Aspergillus oryzae</i>	GU385811.1	597	88	99	++	+++
		B - g	1	<i>Trichosporon asahii</i>	AB369919.1	543	69	100	+	–
		B - h	1	<i>Eurotium amstelodami</i>	HM145963.1	556	75	100	–	+
C	Jeonnam, Gangjin	C - a	3	<i>Lichtheimia corymbifera</i>	HQ285657.1	766	94	99	+	–
		C - b	2	<i>Gibberella intermedia</i>	JQ69083.1	560	71	99	–	+
		C - c	1	<i>Pichia burtonii</i>	AM420292.1	452	59	98	–	++
		C - d	4	<i>Pichia burtonii</i>	EU714323.1	449	69	99	–	++
		C - e	2	<i>Aspergillus flavus</i>	JQ269826.1	596	82	100	++	+++
		C - f	1	<i>Aspergillus flavus</i>	JQ269826.1	597	87	99	++	+++
		C - g	3	<i>Aspergillus oryzae</i>	JN561266.1	597	82	100	++	+++
		C - h	2	<i>Aspergillus flavus</i>	JQ269826.1	597	82	100	++	+++
		C - i	2	<i>Aspergillus flavus</i>	HQ340109.1	596	74	100	++	+++
		C - j	2	<i>Aspergillus flavus</i>	HQ010119.1	596	86	100	++	+++
D	Gangwon, Yeongwol	D - a	1	<i>Absidia corymbifera</i>	DQ658188.1	771	94	99	–	–
		D - b	11	<i>Aspergillus flavus</i>	JQ269826.1	597	82	100	++	+++
		D - c	1	<i>Rhizomucor pusillus</i>	AB369914.1	629	86	99	+	+++
		D - d	4	<i>Scopulariopsis brevicaulis</i>	KC311514.1	631	99	98	++	+
		D - e	2	<i>Scopulariopsis brevicaulis</i>	JN942890.1	630	92	98	++	+
		D - f	2	<i>Eupenicillium</i> sp.	JQ828843.1	616	84	86	++	++

ITS, internal transcribed spacer.


<sup>a</sup>Enzymatic activity was assigned based on distance of the clearance zone from the colony margin (+++, > 10 mm; ++, 5~10 mm; +, < 5 mm; –, absent).

water, vigorously vortexed for 30 sec, and maintained at room temperature for 30 min. Serial dilutions of  $10^{-4}$ ,  $10^{-5}$ , and  $10^{-6}$  were prepared and 0.3 mL of each of these dilutions was spread onto the surface of potato dextrose agar medium (PDA) containing 17 mg/L chloramphenicol.


The plates were incubated for 2~5 days at 30°C, the isolated fungal colonies were transferred to a new PDA plate and incubated for 3~7 days at 30°C to obtain a pure culture. Agar blocks containing actively growing mycelia were stored as previously described for further study [10].

The whole of the four meju samples yielded a total of 112 distinctive fungal colonies. Based on characteristics of colony morphology, growth rate, and pigmentation, 41, 28, 22, and 21 isolates from samples A~D could be assigned to 10, 8, 10, and 6 groups, respectively. Each isolates were identified in genus standard methods [11]. At least two representative isolates for each type (except for those with a single isolate) were classified by sequence comparison of the internal transcribed spacer (ITS) region of the nuclear ribosomal gene (rDNA). The ITS region was amplified using with primers ITS1 (5'-CTAGCGGAGACGGGCTGC-3') and ITS4 (5'-GGGTCTCTTGGTATGGTC-3'). The resulting PCR amplicon was cloned into the pGEM-T Easy vector (Promega, Madison, WI, USA) and four clones per one isolate were sequenced from both directions using the dideoxynucleotide method with universal and synthetic oligonucleotide primers. A total of 112 fungal isolates were identified as belonging to 13 genera, 19 species, and four *Lichtheimia* and two *Eupenicillium* isolates of unclassified species. Seven, seven, five, and five species were identified for samples A~D, respectively (Table 1). Even though some representative isolates showed slightly different cultural characteristics of colony morphology, growth rate, and pigmentation, they were identified as the same species. Among classified isolates, the genera *Aspergillus* and *Mucor*

were the predominant fungi; 41 and 10 isolates of *A. flavus* and *A. oryzae*, respectively, and 16 and 9 isolates of *M. racemosus* and *M. circinelloides*, respectively, were obtained (Fig. 1). The mycoflora of meju fermentation showed a



**Fig. 1.** Venn diagram summarizing the number of unique and shared fungal species in the four meju samples from four well-separated geographic origins. A, B, C, and D, as indicated in Table 1. Numbers in parentheses represent the number of identified fungal isolates from each sample. Parentheses of shared species are shown according to the relative sample locations. Note that only one species was found in all four meju samples and six species were present in two different samples.



**Fig. 2.** Neighbor-joining tree resulting from the gene sequences of the internal transcribed spacer region of rDNA in 19 identified fungal strains derived from meju, starter material for fermented soybean products. Bootstrap values supporting branch points are expressed as the percentages and confidence values for individual branches were determined by 1,000 replication.

different pattern in that common species were isolated from different samples. For example, *A. flavus* was isolated from all four meju samples, *A. oryzae* was isolated from samples B and C, and *M. circinelloides* was found in samples A and B (Fig. 1). All identified species had been observed in previous studies [4, 6, 8, 12, 13] except for *Trichosporon asahii* and *Scopulariopsis brevicaulis*. Although *S. chartarum* and *S. trigonospora* have been reported in a previous study [8], *S. brevicaulis* has not previously been isolated. In addition, this is the first report of the genus *Trichosporon* in the mycoflora of a fermented soybean product.


The phylogenetic relationships of all identified isolates were presented (Fig. 2). A clade with significant support by bootstrap analysis included species from different meju samples. These results suggested that common origin of fungal flora in soybean products irrespective of the separated geographic origin and fermentation method.

Extracellular enzymatic characteristics that are believed to be important for traditional fermentation, such as protease and amylase activities, were analyzed for all isolated fungi (Table 1). To measure protease and amylase activities, agar blocks containing actively growing mycelia from 10-day-old cultures were transferred to the center of petri dishes containing casein [14] and starch media [15], respectively. The dishes in triplicate were maintained at 30°C for 7 days. To confirm the presence of true proteolytic and amylolytic activities, an acid solution of mercury chloride and Gram's iodine solution, respectively, were added to each plate. Extracellular enzymatic activity was certificated by the formation of a transparent halo zone. These procedures revealed differences hydrolysis region around the colony which indicated the protease and amylase producing ability of the fungal strains [14, 15]. All *Aspergillus* isolates showed high levels of both enzymes. In addition, most of the isolates from the meju showed at least one of the enzymatic activities except for *A. corymbifera* and *R. stolonifer*. *A. corymbifera*, which is known as a common causal agent of mucormycosis requiring high enzymatic activities necessary for food processing and pathogenesis and also found in another meju study [13], did show no enzymatic activity under the current culture conditions. Therefore, although it has not been detected in this study, it will be of great interest to analyze the enzyme activities of *A. corymbifera* on sterile culture and different culture conditions. Although several studies have shown a broad spectrum of enzymatic activity from *R. stolonifer*, including starch utilization [16], our assay did not successfully detect amylase activity, possibly because of differences in the assay conditions or culture conditions.

Mycotoxin contamination of fermented food has attracted worldwide attention because of its adverse effects on human health and livestock [17, 18]. It is possible for meju to be contaminated with aflatoxins (AFs; Sigma, St. Louis, MO, USA) produced by *A. flavus* or *A. parasiticus*, which have similar morphological and biochemical properties to *A. oryzae* or *A. sojae* [19-21]. In addition, 3 strains among

51 isolates of the *A. flavus* and *A. oryzae* group present in meju were previously shown to be aflatoxinogenic [4]. Since we found 51 isolates of the *A. flavus* and *A. oryzae* group, we analyzed for the production of mycotoxin from all isolates. To determine production of sterigmatocystin (ST; Sigma) or AF,  $1 \times 10^6$  conidia/mL for each isolates were cultured in the fungal complete media 3 mL under constant dark condition at 30°C for 7 days without agitation. The whole mycelia and culture filtrate were then extracted by 1 mL chloroform thoroughly. The ST/AF in 50  $\mu$ L of chloroform-extract was separated by silica gel TLC plate (Merck, Darmstadt, Germany) in toluene:ethyl acetate:acetic acid (80:10:10 [v/v/v]) solvent system, followed by treatment of 10% aluminum chloride in 95% ethanol to enhance ST/AF fluorescence upon exposure to a longwave UV light [22]. As shown in Fig. 3, we did not detect the presence of AF or ST from any of the isolates in our culture conditions.

In conclusion, the current study examined the mycoflora from the starting material for fermented soybean products, meju in four well-separated geographic origins. From four meju samples, a total of 112 fungal isolates were obtained and identified as belonging to 13 genera and 19 species. There were four isolates of *Lichtheimia* and two isolates of



**Fig. 3.** Thin layer chromatography (TLC) analysis of sterigmatocystin and aflatoxin from culture extracts of *Aspergillus flavus* and *A. oryzae* strains in this study. A representative TLC plate of the organic extracts from culture filtrate of 12 *A. flavus* and *A. oryzae* strains is shown. Strains are indicated under the panel by the lane number. Lanes 1~6 contain extracts from *A. flavus* strains and lanes 7~11 contain extracts from *A. oryzae* strains. Lanes labeled S, A, C, and N represent sterigmatocystin standard, aflatoxin standard, culture medium alone, and a culture extract of no aflatoxinogenic *A. oryzae* strain (KACC 44823) [23] as a negative control, respectively.

*Eupenicillium* of unclassified species. Biochemical analysis of protease and amylase activities, which are closely related to fermentation, indicated that extracellular enzymatic activity was observed in most isolated strains and was particularly high in general fungal species of *Aspergillus*. No aflatoxigenic fungal isolates were detected in this study.

## ACKNOWLEDGEMENTS

This work was supported by the Bio-industry Technology Development Program, Ministry for Food, Agriculture, Forestry and Fisheries, Republic of Korea and in part by the NRF grants by MSIP (No. 2008-0061897 and NRF-2013R1A1A2012433). This research was also supported in part by Korea Research Council of Fundamental Science & Technology (Joint Degree and Research Center for Biorefinery) and “Cooperative Research Program for Agricultural Science & Technology Development (Project No. PJ009999801), Rural Development Administration, Republic of Korea. We thank the Institute of Molecular Biology and Genetics at Chonbuk National University for kindly providing the facilities for this research. J.H. Baek, K.K. So, and Y.H. Ko were supported by BK21 PLUS program in the Department of Bioactive Material Sciences.

## REFERENCES

1. Lee SS. Meju fermentation for a raw material of Korean traditional soy products. *Korean J Mycol* 1995;23:161-75.
2. An BJ, Son GM, Choi C. Changes in protein and amino acid composition of native meju during fermentation. *J Korean Soc Food Nutr* 1986;15:152-7.
3. Kim JH, Yoo JS, Lee CH, Kim SY, Lee SK. Quality properties of soybean pastes made from meju with mold producing protease isolated from traditional meju. *J Korean Soc Appl Biol Chem* 2006;49:7-14.
4. Jung YJ, Chung SH, Lee HK, Chun HS, Hong SB. Isolation and identification of fungi from a meju contaminated with aflatoxins. *J Microbiol Biotechnol* 2012;22:1740-8.
5. Hong SB, Kim DH, Lee M, Baek SY, Kwon SW, Samson RA. Taxonomy of *Eurotium* species isolated from meju. *J Microbiol* 2011;49:669-74.
6. Kim DH, Kim SH, Kwon SW, Lee JK, Hong SB. Fungal diversity of rice straw for meju fermentation. *J Microbiol Biotechnol* 2013;23:1654-63.
7. Cho DH, Lee WJ. Microbiological studies of Korean native soy-sauce fermentation: a study on the microflora of fermented Korean maeju loaves. *J Korean Agric Chem Soc* 1970;13:35-42.
8. Kim DH, Kim SH, Kwon SW, Lee JK, Hong SB. Mycoflora of soybeans used for meju fermentation. *Mycobiology* 2013;41:100-7.
9. Hong SB, Lee M, Kim DH, Chung SH, Shin HD, Samson RA. The proportion of non-aflatoxigenic strains of the *Aspergillus flavus/oryzae* complex from meju by analyses of the aflatoxin biosynthetic genes. *J Microbiol* 2013;51:766-72.
10. Lee JK, Kim BT, Kim JA, Chung HJ, Park SM, Yang MS, Hwang KJ, Kim DH. Cultural characteristics and extraction of the fungal pigment phleochrome from the phytopathogenic fungus *Cladosporium phlei*. *Biotechnol Bioprocess Eng* 2007;12:508-15.
11. Pitt JI, Hocking AD. *Fungi and food spoilage*. 3rd ed. New York: Springer; 2009.
12. Hong SB, Kim DH, Lee M, Baek SY, Kwon SW, Houbraken J, Samson RA. *Zygomycota* associated with traditional meju, a fermented soybean starting material for soy sauce and soybean paste. *J Microbiol* 2012;50:386-93.
13. Kim JY, Yeo SH, Baek SY, Choi HS. Molecular and morphological identification of fungal species isolated from bealmijang meju. *J Microbiol Biotechnol* 2011;21:1270-9.
14. Nadeem M, Qazi JI, Baig S. Enhanced production of alkaline protease by a mutant of *Bacillus licheniformis* N-2 for dehairing. *Braz Arch Biol Technol* 2010;53:1015-25.
15. Tripathy SS, Dash S, Gupta N. Screening and selection of some fungi for production of extracellular amylase. *Indian J Fundam Appl Life Sci* 2011;1:131-6.
16. Pothiraj C, Eyini M. Enzyme activities and substrate degradation by fungal isolates on cassava waste during solid state fermentation. *Mycobiology* 2007;35:196-204.
17. Chelkowski J. Mycological quality of mixed feeds and ingredients. In: Chelkowski J, editor. *Cereal grain, mycotoxins, fungi and quality in drying and storage*. Amsterdam, London, New York: Elsevier; 1991. p. 217-27.
18. Hussein HS, Brasel JM. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology* 2001;167:101-34.
19. Lee CH, Lee SS. Cereal fermentation by fungi. *Appl Mycol Biotechnol* 2002;2:151-70.
20. Niessen L. PCR-based diagnosis and quantification of mycotoxin-producing fungi. *Adv Food Nutr Res* 2008;54:81-138.
21. Park JH, Kang SJ, Oh SS, Chung DH. The screening of aflatoxin producing fungi from commercial meju and soy bean paste in western Gyeongnam by immunoassay. *J Food Hyg Saf* 2001;16:274-9.
22. Stack M, Rodricks JV. Method for analysis and chemical confirmation of sterigmatocystin. *J Assoc Off Anal Chem* 1971;54:86-90.
23. Murakami H. Classification of the koji mold. *J Gen Appl Microbiol* 1971;17:281-309.