

Original Article

Effect of high-intensity pulse irradiation with linear polarized near-infrared rays on muscle tone in patients with cerebrovascular disease: a randomized controlled trial

NOBUYUKI TAKEUCHI, RPT, PhD^{1,2)*}, NOBUHIRO TAKEZAKO, RPT³⁾, YUKO SHIMONISHI, RPT²⁾, SHIGERU USUDA, RPT, PhD⁴⁾

¹⁾ Department of Physical Therapy, Faculty of Health Care, Takasaki University of Health and Welfare: 501 Nakaoruicho, Takasaki City, Gunma 370-0033, Japan

²⁾ Department of Rehabilitation, Honjo General Hospital, Japan

³⁾ Department of Rehabilitation, Takasaki University of Health and Welfare Attached Clinic, Japan

⁴⁾ Gunma University Graduate School of Health Sciences, Japan

Abstract. [Purpose] This study evaluated effects of a high-intensity linear polarized near-infrared ray irradiation for mitigation of muscle hypertonia. [Subjects] The subjects were 20 patients with cerebrovascular disease. [Methods] Subjects were randomly allocated to an intervention or control group. The intervention group received irradiation of the triceps surae. Passive range of motion and passive resistive joint torque of ankle dorsiflexion were measured before and after the intervention in knee extended and flexed positions. [Results] In the knee extended position, the mean changes in passive range of motion were 2.70° and -0.50° in the intervention and control groups, respectively, and the mean changes in passive resistive joint torque were -1.42 and -0.26 N·m in the intervention and control groups, respectively. In the knee flexed position, the mean changes in passive range of motion were 3.70° and 0.70° in the intervention and control groups, respectively, and the mean changes in passive resistive joint torque were -2.38 and -0.31 N·m in the intervention and control groups, respectively. In both knee positions, the mean changes in the two indices were greater in the intervention group than in the control group. [Conclusion] High-intensity linear polarized near-infrared ray irradiation increases passive range of motion and decreases passive resistive joint torque.

Key words: Infrared ray, Muscle tone, Cerebrovascular disease

(This article was submitted Aug. 20, 2015, and was accepted Sep. 17, 2015)

INTRODUCTION

Muscle tone influences various movements, and reduction of excessive muscle tone is an important treatment target for physical therapy. There are many therapeutic approaches for hypertonia, and a lot of research has been conducted on this topic. Phototherapy, such as laser therapy or linear polarized near-infrared ray (LPNR) irradiation, is one therapeutic approach that has been shown to be effective^{1,2)}. The equipment required for LPNR irradiation is a phototherapy device that emits infrared rays at deep penetrating wavelengths (600–1,600 nm)^{3–7)}. LPNR irradiation is commonly used in various therapeutic scenarios because of its noninvasiveness, safety, and simplicity^{2, 4–6, 8–10)}.

The depth of penetration by emitted rays in phototherapy

generally depends on the intensity (output), and rays reach deeper regions as the intensity increases, but the burn risk also increases. The maximum output of most LPNR irradiation therapy equipment is about 2,000 mW^{4, 5)}, and continuous irradiation at this intensity may cause burns. Thus, irradiation is typically applied intermittently, with repeated irradiation lasting several seconds followed by a rest interval^{4, 5)}. High-intensity LPNR (Hi-LPNR) irradiation therapy using equipment with a maximum output of 10 W has recently been reported¹¹⁾. Continuous irradiation at this output has a very high risk of burns. The burn risk is reduced and deep penetration at the high-intensity peak output is secured by applying pulse irradiation, which involves repeated irradiation for several tens of milliseconds or hundreds of microseconds followed by a rest interval^{11–15)}. These high-intensity phototherapy devices are already being used clinically, but reports on them are mainly limited to pain relief or treatment of paralysis^{11–17)}.

We have previously reported in Japan the inhibitory effect of LPNR irradiation at an output of 1,440 mW on muscle tone in patients with cerebrovascular disease (CVD)¹⁸⁾. The maximum output of Hi-LPNR is higher than that of LPNR; therefore, the effect on deep tissue may also be greater. An

*Corresponding author. Nobuyuki Takeuchi (E-mail: ntakeuchi@takasaki-u.ac.jp)

inhibitory effect of Hi-LPNR irradiation on muscle tone in patients with CVD is also expected, but no study has been performed to test this. Therefore, we performed a study to clarify the effect of Hi-LPNR irradiation on hypertonia in patients with CVD, and the findings of the study are reported in this paper.

SUBJECTS AND METHODS

The objective and methods were explained to all patients, and written informed consent was obtained. This study was approved by the ethics committee at Honjo General Hospital (approval number 20110907).

The study was conducted at Honjo General Hospital in Japan. The subjects were 20 patients with CVD and hypertonia of the triceps surae muscle on the paretic side. The inclusion criterion was a score ≥ 1 on at least one item of the Ankle Plantar Flexors Tone Scale (APTS)¹⁹. The APTS is a plantar flexor muscle tone evaluation index comprised of three items: stretch reflex, middle-range resistance, and final-range resistance. Each item is evaluated on a 5-point scale ranging from 0 to 4 (Appendix 1). The stretch reflex mainly reflects neurological elements constituting muscle tone, and middle- and final-range resistance mainly reflect non-neurological elements of muscle tone¹⁹. The exclusion criteria were previous surgical treatment or drug therapy for hypertonia, limited range of motion of the ankle joint on the paretic side with a cause other than hypertonia, and the presence of pain or another factor that could influence measurement. The same physical therapist performed all measurements. In addition, one of the two assistants of the physical therapist helped to retain the position of the lower limb during measurement, and they assisted with measurement.

The subjects were randomly allocated to an intervention group treated with Hi-LPNR irradiation or a control group without irradiation (10 patients each, Fig. 1). In the intervention group, the patients received 5-min Hi-LPNR irradiation while side lying with the paretic side up and mild flexion of the bilateral hip and knee joints. In the control group, the patients rested in the supine position for 5 min.

Hi-LPNR irradiation therapy was delivered using the Super Lizer PX Type 1 (Tokyo Iken, Tokyo, Japan). This equipment emits pulse irradiation (3-ms irradiation at 7-ms intervals) with a maximum output of 10 W. Several probes with a changeable irradiation caliper were available. A handheld irradiation probe with a 20-mm diameter was used in this study. Although the maximum output is 10 W, the output was set at 8 W for this study, and the total irradiation time was 5 min. We determined the irradiation conditions after performing a preliminary investigation and confirming that irradiation under these conditions could be applied safely without adverse events.

Four sites on the origin side of the myotendinous junction of the triceps surae muscle on the paretic side were irradiated. These sites were selected based on a study using LPNR irradiation at 1,440 mW that some of us had previously performed¹⁸, in which a muscle tone-inhibitory effect was noted in this region. Each site was consecutively irradiated for 3 s, and this procedure was repeated for 5 min. The tip




Fig. 1. Flow diagram of the intervention and control groups
Hi-LPNR: high-intensity polarized near-infrared ray

of the irradiation probe was lightly touched to the skin and rested there without excessive pressure. The myotendinous junction was confirmed by palpation and visual examination.

The outcome measures were passive range of motion of ankle dorsiflexion (PROM) and passive resistive joint torque of ankle dorsiflexion (PRJT), which were measured in the supine position. For PROM, a line perpendicular to the fibula was regarded as the stationary axis, and the 5th metatarsal bone was regarded as the migrating axis. The maximum dorsiflexion angle was measured using a goniometer with 1° increments. The fibular head, lateral malleolus, and head and base of the 5th metatarsal bone were drawn on the patient's skin with a marker as indices during measurement, and dorsiflexion was slowly applied. The change in PROM from before to after the intervention was calculated and analyzed.

PRJT was measured using a handheld dynamometer (HHD) (Power Track II, JTECH Medical, Salt Lake City, UT, USA). The center of the sensor pad was touched to the head of the second metatarsal bone, and the minimum force required to apply maximum passive dorsiflexion was measured using the HHD. The force was applied to produce dorsiflexion that was as slow as possible. The torque was calculated by multiplying the force measured by the HHD by the distance between the lower end of the medial malleolus, which was the axis of ankle plantar flexion, and the head of the second metatarsal bone. This value was regarded as the PRJT. The change in PRJT torque from before to after the intervention was calculated and analyzed.

PROM and PRJT were measured with the subjects in the supine position. The knee joint was placed in a position of maximum extension or 90° flexion. The order of the knee joint position and PROM and PRJT measurements was randomly set for each subject. Measurement was performed

immediately before and after irradiation in the intervention group, and immediately before and after resting in the control group.

In the intervention group, the presence or absence of such as things as redness and blisters at the irradiated site was examined by visual inspection. In addition, the subjects were interviewed to determine if they felt an excessive sensation of heat or other abnormalities. If redness, blisters, excessive sensation of heat, or other abnormalities were found, Hi-LPNR irradiation was discontinued.

The age, duration of illness, and PROM and PRJT before the intervention were compared between the two groups using the unpaired t-test; the gender distribution, paretic side, and diagnosis were compared between the two groups using the χ^2 test; and the APTS score was compared between the two groups using the Mann-Whitney test. The change in PROM and PRJT from before to after the intervention was compared between the two groups using the unpaired t-test to investigate the effect of Hi-LPNR.

In addition, the 95% confidence interval (CI) of the minimal detectable change (MDC₉₅) was calculated. MDC₉₅ and the standard error of measurement (SEM) necessary to calculate the MDC₉₅ were determined²⁰⁻²² with the equations below using test-retest data from a control group:

$$\text{SEM} = (\text{standard deviation of the difference in the value between test and retest measurements})/\sqrt{2}$$

$$\text{MDC}_{95} = 1.96 \times \sqrt{2} \times \text{SEM}.$$

When the knee was extended, the MDC₉₅ in PROM was 3.36°, and that in PRJT was 1.44N·m. When the knee was flexed, the MDC₉₅ in PROM was 2.45°, and that in PRJT was 1.07N·m. The effect of Hi-LPNR irradiation was investigated by comparing the average change in PROM and PRJT in each group with these MDC₉₅ values. In addition, the number of subjects in whom the change exceeded the MDC₉₅ was compared between the two groups using the χ^2 test.

The significance level was set at 5% in all tests. All analyses were performed using statistical analysis software (IBM SPSS Statistics, Version 23.0 J, IBM Japan, Tokyo, Japan).

RESULTS

Subject characteristics are shown in Table 1, the changes in PROM and PRJT from before to after the intervention are shown in Table 2, and the number of subjects with changes exceeding MDC₉₅ is shown in Table 3.

There was no significant difference in age, gender, diagnosis, paretic side, duration of illness, or APTS score between the two groups ($p > 0.05$; Table 1). No subjects met any of the exclusion criteria. In all subjects, side effects and other abnormalities did not occur. There were no subjects who discontinued Hi-LPNR irradiation. Under the irradiation conditions of the present study, adverse events were not observed, and this suggested that the intervention was safe.

In the knee extended position, the mean changes in PROM were 2.70° (95% CI: 0.95–4.45°) and –0.50° (95% CI: –1.73–0.73°) in the intervention and control groups, respectively. The mean changes in PROM were significantly greater in the intervention group than in the control group ($p = 0.003$). But the mean value of PROM was smaller than

the MDC₉₅ in both groups (Table 2). The mean changes in PRJT were –1.42 N·m (95% CI: –1.83–1.01 N·m) and –0.26 N·m (95% CI: –0.79–0.26 N·m) in the intervention and control groups, respectively, and the change was significantly greater in the intervention group than in the control group ($p = 0.001$). But the mean in PRJT was smaller than the MDC₉₅ in both groups (Table 2). The change in PROM exceeded the MDC₉₅ in four subjects in the intervention group and one subject in the control group (Table 3), but the difference was not significant between the groups ($p = 0.12$). The change in PRJT exceeded the MDC₉₅ in five subjects in the intervention group and one subject in the control group (Table 3), but the difference was not significant between the groups ($p = 0.051$).

In the knee flexed position, the mean changes in PROM were 3.70° (95% CI: 1.82–5.58°) and 0.70° (95% CI: –0.20–1.60°) in the intervention and control groups, respectively, and the change was significantly greater in the intervention group than in the control group ($p = 0.004$). The mean changes in PRJT were –2.38 N·m (95% CI: –3.15–1.60 N·m) and –0.31 N·m (95% CI: –0.70–0.08 N·m) in the intervention and control groups, respectively, and the change was significantly greater in the intervention group than in the control group ($p < 0.001$). The mean changes in PROM and PRJT exceeded the MDC₉₅ in the intervention group, but they were smaller than the MDC₉₅ in the control group (Table 2). The change in PROM exceeded the MDC₉₅ in six subjects in the intervention group and one subject in the control group (Table 3), and the difference was significant between the groups ($p = 0.019$). The change in PRJT exceeded the MDC₉₅ in nine subjects in the intervention group and one subject in the control group (Table 3), and the difference was significant between the groups ($p < 0.001$).

DISCUSSION

The rays emitted by Hi-LPNR therapy equipment have characteristic wavelengths and intensities (output). Hi-LPNR therapy equipment emits complex rays of 600–1,600 nm^{2, 7}. Rays with a wavelength shorter than 600 nm are likely to be absorbed by blood hemoglobin, and those with a wavelength longer than 1,600 nm are likely to be absorbed by water molecules. Accordingly, rays emitted by Hi-LPNR therapy equipment are unlikely to be absorbed by blood (hemoglobin and water molecules) or influenced by heat dispersion through blood flow and thus are superior in terms of heating the irradiated region. The origin of the myotendinous tendon junction of the triceps surae muscle, which was irradiated in this study, is present just under the skin, and Hi-LPNR may have reached the gastrocnemius and soleus muscle. However, there have been no reports of the depth of penetration of Hi-LPNR irradiation into the body, and the details remain to be investigated.

In general, polarized light includes linear, right-handed, and left-handed circularly polarized light, but the effects due to the presence or absence of each type of polarized light and the differences among the types of polarized lights have not been investigated. The influence of differences in polarized light on skin wound healing and spinal cord injury in animals has been reported²³⁻²⁵, but there have been no reports

Table 1. Subject characteristics

	Total (n = 20)	Intervention (n = 10)	Control (n = 10)
Age (years)	77.7 (8.4) [73.7–81.6]	77.7 (8.5) [71.6–83.8]	77.6 (8.8) [71.3–83.9]
Gender (male/female)	11/9	4/6	7/3
Diagnosis			
Cerebral infarction	16	8	8
Intracerebral hemorrhage	4	2	2
Paretic side (right/left)	32/42	4/6	5/5
Duration of illness (d)	2,036.9 (3,369.5) [459.9–3,613.8]	3,127.8 (4,481.6) [–78.2–6,333.8]	945.9 (1,113.2) [149.6–1,742.2]
Ext			
APTS [#] score (0/1/2/3/4)			
SR	8/7/4/1/0	3/4/3/0/0	5/3/1/1/0
MR	5/10/4/1/0	2/6/2/0/0	3/4/2/1/0
FR	0/11/8/1/0	0/5/5/0/0	0/6/3/1/0
Baseline PROM [#] (°)	0.2 (3.9) [–1.7–2.0]	0.0 (4.0) [–2.9–2.9]	0.3 (3.9) [–2.5–3.1]
Baseline PRJT [#] (N·m)	5.8 (1.2) [5.3–6.4]	5.5 (1.1) [4.7–6.2]	6.2 (1.3) [5.3–7.2]
Flex			
APTS [#] score (0/1/2/3/4)			
SR	9/6/3/1/1	3/4/3/0/0	6/2/0/1/1
MR	9/9/2/0/0	4/6/0/0/0	5/3/2/0/0
FR	0/14/6/0/0	0/9/1/0/0	0/5/5/0/0
Baseline PROM [#] (°)	10.4 (6.7) [7.2–13.5]	12.1 (7.6) [6.7–17.6]	8.6 (5.5) [4.6–12.6]
Baseline PRJT [#] (N·m)	5.6 (1.3) [4.9–6.2]	5.1 (1.1) [4.4–5.9]	6.0 (1.4) [5.0–7.0]

Values are either n or mean (standard deviation) [95% confidence interval].

[#]Lower limb paretic side

APTS: Ankle Plantar Flexors Tone Scale; SR: stretch reflex; MR: middle range resistance; FR: final range resistance; Ext: knee extended position; Flex: knee flexed position; PROM: passive range of motion; PRJT: passive resistive joint torque; n.s.: not significant

Table 2. Change in passive range of motion and passive resistive joint torque and minimal detectable change at the 95% confidence level

		PROM (°)		PRJT (N·m)	
		Mean (SD) 95%CI	MDC ₉₅ [†]	Mean (SD) 95%CI	MDC ₉₅ [†]
Ext	Intervention	2.70 (2.45) 0.95 to 4.45	3.36	–1.42 (0.6) –1.83 to 1.01	1.44
	Control	–0.50 (1.72) –1.73 to 0.73		–0.26 (0.74) –0.79 to 0.26	
Flex	Intervention	3.70 (2.63) 1.82 to 5.58	2.45	–2.38 (1.08) –3.15 to –1.60	1.07
	Control	0.70 (1.25) –0.20 to 1.60		–0.31 (0.54) –0.70 to 0.08	

Ext: knee extended position; Flex: knee flexed position; PROM: passive range of motion; PRJT: passive resistive joint torque; CI: confidence interval

[†]MDC₉₅ was calculated using data from a control group.

**p<0.001

Table 3. Number of subjects that showed a change exceeding the minimal detectable change at the 95% confidence level

		PROM				PRJT			
		\geq MDC ₉₅	$<$ MDC ₉₅			\geq MDC ₉₅	$<$ MDC ₉₅		
Ext	Intervention	4	6]	n.s.	5	5]	n.s.
	Control	1	9			1	9		
Flex	Intervention	6	4]	*	9	1]	**
	Control	1	9			1	9		

MDC₉₅: Minimal detectable change at the 95% confidence level; PROM: passive range of motion; PRJT: passive resistive joint torque; Ext: knee extended position; Flex: knee flexed position, n.s.: not significant * $p < 0.05$, ** $p < 0.001$

on the effect of the presence or absence of polarization and differences among polarized lights on human muscle tone. We could not describe the effect of linearly polarized light on muscle tone, and it remains important to clarify this in detail.

The ankle plantar flexor muscle is comprised of biarticular muscles (gastrocnemius and plantaris muscles) and monoarticular muscles (the soleus, tibialis posterior, long fibular, and short fibular muscles). The main muscles are the gastrocnemius and soleus muscles, which account for 80% of the plantar flexion muscle force²⁶). Accordingly, hypertonia of the gastrocnemius and soleus muscles may have a marked influence on overall plantar flexor muscle tone.

The gastrocnemius originates in the medial and lateral epicondyles of the femur, and the soleus muscle originates in the posterior tibial soleal line and fibular head. Both muscles become the Achilles tendon and end at the calcaneal tuberosity. The gastrocnemius is a biarticular muscle that crosses the knee and ankle joints, and the soleus muscle is a monoarticular muscle that crosses only the ankle joint. When the knee joint is extended with a fixed ankle joint position, the biarticular muscle is lengthened, but the monoarticular muscle length is not changed; therefore, the tone of the plantar flexors in this position reflects the tone of both the soleus and the gastrocnemius. On the other hand, when the knee joint is flexed, the tone of the plantar flexors mainly reflects the tone of the monoarticular muscle, because the biarticular muscle is relaxed.

In both knee positions, the mean changes in PROM and PRJT were greater in the intervention group than in the control group. In addition, in the intervention group, the mean changes in PROM and PRJT in the knee flexed position exceeded the MDC₉₅. Furthermore, the number of subjects with changes exceeding the MDC₉₅ was significantly greater in the intervention group than in the control group. These findings suggest that Hi-LPNR irradiation applied to the posterior surface of the crus improved muscle extensibility and reduced the resistance to passive muscle stretch for the monoarticular plantar flexor muscles, mainly the soleus muscle.

The MDC represents the marginal field for which the change between two measured values in repeated measurement, such as test-retest, is due to measurement error, and the 95% CI of the MDC, MDC₉₅, is generally used²²). Changes smaller than the MDC₉₅ could be due to measurement error and therefore cannot be regarded as true changes²⁰). Changes

exceeding the MDC₉₅ can be interpreted as true changes exceeding the error range, i.e., changes due to the effect of the intervention.

The main factor limiting ankle dorsiflexion is limited stretch of the ankle plantar flexor muscles, and PROM reflects the maximum stretch of the ankle plantar flexor muscles. In both knee positions, the mean change in PROM was significantly greater in the intervention group than in the control group. However, in the knee extended position, the mean change in PROM was smaller than the MDC₉₅ in both groups, and there was no significant difference in the number of subjects exceeding the MDC₉₅. By contrast, in the knee flexed position, the mean change in PROM exceeded the MDC₉₅ in the intervention group, and the number of subjects exceeding the MDC₉₅ was significantly higher in the intervention group than in the control group. Therefore, we suggest that the increase in PROM in the knee flexed position in the intervention group was due to Hi-LPNR irradiation, but that the increase in PROM in the knee extended position was due to measurement error, and we therefore suggest that Hi-LPNR irradiation increased ankle plantar flexor muscle extensibility and increased PROM in the knee flexed position.

PRJT reflects the resistance to maximum passive muscle stretch. In both knee positions, the mean change in PRJT was significantly greater in the intervention group than in the control group. The mean change was greater than the MDC₉₅ in the knee flexed position in the intervention group but smaller than the MDC₉₅ in both knee positions in the control group, and the number of subjects with changes exceeding the MDC₉₅ in the knee flexed position was greater in the intervention group than in the control group. These results suggest that Hi-LPNR irradiation increased ankle plantar flexor muscle extensibility and reduced resistance to maximum passive muscle stretch in the knee flexed position, decreasing PRJT.

Muscle tone involves a non-neural component that is based on the flexibility of fibrous tissue, such as muscle fibers and fascia, and a neural component that is based on reflex muscle contraction, such as the stretch reflex. In disorders of upper motor neurons, including CVD, the stretch reflex is enhanced due to impairment of the descending inhibitory system associated with central nervous system abnormality. Spasticity occurs due to an exaggerated stretch reflex resulting from upper motor neuron lesions and is considered velocity-dependent muscle hypertonia²⁷⁻²⁹). When spasticity

persists for a prolonged time without appropriate treatment, the frequency of muscle stretch decreases, and this leads to reduction of muscle fiber and fascial flexibility, which are non-neural components of muscle tone. The threshold of muscle spindles decreases with decreases in the non-neural component of muscle tone, thus increasing the stimulation of muscle spindles and enhancing the stretch reflex, leading to a vicious cycle of hypertonia. The neural and non-neural components of muscle tone are therefore different but closely associated with each other. In this study, PROM and PRJT were measured while the ankle plantar flexor muscles were slowly stretched. Thus, these measures may have captured mainly the non-neural component of muscle tone, but they would also have been influenced by the neural component if reflex muscle contraction occurred during measurement (i.e., during passive stretch). Therefore, PROM and PRJT were measured in the presence of both components of muscle tone. It is difficult to measure the muscle tone while strictly separating the neural and non-neural components in the procedure, and this is a limitation of this study.

As a muscle is heated, there is a reduction of stretch receptivity of muscle spindles and inhibition of muscle tone (Ib inhibition) caused by an increase in Golgi tendon organ activation-induced discharge by Ib fibers³⁰, suggesting that Hi-LP NR irradiation may reduce the stretch receptivity of muscle spindles and subsequently reduce the excitation of the secondary terminals, thus inhibiting the neural component of muscle tone. Heat stimulation also increases extensibility of muscle fibers and fascia³¹. It has been suggested that heat increases the flexibility of collagen in fasciae and tendons and the extensibility of muscle fibers, thus improving the non-neural component of muscle tone.

We acknowledge that this study had some limitations. The rater performing the measurements of PROM and PRJT and the Hi-LP NR irradiation was not blind to group allocation. In addition, the present study only included 20 subjects, and in the future, it will be necessary to increase the number of subjects. In conclusion, there was a significant difference in the changes in PROM and PRJT between the intervention and control group. PROM in a flexed knee position and PRJT in both a flexed and extended knee position changed more than the MDC₉₅ in the intervention group. In conclusion, Hi-LP NR irradiation has an inhibitory effect on muscle tone in patients with CVD.

ACKNOWLEDGEMENTS

This research was supported by a Grant-in-Aid for Young Scientists (B) from the Japan Society for the Promotion of Science (Research Number 24700556). The funding source had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors declare that they have no conflicts of interest.

REFERENCES

- Asagai Y: Application of low reactive-level laser therapy (LLLT) in the functional training of cerebral palsy patients. *Laser Ther*, 1994, 6: 195–202. [CrossRef]
- Demura S, Yamaji S, Ikemoto Y: Effect of linear polarized near-infrared light irradiation on flexibility of shoulder and ankle joints. *J Sports Med Phys Fitness*, 2002, 42: 438–445. [Medline]
- Demura T, Demura S, Aoki H, et al.: Effect of linear polarized near-infrared light irradiation and light exercise on muscle performance. *J Physiol Anthropol*, 2011, 30: 91–96. [Medline] [CrossRef]
- Yokoyama K, Oku T: Rheumatoid arthritis-affected temporomandibular joint pain analgesia by linear polarized near infrared irradiation. *Can J Anaesth*, 1999, 46: 683–687. [Medline] [CrossRef]
- Yokoyama K, Sugiyama K: Temporomandibular joint pain analgesia by linearly polarized near-infrared irradiation. *Clin J Pain*, 2001, 17: 47–51. [Medline] [CrossRef]
- Huang D, Gu YH, Liao Q, et al.: Effects of linear-polarized near-infrared light irradiation on chronic pain. *Scientific World Journal*, 2012. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3349215/pdf/TSWJ2012-567496.pdf> (Accessed Apr. 27, 2015)
- Sasaki A, Touma Y, Ishino Y, et al.: Linear polarized near-infrared irradiation stimulates mechanical expansion of the rat sagittal suture. *Luminescence*, 2003, 18: 58–60. [Medline] [CrossRef]
- Kashima K, Watanabe N, Higashinaka S, et al.: Changes in sensory and pain perception thresholds after linear polarized near-infrared light radiation in the trigeminal region. *Cranio*, 2005, 23: 174–178. [Medline]
- Hwang H, Ha K: The effect of blood velocity change after SUPER LIZER on rotator cuff pain. *J Phys Ther Sci*, 2012, 24: 851–853. [CrossRef]
- Muneshige H, Toda K, Ma D, et al.: Antinociceptive effect of linear polarized 0.6 to 1.6 microm irradiation of lumbar sympathetic ganglia in chronic constriction injury rats. *J Rehabil Res Dev*, 2006, 43: 565–572. [Medline] [CrossRef]
- Momota Y, Kani K, Takano H, et al.: High-wattage pulsed irradiation of linearly polarized near-infrared light to stellate ganglion area for burning mouth syndrome. *Case Rep Dent*, 2014: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4216686/pdf/CRID2014-171657.pdf> (Accessed Jun. 23, 2015)
- Alayat MS, Elsodany AM, El Fiky AA: Efficacy of high and low level laser therapy in the treatment of Bell's palsy: a randomized double blind placebo-controlled trial. *Lasers Med Sci*, 2014, 29: 335–342. [Medline] [CrossRef]
- Fiore P, Panza F, Cassatella G, et al.: Short-term effects of high-intensity laser therapy versus ultrasound therapy in the treatment of low back pain: a randomized controlled trial. *Eur J Phys Rehabil Med*, 2011, 47: 367–373. [Medline]
- Santamato A, Solfrizzi V, Panza F, et al.: Short-term effects of high-intensity laser therapy versus ultrasound therapy in the treatment of people with subacromial impingement syndrome: a randomized clinical trial. *Phys Ther*, 2009, 89: 643–652. [Medline] [CrossRef]
- Mirshahi A, Lashay A, Roozbahani M, et al.: Pain score of patients undergoing single spot, short pulse laser versus conventional laser for diabetic retinopathy. *Graefes Arch Clin Exp Ophthalmol*, 2013, 251: 1103–1107. [Medline] [CrossRef]
- Kheshe AR, Alayat MS, Ali MM: High-intensity versus low-level laser therapy in the treatment of patients with knee osteoarthritis: a randomized controlled trial. *Lasers Med Sci*, 2014, 29: 1371–1376. [Medline] [CrossRef]
- Alayat MS, Atya AM, Ali MM, et al.: Long-term effect of high-intensity laser therapy in the treatment of patients with chronic low back pain: a randomized blinded placebo-controlled trial. *Lasers Med Sci*, 2014, 29: 1065–1073. [Medline] [CrossRef]
- Takeuchi N, Tanaka E, Kuwabara T, et al.: Effects of linear polarized near-infrared ray irradiation on tibial nerve and gastrocnemius of hemiplegic patients: a randomized controlled trial. *Rigaku Ryohogaku*, 2008, 35: 13–22 [in Japanese].
- Takeuchi N, Kuwabara T, Usuda S: Development and evaluation of a new measure for muscle tone of ankle plantar flexors: the ankle plantar flexors tone scale. *Arch Phys Med Rehabil*, 2009, 90: 2054–2061. [Medline] [CrossRef]
- Faber MJ, Bosscher RJ, van Wieringen PC: Clinimetric properties of the performance-oriented mobility assessment. *Phys Ther*, 2006, 86: 944–954. [Medline]
- Weir JP: Quantifying test-retest reliability using the intraclass correlation coefficient and the SEM. *J Strength Cond Res*, 2005, 19: 231–240. [Medline]
- Lin KC, Fu T, Wu CY, et al.: Assessing the stroke-specific quality of life for outcome measurement in stroke rehabilitation: Minimal detectable change and clinically important difference. *Health Qual Life Outcomes*: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3034658/pdf/1477-7525-9-5.pdf>. (Accessed Apr. 27, 2015)
- Tada K, Ikeda K, Tomita K: Effect of polarized light emitting diode irradiation on wound healing. *J Trauma*, 2009, 67: 1073–1079. [Medline]

- [CrossRef]
- 24) Ribeiro MS, Da Silva DF, De Araújo CE, et al.: Effects of low-intensity polarized visible laser radiation on skin burns: a light microscopy study. *J Clin Laser Med Surg*, 2004, 22: 59–66. [Medline] [CrossRef]
 - 25) Ando T, Sato S, Kobayashi H, et al.: Low-level laser therapy for spinal cord injury in rats: effects of polarization. *J Biomed Opt*, 2013, 18: 098002. [Medline] [CrossRef]
 - 26) Murray MP, Guten GN, Baldwin JM, et al.: A comparison of plantar flexion torque with and without the triceps surae. *Acta Orthop Scand*, 1976, 47: 122–124. [Medline] [CrossRef]
 - 27) Satkunam LE: Rehabilitation medicine: 3. Management of adult spasticity. *CMAJ*, 2003, 169: 1173–1179. [Medline]
 - 28) Lance JW: Symposium synopsis. Spasticity: In *Disordered Motor Control*. Chicago: Yearbook Medical, 1980, pp 487–489.
 - 29) Takeuchi N, Kuwabara T, Usuda S: Association between the tones of the ankle plantar flexors and duration of illness in patients with cerebrovascular disease. *J Phys Ther Sci*, 2012, 24: 1289–1294. [CrossRef]
 - 30) Mense S: Effects of temperature on the discharges of muscle spindles and tendon organs. *Pflugers Arch*, 1978, 374: 159–166. [Medline] [CrossRef]
 - 31) Cameron MH: Thermal Agents: Cold and Heat. In: *Physical Agents in Rehabilitation—From Research to Practice*, 3rd ed. Missouri: Saunders Elsevier, 2009, pp 131–138.

Appendix 1. Ankle Plantar Flexors Tone Scale¹⁹⁾

Stretch reflex

- 0: No twitch.
 - 1: Twitch and no clonus.
 - 2: Mild clonus, persisting <3 s.
 - 3: Moderate clonus, persisting 3–10 s.
 - 4: Severe clonus, persisting >10 s.
-

Middle range resistance

- 0: No resistance.
 - 1: Mild resistance, slight increase in resistance.
 - 2: Moderate resistance, greater increase in resistance.
 - 3: Severe resistance, considerable increase in resistance, but able to achieve passive movement.
 - 4: Passive movement is difficult.
-

Final range resistance

- 0: No resistance.
 - 1: Mild resistance, slight increase in resistance.
 - 2: Moderate resistance, greater increase in resistance.
 - 3: Severe resistance, considerable increase in resistance, but able to maintain final position.
 - 4: Unable to maintain final position or passive movement is difficult.
-