

# Characteristics, biosynthesis, decomposition, metabolism and functions of the garlic odour precursor, *S*-allyl-*L*-cysteine sulfoxide (Review)

YUSUKE YAMAGUCHI and HITOMI KUMAGAI

Department of Chemistry and Life Science, Nihon University, Fujisawa-shi 252-0880, Japan

Received September 30, 2019; Accepted November 6, 2019

DOI: 10.3892/etm.2019.8385

**Abstract.** *S*-Allyl-*L*-cysteine sulfoxide (ACSO) is an odour precursor in garlic bulbs. One plausible pathway for the biosynthesis of ACSO involves *S*-2-carboxypropyl glutathione produced from glutathione and methacrylic acid via valine or from  $\gamma$ -glutamyl cysteine. The elimination of glycine and glutamic acid from *S*-2-carboxypropyl glutathione produces *S*-2-carboxypropyl cysteine, which is converted to *S*-allyl cysteine by decarboxylation and oxidation. *S*-Allyl cysteine is also biosynthesized via the elimination of glutamic acid from  $\gamma$ -glutamyl *S*-allyl cysteine by  $\gamma$ -glutamyl transpeptidase. The sulfur oxidation of *S*-allyl cysteine by flavin-containing monooxygenase forms ACSO. When cells are damaged by slicing or grating, ACSO in the cytoplasm or cytoplasmic vesicle is immediately converted to allylsulfenic acid, pyruvic acid, and ammonia by alliinase (C-S lyase), which is located in the vacuoles of vascular bundle sheath cells. Two molecules of allylsulfenic acid form diallyl thiosulfinate (allicin), which exhibits potent antimicrobial activity. Allicin eventually yields garlic odour compounds, such as diallyl disulfide (DADS) and diallyl trisulfide (DATS). Although these sulfides are known to exert various physiological functions, their strong odour limits their use in foods. On the other hand, ACSO is water-soluble and odourless and enhances sweet, salty, and umami tastes, characteristics of which are desirable for food additives. Upon consumption, ACSO is primarily absorbed from the small intestine in the intact form, but is also partly decomposed to allylsulfenic acid, pyruvic acid and ammonia. Allylsulfenic acid is then further converted to DADS and diallyl monosulfide (DAS). ACSO has numerous *in vivo* functions, such as the prevention of diabetes, myocardial ischaemia, hepatic injury, platelet aggregation and

blood ethanol elevation. Although some of these effects may be attributed to its metabolites, ACSO itself contributes to many of these physiological functions.

## Contents

1. Introduction
2. Characteristics of ACSO
3. Biosynthesis of ACSO
4. Content of ACSO and related compounds in garlic
5. Decomposition of ACSO and other sulfoxides
6. Metabolism of ACSO
7. Physiological functions of ACSO
8. Conclusions

## 1. Introduction

Garlic, which is well-known for its characteristic odour, has various biological functions, such as antioxidant and antimicrobial effects (1-3). It is also known to be able to prevent atherosclerosis (4), diabetes (5), cardiac hypertrophy (5), thrombosis, inflammation, hypertension and even cancer (6-9). Garlic itself does not have any potent odour; however, when garlic bulbs are cut or crushed, a sulfuric odour is instantaneously produced. The major odour precursor is *S*-(+)-allyl-*L*-cysteine sulfoxide (*S*-2-propenyl cysteine sulfoxide; ACSO or alliin). When cells are damaged, ACSO is decomposed to produce garlic odour compounds, such as diallyl disulfide (di-2-propenyl disulfide; DADS) and diallyl trisulfide (di-2-propenyltrisulfide; DATS). These sulfides are the functional compounds that are responsible for the beneficial physiological activities of garlic (10); however, as they are lipophilic, volatile and have a strong odour, their use in foods is limited. On the other hand, ACSO is hydrophilic and odourless, rendering it useful as an additive in a wide range of foods. In light of recent findings, this review summarizes the characteristics and physiological functions of ACSO *in vivo*.

## 2. Characteristics of ACSO

ACSO is an amino acid that contains an amino group and carboxyl group in its structure with a molecular weight

---

*Correspondence to:* Professor Hitomi Kumagai, Department of Chemistry and Life Science, Nihon University, 1866 Kameino, Fujisawa-shi 252-0880, Japan  
E-mail: kumagai.hitomi@nihon-u.ac.jp

**Key words:** garlic, odor precursor, *S*-allyl-*L*-cysteine sulfoxide, alliin, sulfide, biosynthesis, decomposition, metabolism, physiological function




Figure 1. Putative pathways for the biosynthesis of ACSO. ACSO, *S*-allyl-L-cysteine sulfoxide.

of 177.22. According to the substance information in Chemical Abstracts (SciFinder Scholar), the physicochemical properties of ACSO are estimated. The density, pKa, and boiling point of ACSO are predicted to be 1.354 g/ml at 20°C, 1.88 at 25°C and 416°C, respectively, according to calculations of the Advanced Chemistry Development (ACD/Labs) software v11.02. Furthermore, the melting point is 165°C according to PhysProp data obtained from the Syracuse Research Corporation. The mass solubilities at pH 1, 2, 3-6, 7, 8, 9 and 10, as predicted by the ACD/Labs Software v11.02, are 120, 35, 27, 32, 76, 535 and 1,000 g/l, respectively, and ACSO is quite soluble in solutions above pH 9. ACSO itself is odourless and enhances sweet, salty and umami tastes (11). Therefore, ACSO may be one of the key components that produce a rich flavour in garlic-containing foods.

### 3. Biosynthesis of ACSO

The biosynthesis of ASCO is presented in Fig. 1. Although several biosynthetic pathways for ACSO have been proposed, its biosynthesis remains under debate (12-14). Tracing the radioactivity of labelled valine reveals that the radioactivity is located in the carboxypropyl group of *S*-2-carboxypropyl glutathione and *S*-2-carboxypropyl cysteine (15). *S*-2-Carboxypropyl glutathione is biosynthesized from glutathione (a tripeptide comprised of glycine, cysteine, and glutamic acid) with methacrylic acid produced from valine. However, two possible pathways forming *S*-2-carboxypropyl cysteine have been reported. One mechanism involves  $\gamma$ -glutamyl *S*-2-carboxypropyl cysteine produced from *S*-2-carboxypropyl glutathione via




Figure 2. Localization of compounds and enzymes involved in biosynthesis of ACSO in a cell (13,17). ACSO, *S*-allyl-L-cysteine sulfoxide.

elimination of glycine, while the other involves generation from cysteine and methacrylic acid without passing through glutathione. These routes have also been proposed for the biosynthesis of *trans*-1-propenyl cysteine in onion (12). *S*-2-Carboxypropyl glutathione may also be produced from cysteine, as  $\gamma$ -glutamyl cysteine is detected in garlic (16). As decarboxylation and oxidation of the carboxypropyl group produce an allyl group (13),  $\gamma$ -glutamyl *S*-allyl cysteine can be produced from  $\gamma$ -glutamyl *S*-2-carboxypropyl cysteine, and *S*-allyl cysteine can be produced from *S*-2-carboxypropyl cysteine. The *S*-oxidation of  $\gamma$ -glutamyl *S*-allyl cysteine forms  $\gamma$ -glutamyl *S*-allyl cysteine sulfoxide, while that of *S*-allyl cysteine forms ACSO. Flavin-containing monooxygenase (FMO) in the cytosol is responsible for the *S*-oxidation in the biosynthesis of ACSO in the presence of nicotinamide adenine dinucleotide phosphate (NADPH) and flavin adenine dinucleotide (FAD) (17). On the other hand, the recombinant  $\gamma$ -glutamyl transpeptidase (GGT) prepared by cloning genes from garlic exhibits a high specificity for  $\gamma$ -glutamyl *S*-allyl cysteine, but not for  $\gamma$ -glutamyl *S*-allyl cysteine sulfoxide (18). Therefore, *S*-allyl cysteine produced from the deglutamylation of  $\gamma$ -glutamyl *S*-allyl cysteine forms ACSO via its *S*-oxidation, and the pathway via  $\gamma$ -glutamyl *S*-allyl cysteine sulfoxide to ACSO may not exist in garlic. The observations that ACSO can be produced with the addition of *S*-allyl cysteine in garlic callus (14) and that *S*-allyl cysteine is found in garlic (19,20), also support this biosynthetic route from *S*-allyl cysteine to ACSO. As the addition of serine and allyl thiol produces ACSO (12), the pathway from serine via *S*-allyl cysteine to form ACSO is also probable.

The location of ACSO biosynthesis has not yet been convincingly identified.  $\gamma$ -Glutamyl peptides reside in the cytoplasm and cysteine sulfoxides in the cytoplasm or cytoplasmic vesicle, while  $\gamma$ -glutamyl cysteine resides in the chloroplast (13) (Fig. 2). As for the enzymes required for biosynthesis,  $\gamma$ -glutamyl cysteine synthetase is located in the chloroplast,

and  $\gamma$ -glutamyl transpeptidase and flavin-containing monooxygenase are located in the cytoplasm (13).

#### 4. Content of ACSO and related compounds in garlic

The majority of the organosulfur compounds in garlic are  $\gamma$ -glutamyl *S*-alkenyl cysteines and cysteine sulfoxides (21,22), with ACSO being the major component. The ACSO content in fresh garlic is 3-14 mg/g fresh weight (20,23,24) and 20-30 mg/g dry weight (25,26). As regards other cysteine sulfoxides, the contents of *S*-methyl-L-cysteine sulfoxide (MCSO or methiin), *trans*-*S*-1-propenyl-L-cysteine sulfoxide (PeCSO or isoalliin) and cycloalliin are 0.5-2, 0.1-1.2 and 0.5-1.5 mg/g fresh weight, respectively (24).  $\gamma$ -L-Glutamyl-*S*-allyl-L-cysteine content in garlic bulbs is 2-7 mg/g fresh weight (21,24) and 3-20 mg/g dry weight (26), while the  $\gamma$ -L-glutamyl-*S*-(*trans*-1-propenyl)-L-cysteine content is 3-9 mg/g fresh weight (21,24) and 5-15 mg/g dry weight (26).  $\gamma$ -L-Glutamyl-*S*-methyl-L-cysteine is present at 0.1-0.4 mg/g fresh weight (24).

Sulfur fertilization increases the ACSO content in garlic cloves, while nitrogen fertilization decreases this content (27). Irrespective of the fertilization conditions, the ACSO content increases during storage at 20°C and decreases during storage at temperatures >35°C (28). Drying at 60°C does not affect the ACSO content (29).

ACSO accumulates in leaves during early growth stages and translocates to the bulbs during bulb formation (30). Approximately 85% of ACSO is present in the bulbs, with approximately 12% in the leaves and 2% in the roots, while  $\gamma$ -glutamyl cysteines are found only in the bulbs (24). The  $\gamma$ -glutamyl-cysteine content in the bulbs increases a few weeks prior to harvest (25,31). The ACSO content in the bulbs also begins to increase a few weeks prior to harvest (25,31) but also further increases during the curing process (drying process after harvest for storage) (21). The content of ACSO in the leaves, however, remains relatively stable from 6 weeks to 1 week prior to harvest and subsequently decreases.




Figure 3. Formation of volatile compounds from ACSO. ACSO, *S*-allyl-L-cysteine sulfoxide.




Figure 4. Formation of volatile compounds from ACSO and MCSO. ACSO, *S*-allyl-L-cysteine sulfoxide; MCSO, *S*-methyl-L-cysteine sulfoxide.

## 5. Decomposition of ACSO and other sulfoxides

The decomposition of ACSO and other sulfoxides is illustrated in Figs. 3 and 4. When garlic bulbs are sliced or ground during

cooking, ACSO that resides in the cytoplasm or cytoplasmic vesicle of all cells (13) binds alliinase (C-S lyase; EC 4.4.1.4) that localizes in the vacuoles of vascular bundle sheath cells located around the veins or phloem (32). Alliinase belongs

to a class 1 family of pyridoxal-5'-phosphate-dependent enzymes (2) and is a homodimeric glycoprotein, consisting of 448-amino-acid subunits with a molecular weight of 51,500 Da each (2). Two alliinase isozymes have been identified in garlic: One is specific for ACSO and PeCSO, and the other is specific for MCSO (22). ACSO is converted to allylsulfenic acid (2-propene-1-sulfenic acid;  $\text{CH}_2=\text{CHCH}_2\text{SOH}$ ) (33) and aminoacrylic acid by alliinase. Aminoacrylic acid is then spontaneously converted to pyruvic acid and ammonia. Two molecules of allylsulfenic acid produce diallyl thiosulfinate (*S*-prop-2-en-1-yl prop-2-ene-1-sulfinothioate; allicin), which exerts strong antimicrobial activity (34-37). The amount of allicin produced is approximately 0.1 to 7 mg/g fresh weight due to the abundance of ACSO in garlic (7,38). The incorporation of an additional allylsulfenic acid and the removal of allyl alcohol yield 1,3-di(prop-2-en-1-yl)trisulfane 1-oxide. When garlic bulbs are homogenized in water or in an alcohol solution, DADS and DATS are mainly produced from the diallyl thiosulfinate and 1,3-di(prop-2-en-1-yl)trisulfane 1-oxide. Volatile oil obtained from garlic products by steam distillation contains DADS and DATS as the major components (38). These sulfides are known to have various physiological activities (10,39); however, they are lipophilic and have a potent garlic odour. On the other hand, when garlic bulbs are homogenized in oil or in non-polar solvent, dithiins become dominant, 2-vinyl-4*H*-1,3-dithiin and 3-vinyl-4*H*-1,2-dithiin being about 0.4 mg/g and 0.17 mg/g, respectively (38). Other components produced are (*E*)-ajoene (0.07 mg/g), (*Z*)-ajoene (0.04 mg/g), DATS (0.07 mg/g), methyl allyl trisulfide (MATS) (0.06 mg/g) and DADS (0.03 mg/g).

As MCSO is the second most abundant cysteine sulfoxide in garlic, representing approximately 10% of the total cysteine sulfoxides, the compounds produced from MCSO contribute to the composition of volatile oil. MCSO is also decomposed by alliinase when garlic is sliced or grated to yield methylsulfenic acid (Fig. 4). Allylsulfenic acid produced from ACSO and methylsulfenic acid produced from MCSO form *S*-methyl prop-2-ene-1-sulfinothioate and *S*-prop-2-en-1-yl methanesulfinothioate, the latter being twice as abundant as the former, though allicin accounts for 60-90% of the total thiosulfates. The *S*-methyl prop-2-ene-1-sulfinothioate and *S*-prop-2-en-1-yl methanesulfinothioate form methyl allyl disulfide (MADS) and MATS. Importantly, the ratio of allyl groups and methyl groups in total thiosulfates almost directly coincides with that in total *S*-alk(en)yl-L-cysteine sulfoxides in garlic (40). Among the sulfides, DATS (0.8-1.1 mg/g) and DADS (0.5-1 mg/g) are the most abundant, with MATS (0.2-0.7 mg/g) and MADS (0.08-0.5 mg/g) as the next most abundant. Dimethyl trisulfide (DMTS; 0.01-0.1 mg/g) and dimethyl disulfide (DMDS; 0.001-0.06 mg/g) occur in smaller quantities.

## 6. Metabolism of ACSO

The metabolism of ACSO is illustrated in Fig. 5. When 10 ml of 100 mM ACSO/kg body weight is injected into the ligated loop of the small intestine of rats, both ACSO and pyruvic-acid concentrations in the portal vein increase to approximately 4 and 0.2 mM, respectively, after 30 min (41), indicating that ACSO is mostly absorbed from the small intestine into the

blood in the intact form; however, it is also partly converted to allylsulfenic acid, pyruvic acid and ammonia. In addition, mixing ACSO with crude proteins extracted from the small intestine of rats yields DADS and DAS as volatile compounds (41), supporting the assumption that ACSO is decomposed to allylsulfenic acid and two molecules of allylsulfenic acid produce DADS and DAS via allicin. Therefore, ACSO may follow the same fate as allicin and DADS in the body. When garlic oil is consumed, DADS, DAS and DMDS are detected in urine (42). When garlic extract is consumed, *N*-acetyl-*S*-allyl-L-cysteine (allylmercapturic acid) is identified in urine (43,44). *N*-Acetyl-L-cysteine (mercapturic acid; MA) conjugates are commonly formed from glutathione conjugates. The glutathione conjugates are first transformed into L-cysteine conjugates via the elimination of glutamic acid by  $\gamma$ -glutamyl transpeptidase and glycine by cysteinylglycinase. Subsequently, L-cysteine conjugates are *N*-acetylated by cysteine-*S*-conjugate *N*-acetyltransferase to form MA conjugates (43). Allicin reacts with glutathione to produce *S*-allyl-mercaptogluthathione or with L-cysteine to produce *S*-allyl-mercaptocysteine *in vivo* (45). DADS undergoes nucleophilic substitution at the  $\alpha$ -carbon in a reaction with glutathione to yield *S*-allyl-gluthathione and allyl perthiol. This *S*-allyl-gluthathione is then converted to *S*-allyl cysteine by  $\gamma$ -glutamyl transpeptidase and cysteinylglycinase. The *S*-allyl cysteine is further transformed into *N*-acetyl-*S*-allyl-L-cysteine via *N*-acetylation by the cysteine-*S*-conjugate *N*-acetyltransferase (46).

## 7. Physiological functions of ACSO

Orally administered ACSO ameliorates the diabetic condition (47,48). When diabetes is induced in rats by an injection of alloxan monohydrate, fasting blood glucose levels are reduced, and insulin secretion is increased by the daily oral administration of 200 mg ACSO/kg body weight. In addition, orally administered ACSO reduces the generation of reactive aldehydes produced by lipid peroxidation, increasing catalase activity and glutathione content (47). Similar effects have been observed in mice with diet-induced obesity (DIO) (48). When these mice are fed a high-fat (60% kcal from fat) diet for 8 weeks and provided with water containing 0.1 mg ACSO/ml for a further 8 weeks, glucose homeostasis and insulin sensitivity are enhanced. The increase in blood glucose levels following the injection of glucose is suppressed, and the decrease in blood glucose level following the injection of insulin is promoted. In addition, ACSO administration improves the lipid profile and liver functions of the mice with DIO. ACSO treatment reduces the levels of triglycerides, low-density lipoprotein and non-esterized free fatty acid. It also decreases the activities of aspartate aminotransferase and alkaline phosphatase, and the levels of total protein and albumin in the blood (48).

Furthermore, ACSO has been shown to prevent myocardial ischaemia induced by isoproterenol in rats (49). When 40 or 80 mg ACSO/kg body weight is administered to rats for 5 weeks, the increase in the activities of creatine kinase, lactate dehydrogenase, aspartate transaminase and alanine transferase in serum, and their decrease in the heart due to the induction of myocardial ischaemia are both suppressed. In addition, ACSO administration suppresses the increase in 3-hydroxy 3-methyl glutaryl coenzyme A (HMG CoA)




Figure 5. Putative pathways for metabolism of ACSO. ACSO, S-allyl-L-cysteine sulfoxide.

reductase activity in the liver and the decrease in lecithin cholesterol acyl transferase activity in the liver and heart in these isoproterenol-treated rats (49). The increase in the levels of plasma thiobarbituric acid reactive substances and hydroperoxide in mice with myocardial-ischaemia-induced is also suppressed by ACSO administration.

ACSO has also been shown to suppress oxidative stress in rats with carbon tetrachloride ( $\text{CCl}_4$ )-induced hepatic injury (41). In rats with hepatic injury, following the oral administration of 50  $\mu\text{mol}$  ACSO for 7 days, the increased aspartate transaminase, alanine transaminase and lactate dehydrogenase activities as well as the amount of thiobarbituric acid reactive substances are suppressed. In addition, the decrease in glutathione levels and the activities of glutathione S-transferase and glutathione peroxidase are also suppressed. These observations indicate that ACSO prevents hepatic injury by enhancing the activities of phase II detoxifying enzymes

and reducing oxidative stress. As MCSO, S-ethyl-L-cysteine sulfoxide (ECSO) and S-allyl-L-cysteine (ACS) are not effective for the prevention of hepatic injury, the allyl and sulfoxide groups in ACSO are essential for this preventive effect. Moreover, ACSO promotes nuclear factor erythroid 2-related factor 2 (Nrf2) nuclear translocation in HepG2 cells, leading to the induction of antioxidant and detoxifying enzymes, and contributing to these preventive effects (41).

In garlic, some of the sulfides, such as MATS and DMTS, exhibit inhibitory activity against platelet aggregation *in vitro* (50). As thrombosis is triggered by platelet aggregation, these sulfides may be the factors responsible for the prevention of thrombosis following garlic consumption. The administration of ACSO to rats, however, results in greater inhibitory activity against platelet aggregation than DADS (51). The effect of ACSO is most potent at a dosage of 10 mg/kg body weight, and the duration of the effect persists for 16-24 h

following administration. Although DADS is produced during the absorption of ACSO from the small intestine (41), the functional component is assumed to be different, suggesting the *in vivo* production of potent ACSO metabolites that confer inhibitory activity against platelet aggregation.

Finally, ACSO suppresses the increase in blood ethanol concentration following the oral administration of ethanol and ACSO together (52). Although the supply of NAD<sup>+</sup> is the rate-limiting factor for the metabolism of ethanol to acetaldehyde, and NAD<sup>+</sup> is produced from NADH during the conversion of pyruvic acid to lactic acid by lactate dehydrogenase, pyruvate production during the absorption of ACSO from the small intestine is not responsible for this suppressive effect on blood ethanol elevation. The administration of pyruvic acid at levels equivalent to those of ACSO does not exert the suppressive effect on blood ethanol elevation. The mechanisms underlying the reduction in blood ethanol concentrations by ACSO involve the promotion of ethanol metabolism in the liver and the suppression of ethanol absorption from the gut (52). Specifically, ACSO induces the activities of alcohol dehydrogenase, which converts ethanol into acetaldehyde, and aldehyde dehydrogenase, which converts acetaldehyde into acetic acid. In addition, the injection of ACSO into the ligated stomach or small intestine together with ethanol results in the suppression of the increase in ethanol concentration in the portal vein, indicating overall suppression of ethanol absorption. Garlic extract rich in ACSO (Garlic-H) exhibits an effect similar to ACSO itself, while garlic extract that does not contain ACSO (Garlic-N) exerts less-potent effects compared to Garlic-H (52).

## 8. Conclusions

On the whole, as discussed in this review article, the physiological functions of ACSO are extensive and can be applied to promote human health. As ACSO is water-soluble and odourless and enhances taste, the addition of this garlic component to food products will provide both a rich flavour and may confer numerous health benefits.

## Acknowledgements

Not applicable.

## Funding

No funding was received.

## Availability of data and materials

Not applicable.

## Authors' contributions

YY and HK conceived and designed this review article and wrote the manuscript. Both authors have read and approved the final manuscript.

## Ethics approval and consent to participate

Not applicable.

## Patient consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References

- Goncagul G and Ayaz E: Antimicrobial effect of garlic (*Allium sativum*) and traditional medicine. *J Anim Vet Adv* 9: 1-4, 2010.
- Touloupakis E and Ghanotakis DF: Nutraceutical use of garlic sulfur-containing compounds. *Adv Exp Med Biol* 698: 110-121, 2010.
- Rahman MS, Al-Sheibani HI, Al-Riziqi MH, Mothershaw A, Guizani N and Bengtsson G: Assessment of the anti-microbial activity of dried garlic powders produced by different methods of drying. *Int J Food Prop* 9: 503-513, 2006.
- Durak I, Aytac B, Atmaca Y, Devrim E, Avci A, Erol C and Oral D: Effects of garlic extract consumption on plasma and erythrocyte antioxidant parameters in atherosclerotic patients. *Life Sci* 75: 1959-1966, 2004.
- Padiya R, Chowdhury D, Borkar R, Srinivas R, Pal Bhadra M and Banerjee SK: Garlic attenuates cardiac oxidative stress via activation of PI3K/AKT/Nrf2-Keap1 pathway in fructose-fed diabetic rat. *PLoS One* 9: e94228, 2014.
- Bongiorno PB, Fratellone PM and LoGiudice P: Potential health benefits of garlic (*Allium sativum*): A narrative review. *J Complement Integr Med* 5: 1-26, 2008.
- Rose P, Whiteman M, Moore PK and Zhu YZ: Bioactive S-alk(en)yl cysteine sulfoxide metabolites in the genus *Allium*: The chemistry of potential therapeutic agents. *Nat Prod Rep* 22: 351-368, 2005.
- Zawistowski J, Kopec A, Jedrszczyk E, Francik R and Bystrowska B: Garlic grown from air bulbils and its potential health benefits. *ACS Symp Ser* 1286: 315-328, 2018.
- Mithen R: Sulphur-containing compounds. In: Plant secondary metabolites: occurrence, structure and role in the human diet. Crozier A, Clifford N and Ashida H (eds). Blackwell Publishing, Hoboken, NJ, pp25-46, 2006.
- Palani S, Joseph NM, Tegene Y and Zacharia A: Medical properties of garlic - A concise review. *Curr Res Pharm Sci* 4: 92-98, 2014.
- Ueda Y, Sakaguchi M, Hirayama K, Miyajima R and Kimizuka A: Characteristic flavor constituents in water extract of garlic. *Agric Biol Chem* 54: 163-169, 1990.
- Granroth B: Biosynthesis and decomposition of cysteine derivatives in onion and other *Allium* species. *Ann Acad Sci Fenn, Ser A2, Chem* 154: 1-71, 1970.
- Jones MG, Hughes J, Tregova A, Milne J, Tomsett AB and Collin HA: Biosynthesis of the flavour precursors of onion and garlic. *J Exp Bot* 55: 1903-1918, 2004.
- Hughes J, Tregova A, Tomsett AB, Jones MG, Cosstick R and Collin HA: Synthesis of the flavour precursor, alliin, in garlic tissue cultures. *Phytochemistry* 66: 187-194, 2005.
- Suzuki T, Sugii M and Kakimoto T: Metabolic incorporation of L-valine-[<sup>14</sup>C] into S-(2-carboxypropyl) glutathione and S-(2-carboxypropyl) cysteine in garlic. *Chem Pharm Bull (Tokyo)* 10: 328-331, 1962.
- Lancaster JE and Shaw ML:  $\gamma$ -Glutamyl peptides in the biosynthesis of S-alk(en)yl-L-cysteine sulfoxides (flavor precursors) in *Allium*. *Phytochemistry* 28: 455-460, 1989.
- Yoshimoto N, Onuma M, Mizuno S, Sugino Y, Nakabayashi R, Imai S, Tsuneyoshi T, Sumi S and Saito K: Identification of a flavin-containing S-oxygenating monooxygenase involved in alliin biosynthesis in garlic. *Plant J* 83: 941-951, 2015.
- Yoshimoto N, Yabe A, Sugino Y, Murakami S, Sai-Ngam N, Sumi S, Tsuneyoshi T and Saito K: Garlic  $\gamma$ -glutamyl transpeptidases that catalyze deglutamylation of biosynthetic intermediate of alliin. *Front Plant Sci* 5: 758, 2015.
- Suzuki T, Sugii M, Kakimoto T and Tsuboi N: Isolation of (-)-S-allyl-L-cysteine from garlic. *Chem Pharm Bull (Tokyo)* 9: 251-252, 1961.
- Ziegler SJ and Sticher O: HPLC of S-alk(en)yl-L-cysteine derivatives in garlic including quantitative determination of (+)-S-allyl-L-cysteine sulfoxide (alliin). *Planta Med* 55: 372-378, 1989.

21. Lawson LD, Wang Z-Y and Hughes BG:  $\gamma$ -Glutamyl-S-alkylcysteines in garlic and other *Allium* spp.: Precursors of age-dependent *trans*-1-propenyl thiosulfinates. *J Nat Prod* 54: 436-444, 1991.
22. Farooqui AA and Farooqui T (eds): *Garlic and its effects in neurological disorders*. In: *Neuroprotective effects of phytochemicals in neurological disorders*. John Wiley & Sons, Hoboken, NJ, pp113-131, 2017.
23. Hirata S, Abdelrahman M, Yamauchi N and Shigyo M: Characteristics of chemical components in genetic resources of garlic *Allium sativum* collected from all over the world. *Genet Resour Crop Evol* 63: 35-45, 2016.
24. Lawson LD: Garlic: A review of its medicinal effects and indicated active compounds. *ACS Symp Ser* 691: 176-209, 1998.
25. Ueda Y, Kawajiri H, Miyamura N and Miyajima R: Content of some sulfur-containing components and free amino acids in various strains of garlic. *Nippon Shokuhin Kogyo Gakkaishi (J Jpn Soc Food Sci Technol)* 38: 429-434, 1991.
26. Mütsch-Eckner M, Sticher O and Meier B: Reversed-phase high-performance liquid chromatography of S-alk(en)yl-L-cysteine derivatives in *Allium sativum* including the determination of (+)-S-allyl-L-cysteine sulphoxide,  $\gamma$ -L-glutamyl-S-allyl-L-cysteine and  $\gamma$ -L-glutamyl-S-(*trans*-1-propenyl)-L-cysteine. *J Chromatogr A* 625: 183-190, 1992.
27. Bloem E, Haneklaus S and Schnug E: Storage life of field-grown garlic bulbs (*Allium sativum* L.) as influenced by nitrogen and sulfur fertilization. *J Agric Food Chem* 59: 4442-4447, 2011.
28. Yamazaki Y and Okuno T: Accumulation of S-allyl-L-cysteine in garlic bulbs by warming. *Nippon Shokuhin Kagaku Kogaku Kaishi* 55: 410-415, 2008.
29. Rahman MS: Allicin and other functional active components in garlic: Health benefits and bioavailability. *Int J Food Prop* 10: 245-268, 2007.
30. Martins N, Petropoulos S and Ferreira ICFR: Chemical composition and bioactive compounds of garlic (*Allium sativum* L.) as affected by pre- and post-harvest conditions: A review. *Food Chem* 211: 41-50, 2016.
31. Matsuura H, Inagaki M, Maeshige K, Ide N, Kajimura Y and Itakura Y: Changes in contents of  $\gamma$ -glutamyl peptides and fructan during growth of *Allium sativum*. *Planta Med* 62: 70-71, 1996.
32. Ellmore GS and Feldberg RS: Allin lyase localization in bundle sheaths of the garlic clove (*Allium sativum*). *Am J Bot* 81: 89-94, 1994.
33. Penn RE, Block E and Revelle LK: Flash vacuum pyrolysis studies. 5. Methanesulfenic acid. *J Am Chem Soc* 100: 3622-3623, 1978.
34. Cavallito CJ, Buck JS and Suter CM: Allicin, the antibacterial principle of *Allium sativum*. II. Determination of the chemical structure. *J Am Chem Soc* 66: 1952-1954, 1944.
35. Borlinghaus J, Albrecht F, Gruhlke MC, Nwachukwu ID and Slusarenko AJ: Allicin: Chemistry and biological properties. *Molecules* 19: 12591-12618, 2014.
36. Choi MK: CHae K-Y, Lee J-Y and Kyung KH: Antimicrobial activity of chemical substances derived from S-alk(en)yl-L-cysteine sulfoxide (alliin) in garlic, *Allium sativum* L. *Food Sci Biotechnol* 16: 1-7, 2007.
37. Rabinkov A, Miron T, Konstantinovski L, Wilchek M, Mirelman D and Weiner L: The mode of action of allicin: Trapping of radicals and interaction with thiol containing proteins. *Biochim Biophys Acta* 1379: 233-244, 1998.
38. Lawson LD, Wang ZJ and Hughes BG: Identification and HPLC quantitation of the sulfides and dialk(en)yl thiosulfinates in commercial garlic products. *Planta Med* 57: 363-370, 1991.
39. Singh VK and Singh DK: Pharmacological effects of garlic (*Allium sativum* L.). *Ann Rev Biomed Sci* 10: 6-26, 2008.
40. Block E, Naganathan S, Putman D and Zhao SH: *Allium* chemistry: HPLC analysis of thiosulfinates from onion, garlic, wild garlic (ramsoms), leek, scallion, shallot, elephant (great-headed) garlic, chive, and Chinese chive. Uniquely high allyl to methyl ratios in some garlic samples. *J Agric Food Chem* 40: 2418-2430, 1992.
41. Yamaguchi Y, Honma R, Yazaki T, Shibuya T, Sakaguchi T, Uto-Kondo H and Kumagai H: Sulfuric odor precursor S-allyl-L-cysteine sulfoxide in garlic induces detoxifying enzymes and prevents hepatic injury. *Antioxidants* 8: 385, 2019.
42. Bartzatt R, Blum D and Nagel D: Isolation of Garlic Derived Sulfur Compounds from Urine. *Anal Lett* 25: 1217-1224, 1992.
43. de Rooij BM, Boogaard PJ, Rijkssen DA, Commandeur JN and Vermeulen NP: Urinary excretion of N-acetyl-S-allyl-L-cysteine upon garlic consumption by human volunteers. *Arch Toxicol* 70: 635-639, 1996.
44. Jandke J and Spittler G: Unusual conjugates in biological profiles originating from consumption of onions and garlic. *J Chromatogr A* 421: 1-8, 1987.
45. Trio PZ, You S, He X, He J, Sakao K and Hou D-X: Chemopreventive functions and molecular mechanisms of garlic organosulfur compounds. *Food Funct* 5: 833-844, 2014.
46. Verhagen H, Hageman GJ, Rauma A-L, Versluis-de Haan G, van Herwijnen MHM, de Groot J, Törrönen R and Mykkänen H: Biomonitoring the intake of garlic via urinary excretion of allyl mercapturic acid. *Br J Nutr* 86 (Suppl 1): S111-S114, 2001.
47. Augusti KT and Sheela CG: Antiperoxide effect of S-allyl cysteine sulfoxide, an insulin secretagogue, in diabetic rats. *Experientia* 52: 115-120, 1996.
48. Zhai B, Zhang C, Sheng Y, Zhao C, He X, Xu W, Huang K and Luo Y: Hypoglycemic and hypolipidemic effect of S-allyl-cysteine sulfoxide (alliin) in DIO mice. *Sci Rep* 8: 3527, 2018.
49. Sangeetha T and Darlin Quine S: Preventive effect of S-allyl cysteine sulphoxide (Alliin) on mitochondrial dysfunction in normal and isoproterenol induced cardiotoxicity in male Wistar rats: A histopathological study. *Mol Cell Biochem* 328: 1-8, 2009.
50. Nishimura H and Ariga T: Vinylidithiols in garlic and Japanese domestic *Allium* (*A. victorialis*). *CS Symp Ser* 546: 128-143, 1993.
51. Akao M, Shibuya T, Shimada S, Sakurai H and Kumagai H: In vivo production of bioactive compounds from S-allyl-L-cysteine sulfoxide, garlic odor precursor, that inhibit platelet aggregation. *J Clin Biochem Nutr (Suppl 43)*: 1-3, 2008.
52. Uto-Kondo H, Hase A, Yamaguchi Y, Sakurai A, Akao M, Saito T and Kumagai H: S-Allyl-L-cysteine sulfoxide, a garlic odor precursor, suppresses elevation in blood ethanol concentration by accelerating ethanol metabolism and preventing ethanol absorption from gut. *Biosci Biotechnol Biochem* 82: 724-731, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.