

Recent Shift of Body Mass Index Distribution in Korea: a Population-based Korea National Health Insurance Database, 2002–2013

Yeong Sook Yoon¹ and Sang Woo Oh²

¹Department of Family Medicine, Inje University Ilsan Paik Hospital, Goyang, Korea; ²Department of Family Medicine, Center for Obesity, Metabolism, and Nutrition, Dongguk University Ilsan Hospital, Goyang, Korea

Received: 27 September 2016
Accepted: 19 November 2016

Address for Correspondence:

Sang Woo Oh, MD, PhD
Department of Family Medicine, Center for Obesity, Metabolism, and Nutrition, Dongguk University Ilsan Hospital, 26 Dongguk-ro, Ilsandong-gu, Goyang 10326, Republic of Korea
E-mail: osw6021@yahoo.co.kr

We investigated trends of obesity prevalence among adult Koreans. Data from 110,280,952 participants who underwent medical examination provided by the National Health Insurance System from 2002–2013 were analyzed. The prevalence of moderate obesity (body mass index [BMI] 25.0–29.9 kg/m²) was slightly increased from 2002 (30.7%) through 2010 (33.1%) and reached a plateau since 2010 among men but continuously decreased from 22.5% in 2002 to 20.9% in 2013 among women. However, severe obesity (BMI ≥ 30.0 kg/m²) was rapidly increased and the prevalence in 2013 (4.2%) was almost doubled the prevalence in 2002 (2.5%). We observed a steep rise of severe obesity as well as potentially right shifting of the obese category.

Keywords: Obesity; Prevalence; Ethnicity; Weight Status

INTRODUCTION

Obesity is the most important public health problem that increases the risk of type 2 diabetes, hypertension, cardiovascular disease, cancer, and premature death; and the subsequent financial burden on healthcare systems (1).

In the developed European countries and US, obesity prevalence was remarkably increased in the several decades between the 1970s and 1990s but recently stabilized through the last decade (2–4). Moreover, based on the trajectory of prevalence of obesity in US and European society, recent studies have indicated an even more significant increase in the prevalence of severe obesity (body mass index [BMI] ≥ 35.0 kg/m²) (5) and this reflects in a rightward shift of the BMI distribution.

On the other hand, the increasing trend in prevalence of obesity in Asian countries is becoming more apparent due to the rapid changes in dietary habits and lifestyles towards a developed country (3,6). Obesity (BMI ≥ 25.0 kg/m²) prevalence varied differently in each Asia Pacific country ranging from less than 5% in India to 60% in Australia, and the rapid rate of increase in obesity was observed in China and Japan (6). Though it is apparent that obesity is endemic in much of the Asia Pacific region, the recent trends of obesity epidemic has not been well known in each Asian country.

Hence, we performed this study to describe the prevalence of obesity among the adult population in Korea using representative data and to obtain time-trend data over the last 12 years.

MATERIALS AND METHODS

National Health Insurance System (NHIS) in South Korea is a compulsory social insurance system. This NHIS insures about 97% of the population and the remaining population is covered by Medical Aid. The Korean NHIS database consists of Qualification DB, Health Check-up DB, and Claim DB and represents the entire Korean population (7). Among sub-datasets of the NHIS database, we used Qualification DB and Health Check-up DB. Health Check-up DB includes the results from biennial medical examinations of all NHI members older than 40 years and of those who are employed regardless of age (8). The proportion of complete health check-ups was approximately 40% in 2002, whereas it increased up to 68% in 2013 (7). Additional details of the Korean NHIS Database are described elsewhere (9).

The population of this study consisted of Korean adults (≥ 20 years old) who underwent medical examination provided by NHIS from 2002 through 2013. A total of 110,280,952 participants was available in the Health Check-up DB, which included anthropometric data for each participant. The distribution of our study population by gender and age were presented in the Supplementary Table 1.

Anthropometric measurements were conducted with the same protocols based on the Health Screening Implementation Standards provided by Ministry of Health and Welfare. BMI was calculated as measured weight (in kg) divided by measured height (in metres) squared. We used World Health Organization

Table 1. Trends in the prevalence of underweight, normal weight, and obesity for adults aged 20 years and older: KNHID, 2002–2013

BMI category, kg/m ²	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
All												
< 18.5	4.89	4.82	4.80	4.72	4.63	4.67	4.70	4.73	4.87	4.75	4.84	4.81
18.5–24.9	66.11	65.82	66.46	66.10	65.91	65.49	65.17	64.84	64.70	64.27	64.26	63.84
25.0–29.9	26.47	26.64	26.14	26.40	26.62	26.79	26.92	26.99	26.90	27.17	26.99	27.13
30.0–34.9	2.36	2.52	2.41	2.57	2.63	2.80	2.94	3.13	3.20	3.43	3.50	3.73
≥ 35.0	0.17	0.20	0.19	0.21	0.21	0.25	0.27	0.31	0.33	0.38	0.41	0.49
Men												
< 18.5	2.83	2.92	2.70	2.72	2.43	2.49	2.37	2.43	2.32	2.24	2.25	2.22
18.5–24.9	64.13	64.08	63.69	63.47	62.65	62.41	61.53	61.35	60.76	60.13	59.88	59.32
25.0–29.9	30.70	30.46	31.02	31.03	32.01	31.97	32.67	32.60	33.09	33.45	33.52	33.72
30.0–34.9	2.21	2.38	2.43	2.60	2.72	2.92	3.17	3.34	3.52	3.82	3.95	4.26
≥ 35.0	0.13	0.16	0.16	0.18	0.19	0.22	0.25	0.28	0.31	0.36	0.40	0.47
Women												
< 18.5	6.84	6.62	6.78	6.61	6.71	6.75	6.90	6.91	7.28	7.12	7.29	7.27
18.5–24.9	67.99	67.48	69.09	68.59	69.01	68.41	68.62	68.15	68.43	68.20	68.41	68.12
25.0–29.9	22.46	23.02	21.52	22.02	21.51	21.88	21.47	21.67	21.05	21.22	20.80	20.88
30.0–34.9	2.50	2.65	2.40	2.54	2.54	2.69	2.72	2.93	2.89	3.05	3.06	3.24
≥ 35.0	0.21	0.24	0.21	0.24	0.24	0.28	0.29	0.34	0.36	0.41	0.43	0.50

Values are presented as percentage.
KNHID = Korea National Health Insurance Database, BMI = body mass index.

(WHO) BMI cutoffs for adult Asians (10): underweight was defined as BMI < 18.5 kg/m², normal weight as BMI 18.5–24.9 kg/m², and obesity as BMI ≥ 25.0 kg/m². We categorized obesity into the following severity categories: obesity class I: 25.0–29.9 kg/m²; obesity class II: 30.0–34.9 kg/m²; and obesity class III: ≥ 35.0 kg/m². We mentioned class I obesity as moderate obesity, class II and III as severe obesity, and class III obesity as extreme obesity to prevent confusion.

Analyses were conducted by sex and 10-year age groups. To analyze changes in the age-adjusted prevalence of obesity, we applied the direct standardization method using the 2010 Korean Census population as the standard population. All statistical analyses were conducted using STATA 12 (StataCorp LP, College Station, TX, USA).

Ethics statement

This study was approved by the Institutional Review Boards at the Dongguk University Ilsan Hospital (IRB File No.: 2016-45), and written informed consent was exempted.

RESULTS

Trends in the age-standardized prevalence of obesity from 2002 to 2013 by sex are presented in Table 1. Among men aged 20 years and older, the prevalence of class I obesity was slightly increased through 2002 to 2010 (30.70% vs. 33.09%) and reached a plateau since 2010. However, class II and class III obesity was rapidly increased and prevalence in 2013 was more than twice the prevalence in 2002. Women also showed rapid increase of class II and III obesity since 2007. However, the prevalence of class I obesity among women, unlike men, continuously decreased from 22.46% in 2002 to 20.88% in 2013. The prevalence of

class I obesity was higher in men than in women but the prevalence of class III obesity was higher in women than in men.

Fig. 1 showed trends in obesity from 2002 to 2013 in both men and women by age groups. Over 12 years, the prevalence of class I obesity among men was highest in the 50s age groups but those of class II and III obesity was highest in the 30s and 20s age groups, respectively. On the other hand, women in their 60s showed highest prevalence of class I and II obesity from 2002 to 2013. Class III obesity prevalence was highest in women aged 50–59 in 2002 but was highest in women aged 30–39 and 20–29 in 2013. A clear, rapid rise in class II and III of obesity continued from 2002 through 2013 in both men and women in their 20s and 30s.

DISCUSSION

Our study showed that plateaued trends in the prevalence of moderate obesity and dramatic increasing trends in severe obesity over 12 years in Korean adults.

In this study, the prevalence of obesity among men might have slightly increased from 2002 to 2010 and then stabilized from 2010 to 2013, while the prevalence of obesity among women generally decreased across all the years examined. These trends were similar to results from the study using the Korea Nation Health and Nutrition Examination Survey (KNHANES) 1998, 2005, and 2007 to 2009 (11,12). When compared with obesity trends in the East Asia countries, our result was very similar to China (13) but differed from Japan (14). Lao et al. (13) reported the prevalence of obesity in China changed very little from 2002 to 2010, although earlier studies using data before 2000 suggested strongly increasing trends in Chinese adults (2,15). However, Japan showed that the percentages of obesity are increasing in men but not in women in all age groups (14).




Fig. 1. Age-stratified trends in the prevalence of class I, II, and III obesity for adults aged 20 years and older by sex: KNHID, 2002–2013. Obesity was categorized into the following severity categories: obesity class I: 25.0–29.9 kg/m²; obesity class II: 30.0–34.9 kg/m²; and obesity class III: ≥ 35.0 kg/m². KNHID = Korea National Health Insurance Database, BMI = body mass index.

It is noteworthy that severe obesity increased almost 1.5 times in men and tripled in women from 2002 through 2013 in this

study. The obesity epidemic in developed countries, such as Australia, Europe, Russia, and the USA has slowed down or sta-

bilized since the 2000s but severe obesity has increased very fast (16). In Asian studies in the early 2000s, prevalence of severe obesity was 2.4% of men and 3.4% of women in China, 2.3% and 3.4% in Japan, and 6.3% and 7.4% in Singapore, respectively (6). Though the overall prevalence of morbid obesity remains low in comparison with the USA, we should take note of a dramatic rise of severe obesity. Patients with morbid obesity have had a higher risk for type 2 diabetes and coronary heart disease than patients with mild obesity (17). Therefore, increasing severe obesity at a faster rate is of importance for future burden of obesity-related morbidity and mortality.

Another remarkable finding in our study was faster rise of morbid obesity in younger adults. Moreover, the prevalence of extreme obesity in 2013 was highest in ages 20 to 29 and ages 30 to 39 among both men and women. In the USA, adults ages 40 to 59 have higher obesity as well as severe obesity than other age groups, but increasing trends of severe obesity at a faster rate in young adults were quite similar to Korea. An increased trend in morbid obesity in young Korean adults may be explained by the fact that adolescents with obesity who were exposed to a dramatic increase in fast food consumption (18) and in snacks high in fat and sugar (19) and to sedentary lifestyle and sleep deprivation (20) due to excessive private education (21) and increase in automobile use now became adults in their 20s and 30s. Therefore, public awareness, intervention, and prevention of obesity should be focused on adolescents and young adults.

The present study has several limitations. We failed to provide obesity trends before 2002. Second, the possibility of selection bias associated with relatively lower proportion of complete health check-ups in 2002 should be considered. Additionally, general health check-up was limited to employee subscribers in their 20s and 30s. Therefore, it is difficult to argue that they are completely representative of individuals in their 20s and 30s, although our study subjects aged 40 years and older are representative.

However, our study also has several strengths. First, we used nationally representative data over 12 years, which included consistent anthropometric measurements of height and weight by the same protocols. Second, to the best of our knowledge, this is the largest and most recent study to investigate the trends of obesity in Korea.

In summary, we investigated obesity trends in Korean adults from 2002 to 2013 and observed that moderate obesity was stabilized but severe obesity was rapidly increased, especially in young adults. Steep rise of severe obesity as well as potentially right shifting of the obese category in Korea seems to take after obesity trends in the USA. Therefore, there should be wider discussion of whether existing diagnostic criteria for obesity in Asians remain unchanged. In addition, considering fast rise in prevalence of morbid obesity in young adults, further studies exploring factors causing these trends are needed to establish a long-

term policy for lowering future burden on health care system.

DISCLOSURE

The authors have no potential conflicts of interest to disclose.

AUTHOR CONTRIBUTION

Conceptualization: Yoon YS, Oh SW. Data curation: Yoon YS, Oh SW. Writing - original draft: Yoon YS. Writing - review & editing: Oh SW.

ORCID

Yeong Sook Yoon <http://orcid.org/0000-0002-9249-2940>

Sang Woo Oh <http://orcid.org/0000-0002-5623-6720>

REFERENCES

1. Hoque ME, Mannan M, Long KZ, Al Mamun A. Economic burden of underweight and overweight among adults in the Asia-Pacific region: a systematic review. *Trop Med Int Health* 2016; 21: 458-69.
2. Wang Y, Beydoun MA. The obesity epidemic in the United States--gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiol Rev* 2007; 29: 6-28.
3. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, Mullany EC, Biryukov S, Abbafati C, Abera SE, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014; 384: 766-81.
4. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999-2010. *JAMA* 2012; 307: 491-7.
5. Ricci MA, De Vuono S, Scavizzi M, Gentili A, Lupattelli G. Facing morbid obesity: how to approach it. *Angiology* 2016; 67: 391-7.
6. Asia Pacific Cohort Studies Collaboration. The burden of overweight and obesity in the Asia-Pacific region. *Obes Rev* 2007; 8: 191-6.
7. Lee YH, Han K, Ko SH, Ko KS, Lee KU; Taskforce Team of Diabetes Fact Sheet of the Korean Diabetes Association. Data analytic process of a nationwide population-based study using national health information database established by National Health Insurance Service. *Diabetes Metab J* 2016; 40: 79-82.
8. Lee H, Cho J, Shin DW, Lee SP, Hwang SS, Oh J, Yang HK, Hwang SH, Son KY, Chun SH, et al. Association of cardiovascular health screening with mortality, clinical outcomes, and health care cost: a nationwide cohort study. *Prev Med* 2015; 70: 19-25.
9. Song SO, Jung CH, Song YD, Park CY, Kwon HS, Cha BS, Park JY, Lee KU, Ko KS, Lee BW. Background and data configuration process of a nationwide population-based study using the Korean national health insurance system. *Diabetes Metab J* 2014; 38: 395-403.
10. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004; 363: 157-63.

11. Oh SW. Obesity and metabolic syndrome in Korea. *Diabetes Metab J* 2011; 35: 561-6.
12. Rhee SY, Park SW, Kim DJ, Woo J. Gender disparity in the secular trends for obesity prevalence in Korea: analyses based on the KNHANES 1998-2009. *Korean J Intern Med* 2013; 28: 29-34.
13. Lao XQ, Ma WJ, Sobko T, Zhang YH, Xu YJ, Xu XJ, Yu DM, Nie SP, Cai QM, Xia L, et al. Overall obesity is leveling-off while abdominal obesity continues to rise in a Chinese population experiencing rapid economic development: analysis of serial cross-sectional health survey data 2002-2010. *Int J Obes (Lond)* 2015; 39: 288-94.
14. Nishi N. Monitoring obesity trends in health Japan 21. *J Nutr Sci Vitaminol (Tokyo)* 2015; 61 Suppl: S17-9.
15. Xi B, Liang Y, He T, Reilly KH, Hu Y, Wang Q, Yan Y, Mi J. Secular trends in the prevalence of general and abdominal obesity among Chinese adults, 1993-2009. *Obes Rev* 2012; 13: 287-96.
16. Rokholm B, Baker JL, Sørensen TI. The levelling off of the obesity epidemic since the year 1999—a review of evidence and perspectives. *Obes Rev* 2010; 11: 835-46.
17. Todd Miller M, Lavie CJ, White CJ. Impact of obesity on the pathogenesis and prognosis of coronary heart disease. *J Cardiometab Syndr* 2008; 3: 162-7.
18. Shin HJ, Cho E, Lee HJ, Fung TT, Rimm E, Rosner B, Manson JE, Wheelan K, Hu FB. Instant noodle intake and dietary patterns are associated with distinct cardiometabolic risk factors in Korea. *J Nutr* 2014; 144: 1247-55.
19. Ha A. Obesity and its association with diets and sedentary life style among school children in Seoul, Korea: compliance with dietary references intakes for Koreans food guides. *Nutr Res Pract* 2007; 1: 212-7.
20. Kong IG, Lee HJ, Kim SY, Sim S, Choi HG. Physical activity, study sitting time, leisure sitting time, and sleep time are differently associated with obesity in Korean adolescents: a population-based study. *Medicine (Baltimore)* 2015; 94: e1965.
21. Korean Statistical Information Service. The private education expenditures survey in 2007-2015 [Internet]. Available at http://kosis.kr/eng/statisticsList/statisticsList_01List.jsp?vwcd=MT_ETITLE&parmTabId=M_01_01 [accessed on 18 November 2016].