

POSTER PRESENTATION

Open Access

P73. Functional characterisation of HBV-specific T cell receptors for redirection of T cells against HBV infected hepatocytes

K Krebs^{1*}, K Metzger¹, L Weigand², C Dargel¹, E Kieback³, W Uckert³, D Busch⁴, A Krackhardt², U Protzer¹

From 1st Immunotherapy of Cancer Conference (ITOC1)
Munich, Germany. 12-14 March 2014

Chronic HBV infection, which is accompanied by a weak and oligoclonal T cell response, is the most common cause of hepatocellular carcinoma (HCC). Current antiviral therapies do not eliminate the virus, but T cell therapy will very likely do so. From PBMCs of two HLA-A2⁺ acutely infected patients and a donor who cleared HBV infection we have established several HBV-specific monoclonal T cell lines. Thereof we isolated 11 different T cell receptors (TCR) that are specific for the HBV S-protein derived peptides S20 (FLLTRILT1) and S172 (WLSLLVPFV) or for the C18 core-peptide (FLPSDFPSV). The aim of this study was a functional comparison of our set of HBV-specific TCRs in order to identify TCRs with optimal recognition of HBV peptides presented on HLA-A2.

By murinization and codon-optimisation of gene sequences of TCR a and b chains, fused by a P2A element for polycistronic expression, TCR expression after retroviral transduction was increased 2-fold to 60% of PBMCs expressing an HBV-specific TCR.

PBMCs transduced with the 11 optimised HBV-specific TCRs were compared in killing assays using peptide-pulsed T2 cells, LCLs and HBV-replicating HepG2.2.15 cells as targets. CD8⁺ T cells transduced with the core-specific TCRs killed target cells loaded with 0.01 nM of peptide. Cells specific for the S20 and S172 peptide were less sensitive with a specific lysis as low as 0.1 nM. Expression of most of the HLA-A2 restricted HBV-specific TCRs in CD4⁺ T cells also led to specific cytotoxicity, which was 10-fold reduced in sensitivity compared to CD8⁺ T cells and independent of CD8 co-receptor binding. Notably,

our HBV-specific TCRs recognised peptide presented on various different HLA-A2 subtypes.

CD8⁺ T cells transduced with HBV-specific TCRs were also able to recognise endogenously processed peptides and specifically kill HBV-replicating hepatoma cells and strongly reduce cccDNA levels in HBV-infected HepaRG cells.

In addition, intracellular cytokine staining after stimulation showed that the TCR-transduced CD8⁺ T cells were polyfunctional, secreting INF- γ , TNF- α and IL-2, whereas CD4⁺ T cells produced mainly TNF- α and/or IL-2.

We will further analyse our HBV-specific TCRs in HBV/HLA-A2 transgenic mice in order to identify the TCR that confers best antiviral activity. Our HBV-specific TCRs may be used for elucidating specific anti-HBV mechanisms exerted by T cells, and most importantly, for adoptive T cell therapy of chronic hepatitis B and HBV-induced HCC.

Authors' details

¹TU Muenchen / Helmholtz Zentrum Muenchen, Institute of Virology, Muenchen, Germany. ²Klinikum rechts der Isar, III. Medizinische Klinik, Muenchen, Germany. ³Max-Delbrück Center for Molecular Medicine, Berlin, Germany. ⁴TU Muenchen, Institute for Medical Microbiology Immunology and Hygiene, Muenchen, Germany.

Published: 12 March 2014

doi:10.1186/2051-1426-2-S2-P47

Cite this article as: Krebs et al.: P73. Functional characterisation of HBV-specific T cell receptors for redirection of T cells against HBV infected hepatocytes. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 2):P47.

¹TU Muenchen / Helmholtz Zentrum Muenchen, Institute of Virology, Muenchen, Germany

Full list of author information is available at the end of the article