

Research



Cite this article: Khalil S, Welklin JF, McGraw KJ, Boersma J, Schwabl H, Webster MS, Karubian J. 2020 Testosterone regulates *CYP2J19*-linked carotenoid signal expression in male red-backed fairywrens (*Malurus melanocephalus*). *Proc. R. Soc. B* **287**: 20201687.
<http://dx.doi.org/10.1098/rspb.2020.1687>

Received: 14 July 2020

Accepted: 19 August 2020

Subject Category:

Evolution

Subject Areas:

evolution, physiology, genetics

Keywords:

ketocarotenoid, pigmentation, ornamental coloration, androgens, gene expression

Author for correspondence:

Sarah Khalil

e-mail: skhalil@tulane.edu

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5106138>.

Testosterone regulates *CYP2J19*-linked carotenoid signal expression in male red-backed fairywrens (*Malurus melanocephalus*)

Sarah Khalil¹, Joseph F. Welklin^{2,3}, Kevin J. McGraw⁴, Jordan Boersma⁵, Hubert Schwabl⁵, Michael S. Webster^{2,3} and Jordan Karubian¹

¹Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, LA, USA

²Macaulay Library, Cornell Laboratory of Ornithology, Ithaca, NY, USA

³Department of Neurobiology and Behavior, Cornell University, Ithaca, NY, USA

⁴School of Life Sciences, Arizona State University, Tempe, AZ, USA

⁵School of Biological Sciences, Washington State University, Pullman, WA, USA

id SK, 0000-0002-6929-4540; JFW, 0000-0002-7534-8795; KJM, 0000-0001-5196-6620; JB, 0000-0001-8355-2027; HS, 0000-0001-8253-0133; MSW, 0000-0001-7585-4578; JK, 0000-0001-8201-9992

Carotenoid pigments produce most red, orange and yellow colours in vertebrates. This coloration can serve as an honest signal of quality that mediates social and mating interactions, but our understanding of the underlying mechanisms that control carotenoid signal production, including how different physiological pathways interact to shape and maintain these signals, remains incomplete. We investigated the role of testosterone in mediating gene expression associated with a red plumage sexual signal in red-backed fairywrens (*Malurus melanocephalus*). In this species, males within a single population can flexibly produce either red/black nuptial plumage or female-like brown plumage. Combining correlational analyses with a field-based testosterone implant experiment and quantitative polymerase chain reaction, we show that testosterone mediates expression of carotenoid-based plumage in part by regulating expression of *CYP2J19*, a ketolase gene associated with ketocarotenoid metabolism and pigmentation in birds. This is, to our knowledge, the first time that hormonal regulation of a specific genetic locus has been linked to carotenoid production in a natural context, revealing how endocrine mechanisms produce sexual signals that shape reproductive success.

1. Introduction

Carotenoid pigments, which provide many of the vivid red, orange and yellow colours observed in vertebrates, have long captured the attention of behavioural ecologists interested in the evolution of social signals. Among vertebrates, carotenoid-based colours can serve as honest indicators of quality because of costs associated with obtaining and producing them, inspiring classic research surrounding the adaptive benefits of sexual signalling [1,2]. However, many aspects of the underlying mechanisms of carotenoid colour production remain unclear [3] (though see [4,5]), even though many assumptions surrounding honest signalling rely on understanding these proximate mechanisms. For example, carotenoids can be important for both antioxidant and immune functions [6,7], which suggests a trade-off between using carotenoids for physiological maintenance versus signal production [8,9]. However, recent work has cast doubt on the degree to which carotenoids function in physiological defences [10] or trade-off with somatic maintenance [11], sparking debate about the costs associated with production or maintenance of

carotenoid-based signals [12,13]. Resolving the underlying proximate mechanisms involved in carotenoid production can help inform this debate by improving our ability to interpret how multiple physiological pathways interact to shape and maintain carotenoid signals [3].

Once acquired from the diet, carotenoids can be modified via endogenous enzymatic processes that modulate carotenoid-based signal expression [14–17]. In contrast to some endogenously synthesized pigments such as eumelanin [18,19], researchers have only recently begun describing and characterizing the enzymes, genes and pathways involved in carotenoid metabolism and deposition [15,20,21]. Carotenoid ketolation, which includes the metabolism of yellow dietary carotenoids into red ketocarotenoids [22], is an important innovation in vertebrate evolution and colour diversification [23–27]. In birds in particular, many well-known examples of sexually selected visual displays involve bright-red ketocarotenoid-based coloration in the plumage or bare parts (e.g. [28–31]). Two recent studies have independently identified and described the locus *CYP2J19* as a gene that codes for the production of a putative ketolase enzyme (from the cytochrome P450 family) that underlies ketocarotenoid pigmentation in birds [14,15]. Though the discovery of this ketolase has important implications for understanding carotenoid signalling, the generality of this mechanism remains unclear since these studies have so far linked *CYP2J19* to red coloration in aberrantly coloured domesticated birds: the mutant ‘yellowbeak’ zebra finch (*Taeniopygia guttata*) and a hybrid breeding line (red-factor) of canary (*Serinus canaria*). However, recent work has suggested that *CYP2J19* expression may account for interspecific variation in red coloration—for example, *CYP2J19* expression was consistently higher in the liver of species of weaverbirds (Ploceidae) with red plumage compared to species with yellow plumage [24]. However, to date to our knowledge, no empirical study has linked *CYP2J19* expression to intraspecific variation in expression of red coloration in a wild species, limiting our ability to evaluate potential adaptive consequences of this gene-regulatory mechanism to sexually selected traits [32].

Endocrine regulation often underlies the development and expression of sexually selected traits [33]. Androgens, such as testosterone, play a fundamental role in mediating gene expression and resulting phenotype [34,35]. In many animals, testosterone is involved in phenotypic integration at the level of the individual, including determining breeding phenotype [34,35]. For example, in species characterized by complex social hierarchies, such as cooperative breeders [36,37], dominant breeding males often have elevated testosterone relative to subordinate ‘helper’ males [38–41]. Testosterone regulates changes in physiology and phenotype to help match organisms to their social environment, including changes in social rank and breeding status [42,43]. At the same time, studies in other systems provide little support for the phenotypic-integration role of steroid hormones [44,45], revealing limitations in our understanding of these relationships. Indeed, many key components of the endocrine-genomic mechanisms involved in regulating signal expression remain unresolved [46,47], especially for carotenoid-based signalling systems [48–50], highlighting the need for additional research into how phenotypic integration operates on a mechanistic level.

The red-backed fairywren (*Malurus melanocephalus*) provides a useful study system for linking these two lines of inquiry and assessing how *CYP2J19* and testosterone regulation may interact to control intraspecific variation in the expression of a

carotenoid-based signal. Within a population of this cooperatively breeding species of bird, males can express either ornamental black body plumage with a carotenoid-based red dorsal feather patch, which is displayed in courtship, or female-like unornamented brown plumage [51,52]. Individual males can flexibly express unornamented or ornamented plumage depending on several factors including (i) breeding status: nearly all males moult into ornamented plumage during the non-breeding season (pre-alternate moult) in preparation for breeding, though timing and duration of this phase vary, and most males moult back into unornamented plumage after the breeding season (pre-basic moult); (ii) age: approximately 15% of males in our study population moult into ornamented plumage in their first breeding season, 90% by 2 years of age, and nearly all males by their third year; and (iii) physiological condition: males in better condition are more likely to moult into ornamented plumage [53]. Ornamented males have higher reproductive success than unornamented males, driven by higher rates of extrapair paternity in ornamented males [54]. In addition, there is experimental evidence for strong female preference for males with redder dorsal plumage [55], suggesting that this is a sexually selected trait in this species, which in turn appears to be driving introgression of redder plumage between subspecies [56]. Previous work also indicates that androgens (in particular testosterone) are important in signal acquisition: ornamented male red-backed fairywrens have higher levels of circulating androgens than unornamented males [57], unornamented males experimentally implanted with testosterone moult into the ornamental red/black plumage [58], and younger non-breeding helper males rapidly increase androgen levels and develop carotenoid-based plumage when experimentally provided with breeding opportunities [59]. In addition, testosterone treatment of females can induce some red plumage coloration in normally brown females that would otherwise have low circulating androgens [60]. Revealing how these endocrine processes may regulate gene expression associated with sexually selected carotenoid coloration has the potential to advance our understanding of the evolutionary origins and trajectories of carotenoid ornaments.

In this study, we test the hypothesis that testosterone regulates carotenoid-based plumage expression at the level of gene expression in male red-backed fairywrens. Specifically, we investigate the relationship between testosterone and carotenoid-based signal production by assessing the correlation between circulating ketocarotenoid concentration and plumage phenotype (i.e. expression of the carotenoid-based plumage patch), as well as experimentally test for the role of testosterone in regulating *CYP2J19* expression. We first use high-performance liquid chromatography (HPLC) to determine circulating carotenoid levels of ornamented red/black males, unornamented brown males, and females to identify the degree to which variation in metabolized ketocarotenoid circulation explains differences in carotenoid-based signal expression among these phenotypes. To evaluate the genomic regulation of these ketocarotenoid differences, we measure relative expression of *CYP2J19* in the liver (a central site for carotenoid metabolism [14,15,61,62]) in each phenotype. Finally, we experimentally test for the role of testosterone in regulating *CYP2J19* expression by measuring expression levels in testosterone-implanted unornamented males that developed carotenoid-based plumage in response to testosterone treatment. Our results are consistent with the hypothesis that, in red-backed fairywrens, testosterone mediates

expression of carotenoid-based plumage by regulating expression of *CYP2J19*.

2. Material and methods

(a) Plasma sample collection and quantifying circulating carotenoids

We collected samples and conducted experiments (below) on free-living red-backed fairywrens captured in mist nets at our long-term study site in Samsonvale, Queensland, Australia (27°27' S, 152°85' E). We collected blood samples (20–70 µl) from the wing vein using heparinized microcapillary tubes from May to August 2017 and 2018 during the non-breeding season, a period when most males were actively moulting into their breeding season plumage. Blood was centrifuged for 5 min at 10 000 r.p.m., after which the plasma was separated from the packed cells and stored at –20°C until transport to the USA where samples were stored at –80°C until HPLC analysis.

Plumage score was recorded at time of capture following prior methods [63], yielding total ornamentation scores that ranged from 0 (brown) to 100 (red/black). Based on this score, males were labelled as either ‘unornamented’ (brown plumage, score < 33), ‘intermediate’ (mixed plumage, score between 33 and 66), or ‘ornamented’ (red/black plumage, score > 66). Females always have completely brown plumage (plumage score = 0) and are therefore considered unornamented. Timing of moult into ornamented plumage is variable (J. F. Welklin, S. M. Lantz, S. Khalil, J. Karubian, M. S. Webster 2020, unpublished data), similar to other recorded *Malurus* species [64], meaning that male plumage at time of capture and sample collection may differ from the ‘final’ plumage colour score the male expressed later in the breeding season. Males can breed in unornamented or ornamented plumage or serve as auxiliaries with unornamented plumage (helpers) at the nest [53]. Because we were interested in differences between unornamented and ornamented plumage, we documented the ‘final’ plumage score of colour-banded individuals on 1 November (the approximate mid-point of the breeding season) and used this ‘final’ score for analyses of unornamented males, ornamented males, and females; we excluded from analysis the relatively small subset of birds whose ‘final’ plumage score was intermediate ($n = 11$). We assigned either minimum or known age (age range 1–7 years, 77%, 123 of 160, were of known age) to all birds at the time of sample collection using nestling banding records or extent of skull ossification (ossification scale modified from [65], and we have validated this scale within this species multiple times). Qualitatively similar results were obtained in analyses run with these age criteria, or using only known-age birds (see the electronic supplementary material, table S1).

We used HPLC to identify and quantify the concentration of carotenoids in the plasma, following the methods of Rowe & McGraw [51]. We analysed carotenoids in 160 plasma samples ($n = 42$ females, $n = 29$ unornamented males, $n = 89$ ornamented males). To assess the relationship between circulating ketocarotenoid levels and plumage phenotype, we ran a linear mixed-effect model with the *lme* function in the R package *nlme* [66], R v. 3.6.0 [67]. The model included total circulating ketocarotenoid concentration (i.e. the sum of alpha-doradexanthin, astaxanthin, adonirubin and canthaxanthin concentrations) as the response variable and the following predictor variables: (i) phenotype (female versus unornamented male versus ornamented male); (ii) age (as a continuous variable); (iii) year of sample collection; and (iv) the interaction between phenotype and age. To control for repeated measures of the same individual across years, we added individual as a random effect ($n = 13$ individuals sampled both years). Residuals were inspected visually for homoscedasticity, and we used the *varIdent* function to control for heterogeneity of variance between groups. Year of sample collection did not improve model

fit (i.e. it did not improve Akaike information criteria by more than 2 and the p -value of the year variable was greater than 0.05) and was therefore dropped. We tested the model for significance of phenotype with a Tukey’s posthoc test using the *glht* function in the R package *multcomp* [68].

(b) Testosterone implantation and liver sample collection

We collected liver samples from breeding, but not auxiliary helper, red-backed fairywrens in November 2017, to control for potentially confounding underlying differences in endocrine or genetic profiles that may exist between auxiliary non-breeding versus breeding individuals [57]. First, three breeding unornamented males were implanted with testosterone. At time of initial capture and implantation, around 10 feathers were plucked from the centre of the back to induce feather replacement at that location. Implants were composed of beeswax (73% by weight; Sigma-Aldrich, St. Louis, MO, USA) and hardened frozen peanut oil (24% by weight; ACROS Organics, NJ, USA) that were mixed in a water bath at 67°C. Once the beeswax/peanut oil mixture was melted, crystalline testosterone (3% by weight; Sigma-Aldrich, St. Louis, MO, USA) was dissolved in 2.5 µl of 200 proof ethanol (Fisher Bioreagents™), and the testosterone suspension was then added to the wax mixture and stirred. The implants were formed by feeding partially solidified wax through the tip of a syringe that was cut so the diameter was 2 mm, resulting in implants of 2 × 3.2 mm weighing between 19.8 and 20.7 mg. Testosterone concentration in the beeswax carrier was scaled to produce high physiological concentrations found in circulation during the breeding season [57]. Implants were inserted subcutaneously using forceps above the thigh into a small (2–3 mm) skin incision that was sealed with veterinary skin adhesive. After confirming that the incision was completely sealed and the bird was in good condition, the bird was released. Three unornamented males were implanted with sham controls (beeswax/peanut oil implant with no testosterone) and also had around 10 back feathers plucked.

Implanted birds were recaptured 10–12 days post-implantation for liver sample collection, a time period that allowed for growth of pin feathers in the plucked plumage patches (red pins in testosterone-implanted males, and a mix of red and brown pins in the sham-implanted males, consistent with what had been observed in another feather-plucking study in this species [59]). We were unable to recapture one of the sham-implanted birds after implantation. Instead, we captured and obtained samples from one additional breeding unornamented male, who was not implanted, to include in our control group. Following recapture on territories in mist nets, birds were immediately sacrificed by cervical dislocation. Body dissection was performed in the field, and the right lower lobe of the liver was removed and stored in 1 ml of RNAlater storage buffer (ThermoFisher Scientific), and immediately placed on dry ice. Samples were stored at –80°C until RNA extraction. In addition to the three testosterone-implanted unornamented males and the three control unornamented males (two with sham implants, one without an implant), we also collected liver samples for three ornamented breeding males and three breeding females (without implants). All birds sacrificed were seen paired with a male or female within two weeks prior to sample collection, and all samples were collected within a period of 9 days. Circulating androgens were measured using an established radioimmunoassay protocol for this species (full methods in [57,69]); the intra-assay coefficient of variation was 8.68%. Testosterone-implanted birds were confirmed to have high concentrations of circulating androgens at time of collection (mean = 3027 pg ml⁻¹, range = 2198–4065 pg ml⁻¹), which is within the natural range of androgens for breeding ornamented males in this species [57]. Ornamented males also had similarly high levels of androgens at time of

collection (mean = 1834 pg ml⁻¹, range = 1124–2925 pg ml⁻¹). We were unable to obtain samples to assay testosterone concentrations for control unornamented males or females.

(c) Quantifying relative expression of *CYP2J19*

To extract messenger RNA (mRNA), we removed liver tissue from the RNAlater buffer and homogenized it in a Qiagen TissueRuptor. We used a Qiagen RNeasy mini-kit, following manufacturer's instructions, and reverse transcribed the mRNA to complementary DNA (cDNA) with a Superscript IV first strand synthesis kit (Invitrogen). All quantitative polymerase chain reaction (qPCR) reactions were run on CFX96 Touch™ Real-Time PCR Detection System (BioRad) with CFX Maestro Software (BioRad), using PowerUp SYBR Green Master Mix (ThermoFisher Scientific). For measurements of *CYP2J19* expression, we used qPCR primers CYP2J2-2F and CYP2J2-2R [14]. We assayed gene expression in triplicate for each sample (except for one female sample where *CYP2J19* was assayed in duplicate) and normalized the data using the housekeeping gene *GAPDH*, using primers Gg_GAPDH_qPCR_F and Gg_GAPDH_qPCR_R [15]. Reaction conditions for qPCR were tested and optimized using a standard curve produced by creating a serial dilution of a pool of all cDNA samples. Efficiencies ranged from 95% to 105%, and we analysed qPCR data using the delta-delta Ct method [70], further described in the electronic supplementary material, Methods.

We found no effect of the presence or the absence of the sham implant on gene expression within unornamented males (see the electronic supplementary material, Methods), and no evidence of homoscedasticity (Breusch Pagan test, $p > 0.05$). We tested for statistical differences in liver *CYP2J19* expression (log fold change) between phenotypes with an ANOVA, using the aov function in R, followed by a Tukey's posthoc test using the Tukey HSD function in R.

3. Results

(a) Circulating carotenoid concentration was associated with sex and plumage phenotype

We identified six different circulating carotenoids in fairywrens: two dietary xanthophylls, lutein and zeaxanthin, and four red metabolized ketocarotenoids previously identified in the red dorsal feathers of red-backed fairywrens [51]: alpha-doradexanthin, astaxanthin, adonirubin and canthaxanthin. Plumage phenotype was a significant predictor of total circulating ketocarotenoid concentration ($F_{2,48} = 240.2$, $p < 0.0001$; figure 1), where ornamented (red/black) males had higher concentrations of circulating ketocarotenoids than either unornamented (brown) males (Tukey's posthoc, $p_{\text{adj}} < 0.001$) or females (brown) ($p_{\text{adj}} < 0.001$). There was also a significant effect of the phenotype \times age interaction ($F_{2,48} = 24.7$, $p < 0.0001$) but no effect of age alone (electronic supplementary material, table S2). Specifically, ages differed in ketocarotenoid circulation only for unornamented males, where 2-year-old unornamented males had higher concentrations of circulating ketocarotenoids than their 1-year-old counterparts (figure 1 and the electronic supplementary material, figure S1).

(b) *CYP2J19* expression in the liver was associated with expression of red plumage

Liver expression of *CYP2J19* differed significantly among plumage phenotypes ($F_{3,8} = 114$, $p < 0.0001$; figure 2). Ornamented males had higher relative expression of *CYP2J19*

than did unornamented males (Tukey's posthoc, $p_{\text{adj}} = 0.038$), and unornamented males had higher relative expression of *CYP2J19* than did females ($p_{\text{adj}} < 0.0001$).

(c) Testosterone upregulated *CYP2J19* expression

Testosterone-implanted unornamented males had significantly higher expression of *CYP2J19* in liver tissue than did control unornamented males (Tukey's posthoc, $p_{\text{adj}} = 0.012$; figure 2) or females (Tukey's posthoc, $p_{\text{adj}} < 0.0001$), with no difference in expression between testosterone-implanted unornamented males and ornamented males (Tukey's posthoc, $p_{\text{adj}} = 0.83$).

4. Discussion

By combining field-based observational and experimental investigations with gene expression and biochemical analyses, we found that testosterone regulates gene expression implicated in the production of sexually selected red plumage in male red-backed fairywrens. Ornamented males with carotenoid-based red plumage ornamentation had higher levels of circulating ketocarotenoids than either females or unornamented males (both with brown plumage). Carotenoid metabolism is an endogenous process, and the ketolase enzyme encoded by *CYP2J19* has previously been demonstrated to convert dietary carotenoids into red ketocarotenoids in birds [14,15]. Along with unmanipulated ornamented fairywren males having higher expression of *CYP2J19* in the liver than either unornamented males or females, our experiment confirmed that elevated testosterone levels resulted in increased *CYP2J19* expression in unornamented males. These findings are supported by field studies showing that ornamented plumage is testosterone-dependent in red-backed fairywrens [57,58]. Taken together, these results are consistent with a mechanistic hypothesis (figure 3), whereby circulating testosterone levels change in males in response to intrinsic or extrinsic cues (breeding status, age, physiological condition [53,57]), which in turn modifies expression of *CYP2J19* in the liver and thus increases concentration of metabolized ketocarotenoids in the plasma, which is associated with red plumage ornamentation. Though we did not test this full mechanistic hypothesis within the same individuals (i.e. it is based on two separate datasets, one for circulating ketocarotenoids and one for the testosterone-gene expression experiment), this proximate pathway for carotenoid-based signal production establishes a link between the endocrine and gene-regulatory system and avian coloration important for sexual signalling and elevated male reproductive success.

Testosterone has been shown in several studies to be an important activator of male ornamentation [33,71,72], yet there is little work investigating how testosterone and ornament production are linked [73,74]. Mutations and expression differences in *CYP2J19* have been implicated in driving carotenoid trait divergence between species and subspecies [14,15,23,24,75,76], but it has not previously been shown whether *CYP2J19* expression levels influences intra-population variation in ornamentation. Our study design focused on characterizing differences in plumage phenotypes within a wild population, revealing how hormonal regulation of *CYP2J19* expression generates variation in a polymorphic sexual signal. Past studies combining endocrine and gene expression analysis in other natural systems have revealed how flexibility in behaviour is mediated within a species




Figure 1. Ornamented male red-backed fairywrens have higher concentrations of circulating metabolized ketocarotenoids than do either unornamented males or females. (a) Scatterplot points show ketocarotenoid concentration as a function of breeding phenotype and age. Lines represent fitted lines from the linear mixed-effect model described in text. (b) Total plasma carotenoid concentration for the different plumage phenotypes are presented as boxplots, indicating the median and quartiles with whiskers reaching up to 1.5 times the interquartile range. The half-violin plot outlines show kernel probability density. The scatterplot points show the value of the total carotenoid concentration for each sample, and the colour of the point represents the concentration of metabolized ketocarotenoids in that sample, represented as a scaled gradient where yellow points have low metabolized ketocarotenoid concentration and red points have high metabolized ketocarotenoid concentration (see inset legend). (Online version in colour.)

[77,78], and how phenotypic differences may evolve between species [79], improving our understanding of hormone-mediated phenotypic evolution. The current study adds to this body of work by suggesting that incorporating endocrine control of gene expression to studies of colour production may be a rewarding research avenue for assessing signal evolution. Specifically, our findings show support for a role of testosterone as a transcriptional regulator of sexually selected phenotypes, which may also be important in other species with strong sexual selection for ornamented phenotypes.

Testosterone may also help explain two unexpected results from this study. First, why would unornamented males have higher expression of *CYP2J19* than females, despite both of them having similar brown plumage? Unornamented male red-backed fairywrens that shift from being an auxiliary helper to being a breeder during the breeding season exhibit rapid increases in their circulating testosterone levels [59], and in general male breeders have higher levels of circulating testosterone than do helpers [57]. These higher concentrations of testosterone probably benefit unornamented breeders by

mediating observed increases in breeding-specific behaviours such as territory defence [63,80], but also probably trigger increased *CYP2J19* expression. Whether this increase in *CYP2J19* expression is merely a by-product of testosterone or whether it somehow benefits these males is unclear, but one possibility is that it could allow newly paired unornamented males to quickly start depositing red ketocarotenoids into their developing feathers if they are still in the moulting window. Second, why do we find an age effect on circulating ketocarotenoids among unornamented males, where 2-year-olds have higher concentration than 1-year-olds? Here again, testosterone may mediate differences in *CYP2J19* expression that underlie these results. Specifically, 1-year-olds are more likely to serve as helpers in their natal group and hence have lower levels of circulating testosterone, whereas most 2-year-olds transition into a breeding role, even after having completed moult into unornamented plumage [63,81], which may explain this shift in carotenoid concentration between the two age groups. Future work with greater sampling will reveal how tightly linked age, testosterone and gene




Figure 2. Testosterone implantation upregulates the expression of the *CYP2J19* gene in the liver of unornamented males. Shown are qPCR measurements of expression of *CYP2J19* relative to housekeeping gene *GAPDH* in ornamented males, testosterone (T)-implanted unornamented males, control unornamented males, and females of the red-backed fairywren. Points represent samples from individual birds, and boxplots indicate the median and quartiles with whiskers reaching up to 1.5 times the interquartile range. Different letters above plots indicate significant differences at $p < 0.05$ by Tukey's HSD.




Figure 3. A working mechanistic model for the acquisition of carotenoid-based plumage in male red-backed fairywrens (modified from [53]). Under this model, extrinsic and intrinsic cues interact to affect testosterone levels. In turn, testosterone levels mediate expression of the *CYP2J19* gene (and probably other genes), which appears to elevate circulating ketocarotenoid levels in plasma, which in turn are linked to plumage phenotype. Solid arrows depict connections supported by this study, whereas dashed arrows depict connections supported by previous work in this system. Illustrations of fairywrens by Allison Johnson. (Online version in colour.)

expression are to each other, and how often ketocarotenoids are produced even when not deposited into the plumage.

Our results corroborate a growing number of studies suggesting the importance of *CYP2J19* for red coloration in birds [23,24,82,83] and other animals [84,85]. The genetic basis of carotenoid-based coloration is not yet well resolved [3,32], but a handful of genes with large effects have been identified in recent years that are associated with carotenoid coloration (e.g. *BCO2* [86–88], *SCARB1* [16], *CYP2J19* [14,15] and other cytochrome P450's in other taxa [89,90]). There are probably many more genes involved in processes such as carotenoid uptake, processing and transport [91], all of which might influence coloration to varying degrees. Yet, without a clearer understanding of how these genes and other underlying processes interact to produce carotenoid phenotypes, our ability to identify and interpret the selective forces and

evolutionary processes that maintain these phenotypes remains limited. One promising route to identifying genes associated with carotenoids is to use RNA sequencing, especially in the context of experimental manipulation of phenotypes as done here with hormones, to identify gene pathways that underlie red coloration. We note the small sample size for our gene expression experiment ($n = 3$ per phenotype, owing to permitting restrictions), and we attempted to minimize any effect of this by only using breeding individuals to reduce phenology differences as well as collecting all birds in a relatively small time period (described in methods). Additionally, future work might test for the role of *CYP2J19* activity in other tissues relevant for colour signal production, including feather follicles and other integumentary tissue (e.g. bill or leg keratinocytes) [92], and investigate finer-grained relationships between *CYP2J19* expression and hue of red

plumage to extend beyond the presence/absence-of-coloration approach we employ in the current study. An integrative approach to studying colour signals by combining new research on proximate mechanisms of signal production with our current understanding of ultimate explanations for these signals (e.g. importance in mate choice and reproductive success), will improve our broader understanding of how such traits are regulated and shaped by selection.

Ethics. All procedures in this study were approved by the Tulane University Institutional Animal Care and Use Committee (IACUC 2019-1715), Cornell University IACUC (2009-0105), Washington State University IACUC (ASAF no. 04573), the James Cook University Animal Ethics Committee (A2100) and under a Queensland Government Department of Environment and Heritage Protection Scientific Purposes Permit (WISP15212314).

Data accessibility. Data available from the Dryad Digital Repository: <https://dx.doi.org/10.5061/dryad.pnvx0k6jp> [93].

Authors' contributions. S.K. contributed to conceptualization, writing, editing, data collection and funding acquisition, and carried out qPCR

laboratory work, formal statistical analysis and visualization. J.F.W. contributed to data collection and editing. K.J.M. carried out HPLC laboratory work and contributed to editing. J.B. made testosterone implants, carried out testosterone assays and contributed to editing. H.S. contributed to conceptualization, data collection and editing. M.S.W. contributed to conceptualization and editing. J.K. contributed to conceptualization, supervision, editing and funding acquisition.

Competing interests. We declare we have no competing interests.

Funding. This work was supported by the National Science Foundation (IOS-1354133 and IRES-1460048 to J.K.) and Tulane University Department of Ecology and Evolution (to S.K.). S.K. was supported by an NSF Graduate Research Fellowship during part of this work.

Acknowledgements. We gratefully acknowledge field technicians Mary Margaret Ferraro, Maria Smith, David Weber, Sarah Duff and Malcom Moniz for their assistance in collecting samples in the field, as well as William Feeney and Matthew Marsh for assistance in the field. We also thank Southeast Queensland Water for access to land where we performed our field research. This manuscript was greatly improved by discussion and feedback from E.D. Enbody, N.R. Hofmeister, J. Walsh, A.C. Demery, I.J. Lovette, A.R. Gunderson and K.G. Ferris, as well as the Karubian laboratory. Illustrations of fairywrens by Allison Johnson.

References

- Hill GE, McGraw KJ. 2006 *Bird coloration, volume 2: function and evolution*. Cambridge, MA: Harvard University Press.
- Svensson APA, Wong BBM. 2011 Carotenoid-based signals in behavioural ecology: a review. *Behaviour* **148**, 131–189. (doi:10.1163/000579510X548673)
- Toews DPL, Hofmeister NR, Taylor SA. 2017 The evolution and genetics of carotenoid processing in animals. *Trends Genet.* **33**, 171–182. (doi:10.1016/j.tig.2017.01.002)
- Badyaev AV, Morrison ES, Belloni V, Sanderson MJ. 2015 Tradeoff between robustness and elaboration in carotenoid networks produces cycles of avian color diversification. *Biol. Direct* **10**, 1–22. (doi:10.1186/s13062-015-0073-6)
- Morrison ES, Badyaev AV. 2018 Structure versus time in the evolutionary diversification of avian carotenoid metabolic networks. *J. Evol. Biol.* **31**, 764–772. (doi:10.1111/jeb.13257)
- Chew BP, Park JS. 2004 Carotenoid action on the immune response. *J. Nutr.* **134**, 2575–2615. (doi:10.1093/jn/134.1.2575)
- Surai PF, Speake BK, Sparks NHC. 2009 Carotenoids in avian nutrition and embryonic development. 2. Antioxidant properties and discrimination in embryonic tissues. *J. Poult. Sci.* **38**, 117–145. (doi:10.2141/jpsa.38.117)
- Weaver RJ, Koch RE, Hill GE. 2017 What maintains signal honesty in animal colour displays used in mate choice? *Phil. Trans. R. Soc. B* **372**, 20160343. (doi:10.1098/rstb.2016.0343)
- Koch RE, Hill GE. 2018 Do carotenoid-based ornaments entail resource trade-offs? An evaluation of theory and data. *Funct. Ecol.* **32**, 1908–1920. (doi:10.1111/1365-2435.13122)
- Koch RE, Kavazis AN, Hasselquist D, Hood WR, Zhang Y, Toomey MB, Hill GE. 2018 No evidence that carotenoid pigments boost either immune or antioxidant defenses in a songbird. *Nat. Commun.* **9**, 491. (doi:10.1038/s41467-018-02974-x)
- Koch RE, Staley M, Kavazis AN, Hasselquist D, Toomey MB, Hill GE. 2019 Testing the resource tradeoff hypothesis for carotenoid-based signal honesty using genetic variants of the domestic canary. *J. Exp. Biol.* **222**, jeb.188102. (doi:10.1242/jeb.188102)
- Costantini D, Møller AP. 2008 Carotenoids are minor antioxidants for birds. *Funct. Ecol.* **22**, 367–370. (doi:10.1111/j.1365-2435.2007.01366.x)
- Hartley RC, Kennedy MW. 2004 Are carotenoids a red herring in sexual display? *Trends Ecol. Evol.* **19**, 353–354. (doi:10.1016/j.tree.2004.04.002)
- Mundy NII *et al.* 2016 Red carotenoid coloration in the zebra finch is controlled by a cytochrome P450 gene cluster. *Curr. Biol.* **26**, 1435–1440. (doi:10.1016/j.cub.2016.04.047)
- Lopes RJ *et al.* 2016 Genetic basis for red coloration in birds. *Curr. Biol.* **26**, 1427–1434. (doi:10.1016/j.cub.2016.03.076)
- Toomey MB *et al.* 2017 High-density lipoprotein receptor SCARB1 is required for carotenoid coloration in birds. *Proc. Natl Acad. Sci. USA* **114**, 5219–5224. (doi:10.1073/pnas.1700751114)
- Våge DI, Boman IA. 2010 A nonsense mutation in the beta-carotene oxygenase 2 (BCO2) gene is tightly associated with accumulation of carotenoids in adipose tissue in sheep (*Ovis aries*). *BMC Genet.* **11**, 10. (doi:10.1186/1471-2156-11-10)
- Meredith P, Sarna T. 2006 The physical and chemical properties of eumelanin. *Pigment Cell Res.* **19**, 572–594. (doi:10.1111/j.1600-0749.2006.00345.x)
- D'Alba L, Shawkey MD. 2019 Melanosomes: biogenesis, properties, and evolution of an ancient organelle. *Physiol. Rev.* **99**, 1–19. (doi:10.1152/physrev.00059.2017)
- Walsh N, Dale J, McGraw KJ, Pointer MA, Mundy NI. 2012 Candidate genes for carotenoid coloration in vertebrates and their expression profiles in the carotenoid-containing plumage and bill of a wild bird. *Proc. R. Soc. B* **279**, 58–66. (doi:10.1098/rspb.2011.0765)
- Brelsford A, Toews DPL, Irwin DE. 2017 Admixture mapping in a hybrid zone reveals loci associated with avian feather coloration. *Proc. R. Soc. B* **284**, 20171106. (doi:10.1098/rspb.2017.1106)
- Brush AH. 1990 Metabolism of carotenoid pigments in birds. *FASEB J.* **4**, 2969–2977. (doi:10.1096/fasebj.4.12.2394316)
- Twyman H, Andersson S, Mundy NI. 2018 Evolution of *CYP2J19*, a gene involved in colour vision and red coloration in birds: positive selection in the face of conservation and pleiotropy. *BMC Evol. Biol.* **18**, 22. (doi:10.1186/s12862-018-1136-y)
- Twyman H, Prager M, Mundy NI, Andersson S. 2018 Expression of a carotenoid-modifying gene and evolution of red coloration in weaverbirds (Ploceidae). *Mol. Ecol.* **27**, 449–458. (doi:10.1111/mec.14451)
- Prager M, Andersson S. 2010 Convergent evolution of red carotenoid coloration in widowbirds and bishops (*Euplectes* spp.). *Evolution* **64**, 3609–3619. (doi:10.1111/j.1558-5646.2010.01081.x)
- Friedman NR, McGraw KJ, Omland KE. 2014 Evolution of carotenoid pigmentation in caciques and meadowlarks (Icteridae): Repeated gains of red plumage coloration by carotenoid C4-oxygenation. *Evolution* **68**, 791–801. (doi:10.1111/evo.12304)
- Thomas DB, McGraw KJ, Butler MW, Carrano MT, Madden O, James HF. 2014 Ancient origins and multiple appearances of carotenoid-pigmented feathers in birds. *Proc. R. Soc. B* **281**, 20140806. (doi:10.1098/rspb.2014.0806)
- Pryke SR, Andersson S, Lawes MJ, Piper SE. 2002 Carotenoid status signaling in captive and wild red-collared widowbirds: independent effects of badge size and color. *Behav. Ecol.* **13**, 622–631. (doi:10.1093/beheco/13.5.622)
- Searcy WA, Yasukawa K. 1983 Sexual selection and red-winged blackbirds. *Am. Sci.* **71**, 166–174.

30. Hill GE. 1990 Female house finches prefer colourful males: sexual selection for a condition-dependent trait. *Anim. Behav.* **40**, 563–572. (doi:10.1016/S0003-3472(05)80537-8)
31. Burley N, Coopersmith CB. 1987 Bill color preferences of zebra finches. *Ethology* **76**, 133–151. (doi:10.1111/j.1439-0310.1987.tb00679.x)
32. Orteu A, Jiggins CD. 2020 The genomics of coloration provides insights into adaptive evolution. *Nat. Rev. Genet.* **21**, 461–475. (doi:10.1038/s41576-020-0234-z)
33. Hau M. 2007 Regulation of male traits by testosterone: implications for the evolution of vertebrate life histories. *Bioessays* **29**, 133–144. (doi:10.1002/bies.20524)
34. Ketterson ED, Nolan V. 1992 Hormones and life histories: an integrative approach. *Am. Nat.* **140**, S33–S62. (doi:10.1086/285396)
35. Cox RM, McGlothlin JW, Bonier F. 2016 Hormones as mediators of phenotypic and genetic integration: an evolutionary genetics approach. *Integr. Comp. Biol.* **56**, 126–137. (doi:10.1093/icb/icw033)
36. Young AJ, Carlson AA, Monfort SL, Russell AF, Bennett NC, Clutton-Brock T. 2006 Stress and the suppression of subordinate reproduction in cooperatively breeding meerkats. *Proc. Natl Acad. Sci. USA* **103**, 12 005–12 010. (doi:10.1073/pnas.0510038103)
37. Nelson-Flower MJ, Ridley AR. 2015 Male-male competition is not costly to dominant males in a cooperatively breeding bird. *Behav. Ecol. Sociobiol.* **69**, 1997–2004. (doi:10.1007/s00265-015-2011-0)
38. Pikus AE, Guindre-Parker S, Rubenstein DR. 2018 Testosterone, social status and parental care in a cooperatively breeding bird. *Horm. Behav.* **97**, 85–93. (doi:10.1016/j.yhbeh.2017.10.008)
39. Poiani A, Fletcher T. 1994 Plasma levels of androgens and gonadal development of breeders and helpers in the bell miner (*Manorina melanophrys*). *Behav. Ecol. Sociobiol.* **34**, 31–41. (doi:10.1007/BF00175456)
40. Peters A, Astheimer LB, Cockburn A. 2001 The annual testosterone profile in cooperatively breeding superb fairy-wrens, *Malurus cyaneus*, reflects their extreme infidelity. *Behav. Ecol. Sociobiol.* **50**, 519–527. (doi:10.1007/s002650100403)
41. Schoech SJ, Mumme RL, Moore MC. 1991 Reproductive endocrinology and mechanisms of breeding inhibition in cooperatively breeding Florida scrub jays (*Aphelocoma c. coerulescens*). *Condor* **93**, 354–364. (doi:10.2307/1368951)
42. Oliveira RF, Hirschenhauser K, Carneiro LA, Canario AVM. 2002 Social modulation of androgen levels in male teleost fish. *Comp. Biochem. Physiol. B* **132**, 203–215. (doi:10.1016/S1096-4959(01)00523-1)
43. Ryder TB, Dakin R, Vernasco BJ, Evans BS, Horton BM, Moore IT. 2019 Testosterone modulates status-specific patterns of cooperation in a social network. *Am. Nat.* **195**, 82–94. (doi:10.1086/706236)
44. Garamszegi LZ, Rosivall B, Rettenbacher S, Markó G, Zsebok S, Szöllösi E, Eens M, Potti J, Török J. 2012 Corticosterone, avoidance of novelty, risk-taking and aggression in a wild bird: no evidence for pleiotropic effects. *Ethology* **118**, 621–635. (doi:10.1111/j.1439-0310.2012.02049.x)
45. Lipshutz SE, George EM, Bentz AB, Rosvall KA. 2019 Evaluating testosterone as a phenotypic integrator: from tissues to individuals to species. *Mol. Cell. Endocrinol.* **496**, 110531. (doi:10.1016/j.mce.2019.110531)
46. Ketterson ED, Atwell JW, McGlothlin JW. 2009 Phenotypic integration and independence: hormones, performance, and response to environmental change. *Integr. Comp. Biol.* **49**, 365–379. (doi:10.1093/icb/icp057)
47. Mank JE. 2017 The transcriptional architecture of phenotypic dimorphism. *Nat. Ecol. Evol.* **1**, 0006. (doi:10.1038/s41559-016-0006)
48. Blas J, Perez-Rodriguez L, Bortolotti GR, Vinuela J, Marchant TA. 2006 Testosterone increases bioavailability of carotenoids: insights into the honesty of sexual signaling. *Proc. Natl Acad. Sci. USA* **103**, 18 633–18 637. (doi:10.1073/pnas.0609189103)
49. Peters A, Roberts ML, Kurvers RHJM, Delhey K. 2012 Testosterone treatment can increase circulating carotenoids but does not affect yellow carotenoid-based plumage colour in blue tits *Cyanistes caeruleus*. *J. Avian Biol.* **43**, 362–368. (doi:10.1111/j.1600-048X.2012.05713.x)
50. Peters A. 2007 Testosterone and carotenoids: an integrated view of trade-offs between immunity and sexual signalling. *Bioessays* **29**, 427–430. (doi:10.1002/bies.20563)
51. Rowe M, McGraw KJ. 2008 Carotenoids in the seminal fluid of wild birds: interspecific variation in fairy-wrens. *Condor* **110**, 694–700. (doi:10.1525/cond.2008.8604)
52. Rowley I, Russel E. 1997 *Fairy-wrens and grasswrens: Maluridae*. Oxford, UK: Oxford University Press.
53. Webster MS, Karubian J, Schwabl H. 2010 Dealing with uncertainty: flexible reproductive strategies by a tropical passerine bird in an unstable ecological and social environment. *Adv. Study Behav.* **42**, 123–153. (doi:10.1016/S0065-3454(10)42004-5)
54. Webster MS, Varian CW, Karubian J. 2008 Plumage color and reproduction in the red-backed fairy-wren: why be a dull breeder? *Behav. Ecol.* **19**, 517–524. (doi:10.1093/beheco/arn015)
55. Baldassarre DT, Webster MS. 2013 Experimental evidence that extra-pair mating drives asymmetrical introgression of a sexual trait. *Proc. R. Soc. B* **280**, 20132175. (doi:10.1098/rspb.2013.2175)
56. Baldassarre DT, White TA, Karubian J, Webster MS. 2014 Genomic and morphological analysis of a semipermeable avian hybrid zone suggests asymmetrical introgression of a sexual signal. *Evolution* **68**, 2644–2657. (doi:10.1111/evo.12457)
57. Lindsay WR, Webster MS, Varian CW, Schwabl H. 2009 Plumage colour acquisition and behaviour are associated with androgens in a phenotypically plastic tropical bird. *Anim. Behav.* **77**, 1525–1532. (doi:10.1016/j.anbehav.2009.02.027)
58. Lindsay WR, Webster MS, Schwabl H. 2011 Sexually selected male plumage color is testosterone dependent in a tropical passerine bird, the red-backed fairy-wren (*Malurus melanocephalus*). *PLoS ONE* **6**, e0026067. (doi:10.1371/journal.pone.0026067)
59. Karubian J, Lindsay WR, Schwabl H, Webster MS. 2011 Bill coloration, a flexible signal in a tropical passerine bird, is regulated by social environment and androgens. *Anim. Behav.* **81**, 795–800. (doi:10.1016/j.anbehav.2011.01.012)
60. Lindsay WR, Barron DG, Webster MS, Schwabl H. 2016 Testosterone activates sexual dimorphism including male-typical carotenoid but not melanin plumage pigmentation in a female bird. *J. Exp. Biol.* **219**, 3091–3099. (doi:10.1242/jeb.135384)
61. Del Val E, Senar JC, Garrido-Fernández J, Jarén M, Borràs A, Cabrera J, Negro JJ. 2009 The liver but not the skin is the site for conversion of a red carotenoid in a passerine bird. *Naturwissenschaften* **96**, 797–801. (doi:10.1007/s00114-009-0534-9)
62. Ge Z, Johnson JD, Cobine PA, McGraw KJ, Garcia R, Hill GE. 2015 High concentrations of ketocarotenoids in hepatic mitochondria of *Haemorrhous mexicanus*. *Physiol. Biochem. Zool.* **88**, 444–450. (doi:10.1086/681992)
63. Karubian J. 2002 Costs and benefits of variable breeding plumage in the red-backed fairy-wren. *Evolution* **56**, 1673–1682. (doi:10.1111/j.0014-3820.2002.tb01479.x)
64. Mulder RA, Magrath MJL. 1994 Timing of pre-nuptial molt as a sexually selected indicator of male quality in superb fairy-wrens (*Malurus cyaneus*). *Behav. Ecol.* **5**, 393–400. (doi:10.1093/beheco/5.4.393)
65. Pyle P, Howell SNG, Yunick R, Desante D. 1987 *Identification guide to North American passerines*. Bolinas, CA: Slate Creek Press.
66. Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. 2018 nlme: Linear and nonlinear mixed effects models. R package version 3.1-137. See <https://CRAN.R-project.org/package=nlme>.
67. R Core Team. 2019 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
68. Hothorn T, Bretz F, Westfall P. 2008 Simultaneous inference in general parametric models. *Biom. J.* **50**, 346–363. (doi:10.1002/bimj.200810425)
69. Barron DG, Webster MS, Schwabl H. 2015 Do androgens link morphology and behaviour to produce phenotype-specific behavioural strategies? *Anim. Behav.* **100**, 116–124. (doi:10.1016/j.anbehav.2014.11.016)
70. Livak KJ, Schmittgen TD. 2001 Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCT} method. *Methods* **25**, 402–408. (doi:10.1006/meth.2001.1262)
71. Fernald RD. 1976 The effect of testosterone on the behavior and coloration of adult male cichlid fish (*Haplochromis burtoni*, Günther). *Horm. Res. Paediatr.* **7**, 172–178. (doi:10.1159/000178726)
72. Mougeot F, Irvine JR, Seiwright L, Redpath SM, Pierny S. 2004 Testosterone, immunocompetence, and honest sexual signaling in male red grouse. *Behav. Ecol.* **15**, 930–937. (doi:10.1093/beheco/arh087)

73. McGraw KJ, Correa SM, Adkins-Regan E. 2006 Testosterone upregulates lipoprotein status to control sexual attractiveness in a colorful songbird. *Behav. Ecol. Sociobiol.* **60**, 117–122. (doi:10.1007/s00265-005-0135-3)
74. Evans MR, Goldsmith AR, Norris SRA. 2000 The effects of testosterone on antibody production and plumage coloration in male house sparrows (*Passer domesticus*). *Behav. Ecol. Sociobiol.* **47**, 156–163. (doi:10.1007/s002650050006)
75. Sin SYW, Lu L, Edwards SV. 2020 De novo assembly of the northern cardinal (*Cardinalis cardinalis*) genome reveals candidate regulatory regions for sexually dichromatic red plumage coloration. *bioRxiv*, 1–28. (doi:10.1101/2020.05.12.092080)
76. Kirschel A, Nwankwo E, Pierce D, Moysi M, Ogolowa B, Monadjem A, Brelsford A. 2020 *CYP2J19* mediates carotenoid colour introgression across a natural avian hybrid zone. (doi:10.22541/au.158880215.54508683/v2)
77. Bentz AB, Dossey EK, Rosvall KA. 2019 Tissue-specific gene regulation corresponds with seasonal plasticity in female testosterone. *Gen. Comp. Endocrinol.* **270**, 26–34. (doi:10.1016/j.ygcen.2018.10.001)
78. Schuppe ER, Fuxjager MJ. 2019 Phenotypic variation reveals sites of evolutionary constraint in the androgenic signaling pathway. *Horm. Behav.* **115**, 104538. (doi:10.1016/j.yhbeh.2019.06.002)
79. Rosvall KA, Bergeon Burns CM, Jayaratna SP, Ketterson ED. 2016 Divergence along the gonadal steroidogenic pathway: implications for hormone-mediated phenotypic evolution. *Horm. Behav.* **84**, 1–8. (doi:10.1016/j.yhbeh.2016.05.015)
80. Dowling J, Webster MS. 2017 Working with what you've got: unattractive males show greater mate-guarding effort in a duetting songbird. *Biol. Lett.* **13**, 20160682. (doi:10.1098/rsbl.2016.0682)
81. Karubian J. 2008 Changes in breeding status are associated with rapid bill darkening in male red-backed fairy-wrens *Malurus melanocephalus*. *J. Avian Biol.* **39**, 81–86. (doi:10.1111/j.0908-8857.2008.04161.x)
82. Hooper DM, Griffith SC, Price TD. 2019 Sex chromosome inversions enforce reproductive isolation across an avian hybrid zone. *Mol. Ecol.* **28**, 1246–1262. (doi:10.1111/mec.14874)
83. Hill GE *et al.* 2019 Plumage redness signals mitochondrial function in the house finch. *Proc. R. Soc. B* **286**, 20191354. (doi:10.1098/rspb.2019.1354)
84. Twyman H, Valenzuela N, Literman R, Andersson S, Mundy NI. 2016 Seeing red to being red: conserved genetic mechanism for red cone oil droplets and co-option for red coloration in birds and turtles. *Proc. R. Soc. B* **283**, 20161208. (doi:10.1098/rspb.2016.1208)
85. Weaver RJ, Gonzalez BK, Santos SR, Havird JC. 2020 Red coloration in an anchialine shrimp: carotenoids, genetic variation, and candidate genes. *Biol. Bull.* **238**, 119–130. (doi:10.1086/708625)
86. Eriksson J *et al.* 2008 Identification of the Yellow skin gene reveals a hybrid origin of the domestic chicken. *PLoS Genet.* **4**, e1000010. (doi:10.1371/journal.pgen.1000010)
87. Gazda MA *et al.* 2020 Genetic basis of de novo appearance of carotenoid ornamentation in bare parts of canaries. *Mol. Biol. Evol.* **37**, 1317–1328. (doi:10.1093/molbev/msaa006)
88. Gazda MA, Toomey MB, Araújo PM, Lopes RJ, Afonso S, Myers CA. 2020 A genetic mechanism for sexual dichromatism in birds. *Science* **12**, 1–6. (doi:10.1126/science.aba0803)
89. Wybouw N *et al.* 2019 Convergent evolution of cytochrome P450s underlies independent origins of keto-carotenoid pigmentation in animals. *Proc. R. Soc. B* **286**, 20191039. (doi:10.1098/rspb.2019.1039)
90. Twomey E, Johnson JD, Castroviejo-Fisher S, Van Bocxlaer I. 2020 A ketocarotenoid-based color polymorphism in the Sira poison frog *Ranitomeya sirensis* indicates novel gene interactions underlying aposematic signal variation. *Mol. Ecol.* **29**, 2004–2015. (doi:10.1111/mec.15466)
91. McLean CA, Lutz A, Rankin KJ, Elliott A, Moussalli A, Stuart-Fox D. 2019 Red carotenoids and associated gene expression explain colour variation in frillneck lizards. *Proc. R. Soc. B* **286**, 20191172. (doi:10.1098/rspb.2019.1172)
92. McGraw KJ. 2004 Colorful songbirds metabolize carotenoids at the integument. *J. Avian Biol.* **35**, 471–476. (doi:10.1111/j.0908-8857.2004.03405.x)
93. Khalil S, Welklin JF, McGraw KJ, Boersma J, Schwabl H, Webster MS, Karubian J. 2020 Data from: Testosterone regulates *CYP2J19*-linked carotenoid signal expression in male red-backed fairywrens (*Malurus melanocephalus*). Dryad Digital Repository. (<https://doi.org/10.5061/dryad.pnvx0k6jp>)