

SCIENTIFIC REPORTS

Corrigendum: Structural basis of a nucleosome containing histone H2A.B/H2A.Bbd that transiently associates with reorganized chromatin

Yasuhiro Arimura, Hiroshi Kimura, Takashi Oda, Koichi Sato, Akihisa Osakabe, Hiroaki Tachiwana, Yuko Sato, Yasuha Kinugasa, Tsuyoshi Ikura, Masaaki Sugiyama, Mamoru Sato & Hitoshi Kurumizaka

Scientific Reports 3:3510; doi: 10.1038/srep03510; published online 16 December 2013; updated on 13 July 2015

The DLS data shown in Figure 5b was obtained using the 146 bp α -satellite DNA derivative, rather than the 145 bp 601 DNA fragment.

Under the subheading ‘Solution structure of the H2A.B nucleosome’:

“In this analysis, we used the 145 bp 601 DNA fragment, which reportedly forms a stably positioned nucleosome³. Consistent with previous electron and atomic force microscopic analyses²⁹, the mean particle size of the H2A.B octasome with the 145 bp 601 DNA fragment was clearly larger than that of the canonical H2A octasome (Fig. 5a and b).”

should read:

“In this analysis, we used a 146 bp DNA fragment, containing the α -satellite sequence, which reportedly forms a stably positioned nucleosome¹. Consistent with previous electron and atomic force microscopic analyses²⁹, the mean particle size of the H2A.B octasome was clearly larger than that of the canonical H2A octasome (Fig. 5a and b).”

In the legend of Figure 5b:

“Particle size distribution profiles in the DLS analysis of the H2A and H2A.B octasomes with a 145 bp 601 DNA fragment”

should read:

“Particle size distribution profiles in the DLS analysis of the H2A and H2A.B octasomes with a 146 bp DNA fragment.”

In the Methods section, under the subheading ‘Dynamic light scattering (DLS) measurements’:

“For the DLS experiments, the 145 bp 601 DNA was used.”

should read:

“For the DLS experiments, a 146 bp DNA containing the α -satellite sequence was used.”